Published in final edited form as:

Immunol Res. 2007 ; 37(1): 61–78.

Intestinal Innate Immunity and the Pathogenesis of *Salmonella* Enteritis

Chittur V. Srikanth and

Mucosal Immunology Laboratory, Division of Pediatric Gastroenterology and Nutrition,
Massachusetts General Hospital and Harvard Medical School, Charlestown, MA 02129

Bobby J. Cherayil

Mucosal Immunology Laboratory, Division of Pediatric Gastroenterology and Nutrition, Room
3400, Massachusetts General Hospital East, Building 114, 16th Street, Charlestown, MA 02129.
cherayil@helix.mgh.harvard.edu

Abstract

Acute gastroenteritis caused by *Salmonella typhimurium* infection is a clinical problem with significant public health impact. The availability of several experimental models of this condition has allowed detailed investigation of the cellular and molecular interactions involved in its pathogenesis. Such studies have shed light on the roles played by bacterial virulence factors and host innate immune mechanisms in the development of intestinal inflammation.

Keywords

Salmonella; Enteritis; Innate immunity; Mucosa

Introduction and Overview

The Gram-negative bacterial pathogen *Salmonella typhimurium* is one of the most frequent causes of acute gastroenteritis in humans, and is responsible for significant morbidity, and some mortality, worldwide (1). It is also one of the few microorganisms that has been used in a well-documented attempt at causing widespread illness for political purposes (2). This incident, involving a religious cult in Oregon, illustrates the potential use of *Salmonella* in bioterrorism, and provides additional motivation for understanding the pathogenesis of the diseases caused by this organism.

Salmonella enteritis is the outcome of the acute intestinal inflammation that results from the triggering of host innate immune responses by the pathogen. These responses are protective, and help to clear the infection over the course of 3–4 d in most cases. However, the inflammation causes disturbances of gastrointestinal function that manifest clinically as abdominal pain, nausea, vomiting, and diarrhea (3). Related serotypes of *Salmonella*, *S. typhi* and *S. paratyphi*, cause the systemic illness typhoid fever rather than gastroenteritis. The pathogenesis of this disease has been reviewed elsewhere (4) and will not be considered further here.

© 2007 Humana Press Inc.

Note Added in Proof

Recent work with TLR5-deficient mice (Uematsu et al., Nat Immunol 2006; 7: 868–874) indicates a dominant role for CD11c⁺ lamina propria cells rather than IECs in TLR5-dependent inflammatory responses to *Salmonella*. TLR5 is also required for the extra-intestinal transport of the bacteria by these cells. The effects of TLR5 deficiency have been described independently by others (Feuillet et al., Proc Natl Acad Sci USA 2006; 103: 12487–12492).

S. typhimurium usually infects the host orally following ingestion of contaminated water or food. Spread of the infection is facilitated by the ability of the organism to survive for several days in groundwater, pond water, or seawater, and for months in contaminated food material like eggs and oysters (4–7). A key aspect of pathogenesis is the active bacterial invasion of intestinal epithelial cells (IECs), particularly those that overlie the Peyer's patches of the small intestine. This process is mediated by *Salmonella* effector proteins that are introduced into the host cells through a specialized secretory apparatus, and is accompanied by the production of several epithelium-derived proinflammatory molecules (8). Once the bacteria have crossed the epithelial monolayer, they interact with resident lamina propria macrophages, and with circulating monocytes and neutrophils that are recruited to the site of infection by the initial invasion of IECs. These cells also respond by secreting cytokines, chemokines, and other soluble mediators that amplify the inflammation. The recruited phagocytes, along with local defense mechanisms, help in the elimination of the bacteria, while other cells such as dendritic cells and lymphocytes initiate the steps leading to adaptive immunity.

In this review, we will discuss what is currently known about the cellular and molecular processes involved in the intestinal innate immune response to *S. typhimurium* and how they contribute to the pathogenesis of acute enteritis. We will focus mainly on the interactions between the bacteria and epithelial cells, macrophages, and neutrophils, because these are the cells that participate in the earliest stages of the response. Before proceeding, however, we will digress briefly to consider the experimental models that have been used to study *Salmonella* enteritis.

Experimental Models of *Salmonella*-Induced Intestinal Inflammation

Apart from volunteer studies conducted to determine the minimal pathogenic dose (9), there have been few, if any, investigations of *Salmonella* pathogenesis in humans at the cellular and molecular level. Researchers have had to rely instead on the use of model systems that are amenable to detailed dissection. In vitro tissue culture studies, involving the infection of cultured cells of various types, including polarized monolayers of the T84 human colonic cell line, have yielded important insights into the functions of the pathogen and host molecules required for cell invasion (10). Such studies have also provided information on the cellular inflammatory mediators that are released in response to infection. However, the reductionist nature of this approach places a significant limitation on understanding the complex interplay of molecules and cells that occurs in vivo. Accordingly, investigators have turned to animal models of salmonellosis. Cattle are natural hosts for *S. typhimurium*, and infected calves develop an acute gastroenteritis that resembles the human disease both clinically and histopathologically (11). Infection of whole animals, or of isolated calf intestinal loops, has thus proved quite useful in examining the role of various *Salmonella* molecules involved in the induction of intestinal inflammation. In general, these studies have confirmed observations made in tissue culture. Unfortunately, the difficulties of large animal husbandry, together with high cost and the inability to manipulate host cellular and immunologic factors, have made the bovine model usable only in restricted circumstances. *Salmonella* enteritis can also be modeled in Rhesus monkeys, but this approach suffers from disadvantages similar to infection in cattle and has not been used extensively (12,13). A recent advance in the field was made when it was discovered that oral streptomycin pre-treatment of C57BL/6 mice followed by oral infection with a streptomycin-resistant strain of *S. typhimurium* resulted in a robust and reproducible large intestinal inflammation (14). This enteritis, which is most marked in the cecum, has several of the pathologic hallmarks of acute *Salmonella* gastroenteritis in humans, including neutrophil and mononuclear infiltration, and epithelial damage. The major difference from the human disease is the lack of a significant secretory response, and the infected mice rarely, if ever, develop diarrhea.

Despite this shortcoming, this model is convenient, relatively inexpensive, and allows the detailed study and manipulation of both bacterial and host factors involved in disease pathogenesis. The ability to use genetically engineered mouse strains deficient in specific aspects of the immune response is a particular advantage of this experimental system. Its usefulness has already been illustrated by several informative studies (15–20) and is likely to make it the model of choice for future investigations.

There is, in general, a satisfying concordance of findings from the various models used to study *Salmonella* infection experimentally. There are discordant results, however, that could be related to variations in methodology, bacterial strain, host cell type, or real differences in biology, and we will attempt to point out these discrepancies where appropriate.

***Salmonella* Epithelial Interactions: The Initiation of the Inflammatory Response**

Colonization of the intestine is an essential pre-requisite to *Salmonella* infection. It takes only 100–1000 colony forming units of *S. typhimurium* to cause disease in humans (9). These bacteria have to survive and proliferate in a microenvironment that is usually hostile to incoming invaders because of the presence of large numbers of commensal microorganisms and the action of host immune defenses. Successful colonization depends on the ability to resist mucosal anti-microbial mechanisms, compete with other bacteria for space and nutrients, and adhere to the apical surface of IECs. The importance of this step of the infection process is suggested by the fact that without the streptomycin pre-treatment step that is routinely used in the mouse model of *Salmonella* enteritis, intestinal colonization by the pathogen is insufficient to cause intestinal inflammation (14). The antibiotic is presumed to alter the commensal flora and create a niche in which *Salmonella* can survive. The existence of *Salmonella* mutants that have abnormal intestinal colonization characteristics also points to specific molecular interactions that must occur for this process to be successful (21–24). Further study of the bacterial and host factors involved in intestinal colonization and their role in enteritis may suggest new ways to interrupt pathogenesis at a very early stage. It should be mentioned that the oral infecting dose required to cause disease in humans is much lower than in mice (9,14), so caution is required when extrapolating from results obtained with the mouse model.

Contact between infecting *Salmonella* and the intestinal epithelium initiates a series of events that represents, in effect, a battle between the host and the pathogen. On the bacterial side, mechanisms for invasion and intracellular survival are activated, while on the epithelial side several defensive measures are deployed, including local production of anti-microbial factors, and the recruitment of back-up troops in the form of phagocytes and other cells of the immune system. Analysis of the molecular details of these interactions is an active area of research, one that has produced a wealth of information on both *Salmonella* biology and mucosal immunity. Broadly speaking, there are two main pathways by which anti-microbial inflammatory responses are activated during *Salmonella*–epithelial interactions. One is a by-product of bacterial invasion, while the other involves specific sensing of microbial components by host pattern recognition receptors (PRRs).

Epithelial Invasion and the Activation of IEC Pro-inflammatory Signals

The ability to actively invade IECs is an important aspect of *Salmonella* virulence. It depends on the concerted action of a battery of bacterial proteins, one key component of which is a specialized secretory apparatus, a type III secretion system (TTSS), encoded by genes located on *Salmonella* pathogenicity island 1 (SPI1) (8). The SPI1 TTSS is a multi-subunit needle-like structure that protrudes from the bacterial surface and penetrates the

plasma membrane of the cell that is being invaded. Several bacterial proteins, many of them also encoded by SPI1 genes, are actively secreted through the needle apparatus into the host cytosol with the assistance of a translo-cone complex made up of the three *Salmonella* proteins, SipB, SipC, and SipD. The main function of the translocated proteins, which are sometimes called effector proteins, is to induce rearrangements of the host actin cytoskeleton at the site of cell–bacterial interaction, an activity in which the bacterial guanine nucleotide exchange factors (GEFs) SopE and SopE2, as well as the actin binding proteins SipA and SipC, play important roles. The cytoskeletal alterations lead to membrane ruffling and the formation of lamellapodial extensions that engulf the bacterium and ultimately encompass it within a membrane-bound intra-cellular compartment known as the *Salmonella*-containing vacuole (SCV). The highly regulated activity of *Salmonella* SptP, a dual function phosphatase and GTPase activating protein (GAP), restores the basal state of the actin cytoskeleton and plasma membrane (25).

Some of the *Salmonella* effectors that are involved in cell invasion also activate intra-cellular signals that result in changes in IEC gene expression and physiology. SopE and SopE2 activate the cellular Rho GTPases Cdc42 and Rac1 to induce actin polymerization, but these interactions also result in the activation of signaling pathways leading to the nuclear translocation of the transcription factor NF- κ B and the resultant upregulation of genes involved in immune and inflammatory responses (26). One outcome is the baso-lateral secretion of various soluble mediators such as IL-8 that recruit neutrophils and other cells from the circulation into the intestinal lamina propria. Experiments carried out in our laboratory suggest that signals triggered by SopE2 act at a step of NF- κ B activation that is downstream of the induced degradation of the inhibitor of this transcription factor, I κ B α , and that such signals may interact synergistically with those initiated by Toll-like receptor (TLR) 5 (see below) (27).

Neutrophils recruited to the intestinal lamina propria are induced to migrate across the epithelium into the gut lumen in response to the effects of hepxilin A3. This prostanoid chemoattractant is secreted apically by IECs as a result of cellular interactions with the *Salmonella* SipA protein (28). Interestingly, this function of SipA does not appear to require translocation into the IEC cytosol, and, instead, may involve stimulation of a cell surface receptor that is yet to be characterized. Activation of protein kinase C is a key step downstream of this putative receptor in SipA-induced production of hepxilin A3 (29). Because migration of neutrophils across the epithelium disrupts the integrity of the monolayer, contributes to IEC dysfunction, and gives rise to the microabscesses that are a characteristic pathologic feature (30,31), further study of the effects of SipA is likely to lead to important insights into the pathogenesis of *Salmonella* enteritis.

The *Salmonella* effector SopB is an inositol phosphatase that increases inositol flux in IEC tissue culture models (32). It has also been shown to induce chloride secretion in such models (32,33) and may contribute thereby to the pathogenesis of diarrhea. In addition, this protein activates the phosphatidylinositol 3-kinase/Akt kinase signaling pathway in epithelial cells (34). We demonstrated recently that this pathway can also be activated in a SopB-independent manner, and that it may have an anti-inflammatory function by inhibiting IL-8 production (35). Such attenuation of IEC inflammatory responses may be necessary for *Salmonella* to obtain a foothold in the gut. Additional anti-inflammatory effects are provided by AvrA, a SPI1 effector that inhibits the activation of MAP kinase-dependent signals, and by an uncharacterized bacterial factor that appears to act by altering neddylation of the enzyme complex that ubiquitinates I κ B α (36,37).

The in vivo role of the various SPI1 effectors has been examined in both the bovine and murine models of enteritis (15,38,39). These studies have confirmed the importance of

SipA, SopB, SopE, and SopE2 in intestinal inflammation, but exactly how these proteins contribute to disease pathogenesis, and, specifically, whether their effects are primarily on cell invasion or on activation of pro-inflammatory signaling pathways, are issues that await clarification. Experiments with the bovine model have also implicated the SopA and SopD effectors in the pathogenesis of gastroenteritis, but very little is known about the molecular function of these proteins in inflammation other than that they are involved in invasion of polarized epithelial cells (38–40). The putative anti-inflammatory function of AvrA has not been substantiated in vivo (38). Table 1 summarizes what is currently known about the role of *Salmonella* effector proteins in the activation of IEC responses and in the pathogenesis of gastroenteritis, and Fig. 1 illustrates some of these concepts.

Epithelial PRRs in the Response to *Salmonella*

The cells of the innate immune system use relatively invariant, non-clonotypic, germ-line encoded receptors to recognize conserved molecules that are expressed by broad groups of microorganisms. An important subset of such PRRs is made up of a family of structurally related transmembrane proteins known as the TLRs (41). There are 11 TLRs that are involved in sensing and responding to various microbial components such as lipopolysaccharide (LPS), lipopeptides, and DNA, and there has been a great deal of recent interest in elucidating the role of these receptors in epithelial responses. TLR5, the receptor for monomeric bacterial flagellin, is expressed on the basolateral aspect of primary intestinal epithelium or polarized IEC lines, a location that is presumed to prevent inappropriate activation by commensal bacteria (42,43). Pathogenic flagellated bacteria have special virulence mechanisms that result in flagellin crossing the epithelial monolayer, or the bacteria may themselves invade or damage the epithelium and come into contact with TLR5. Apical exposure of T84 cells to *Salmonella*, for instance, results in the vesicle-mediated transcytosis of flagellin to the basolateral aspect of the monolayer, a process that depends on molecules encoded by a second *Salmonella* pathogenicity island SPI2 (44). The polarized localization of TLR5 thus allows it to respond selectively to the presence of flagellin on the basolateral aspect of the epithelium, circumstances that would indicate infection with a presumptive pathogen. Flagellin is one of the few TLR ligands that has been demonstrated to associate physically with its receptor (45). The signals transduced by TLR5 following flagellin binding are a major stimulus for the epithelial expression of IL-8, implying an important role in neutrophil recruitment to the intestine (42,43). Studies carried out recently in the mouse enteritis model support the importance of TLRs in the intestinal inflammatory response by demonstrating that *Salmonella*-induced intestinal inflammation was significantly attenuated in mice lacking MyD88, an adaptor protein that is required for signal transduction downstream of most TLRs (17). It is not clear, however, whether the requirement for MyD88 is in IECs or in sub-epithelial cells such as macrophages. These studies also raised some questions about the role of flagellin–TLR5 interactions, and suggested that the attenuated enteritis caused by aflagellate mutants of *Salmonella* may have more to do with decreased motility than a failure to activate TLR5 (16). Definitive resolution of this issue will require investigations in TLR5-deficient mice. See note added in proof.

Recent work has demonstrated that the production of monomeric flagellin, the ligand for TLR5, may depend on exposure of *Salmonella* to lysophospholipids secreted by IECs (46). Because IEC production of the lysophospholipids requires contact with the bacteria, bidirectional cross-talk between host and pathogen appears to be involved in the activation of TLR5. The in vivo significance of these intriguing tissue culture observations is yet to be confirmed.

Other TLRs, such as TLR2, TLR3, TLR4, and TLR9, have been reported to be expressed on IECs (47,48). However, there is some controversy regarding the precise role of these

receptors in the intestinal epithelium, and their in vivo contribution to the activation of IEC inflammatory responses in infectious enteritis has yet to be convincingly demonstrated.

Other Factors Involved in Bacterial Epithelial Interactions

Apart from the SPI1 TTSS and flagellin, other bacterial structures may contribute to the *Salmonella*–IEC interaction. The *Salmonella* genome contains a number of fimbrial operons that carry out crucial roles during the intestinal phase of infection. Fimbrial adhesins are involved in attachment to and invasion of IECs and also act as targets of the adaptive immune response (49). The csgBA *Salmonella* mutant, which carries defects in one of the fimbrial operons, is impaired in eliciting fluid accumulation and induction of Gro α mRNA in bovine ligated ileal loops, possibly because of a failure to activate TLR2 (50).

Epithelial Anti-microbial Peptides in the Control of *Salmonella* Infection

In addition to the production of inflammatory mediators involved in recruiting circulating phagocytic cells to the site of infection, IECs also play an important role in killing bacterial invaders via the secretion of antimicrobial peptides. Paneth cells, one of the lineages that differentiate from epithelial precursors, contribute significantly to this function. These cells, which are located at the bottom of small intestinal crypts, express a number of short cationic peptides, the α -defensins or cryptdins, that are able to disrupt the membranes of a number of microorganisms, including *Salmonella* (51). The defensins are secreted apically into the crypt lumen when Paneth cells are exposed to intact bacteria or bacterial components such as LPS (52). The importance of defensins in protection against enteropathogens is illustrated by the observation that transgenic mice expressing human defensin 5 exhibited a dramatic increase in resistance to *Salmonella* infection (53).

IECs express members of another class of defensive peptide, the cathelicidins, in which a cathelin domain is linked to a peptide with anti-microbial activity (54). Several cathelicidins have been identified in mammals, but of these humans express only one, LL-37 (hCAP18). LL-37/hCAP18 is expressed in the epithelial cells located at the surface and upper crypts of the normal human colon (55). In the mouse, cathelicidin-related anti-microbial peptide (CRAMP) has been shown to inhibit *Salmonella* replication in vivo and in vitro, and to induce the formation of long filamentous bacteria (56).

Developmental Changes in Epithelial Function and the Intestinal Response to *Salmonella* Infection

Systemic spread of *Salmonella* during intestinal infection in normal adult individuals is minimized by the acute intestinal inflammatory response. However, young infants, particularly those under 3 mo age, are prone to extra-intestinal dissemination of the bacteria, leading to complications such as sepsis, osteomyelitis, and meningitis. We observed recently that this age-dependent variation in the response to *Salmonella* infection also occurs in the mouse model (19). One- to 2-wk-old pre-weaned mouse pups had significantly less intestinal inflammation, based on a number of assays, than 5–6-wk-old adults. Correspondingly, the younger animals had larger numbers of *Salmonella* in the mesenteric lymph nodes and spleen than the adults. Using this model, we found that the intestinal expression of a number of genes involved in anti-microbial and inflammatory responses was significantly upregulated with age. The expression of many of these genes, including chemokines such as CXCL9 and CXCL10, and regulators of intracellular infection such as LRG-47 and IIGTP, are under the control of interferon (IFN) γ . Indeed, we found that the age-dependent increase in expression of such genes did not occur in IFN γ -deficient mice. Correspondingly, the response of adult IFN γ -deficient animals to *Salmonella* infection resembled that of wild-type pups, with lower levels of intestinal inflammation and higher numbers of tissue bacteria at systemic sites. Based on these observations, we have suggested

that IFN γ plays an important role in the maturation of intestinal innate immunity, and in regulating *Salmonella*-induced intestinal inflammation.

***Salmonella* Beyond the Epithelium: Amplification of the Inflammatory Response**

Once *Salmonella* has breached the intestinal epithelial barrier, either by transcytosis across the IEC or by paracellular movement through breaks in the epithelium, the bacteria encounter cells of the immune system that are resident in the lamina propria, such as tissue macrophages, dendritic cells, and mast cells, as well as newly recruited neutrophils and monocytes. Resident lamina propria macrophages of the intestine, unlike their newly recruited counterparts, are generally considered to be poorly responsive to bacteria because of low levels of expression of the TLR accessory molecule CD14 and the influence of TGF β in the local microenvironment (57). Nevertheless, they are avidly phagocytic and have been shown to kill internalized *Salmonella* very effectively (58). Mast cells, which are present in large numbers in the gastrointestinal tract, play an important role in bacterial infection by their ability to rapidly release several inflammatory mediators from pre-formed granule stores (59). However, recent experiments with a low-dose intraperitoneal infection model suggest that these cells are not critical to protection against *Salmonella* (60). It is not clear whether this conclusion is also applicable to enteric infection. Dendritic cells are key to the development of adaptive immune responses during *Salmonella* infection and may actively take up the bacteria from the lumen even before epithelial invasion (61). The interactions of these cells with *Salmonella* have been reviewed in detail elsewhere (62).

Influx of neutrophils and monocytes into the intestinal lamina propria and epithelial layer is one of the characteristic histopathologic features of *Salmonella* enteritis. The migration of neutrophils to the site of infection depends on specific chemoattractants such as IL-8, GRO α , GRO γ , and hepxilin A3 that are produced by infected IECs and other cells in the lamina propria (63). The exact chemoattractants involved in recruiting monocyte/macrophages have not been delineated, although epithelial expression of CCL2 has been implicated in both human tissue culture systems and in vivo mouse models of salmonellosis (64,65). Both cell types play important roles in phagocytosing and killing *Salmonella* and in the production of various mediators such as IL-1 and TNF α that amplify the events set in motion by IECs. Some of the molecules secreted by these cells contribute significantly to the tissue damage that is the inevitable accompaniment of the inflammatory response. The molecular details of these processes are discussed below, and are depicted in Fig. 2.

Phagocyte Anti-microbial Mechanisms

Bacteria phagocytosed by macrophages and neutrophils are subjected to intracellular killing mechanisms that include trafficking to the degradative environment of the lysosome, the generation of toxic reactive oxygen and nitrogen intermediates (ROI and RNI) by phagocyte oxidase (phox) and inducible nitric oxide synthase (iNOS), respectively, exposure to bactericidal peptides, and micronutrient deprivation. As a successful intracellular pathogen, *Salmonella* has evolved strategies to resist some of these anti-microbial defenses, allowing it to survive for extended periods inside macrophages both in vitro and in vivo (66). Genes encoded by SPI2 play an important role in intra-macrophage survival, largely by modulating the characteristics of the SCV so that fusion with the lysosome, as well as recruitment of phox and iNOS, are prevented (67). Certain SPI2 defective strains of *Salmonella*, specifically those with mutations in the *sseD* or *ssaT* genes, cause an attenuated form of enteritis in both mouse and bovine models, suggesting that the ability to survive within macrophages influences the inflammatory response (11,17,68). However, inactivation of

another SPI2 gene, *ssrA*, does not result in alteration of fluid secretion in a rabbit ligated ileal loop model (69).

Intracellular survival and multiplication of *Salmonella* are also regulated by the Nramp1 gene, which encodes a macrophage-specific divalent cation transporter that is believed to pump iron out of the SCV, thereby depriving the bacteria of an essential nutrient (70). Although there are no published studies that directly address the role of Nramp1 in *Salmonella*-induced intestinal inflammation, it is reasonable, based on the effects of SPI2 mutations, that changes in the number of intracellular bacteria would influence the enteritis. Furthermore, production of the macrophage anti-inflammatory protein, secretory leukocyte protease inhibitor (SLPI), which inhibits TLR activation, is dependent on Nramp1 (71,72). We have shown recently that another macrophage iron transporter, ferroportin (FPN), influences the intracellular growth of *Salmonella*, probably by reducing iron concentrations within the SCV (73). We are currently in the process of analyzing the effects of altered FPN expression on enteritis in the mouse model.

Phagocyte Inflammatory Mediators

One of the important functions of macrophages newly recruited to the gut during *Salmonella* infection is the secretion of soluble mediators that help to amplify the inflammatory response initiated by the IECs. Two mediators that play important roles in this process are TNF α and IL-1 β (74,75). TNF α promotes the recruitment of additional leukocytes, including neutrophils, by inducing chemokine and adhesion molecule expression on surrounding cells. In addition, it enhances phox- and iNOS-independent microbicidal functions of macrophages, thus aiding in the killing of phagocytosed bacteria. Not surprisingly, mice deficient in TNF receptor I are impaired in their ability to control the growth of several intracellular pathogens including *Salmonella*. IL-1 β has effects similar to TNF α and also acts on the liver to induce production of acute phase reactants and on the hypothalamus to influence body temperature regulation.

The anti-microbial armamentarium of neutrophils includes a number of proteases that are stored in granules (76). Upon appropriate stimulation, these granules fuse with the phagolysosome and discharge their contents into the environment of the internalized bacteria. However, some of the granules also undergo exocytosis, resulting in the release of the proteases outside the cell, with consequent degradation of extracellular matrix components and damage to epithelial cells (31). The importance of this aspect of the inflammatory response is highlighted by the observation that inhibition of neutrophil elastase results in significant amelioration of dextran sulfate sodium (DSS)-induced colitis in mice (77). Interestingly, some neutrophil proteases also have anti-inflammatory effects. Neutrophil elastase and cathepsin G degrade flagellin from several kinds of bacteria, including *Salmonella*, thereby inhibiting flagellin-induced inflammatory responses (78). It will be important and informative to determine the role of neutrophil proteases in *Salmonella* enteritis in vivo.

One of the consequences of the *Salmonella*-induced inflammatory response is the induction of cell death. Intestinal epithelial cells may die from the direct effects of the infection or in response to various inflammatory mediators. Neutrophils are relatively short-lived cells, die once they have fulfilled their anti-bacterial function, and are phagocytosed by macrophages. Macrophages themselves are induced to undergo apoptosis as a result of the actions of the *Salmonella* SipB effector protein (discussed below) or in response to stimulation by LPS (79). Cell death may facilitate the spread of *Salmonella*, but may also lead to the generation of additional inflammatory stimuli as a result of the release of intracellular contents such as uric acid (80).

Phagocyte Sensing of Bacteria

The activation of phagocyte anti-microbial and inflammatory mechanisms occurs in response to microbial triggers that are recognized by specific cellular sensors. The TLRs constitute one important class of such sensors, and both macrophages and neutrophils express several of these receptors (81). Work from our laboratory, as well those of others, has shown that TLR2 and TLR4 play important roles in inducing the production of TNF α and other inflammatory cytokines by macrophages during *Salmonella* infection (82–85). TLR signals also lead to increased expression of IL-1 β mRNA and intracellular protein (74).

In addition to the TLRs, which are expressed on the cell surface, recent work has demonstrated the existence of PRRs in the cytosol. This intracellular sensing system consists of a large family of structurally related proteins that have been given the name NLRs, for NACHT (a domain common to this family)-leucine rich repeats (86). The pattern of expression of these proteins varies, but several NLRs are expressed in macrophages and neutrophils. Their roles in innate immunity are the subject of active, on-going investigations. One important function of a subset of these proteins that has already come to light is the activation of caspase 1. This protease is essential for the proteolytic maturation and secretion of IL-1 β and the related cytokine IL-18, and is also involved in the induction of apoptosis. Various members of the NLR family are required for inducing IL-1 β processing and secretion in response to a number of endogenous and exogenous triggers, including ATP, uric acid crystals, Gram-negative and Gram-positive bacteria, bacterial toxins, and bacterial RNA (87–90). Several laboratories have also reported that *Salmonella*-induced IL- β secretion depends on at least two of the NLR proteins, ASC and IPAF (89,91,92). IPAF has a direct activating interaction with caspase 1, while ASC functions as an adaptor that links other NLRs with the protease. The specific *Salmonella* component involved in IPAF-dependent induction of IL-1 β secretion was found to be soluble flagellin (91,92), and it was demonstrated that the entry of flagellin into the cytosol required the SPI1 TTSS (91). This requirement would help to explain earlier observations indicating a role for SipB in caspase 1 activation and apoptosis, although the significance of the direct binding of SipB to caspase 1 needs further clarification (93). It is not yet clear whether the flagellin is translocated through the needle complex of the SPI1 TTSS, or whether it leaks from the phagosome via the pore made by this structure. The exact roles played by IPAF and ASC in flagellin sensing are also matters of speculation. It is tempting to think that the leucine-rich repeat domain of IPAF may bind to flagellin directly, but there is no evidence yet to support this idea. It has been suggested that ASC may act to stabilize the IPAF–caspase 1 interaction, but it is also possible that it could be part of a separate NLR complex required for the response to cytosolic flagellin. Recent experiments have confirmed the protective role of *Salmonella*-induced caspase 1 activation in vivo (94), in contrast to earlier observations suggesting that caspase 1 activity facilitated colonization of Peyer's patches by the bacteria (95).

Experiments in cultured macrophages have revealed an additional mechanism for recognizing bacteria in the cytosol (96). These studies demonstrated that although the majority of intracellular *Salmonella* reside within the SCV, a small population of the bacteria escape into the cytosol. Such bacteria become marked by the accumulation of polyubiquitinated proteins on their surface and associate with the proteasome. Inhibition of proteasomal activity enhanced intracellular growth of *Salmonella*, suggesting that ubiquitination of cytosolic bacteria may be an important antimicrobial defense mechanism. Elucidation of the molecular pathway that leads to tagging of bacteria with ubiquitin, which may involve the NLRs or other types of intracellular PRRs, will be an important line of further investigation.

Why have cells evolved special systems for responding to bacteria in the cytosol when it would appear that the TLRs would be adequate for bacterial sensing? NLRs may serve

simply as a redundant back-up system, but there may be an advantage to recognizing the presence of bacterial components specifically in the cytoplasm. Pathogens are more likely to end up in the cytosolic compartment, e.g., both *Listeria* and *Shigella* can escape from the phagosome into the cytoplasm. Pathogens are also more likely to have virulence mechanisms such as the SPI1 TTSS of *Salmonella* or the type IV secretory system of *Helicobacter pylori* that are able to facilitate the movement of bacterial molecules into the cytosol. Therefore, triggering of the NLRs and the consequent release of IL-1 β would indicate a potentially more “dangerous” situation than TLR activation alone. The induction of apoptosis by NLRs may represent an extreme response to such a situation by helping to eliminate pathogen-infected cells. Clearly, this is a growth area in inflammation research, and the next few years (if not months) will no doubt reveal the molecular details of NLR function, and the role played by these proteins in a variety of inflammatory processes, including *Salmonella* enteritis.

The Role of IFN γ in the Intestinal Response to *Salmonella*

IFN γ plays a very important role in protection against several intracellular pathogens, including *Salmonella*. This cytokine is typically produced by T cells of the Th1 phenotype and is required for activating various macrophage microbicidal functions. Humans with defects in Th1 development, or in the production of or response to IFN γ , are susceptible to severe mycobacterial and *Salmonella* infections (97). IFN γ can also be produced early in the course of infection from cells of the innate immune system such as macrophages and natural killer (NK) cells, as well as intra-epithelial lymphocytes and memory CD8 $^{+}$ T cells, and can have a major impact on subsequent immunologic events (62). We have shown recently that IFN γ has a significant influence on *Salmonella* enteritis in the mouse model (19). The intestinal inflammation induced by *Salmonella* in adult IFN γ -deficient mice was significantly attenuated, as indicated by fewer histopathological abnormalities and lower levels of inflammatory mediator expression. Some of these effects could be related to the influence of IFN γ on maturation of intestinal innate immune defenses, but direct regulation of IEC and macrophage function with respect to cytokine/ chemokine expression and anti-microbial mechanisms also probably plays a role.

Conclusion

The intestinal inflammation induced by *S. typhimurium* is an integral part of the innate immune response to the organism in which IECs, macrophages, and neutrophils play key roles. It serves to protect the host against spread of the infection, and in cases where the response is effective, the bacteria are confined to the gut and ultimately eliminated. When it is poorly developed, as in young infants or pre-weaned mouse pups, the risk of systemic dissemination of the bacteria is increased. Does the enteritis serve the pathogen in any way? The fact that several of the SPI1 TTSS effector proteins activate IEC inflammatory responses raises this question, and it is possible that the induction of diarrhea and the consequent spread of the organism to the environment may facilitate its transmission to other hosts.

Salmonella enteritis in humans is generally self-limited and is not usually treated except in infants less than 3 mo of age. With increased understanding of the cells and molecules involved in the pathogenesis of this condition, opportunities to minimize even the transient morbidity associated with the infection will become available through the development of novel therapeutic and prophylactic interventions. Targeting cytokines such as TNF α and IL-1 β , or transcription factors such as NF- κ B, is a potential approach to controlling the inflammation, one that can now be tested fairly easily using the mouse model.

One issue that we have not touched on is the resolution of *Salmonella*-induced intestinal inflammation. To our knowledge, there have been no studies on how healing occurs in this condition. This is likely to be an interesting and fruitful line of investigation, but it is hampered by the fact that *Salmonella* infection in the C57BL/6 strain that is typically used in the murine enteritis model results in death in 5–7 d in a large number of animals. Mouse strains such as 129/SvJ that carry the resistant allele of *Nramp1* are able to survive the acute infection (66), and it would be worthwhile to analyze the course of *Salmonella* enteritis in these strains. Insights into the processes involved in the restitution of normal intestinal structure and function could lead to the identification of additional molecular targets for therapeutic manipulation, and could prove beneficial for the control of both *Salmonella*-induced and other types of intestinal inflammation.

Acknowledgments

Work in our laboratory is supported by the NIH through R01 AI48815 and R21 AI065619.

References

1. Rabsch W, Tschape H, Baumler A. Non-typhoidal salmonellosis: emerging problems. *Microbes Infect.* 2001; 3:237–247. [PubMed: 11358718]
2. Torok TJ, Tauxe RV, Wise RP, et al. A large community outbreak of salmonellosis caused by intentional contamination of restaurant salad bars. *JAMA.* 1997; 278:389–395. [PubMed: 9244330]
3. Graham SM. Salmonellosis in children in developing and developed countries and populations. *Curr Opin Infect Dis.* 2002; 15:507–512. [PubMed: 12686884]
4. Bhan MK, Bahl R, Bhatnagar S. Typhoid and paratyphoid fever. *Lancet.* 2005; 366:749–762. [PubMed: 16125594]
5. Murase T, Yamada M, Muto T, Matsushima A, Yamai S. Fecal excretion of *Salmonella enterica* serovar Typhimurium following a food-borne outbreak. *J Clin Microbiol.* 2000; 38:3495–3497. [PubMed: 10970414]
6. Nishio T, Nakamori J, Miyazaki K. Survival of *Salmonella typhi* in oysters. *Zentralbl Bakteriell Mikrobiol Hyg [B].* 1981; 172:415–426.
7. Wait DA, Sobsey MD. Comparative survival of enteric viruses and bacteria in Atlantic Ocean seawater. *Water Sci Technol.* 2001; 43:139–142. [PubMed: 11464742]
8. Galan JE. *Salmonella* interactions with host cells: type III secretion at work. *Annu Rev Cell Dev Biol.* 2001; 17:53–86. [PubMed: 11687484]
9. Blaser MJ, Newman LS. A review of human salmonellosis: I. Infective dose. *Rev Infect Dis.* 1982; 4:1096–1106. [PubMed: 6760337]
10. Hurley BP, McCormick BA. Translating tissue culture results into animal models: the case of *Salmonella typhimurium*. *Trends Microbiol.* 2003; 11:562–569. [PubMed: 14659688]
11. Tsois RM, Adams LG, Ficht TA, Baumler AJ. Contribution of *Salmonella typhimurium* virulence factors to diarrheal disease in calves. *Infect Immun.* 1999; 67:4879–4885. [PubMed: 10456944]
12. Kent TH, Formal SB, Labrec EH. *Salmonella* gastroenteritis in Rhesus monkeys. *Arch Pathol.* 1966; 82:272–279. [PubMed: 4957636]
13. Rout WR, Formal SB, Dammin GJ, Giannella RA. Pathophysiology of *Salmonella* diarrhea in the Rhesus monkey: Intestinal transport, morphological and bacteriological studies. *Gastroenterology.* 1974; 67:59–70. [PubMed: 4210000]
14. Barthel M, Hapfelmeier S, Quintanilla-Martinez L, et al. Pretreatment of mice with streptomycin provides a *Salmonella enterica* serovar Typhimurium colitis model that allows analysis of both pathogen and host. *Infect Immun.* 2003; 71:2839–2858. [PubMed: 12704158]
15. Hapfelmeier S, Ehrbar K, Stecher B, Barthel M, Kremer M, Hardt WD. Role of the *Salmonella* pathogenicity island 1 effector proteins SipA, SopB, SopE, and SopE2 in *Salmonella enterica* subspecies 1 serovar Typhimurium colitis in streptomycin-pretreated mice. *Infect Immun.* 2004; 72:795–809. [PubMed: 14742523]

16. Stecher B, Hapfelmeier S, Muller C, Kremer M, Stallmach T, Hardt WD. Flagella and chemotaxis are required for efficient induction of *Salmonella enterica* serovar Typhimurium colitis in streptomycin-pretreated mice. *Infect Immun*. 2004; 72:4138–4150. [PubMed: 15213159]
17. Hapfelmeier S, Stecher B, Barthel M, et al. The *Salmonella* pathogenicity island (SPI)-2 and SPI-1 type III secretion systems allow *Salmonella* serovar typhimurium to trigger colitis via MyD88-dependent and MyD88-independent mechanisms. *J Immunol*. 2005; 174:1675–1685. [PubMed: 15661931]
18. Hapfelmeier S, Hardt WD. A mouse model for *S. typhimurium*-induced enterocolitis. *Trends Microbiol*. 2005; 13:497–503. [PubMed: 16140013]
19. Rhee SJ, Walker WA, Cherayil BJ. Developmentally regulated intestinal expression of IFN γ and its target genes and the age-specific response to enteric *Salmonella* infection. *J Immunol*. 2005; 175:1127–1136. [PubMed: 16002714]
20. Suar M, Jantsch J, Hapfelmeier S, et al. Virulence of broad- and narrow-host-range *Salmonella enterica* serovars in the streptomycin-pretreated mouse model. *Infect Immun*. 2006; 74:632–644. [PubMed: 16369020]
21. McCormick BA, Stocker BA, Laux DC, Cohen PS. Roles of motility, chemotaxis, and penetration through and growth in intestinal mucus in the ability of an avirulent strain of *Salmonella typhimurium* to colonize the large intestine of streptomycin-treated mice. *Infect Immun*. 1988; 56:2209–2217. [PubMed: 3044995]
22. Murray RA, Lee CA. Invasion genes are not required for *Salmonella enterica* serovar Typhimurium to breach the intestinal epithelium: evidence that *Salmonella* pathogenicity island 1 has alternative functions during infection. *Infect Immun*. 2000; 68:5050–5055. [PubMed: 10948124]
23. Humphries AD, Townsend SM, Kingsley RA, Nicholson TL, Tsolis RM, Baumler AJ. Role of fimbriae as antigens and intestinal colonization factors of *Salmonella* serovars. *FEMS Microbiol Lett*. 2001; 201:121–125. [PubMed: 11470349]
24. Dorsey CW, Laarakker MC, Humphries AD, Weening EH, Baumler AJ. *Salmonella enterica* serotype Typhimurium MisL is an intestinal colonization factor that binds fibronectin. *Mol Microbiol*. 2005; 57:196–211. [PubMed: 15948960]
25. Kubori T, Galan JE. Temporal regulation of *Salmonella* virulence effector function by proteasome-dependent protein degradation. *Cell*. 2003; 115:333–342. [PubMed: 14636560]
26. Hayden MS, Ghosh S. Signaling to NF- κ B. *Genes Dev*. 2004; 18:2195–2224. [PubMed: 15371334]
27. Huang FC, Werne A, Li Q, Galyov EE, Walker WA, Cherayil BJ. Cooperative interactions between flagellin and SopE2 in the epithelial interleukin-8 response to *Salmonella enterica* serovar Typhimurium infection. *Infect Immun*. 2004; 72:5052–5062. [PubMed: 15321998]
28. Mrsny RJ, Gewirtz AT, Siccaldi D, et al. Identification of hepoxilin A3 in inflammatory events: a required role in neutrophil migration across intestinal epithelia. *Proc Natl Acad Sci USA*. 2004; 101:7421–7426. [PubMed: 15123795]
29. Silva M, Song C, Nadeau WJ, Matthews JB, McCormick BA. *Salmonella typhimurium* SipA-induced neutrophil transepithelial migration: involvement of a PKC- α -dependent signal transduction pathway. *Am J Physiol Gastrointest Liver Physiol*. 2004; 286:G1024–1031. [PubMed: 14739142]
30. Zen K, Parkos CA. Leukocyte-epithelial interactions. *Curr Opin Cell Biol*. 2003; 15:557–564. [PubMed: 14519390]
31. Ginzberg HH, Shannon PT, Suzuki T, et al. Leukocyte elastase induces epithelial apoptosis: role of mitochondrial permeability changes and Akt. *Am J Physiol Gastrointest Liver Physiol*. 2004; 287:G286–298. [PubMed: 15194561]
32. Norris FA, Wilson MP, Wallis TS, Galyov EE, Majerus PW. SopB, a protein required for virulence of *Salmonella dublin*, is an inositol phosphate phosphatase. *Proc Natl Acad Sci USA*. 1998; 95:14057–14059. [PubMed: 9826652]
33. Feng Y, Wente SR, Majerus PW. Overexpression of the inositol phosphatase SopB in human 293 cells stimulates cellular chloride influx and inhibits nuclear mRNA export. *Proc Natl Acad Sci USA*. 2001; 98:875–879. [PubMed: 11158563]

34. Steele-Mortimer O, Knodler LA, Marcus SL, et al. Activation of Akt/protein kinase B in epithelial cells by the *Salmonella typhimurium* effector sigD. *J Biol Chem*. 2000; 275:37718–37724. [PubMed: 10978351]
35. Huang FC, Li Q, Cherayil BJ. A phosphatidylinositol-3-kinase-dependent anti-inflammatory pathway activated by *Salmonella* in epithelial cells. *FEMS Microbiol Lett*. 2005; 243:265–270. [PubMed: 15668028]
36. Neish AS, Gewirtz AT, Zeng H, et al. Prokaryotic regulation of epithelial responses by inhibition of I κ B α ubiquitination. *Science*. 2000; 289:1560–1563. [PubMed: 10968793]
37. Collier-Hyams LS, Zeng H, Sun J, et al. Cutting edge: *Salmonella* AvrA effector inhibits the key proinflammatory, anti-apoptotic NF- κ B pathway. *J Immunol*. 2002; 169:2846–2850. [PubMed: 12218096]
38. Zhang S, Santos RL, Tsolis RM, et al. The *Salmonella enterica* serotype Typhimurium effector proteins SipA, SopA, SopB, SopD, and SopE2 act in concert to induce diarrhea in calves. *Infect Immun*. 2002; 70:3843–3855. [PubMed: 12065528]
39. Zhang S, Adams LG, Nunes J, Khare S, Tsolis RM, Baumler AJ. Secreted effector proteins of *Salmonella enterica* serotype Typhimurium elicit host-specific chemokine profiles in animal models of typhoid fever and enterocolitis. *Infect Immun*. 2003; 71:4795–4803. [PubMed: 12874363]
40. Raffatellu M, Wilson RP, Chessa D, et al. SipA, SopA, SopB, SopD, and SopE2 contribute to *Salmonella enterica* serotype Typhimurium invasion of epithelial cells. *Infect Immun*. 2005; 73:146–154. [PubMed: 15618149]
41. Beutler B, Jiang Z, Georgel P, et al. Genetic analysis of host resistance: Toll-like receptor signaling and immunity at large. *Annu Rev Immunol*. 2006; 24:353–389. [PubMed: 16551253]
42. Gewirtz AT, Navas TA, Lyons S, Godowski PJ, Madara JL. Cutting edge: bacterial flagellin activates basolaterally expressed TLR5 to induce epithelial proinflammatory gene expression. *J Immunol*. 2001; 167:1882–1885. [PubMed: 11489966]
43. Gewirtz AT, Simon PO Jr, Schmitt CK, et al. *Salmonella typhimurium* translocates flagellin across intestinal epithelia, inducing a proinflammatory response. *J Clin Invest*. 2001; 107:99–109. [PubMed: 11134185]
44. Lyons S, Wang L, Casanova JE, Sitaraman SV, Merlin D, Gewirtz A. *Salmonella typhimurium* transcytoses flagellin via an SPI2-mediated vesicular transport pathway. *J Cell Sci*. 2004; 117:5771–1580. [PubMed: 15507487]
45. Smith KD, Andersen-Nissen E, Hayashi F, et al. Toll-like receptor 5 recognizes a conserved site on flagellin required for protofilament formation and bacterial motility. *Nat Immunol*. 2003; 4:1247–1253. [PubMed: 14625549]
46. Subramanian N, Qadri A. Lysophospholipid sensing triggers secretion of flagellin from pathogenic *Salmonella*. *Nat Immunol*. 2006; 7:583–589. [PubMed: 16648855]
47. Backhed F, Hornef M. Toll-like receptor 4-mediated signaling by epithelial surfaces: necessity or threat? *Microbes Infect*. 2003; 5:951–959. [PubMed: 12941387]
48. Abreu MT, Fukata M, Arditi M. TLR signaling in the gut in health and disease. *J Immunol*. 2005; 174:4453–4460. [PubMed: 15814663]
49. Humphries AD, Townsend SM, Kingsley RA, Nicholson TL, Tsolis RM, Baumler AJ. Role of fimbriae as antigens and intestinal colonization factors of *Salmonella* serovars. *FEMS Microbiol Lett*. 2001; 20:121–125. [PubMed: 11470349]
50. Tukul C, Raffatellu M, Humphries AD, et al. CsgA is a pathogen-associated molecular pattern of *Salmonella enterica* serotype Typhimurium that is recognized by Toll-like receptor 2. *Mol Microbiol*. 2005; 58:289–304. [PubMed: 16164566]
51. Selsted ME, Ouellette AJ. Mammalian defensins in the antimicrobial immune response. *Nat Immunol*. 2005; 6:551–557. [PubMed: 15908936]
52. Ayabe T, Satchell DP, Wilson CL, Parks WC, Selsted ME, Ouellette AJ. Secretion of microbicidal alpha-defensins by intestinal Paneth cells in response to bacteria. *Nat Immunol*. 2000; 1:113–118. [PubMed: 11248802]

53. Salzman NH, Ghosh D, Huttner KM, Paterson Y, Bevins CL. Protection against enteric salmonellosis in transgenic mice expressing a human intestinal defensin. *Nature*. 2003; 422:522–526. [PubMed: 12660734]
54. Lehrer RI, Ganz T. Cathelicidins: a family of endogenous antimicrobial peptides. *Curr Opin Hematol*. 2002; 9:18–22. [PubMed: 11753073]
55. Hase K, Eckmann L, Leopard JD, Varki N, Kagnoff MF. Cell differentiation is a key determinant of cathelicidin LL-37/human cationic antimicrobial protein 18 expression by human colon epithelium. *Infect Immun*. 2002; 70:953–963. [PubMed: 11796631]
56. Rosenberger CM, Gallo RL, Finlay BB. Interplay between antibacterial effectors: a macrophage antimicrobial peptide impairs intracellular *Salmonella* replication. *Proc Natl Acad Sci USA*. 2004; 101:2422–2427. [PubMed: 14983025]
57. Smith PD, Ochsenbauer-Jambor C, Smythies LE. Intestinal macrophages: unique effector cells of the innate immune system. *Immunol Rev*. 2005; 206:149–159. [PubMed: 16048547]
58. Smythies LE, Sellers M, Clements RH. Human intestinal macrophages display profound inflammatory anergy despite avid phagocytic and bacteriocidal activity. *J Clin Invest*. 2005; 115:66–75. el al. [PubMed: 15630445]
59. Malaviya R, Abraham SN. Mast cell modulation of immune responses to bacteria. *Immunol Rev*. 2001; 179:16–24. [PubMed: 11292019]
60. Chatterjea D, Burns-Guydish SM, Sciuto TE, Dvorak A, Contag CH, Galli SJ. Adoptive transfer of mast cells does not enhance the impaired survival of Kit(W)/Kit(W-v) mice in a model of low dose intraperitoneal infection with bioluminescent *Salmonella typhimurium*. *Immunol Lett*. 2005; 99:122–129. [PubMed: 15894120]
61. Niess JH, Brand S, Gu X, et al. CX3CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance. *Science*. 2005; 307:254–258. [PubMed: 15653504]
62. Sundquist M, Rydstrom A, Wick MJ. Immunity to *Salmonella* from a dendritic point of view. *Cell Microbiol*. 2004; 6:1–11. [PubMed: 14678326]
63. Tukel C, Raffatellu M, Chessa D, Wilson RP, Akcelik M, Baumler AJ. Neutrophil influx during non-typhoidal salmonellosis: who is in the driver's seat? *FEMS Immunol Med Microbiol*. 2006; 46:320–329. [PubMed: 16553804]
64. Jung HC, Eckmann L, Yang SK, et al. A distinct array of proinflammatory cytokines is expressed in human colon epithelial cells in response to bacterial invasion. *J Clin Invest*. 1995; 95:55–65. [PubMed: 7814646]
65. Depaolo RW, Lathan R, Rollins BJ, Karpus WJ. The chemokine CCL2 is required for control of murine gastric *Salmonella enterica* infection. *Infect Immun*. 2005; 73:6514–6522. [PubMed: 16177325]
66. Monack DM, Bouley DM, Falkow S. *Salmonella typhimurium* persists within macrophages in the mesenteric lymph nodes of chronically infected Nrp1^{+/+} mice and can be reactivated by IFN γ neutralization. *J Exp Med*. 2004; 199:231–241. [PubMed: 14734525]
67. Abrahams GL, Hensel M. Manipulating cellular transport and immune responses: dynamic interactions between intracellular *Salmonella enterica* and its host cells. *Cell Microbiol*. 2006; 8:728–737. [PubMed: 16611223]
68. Bispham J, Tripathi BN, Watson PR, Wallis TS. *Salmonella* pathogenicity island 2 influences both systemic salmonellosis and *Salmonella*-induced enteritis in calves. *Infect Immun*. 2001; 69:367–377. [PubMed: 11119526]
69. Everest P, Ketley J, Hardy S, et al. Evaluation of *Salmonella typhimurium* mutants in a model of experimental gastroenteritis. *Infect Immun*. 1999; 67:2815–2821. [PubMed: 10338486]
70. Blackwell JM, Goswami T, Evans CA, et al. SLC11A1 (formerly NRAMP1) and disease resistance. *Cell Microbiol*. 2001; 3:773–784. [PubMed: 11736990]
71. Nakamura A, Mori Y, Hagiwara K, et al. Increased susceptibility to LPS-induced endotoxin shock in secretory leukoprotease inhibitor (SLPI)-deficient mice. *J Exp Med*. 2003; 197:669–674. [PubMed: 12615907]
72. Thuraisingam T, Sam H, Moisan J, Zhang Y, Ding A, Radzioch D. Delayed cutaneous wound healing in mice lacking solute carrier 11a1 (formerly Nrp1): correlation with decreased

- expression of secretory leukocyte protease inhibitor. *J Invest Dermatol.* 2006; 126:890–901. [PubMed: 16470178]
73. Chlosta S, Fishman DS, Harrington L, et al. The iron efflux protein ferroportin regulates the intracellular growth of *Salmonella enterica*. *Infect Immun.* 2006; 74:3065–3067. [PubMed: 16622252]
 74. Dinarello CA. Blocking IL-1 in systemic inflammation. *J Exp Med.* 2005; 201:1355–1359. [PubMed: 15867089]
 75. Hehlhans T, Pfeffer K. The intriguing biology of the tumour necrosis factor/tumour necrosis factor receptor superfamily: players, rules and the games. *Immunology.* 2005; 115:1–20. [PubMed: 15819693]
 76. Pham CT. Neutrophil serine proteases: specific regulators of inflammation. *Nat Rev Immunol.* 2006; 6:541–550. [PubMed: 16799473]
 77. Morohoshi Y, Matsuoka K, Chinen H, et al. Inhibition of neutrophil elastase prevents the development of murine dextran sulfate sodium-induced colitis. *J Gastroenterol.* 2006; 41:318–324. [PubMed: 16741610]
 78. Lopez-Boado YS, Espinola M, Bahr S, Belaouaj A. Neutrophil serine proteinases cleave bacterial flagellin, abrogating its host response-inducing activity. *J Immunol.* 2004; 172:509–515. [PubMed: 14688361]
 79. Guiney DG. The role of host cell death in *Salmonella* infections. *Curr Top Microbiol Immunol.* 2005; 289:131–150. [PubMed: 15791954]
 80. Shi Y, Evans JE, Rock KL. Molecular identification of a danger signal that alerts the immune system to dying cells. *Nature.* 2003; 425:516–521. [PubMed: 14520412]
 81. Parker LC, Whyte MK, Dower SK, Sabroe I. The expression and roles of Toll-like receptors in the biology of the human neutrophil. *J Leukoc Biol.* 2005; 77:886–892. [PubMed: 15728244]
 82. Li Q, Cherayil BJ. Role of Toll-like receptor 4 in macrophage activation and tolerance during *Salmonella enterica* serovar Typhimurium-infection. *Infect Immun.* 2003; 71:4873–4882. [PubMed: 12933828]
 83. Royle M, Totemeyer S, Aldridge LC, Maskell DJ, Bryant CE. Stimulation of Toll-like receptor 4 by lipopolysaccharide during cellular invasion by live *Salmonella typhimurium* is a critical but not exclusive event leading to macrophage responses. *J Immunol.* 2003; 170:5445–5454. [PubMed: 12759420]
 84. Lembo A, Kalis C, Kirschning CJ, et al. Differential contribution of Toll-like receptors 4 and 2 to the cytokine response to *Salmonella enterica* serovar Typhimurium and *Staphylococcus aureus* in mice. *Infect Immun.* 2003; 71:6058–6062. [PubMed: 14500530]
 85. Weiss DS, Raupach B, Takeda K, Akira S, Zychlinsky A. Toll-like receptors are temporally involved in host defense. *J Immunol.* 2004; 172:4463–4469. [PubMed: 15034062]
 86. Martinon F, Tschopp J. NLRs join TLRs as innate sensors of pathogens. *Trends Immunol.* 2005; 26:447–454. [PubMed: 15967716]
 87. Kanneganti TD, Ozoren N, Body-Malapel M, et al. Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature.* 2006; 440:233–236. [PubMed: 16407888]
 88. Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature.* 2006; 440:237–241. [PubMed: 16407889]
 89. Mariathasan S, Weiss DS, Newton K, et al. Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature.* 2006; 440:228–232. [PubMed: 16407890]
 90. Sutterwala FS, Ogura Y, Szczepanik M, et al. Critical role for NALP3/CIA1/Cryopyrin in innate and adaptive immunity through its regulation of caspase-1. *Immunity.* 2006; 24:317–327. [PubMed: 16546100]
 91. Miao EA, Alpuche-Aranda CM, Dors M, et al. Cytoplasmic flagellin activates caspase 1 and secretion of IL-1 β via Ipaf. *Nat Immunol.* 2006; 7:569–575. [PubMed: 16648853]
 92. Franchi L, Amer A, Body-Malapel M, et al. Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1 β in *Salmonella*-infected macrophages. *Nat Immunol.* 2006; 7:576–582. [PubMed: 16648852]

93. Hersh D, Monack DM, Smith MR, Ghori N, Falkow S, Zychlinsky A. The *Salmonella* invasin SipB induces macrophage apoptosis by binding to caspase-1. *Proc Natl Acad Sci USA*. 1999; 96:2396–2401. [PubMed: 10051653]
94. Lara-Tejero M, Sutterwala FS, Ogura Y, et al. Role of the caspase-1 inflammasome in *Salmonella typhimurium* pathogenesis. *J Exp Med*. 2006; 203:1407–1412. [PubMed: 16717117]
95. Monack DM, Hersh D, Ghori N, Bouley D, Zychlinsky A, Falkow S. *Salmonella* exploits caspase-1 to colonize Peyer's patches in a murine typhoid model. *J Exp Med*. 2000; 192:249–258. [PubMed: 10899911]
96. Perrin AJ, Jiang X, Birmingham CL, So NS, Brumell JH. Recognition of bacteria in the cytosol of mammalian cells by the ubiquitin system. *Curr Biol*. 2004; 14:806–811. [PubMed: 15120074]
97. van de Vosse E, Hoeve MA, Ottenhof TH. Human genetics of intracellular infectious diseases: molecular and cellular immunity against mycobacteria and salmonellae. *Lancet Infect Dis*. 2004; 4:739–749. [PubMed: 15567123]
98. Zhou D, Moosekar MS, Galan JE. Role of the *S. typhimurium* actin-binding protein SipA in bacterial internalization. *Science*. 1999; 283:2092–2095. [PubMed: 10092234]
99. Hayward RD, Koronakis V. Direct nucleation and bundling of actin by the SipC protein of invasive *Salmonella*. *EMBO J*. 1999; 18:4926–434. [PubMed: 10487745]
100. Collazo CM, Galan JE. The invasion-associated type III system of *Salmonella typhimurium* directs the translocation of Sip proteins into the host cell. *Mol Microbiol*. 1997; 24:747–756. [PubMed: 9194702]
101. Jones MA, Wood MW, Mullan PB, Watson PR, Wallis TS, Galyov EE. Secreted effector proteins of *Salmonella dublin* act in concert to induce enteritis. *Infect Immun*. 1998; 66:5799–5804. [PubMed: 9826357]
102. Hardt WD, Chen LM, Schuebel KE, Bustelo XR, Galan JE. *S. typhimurium* encodes an activator of Rho GTPases that induces membrane ruffling and nuclear responses in host cells. *Cell*. 1998; 93:815–826. [PubMed: 9630225]
103. Bakshi CS, Singh VP, Wood MW, Jones PW, Wallis TS, Galyov EE. Identification of SopE2, a *Salmonella* secreted protein which is highly homologous to SopE and involved in bacterial invasion of epithelial cells. *J Bacteriol*. 2000; 182:2341–2344. [PubMed: 10735884]
104. Stender S, Friebe A, Linder S, Rohde M, Mirol S, Hardt WD. Identification of SopE2 from *Salmonella typhimurium*, a conserved guanine nucleotide exchange factor for Cdc42 of the host cell. *Mol Microbiol*. 2000; 36:1206–1221. [PubMed: 10931274]
105. Fu Y, Galan JE. A *Salmonella* protein antagonizes Rac-1 and Cdc42 to mediate host-cell recovery after bacterial invasion. *Nature*. 1999; 401:293–297. [PubMed: 10499590]
106. Hardt WD, Galan JE. A secreted *Salmonella* protein with homology to an avirulence determinant of plant pathogenic bacteria. *Proc Natl Acad Sci USA*. 1997; 94:9887–9892. [PubMed: 9275221]

Fig. 1.

Initial interactions between *Salmonella* and IECs. Several effector proteins are translocated into the IEC cytosol via the SPI1 TTSS. Some of these proteins are involved in the actin remodeling and plasma membrane protrusions that are required for cell invasion, and also activate intracellular signals. SipA may also act from an extracellular location to induce hepxilin A3 secretion. Flagellin is transcytosed from the apical surface of the epithelium by vesicular transport, and upon reaching the basolateral aspect is able to activate TLR5. The combined effect of the signals activated by the effector proteins and flagellin/TLR5 interactions results in changes in gene expression and the secretion of inflammatory mediators from either the basolateral or apical surface.

Fig. 2.

Interactions between *Salmonella* and phagocytic cells. Phagocytic cells such as macrophages recognize and respond to the bacteria through cell surface TLRs, which sense microbial components such as LPS, and intracellular NLRs, which sense flagellin translocated into the cytosol through the SPI1 TTSS. These interactions lead to the secretion of inflammatory cytokines and the induction of apoptosis.

Table 1

Summary of the Functions of SPI1 Effector Proteins and Their Roles in Intestinal Inflammation

Effector protein	Molecular function	Role in inflammation	Reference
SipA	Actin crosslinking		98
	Hepoxillin A3 induction	PMN migration/ diarrhea in calves, murine colitis	15,28 38
SipB	Caspase binding, translocase	Diarrhea in calves	11,93
SipC	Actin bundling, nucleation, translocase		99
SipD	Translocase		100
SopA	Unknown	Diarrhea in calves	38
SopB	Inositol phosphatase	Diarrhea in calves	15,32
			38
SopD	Unknown	Diarrhea in calves	38,101
SopE	GEF for Cdc42/Rac1	Murine colitis	15,102
SopE2	GEF for Cdc42/Rac1	Diarrhea in calves, murine colitis	15,38
			103,104
SptP	Phosphatase and GTPase activation	None	38,105
AvrA	Inhibition of NF- κ B activation	None	38,106