Published in final edited form as:

J Org Chem. 2011 November 4; 76(21): 8944–8954. doi:10.1021/jo201658y.

Microwave-Based Reaction Screening: Tandem Retro-Diels-Alder/Diels-Alder Cycloadditions of *ortho*-Quinol Dimers

Suwei Dong[†], Katharine J. Cahill[‡], Moon-Il Kang[§], Nancy H. Colburn[§], Curtis J. Henrich[§], Jennifer A. Wilson[§], John A. Beutler[§], Richard P. Johnson^{‡,*}, and John A. Porco Jr.^{†,*}

[†]Department of Chemistry and Center for Chemical Methodology and Library Development (CMLD-BU), Boston University, Boston, Massachusetts 02215

[‡]Department of Chemistry, University of New Hampshire, Durham, NH 03824

[§]Laboratory of Cancer Prevention and Molecular Targets Laboratory, Center for Cancer Research, NCI-Frederick, Frederick, MD 21702

Abstract

We have accomplished a parallel screen of cycloaddition partners for *ortho*-quinols utilizing a plate-based microwave system. Microwave irradiation improves the efficiency of retro-Diels-Alder/Diels-Alder cascades of *ortho*-quinol dimers which generally proceed in a diastereoselective fashion. Computational studies indicate that asynchronous transition states are favored in Diels-Alder cycloadditions of *ortho*-quinols. Subsequent biological evaluation of a collection of cycloadducts has identified an inhibitor of activator protein-1 (AP-1), an oncogenic transcription factor.

INTRODUCTION

ortho-Quinols are highly reactive 2,4-cyclohexadienones and have been proposed as precursors for a number of natural products. *ortho*-Quinol dimers are common building blocks for preparation of bicyclo[2.2.2]octenones *via* retro-Diels-Alder reaction followed by a Diels-Alder cycloaddition with external dienophiles.¹ We recently reported the synthesis of the natural product chamaecypanone C (**1**)² involving a retro-Diels-Alder/Diels-Alder cascade of 2,4-cyclohexadienone (*o*-quinol) dimer **2** and diaryl enone **3** under thermal, oxidative conditions (Scheme 1). [4+2] Cycloaddition of the derived *ortho*-quinol monomer **4** and 2,4-diaryl cyclopentadienone **5** was found to proceed in a highly regio- and diastereoselective manner. This result, along with reactions with a few dienophiles, prompted us to further evaluate cycloadditions of **4** and related *ortho*-quinols with a range of reaction partners, in order to better understand the mode of reactivity for dienophiles as well as diastereoselectivity with respect to the *o*-quinol.

Retro-[4+2]/[4+2] reactions of dimers derived from *o*-quinols and masked *o*-benzoquinones (MOBs) have been utilized for preparation of reactive 2,4-cyclohexadienones enroute to

*porco@bu.edu & richard.johnson@unh.edu.

Supporting Information Available Experimental procedures and characterization data for all new compounds, including X-ray structure analysis of compound **10e**, X-ray crystallographic data (CIF), and detailed biological methods and results. This material is available free of charge *via* the Internet at <http://pubs.acs.org>.

bicyclo[2.2.2]octenones.^{1,3,4} However, most reported examples involve thermolysis at high temperatures for extended reaction times. In some cases, sealed tubes have been used at temperatures up to 220 °C,^{3b} where safety issues may become a concern. In comparison to conventional heating, microwave heating has been proven to be efficient in many cases leading to a dramatic reduction in reaction time.⁵ In particular, use of thermal susceptors or “sensitizers” including graphite⁶ and silicon carbide (SiC),⁷ has been shown to further improve the efficiency of thermal transformations, as these materials can reach high temperature under microwave irradiation in spite of the nature of reaction solvents.⁸ Recently, Kappe and coworkers have reported the development of sintered SiC microtiter plates for performing parallel synthesis in a multimode microwave reactor.⁹ Considering the reactivity of *ortho*-quinol dimers and the available literature precedents, we wished to utilize SiC plates in a high-throughput reaction partner assessment.¹⁰ In the current study, a broad panel of reaction partners in the retro-[4+2]/[4+2] cascade of *o*-quinol dimers have been evaluated under microwave conditions in an effort to study the scope and limitations of this methodology, to obtain both bicyclo[2.2.2]octenone and *cis*-decalin frameworks, and to understand the factors determining regio- and diastereoselectivity. To explain the observed reactivities and diastereoselectivities, in particular the regioselectivity for cycloadditions, computational studies of Diels-Alder reactions between *o*-quinols and select dienophiles/dienes were also conducted.¹¹ In this paper, we report the results of this study as well as biological screening of the resulting collection of complex cycloadducts.

RESULTS AND DISCUSSION

Reaction Optimization

We first evaluated reactions between the readily accessible dimer (–)-**2**¹² and *N*-phenylmaleimide (**6**) under microwave conditions in various solvents (Table 1).¹³ Although reactions went to completion within 30 min in either DMF (entry 1) or DMA (entry 2) to afford cycloadduct **7**, we noticed that use of mesitylene as solvent afforded cleaner reactions (entry 3). Further experiments (entry 4) indicated that reactions could be completed within 15 min (μW, 180 °C) in comparison to use of conventional heating (entry 5) which required approximately 90 minutes for completion.²

Screening of Reaction Partners

Utilizing the optimized microwave conditions, 57 reaction partners were evaluated in a reaction screen.¹² Representative structures were shown in Figure 1. A number of alkene partners were selected based on retro-[4+2]/[4+2] cycloadditions of *o*-quinol dimers and MOB dimers reported in the literature.^{1,3} Representative dipolarophiles and dienes were also included. Reactions were conducted on a 0.009 mmol scale using 3.0 mg of dimer (–)-**2** as substrate, 10 equiv of the corresponding reaction partner, and 100 μL of mesitylene under microwave irradiation at 188 °C. After 60 min, reaction mixtures were filtered through a silica gel plug, concentrated, and evaluated using ultra performance liquid chromatography (UPLC)-MS which allows for short analytical run times (approx. three minutes).¹⁴ Based on UPLC-MS data, reactions showing major peaks in the HPLC trace were scaled up (0.06 mmol, 20 mg substrate) using a SiC chip as passive heating element in a single mode microwave reactor.¹² The results of the reaction screening indicated that dimer **2** was completely consumed in all cases. Of the 57 reactions, 12 afforded products corresponding to [4+2] adducts between the *o*-quinol and alkene reaction partners, and other reactions afforded product **8** (Scheme 2), likely generated from a dienone-phenol rearrangement of intermediate **4**.¹⁵ The formation of **8** is in accordance with literature reported thermal degradation results for dimer **2**,¹⁶ and suggests that the corresponding reaction partners are unreactive in the cycloaddition event.

As shown in Table 2, *o*-quinol **4** may react with activated alkenes, including the normal-demand dienophiles MVK (**9a**), indene (**9b**), 4-methoxy styrene (**9c**), the inverse-demand dienophiles dihydrofuran (DHF **9d**), vinylene carbonate (**9e**), and the alkynes dimethyl acetylenedicarboxylate (DMAD, **9f**) and phenylacetylene (**9g**). In all cases, Diels-Alder cycloaddition between *o*-quinol **4** and dienophiles **9** proceeded in a highly diastereoselective fashion with [4+2] adducts **10a–g** isolated in moderate to excellent yields. Structure elucidation of products using 2D NMR experiments or X-ray crystallography confirmed that *endo*-[4+2] cycloadducts were the predominate products isolated from the reactions,¹² which is in accordance with both experimental results and theoretical calculations in literature that secondary orbital overlap is believed to govern *endo*-selectivity.¹⁷ Facial selectivity could be explained by either sterics or hyperconjugation effects according to our computational studies (*vide infra*).

In the cases of reaction with dienes (**11**, Table 3), *o*-quinol **4** was found to react as either a 4π or 2π reaction partner to afford bicyclooctenone derivative **12** or *cis*-decalin product **13**. Reaction with 1-phenyl substituted 1,3-butadiene **11a** (entry 1) cleanly afforded a mixture of bicyclooctenones *endo*-**12aa** and *exo*-**12aβ** in a 4:1 ratio favoring the *endo*-adduct. When the 2-substituted-1,3-butadiene β-myrcene (**11b**, entry 2) was used in the reaction, *cis*-decalin **13b** was found to be the major product.¹⁸ Considering the result that a mixture of two compounds was generated at lower reaction temperature (150 °C),² cycloadduct **13b** should be thermodynamically more favored in comparison to its bicyclooctenone isomer obtunone.¹⁹ Similarly, reaction with 2,3-disubstituted-1,3-butadiene **11c** (entry 3) produced *cis*-decalin **13c** as the major product in excellent yield (94%). While reaction with cyclopentadiene dimer **11d** (entry 4) generated cycloadduct **12d** as the sole product in almost quantitative isolated yield, use of cyclohexadiene **11e** (entry 5) as reaction partner led to a more complex product mixture. Structure elucidation confirmed that compounds **12e** and **13e** were the major products in this instance along with a small amount of the rearranged bicyclooctene **14** (approx. 10%).

In order to probe possible interconversion between adducts **12e** and **13e**, pure compound **12e** was resubjected to the reaction conditions (μ W, SiC, mesitylene, 180 °C, 15 min). Product analysis indicated that a mixture of cycloadducts **12e**, **13e**, and **14** was generated in a 2:1:1 ratio. Extended reaction time (1 h) or elevated temperature (200 °C) did not change the ratio in which case further decomposition was observed. It is plausible that compounds **12e** and **13e** may interconvert through Cope rearrangement²⁰ (Figure 2, pathway a) or *via* a retro-[4+2] reaction of **12e** followed by recombination of *o*-quinol and cyclohexadiene (pathway b). To further understand this process, trapping experiments were conducted as shown in Scheme 3. In both experiments, no crossover Diels-Alder cycloadduct was observed; reactions either afforded a mixture of **12e/13e/14** from **12e** or recovered starting materials from cycloadduct **7**. These results support an intramolecular [3,3] sigmatropic rearrangement mechanism for the interconversion of cycloadducts **12e** and **13e**.²¹

The generation of product **14** may be explained by thermal α -ketol rearrangement of hydroxy enone **13e** (Figure 2, pathway c).²² A similar ring contraction has been observed in a related α -methyl- α -hydroxycyclohexanone system under basic conditions.²³ However, use of related basic conditions (LiHMDS/THF or Al₂O₃/hexanes) did not effect ketol rearrangement of **13e**.

Based on the results from our reaction screen and follow-up product analyses, we noticed that activated alkenes or alkynes are more reactive in the reaction with the model *o*-quinol. Moreover, *cis*-alkenes in general showed better reactivity than *trans*-alkenes, 1,1-disubstituted, and trisubstituted alkenes. Acyclic 1,3-dienes were found to react as 4π components only if they lacked substitution at the C1 or C4 positions, and displayed

reactivity as 2π components in the cases of less substituted alkenes. Taken together, these results suggest that Diels-Alder cycloaddition of *o*-quinols is an electronically and sterically demanding process. These observations are also in accordance with literature reported examples of 2,4-cyclohexadienones.^{1,3}

To further evaluate the scope of the microwave-assisted protocol, *ortho*-quinol dimers (+)-**15**,¹¹ **16**,²⁴ **17**, and **18**²⁵ were subjected to optimized reaction conditions (μ W, SiC, mesitylene, 170 °C, 15 min, Table 4). In all cases, good to excellent isolated yield of the desired cycloadducts **19** to **25** were obtained as single diastereomers.

Computational Studies

Previously computational modeling has been employed mainly in studies of MOBs as dienes in [4+2] cycloadditions.^{3a,26} Quideau and coworkers have also utilized calculations to study the dimerization of *ortho*-quinols.²⁷ We wished to further study Diels-Alder cycloadditions of *o*-quinols and dienophiles/dienes computationally to further understand the energetics, regio-, and stereochemical features of the cycloadditions.

In our studies, B3LYP calculations²⁸ first showed (Table 5, entry 1) that cycloreversion of dimer **2** is modestly endothermic, with forward and reverse barriers of 23.9 and 12.3 kcal/mol, respectively. The lowest energy transition state has C_2 symmetry; similar results have been described for related cycloadditions.²³ Predicted reaction energetics support a low transient concentration of dienone **4**. Cycloaddition of **4** with **9a**, **9d**, **9b** as representative dienophiles is predicted in each case (entries 2 – 4) to have a low reaction barrier, with exothermicity favorable to a unidirectional process. For each of these reactions, we optimized all *eight* potential transition states; the observed product was correctly assigned by theory in all cases. In every favored cycloaddition, the shorter transition state bond connects to C-5 of dienone **4** (Figure 3). Two failed reactants, **27** and **28**, showed unexpectedly higher reaction barriers (entries 6 and 7) when compared to structural analogues **9a** and **9b**. It is noteworthy that the barriers for two reaction modes of myrcene (**11b**, entry 5) are nearly equal, consistent with the experimental product distribution (ca. 1:1) at lower reaction temperature (150 °C).² The lower energy **13b** was the sole product generated from the reaction at higher temperature (180 °C, *cf.* Table 3, entry 2), presumably from cycloadduct **26** *via* Cope rearrangement.

The observed ensemble of selectivity and the correlation between theory and experiment are striking. In all cases, cycloaddition stereochemistry favors the usual Alder *endo* rule, with additional facial selectivity *syn* to the less sterically demanding hydroxyl group. Hyperconjugative effects (Cieplak-Fallis model)²⁹ from the methyl group in the transition state may also contribute to facial selectivity.^{24, 30} Dienone **4** has a high lying HOMO (−6.9 eV) and a low-lying LUMO (−2.3 eV) and as a consequence reacts with both electron-rich and electron-poor dienophiles. In every case, the preferred regiochemistry follows initial bonding to C5 of the dienone which maximizes resonance stabilization in the asynchronous transition state. In general, reactions with lower barriers showed more asynchronous transition state structures.³¹

In a side by side comparison of MVK (**9a**) and cyclopentenone **27**, similar dienophiles electronically, the MVK π system shows better secondary-orbital overlap in transition states in reactions with *o*-quinol **4** (Figure 3, TS2). In contrast, the *exo* C=O double bond of cyclic ketone **27** does not align well with the *o*-quinol in the transition state (TS6). Similarly, in TS4 the arene has a more optimal π - π interaction with the 4π partner, while in TS7 the dihydronaphthalene (**28**) aromatic ring is oriented nearly perpendicular to the cyclohexadienone ring, presumably to minimize steric interactions, thereby raising TS energetics by 5.1 kcal/mol in comparison to the indene case (*cf.* Table 5, entries 4 and 7).

Besides the sterics and electronics of dienophiles, these results suggest that secondary-orbital overlap of dienophiles with *ortho*-quinols may be another important factor determining dienophile reactivity.

Competition between cycloreversion, [3,3] sigmatropic shifts¹⁹ and secondary dienone-phenol³² or α -ketol²¹ rearrangements add complexity to these results. For example, DFT calculations show that (–)-**2** should undergo a degenerate and homochiral Cope rearrangement with a barrier of 22.4 kcal/mol which should compete with cycloreversion. Figure 4 summarizes predicted energetics for the dienone **4** + 1,3-cyclohexadiene potential surface. No transition state for [4+2] cycloaddition leading directly to **13e** (TS9) was identified with DFT calculations. Repeated attempts to locate TS9 instead optimized to TS8 which leads to **12e**. Based on these predictions, the most likely pathway to **13e** is through a Cope rearrangement of **12e** which has a predicted barrier of 29.6 kcal/mol. According to their relative energetics, Cope rearrangement of **12e** should be favored over cycloreversion. Our trapping experiments (Scheme 3) supported only a [3,3] sigmatropic rearrangement for interconversion of **12e** and **13e**, consistent with computations.

The rearrangement of **13e** to **14** can proceed by thermal isomerization of neutral **13e** or through catalysis by acid, base, or lithiation. Computational results are summarized in Scheme 4. The predicted thermal barrier (41.1 kcal/mol) is too high to be operative. Protonation, deprotonation, or lithiation of **13e** all decrease the predicted barrier significantly and affect product energetics. Adventitious proton catalysis seems most compatible with our experimental results. For the most favorable case, B3LYP/6-31+G(d) calculations predict that deprotonated **13e** should undergo an exothermic rearrangement with a barrier of only 13.1 kcal/mol. The apparent conflict of this result with our LiHMDS-catalyzed experiments (*vide infra*) was resolved by showing that rearrangement of lithiated **13e** is now endothermic by 3.5 kcal/mol because O-Li-O bridging creates product strain.

Rearrangement of **4** to **8** can proceed by similarly diverse mechanisms.^{21,31} DFT computations for the methyl migration step (Scheme 5) showed low barriers for ionic pathways. Once again, our experimental results are consistent with trace acid catalysis.

Biological Studies

Both enantiomers of synthesized bicyclo[2.2.2]octenones and *cis*-decalins were subjected to screening at 20 μ M for inhibition in three biological assays at the National Cancer Institute (NCI): AP-1 (activator protein-1, an oncogenic transcription factor),³³ TRAIL (Tumor necrosis factor- α -related apoptosis-inducing ligand) sensitization,³⁴ and HIF-2 (hypoxia inducible factor 2).³⁵ Compound (–)-**31** (Figure 5, inset) showed selective inhibition against AP-1 at 4 μ M concentration with luciferase reporter assays in HEK293 cells. TPA (12-O-tetradecanoylphorbol-13-acetate) induced AP-1 activity was repressed by more than 50% after pretreatment with the compound at 8 μ M. However, the compound did not show inhibition of NF- κ B (nuclear factor kappa B) or SRE (serum response element) dependent transcription at the same concentration (Figure 5). SRE was also used as a proliferation control which can measure cytotoxic activity for the compound. Thus, this result suggests that compound (–)-**31** targets the events needed for AP-1 activation selectively, rather than other transcription factors such as HIF-2, NF- κ B, and SRE. As AP-1 is required for tumor promotion and progression,³⁶ identification of novel and specific AP-1 inhibitors such as bicyclooctenone (–)-**31** may be beneficial for cancer prevention and therapy. Although further experiments in the NCI 60-cell screen at 10 μ M showed that (–)-**31** did not pass the threshold for further evaluation of cell growth inhibition, analogues of compound **31** may be pursued in future studies for SAR analysis.

CONCLUSION

We have utilized microwave-assisted parallel screening for evaluation of Diels-Alder reactions of *ortho*-quinol dimers in an effort to identify reactive reaction partners among a broad panel of compounds. Elucidation of products, analysis of the results, and computational studies allowed us to not only confirm the diastereoselectivity of [4+2] cycloadditions of *ortho*-quinols, but also determine that sterics and proper secondary orbital alignment are the major governing factors for dienophile reactivity. The high concordance between experimental and calculated results suggest that theoretical calculations of energetics may be used to pre-screen reaction partners for [4+2] cycloadditions of *ortho*-quinols.³⁷ Furthermore, biological evaluation of a set of compounds generated in the study showed that bicyclo[2.2.2]octenones may be worthwhile scaffolds for development of screening libraries, as novel AP-1 inhibitory activity was observed for one member of the small library. Further studies involving retro-Diels-Alder/Diels-Alder cascade processes to produce natural products and analogues are currently in progress and will be reported in future publications.

EXPERIMENTAL SECTION

Optimization of the retro-[4+2]/[4+2] reaction

A 10 mL microwave tube with a silicon carbide chip was charged with dimer **2** (10.0 mg, 0.030 mmol) and *N*-phenylmaleimide **6** (15.6 mg, 0.150 mmol).¹² 0.5 mL of solvent was added under an argon atmosphere and the reaction was irradiated in a microwave reactor (stir off) at the indicated temperature (*cf.* Table 1). Reactions in DMF (entry 1) and DMA (entry 2) were directly submitted to UPLC-MS analysis, and crude reactions were diluted with EtOAc, washed with water, brine, dried over MgSO₄. The organic fraction was concentrated, dried *in vacuo*, and purified by column chromatography. Reactions in mesitylene were stopped after indicated reaction time, the crude mixture was passed through a short silica gel plug, flushed using hexanes to remove mesitylene, and finally flushed with EtOAc to elute both the product and excess maleimide **6**. The organic fraction was concentrated, dried *in vacuo*, analyzed by UPLC-MS, and purified by column chromatography to afford cycloadduct **7**.

Reaction partner screening

In a microwave reaction vessel (4×48 well SiC block) were charged dimer **2** (3.0 mg, 0.009 mmol), dienophiles/dienes (10 equiv), and 100 μ L mesitylene.¹² The resulting mixtures were heated in a microwave reactor with 48 well SiC block (stir off) at 180 °C (IR temperature) for 1 h. After the reaction, the crude mixture was passed through a short silica gel plug, flushed using hexanes to remove mesitylene, and finally flushed with EtOAc to elute both the product and excess dienophiles/dienes. The organic fraction was finally concentrated and analyzed by UPLC-MS.

General procedure for microwave reaction of *o*-quinol dimers (20–30 mg scale)

A 10 mL microwave tube with silicon carbide chip was charged with *o*-quinol dimer (0.060 mmol) and dienophile (equivalents, *cf.* Tables 2–4). 0.5 mL of mesitylene was added under an argon atmosphere and the reaction was irradiated in a microwave reactor at 170 or 180 °C (IR temperature) for 15 min (stir off). Reactions were stopped after indicated reaction time, the crude mixture was passed through a short silica gel plug, flushed using hexanes to remove mesitylene (and in a few cases excess nonpolar dienophiles), and finally flushed with EtOAc to elute the crude products. The organic fraction was concentrated, dried *in vacuo*, and the residue was purified by silica gel column chromatography to afford the corresponding Diels-Alder adducts.

(1*R*,3*S*,4*S*,7*S*)- 3-Hydroxy-7-(4-methoxy)phenyl-3-methyl-6-(propan-2-yl)bicyclo[2.2.2]oct-5-en-2-one (10c)

Dimer **2** (20.0 mg, 0.060 mmol) and 4-vinylanisole **9c** (81.0 μ L, 0.60 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=10:1 to 8:1) followed by preparative TLC (CH₂Cl₂/EtOAc=10:1) to afford product **10c** (35.5 mg, 98%) as a yellow oil. *R*_f = 0.41 (hexanes/EtOAc=2:1); IR (thin film): ν_{max} 3434, 2959, 1721, 1511, 1248, 1178 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 7.07 (d, *J* = 8.0 Hz, 2H), 6.79 (d, *J* = 8.0 Hz, 2H), 6.18 (d, *J* = 7.2 Hz, 1H), 3.77 (s, 3H), 3.38 (dd, *J* = 8.4, 5.6 Hz, 1H), 3.06 (d, *J* = 0.8 Hz, 1H), 2.94 (ddd, *J* = 6.0, 3.2, 2.4 Hz, 1H), 2.68 (ddd, *J* = 12.8, 9.6, 2.4 Hz, 1H), 2.53 (s, 1H), 1.90 (sept., *J* = 6.8 Hz, 1H), 1.61 (ddd, *J* = 12.8, 6.0, 2.8 Hz, 1H), 1.28 (s, 3H), 0.89 (d, *J* = 6.8 Hz, 3H), 0.73 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 213.0, 158.4, 144.7, 135.9, 128.7 (two carbons overlapping), 125.7, 113.6 (two carbons overlapping), 72.1, 59.0, 55.2, 43.1, 39.9, 33.1, 27.6, 25.9, 20.5, 19.7; HRMS–ESI (*m/z*): [M–H₂O+H]⁺ calcd for C₁₉H₂₃O₂, 283.1698; found, 283.1713. [α]_D²² = +36.4° (*c* = 0.67, CHCl₃).

(1*R*,2*S*,6*R*,7*S*,9*S*)- 9-Hydroxy-9-methyl-11-(propan-2-yl)-3,5-dioxatricyclo[5.2.2.0^{2,6}]undec-10-ene-4,8-dione (10e)

Dimer **2** (20.0 mg, 0.060 mmol) and vinylene carbonate **9e** (77.3 μ L, 1.20 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=3:1) to afford product **10e** (23.1 mg, 76%) as a white solid. *R*_f = 0.11 (hexanes/EtOAc=3:1); mp 135–137 °C; IR (thin film): ν_{max} 3426, 2964, 1795, 1731, 1164, 1057 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 5.97 (d, *J* = 6.4 Hz, 1H), 5.36 (dd, *J* = 8.0, 3.6 Hz, 1H), 5.04 (dd, *J* = 8.0, 3.6 Hz, 1H), 3.64 (dd, *J* = 3.6, 2.0 Hz, 1H), 3.38 (dd, *J* = 6.4, 3.6 Hz, 1H), 2.43 (s, 1H), 2.39 (sept., *J* = 6.8 Hz, 1H), 1.32 (s, 3H), 1.022 (d, *J* = 6.8 Hz, 3H), 1.016 (d, *J* = 6.8 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz): δ 205.5, 154.4, 145.5, 122.2, 74.6, 74.3, 71.9, 54.4, 45.8, 33.1, 25.2, 20.23, 20.18; HRMS–ESI (*m/z*): [2M + Na]⁺ calcd for C₂₆H₃₂NaO₁₀, 527.1893; found, 527.1880. [α]_D²² = +125.6° (*c* = 0.52, CHCl₃).

(1*R*,4*S*,7*S*)-Dimethyl-7-hydroxy-7-methyl-8-oxo-5-(propan-2-yl)bicyclo[2.2.2]octa-2,5-diene-2,3-dicarboxylate (10f)

Dimer **2** (20.0 mg, 0.060 mmol) and dimethyl acetylenedicarboxylate **9f** (74.0 μ L, 0.60 mmol) in 300 μ L of mesitylene were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (CH₂Cl₂/EtOAc=15:1 to 10:1) to afford product **10f** (33.3 mg, 90%) as a yellow oil. *R*_f = 0.30 (CH₂Cl₂/EtOAc=5:1); IR (thin film): ν_{max} 3485, 2959, 1726, 1265 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 6.03 (d, *J* = 6.4 Hz, 1H), 4.35 (d, *J* = 2.0 Hz, 1H), 4.04 (d, *J* = 6.0 Hz, 1H), 3.81 (s, 3H), 3.77 (s, 3H), 2.54 (s, 1H), 2.48 (doublet of sept., *J* = 6.8, 1.2 Hz, 1H), 1.32 (s, 3H), 1.02 (d, *J* = 6.8 Hz, 6H); ¹³C NMR (CDCl₃, 100 MHz): δ 201.9, 166.4, 164.2, 149.8, 143.2, 134.4, 123.7, 67.4, 58.1, 52.7, 52.6, 50.8, 31.9, 26.3, 20.3 (two carbons overlapping); HRMS–ESI (*m/z*): [M + Na]⁺ calcd for C₁₆H₂₀NaO₆, 331.1158; found, 331.1158. [α]_D²² = +58.7° (*c* = 0.51, CHCl₃).

(1*S*,3*S*,4*R*)- 3-Hydroxy-3-methyl-6-phenyl-7-(propan-2-yl)bicyclo[2.2.2]octa-5,7-dien-2-one (10g)

Dimer **2** (20.0 mg, 0.060 mmol) and vinylene carbonate **9g** (65.9 μ L, 1.20 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=8:1) to afford product **14** (25.8 mg, 80%) as a yellow oil. *R*_f = 0.27 (hexanes/EtOAc=3:1); IR (thin film): ν_{max} 3434, 2963, 1724, 1445, 1068 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 7.46–7.44 (m, 2H), 7.36–7.32 (m, 2H), 7.29–7.26 (m,

1H), 6.73 (dd, $J = 6.0, 2.0$ Hz, 1H), 6.08 (dd, $J = 6.4, 2.0$ Hz, 1H), 4.43 (dd, $J = 2.0, 2.0$ Hz, 1H), 3.79 (ddd, $J = 6.4, 6.4, 2.0$ Hz, 1H), 2.53 (sept., $J = 6.8$ Hz, 1H), 2.30 (s, 1H), 1.35 (s, 3H), 1.07 (d, $J = 6.8$ Hz, 3H), 1.05 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 205.3, 149.2, 141.1, 136.0, 129.2, 128.6 (two carbons overlapping), 127.8, 125.3 (two carbons overlapping), 125.1, 69.5, 60.3, 48.5, 32.1, 26.4, 20.7 (two carbons overlapping); HRMS–ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{18}\text{H}_{20}\text{NaO}_2$, 291.1361; found, 291.1363. $[\alpha]_{\text{D}}^{22} = +27.6^\circ$ ($c = 0.81$, CHCl_3).

(1*R*,3*S*,4*S*,7*R*)-3-Hydroxy-3-methyl-7-[(*E*)-2-phenylethenyl]-6-(propan-2-yl)bicyclo[2.2.2]oct-5-en-2-one (12a α)

Dimer **2** (20.0 mg, 0.120 mmol) and 1-phenyl-1,3-butadiene³⁸ (**11a**, 76.1 mg, 1.20 mmol) were thermolyzed following the general procedure. The crude mixture was purified by preparative TLC (hexanes/EtOAc=4:1) to afford product **12a α** (28.8 mg, 81%) as a white solid. $R_f = 0.22$ (hexanes/EtOAc=5:1); mp 68–70 °C; IR (thin film): ν_{max} 3439, 2962, 2932, 1725, 1449, 1126 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 7.27~7.26 (m, 4H), 7.20~7.17 (m, 1H), 6.37 (dd, $J = 16.0, 1.0$ Hz, 1H), 6.11 (ddd, $J = 7.0, 1.5, 1.5$ Hz, 1H), 5.86 (dd, $J = 16.0, 8.5$ Hz, 1H), 3.09 (dd, $J = 2.0, 2.0$ Hz, 1H), 3.00~2.94 (m, 1H), 2.83 (ddd, $J = 7.0, 3.0, 2.5$ Hz, 1H), 2.53 (ddd, $J = 12.5, 8.5, 3.0$ Hz, 1H), 2.50 (bs, 1H), 2.30 (doublet of sept., $J = 7.0, 1.0$ Hz, 1H), 1.26 (s, 3H), 1.21 (ddd, $J = 13.0, 5.0, 2.5$ Hz, 1H), 1.02 (d, $J = 7.0$ Hz, 3H), 0.98 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 212.9, 144.4, 137.1, 132.6, 129.7, 128.5 (two carbons overlapping), 127.2, 126.2, 126.0 (two carbons overlapping), 72.3, 56.6, 42.9, 39.2, 33.1, 27.3, 25.9, 20.9, 20.5; HRMS–ESI (m/z): $[\text{M}-\text{H}_2\text{O}+\text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{23}\text{O}$, 279.1749; found, 279.1748. $[\alpha]_{\text{D}}^{22} = +119.5^\circ$ ($c = 0.91$, CHCl_3).

(1*R*,3*S*,4*S*,7*S*)-3-Hydroxy-3-methyl-7-[(*E*)-2-phenylethenyl]-6-(propan-2-yl)bicyclo[2.2.2]oct-5-en-2-one (12a β)

Utilizing the same protocol as compound **12a α** , product **12a β** (19.8 mg, 97%) was isolated as a white solid. $R_f = 0.32$ (hexanes/EtOAc=5:1); mp 104–106 °C; IR (thin film): ν_{max} 3440, 2962, 2871, 1726, 1449, 1142, 1081 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 7.33~7.26 (m, 4H), 7.21~7.17 (m, 1H), 6.40 (d, $J = 15.6$ Hz, 1H), 6.10 (ddd, $J = 15.6, 8.8, 1.6$ Hz, 1H), 6.04 (dd, $J = 6.8, 1.2$ Hz, 1H), 3.03 (dd, $J = 1.6, 1.6$ Hz, 1H), 2.85~2.82 (m, 1H), 2.65~2.58 (m, 1H), 2.40 (sept., $J = 6.8$ Hz, 1H), 2.30 (s, 1H), 2.12 (ddd, $J = 12.8, 5.2, 2.0$ Hz, 1H), 1.77~1.70 (m, 1H), 1.28 (s, 3H), 1.05 (d, $J = 6.8$ Hz, 3H), 1.04 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 213.4, 147.8, 137.1, 132.1, 130.3, 128.5 (two carbons overlapping), 127.2, 126.3 (two carbons overlapping), 125.5, 73.5, 56.6, 43.1, 42.5, 32.6, 26.4, 26.3, 20.54, 20.50; HRMS–ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{25}\text{O}_2$, 297.1855; found, 297.1867. $[\alpha]_{\text{D}}^{22} = +137.8^\circ$ ($c = 0.57$, CHCl_3).

(1*S*,4*aR*,8*aS*)-1-Hydroxy-1-methyl-6-[4-methylpent-3-en-1-yl]-4-(propan-2-yl)-4a,5,8,8a-tetrahydronaphthalen-2(1*H*)-one (13b)

Dimer **2** (20.0 mg, 0.060 mmol) and β -myrcene **11c** (103.3 μL , 0.60 mmol) were thermolyzed following the general procedure. The crude mixture was purified and **13b** was isolated as a colorless oil (20.8 mg, 57%). ^1H -NMR, ^{13}C -NMR, and IR spectra for compound **13b** were found to be identical to literature reported data.¹⁷ $R_f = 0.38$ (hexanes/EtOAc=5:1); $[\alpha]_{\text{D}}^{22} = +217.9^\circ$ ($c = 1.01$, CHCl_3).

(1*S*,4*aR*,8*aS*)-1-Hydroxy-1,6,7-trimethyl-4-(propan-2-yl)-4a,5,8,8a-tetrahydronaphthalen-2(1*H*)-one (13c)

Dimer **2** (20.0 mg, 0.060 mmol) and 2,3-dimethyl-1,3-butadiene **11c** (67.9 μL , 0.60 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=20:1 to 15:1) to afford product **13c** (27.3 mg, 92%) as

an oil. $R_f = 0.38$ (hexanes/EtOAc=8:1); IR (thin film): $\nu_{\max}$ 3491, 2968, 2926, 1722, 1671, 1242, 1152 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 5.85 (d, $J = 2.8$ Hz, 1H), 3.55 (s, 1H), 2.94~2.91 (m, 1H), 2.38 (sept., $J = 6.8$ Hz, 1H), 2.35~2.30 (m, 3H), 2.07 (dd, $J = 17.6$, 6.8 Hz, 1H), 1.59 (s, 3H), 1.48 (s, 3H), 1.32 (s, 3H), 1.06 (d, $J = 6.8$ Hz, 3H), 1.01 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 203.1, 172.8, 126.1, 123.3, 120.0, 76.6, 44.4, 36.8, 33.3, 30.8, 29.4, 24.4, 22.4, 20.6, 19.0, 18.7; HRMS–ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{24}\text{O}_2$, 249.1855; found, 249.1876. $[\alpha]^{22}_{\text{D}} = +133.6^\circ$ ($c = 0.46$, CHCl_3).

(1S,2R,7S,8R,10S)-10-Hydroxy-10-methyl-12-(propan-2-yl)tricyclo-[6.2.2.0^{2,7}]dodeca-5,11-dien-9-one (12e)

Dimer **2** (60.0 mg, 0.180 mmol) and 1,3-cyclohexadiene **11e** (343.0 μL , 3.60 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel column (hexanes:EtOAc=20:1 to 15:1) to afford **12e** (51.8 mg, 57%) as a white solid. $R_f = 0.32$ (hexanes/EtOAc=5:1); mp 46–48 $^\circ\text{C}$; IR (thin film): $\nu_{\max}$ 3443, 2960, 2928, 1725, 1366, 1137 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 5.89 (d, $J = 7.6$ Hz, 1H), 5.87~5.82 (m, 1H), 5.42 (ddd, $J = 9.6$, 2.4, 2.4 Hz, 1H), 3.00 (dd, $J = 2.0$, 2.0 Hz, 1H), 2.79~2.72 (m, 2H), 2.64~2.62 (m, 1H), 2.52 (s, 1H), 2.22 (doublet of sept., $J = 7.2$, 0.8 Hz, 1H), 1.93~1.86 (m, 1H), 1.84~1.75 (m, 1H), 1.61 (dddd, $J = 12.4$, 10.0, 4.8, 4.8 Hz, 1H), 1.23 (s, 3H), 1.21~1.16 (m, 1H), 0.94 (d, $J = 6.8$ Hz, 3H), 0.92 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 213.6, 144.8, 130.2, 128.4, 125.3, 72.9, 56.3, 49.2, 35.5, 32.9, 31.8, 26.3, 25.9, 23.5, 20.8, 20.5; HRMS–ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{23}\text{O}_2$, 247.1698; found, 247.1721. $[\alpha]^{22}_{\text{D}} = +160.4^\circ$ ($c = 1.17$, CHCl_3).

(1R,2S,3S,7R,8S)-3-Hydroxy-3-methyl-6-(propan-2-yl)tricyclo-[6.2.2.0^{2,7}]dodeca-5,9-dien-4-one (13e)

Utilizing the same protocol for compound **12e**, product **13e** (22.5 mg, 25%) was isolated as a white solid. $R_f = 0.39$ (hexanes: EtOAc = 5:1); mp 52–53 $^\circ\text{C}$; IR (thin film) $\nu_{\max}$: 3463, 2963, 2867, 1673, 1376, 1170 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 6.04 (dd, $J = 3.2$, 3.2 Hz, 1H), 5.84 (s, 1H), 5.77 (dd, $J = 3.2$, 3.2 Hz, 1H), 3.99 (s, 1H), 2.96 (dddd, $J = 4.4$, 3.2, 2.0, 2.0 Hz, 1H), 2.83 (dd, $J = 8.0$, 2.8 Hz, 1H), 2.78 (ddd, $J = 4.0$, 2.0, 2.0 Hz, 1H), 2.44 (sept., $J = 6.8$ Hz, 1H), 2.26 (dd, $J = 8.4$, 1.6 Hz, 1H), 1.64~1.59 (m, 1H), 1.52 (dddd, $J = 12.4$, 9.2, 2.8, 2.8 Hz, 1H), 1.40 (dddd, $J = 12.4$, 12.4, 3.2, 3.2 Hz, 1H), 1.27~1.20 (m, 1H), 1.19 (s, 3H), 1.08 (d, $J = 6.8$ Hz, 3H), 1.07 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 202.5, 170.8, 133.8, 132.7, 119.2, 73.9, 47.9, 45.7, 35.4, 32.8, 32.0, 29.9, 26.1, 24.5, 23.2, 20.9; HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{23}\text{O}_2$, 247.1698; found, 247.1707. $[\alpha]^{22}_{\text{D}} = -128.1^\circ$ ($c = 0.52$, CHCl_3).

1-[(1R,2S,3S,6R,7S)-3-hydroxy-5-(propan-2-yl)tricyclo-[5.2.2.0^{2,6}]-undeca-4,8-dien-3-yl]ethanone (14)

Utilizing the same protocol for compound **12g**, product **14** (8.2 mg, 9%) was isolated as a white solid. $R_f = 0.46$ (hexanes: EtOAc = 5:1); mp 98–100 $^\circ\text{C}$; IR (thin film) $\nu_{\max}$: 3450, 2952, 2936, 2915, 1698, 1367, 1102 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz): δ 6.27 (dd, $J = 7.2$, 7.2 Hz, 1H), 5.89 (dd, $J = 7.2$, 7.2 Hz, 1H), 5.03 (dd, $J = 1.5$, 1.5 Hz, 1H), 3.92 (s, 1H), 2.98 (ddd, $J = 8.5$, 2.2, 2.2 Hz, 1H), 2.77~2.75 (m, 1H), 2.58~2.55 (dd, $J = 2.0$ Hz, 1H), 2.34 (dd, $J = 8.2$, 2.2 Hz, 1H), 2.29 (sept., $J = 7.0$ Hz, 1H), 2.09 (2, 3H), 1.49 (dddd, $J = 12.0$, 9.5, 4.5, 2.2 Hz, 1H), 1.41 (dddd, $J = 12.0$, 9.0, 3.0, 3.0 Hz, 1H), 1.33 (dddd, $J = 11.5$, 11.5, 3.5, 3.5 Hz, 1H), 1.26~1.20 (m, 1H), 1.06 (d, $J = 7.0$ Hz, 3H), 1.04 (d, $J = 7.0$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 210.5, 159.5, 134.0, 129.2, 124.8, 90.8, 53.5, 48.7, 32.7, 30.9, 27.7, 25.1, 24.2, 23.6, 21.7, 20.8; HRMS–ESI (m/z): $[\text{M}-\text{H}_2\text{O}+\text{H}]^+$ calcd for $\text{C}_{16}\text{H}_{21}\text{O}$, 229.1592; found, 229.1597. $[\alpha]^{22}_{\text{D}} = +210.4^\circ$ ($c = 0.30$, CHCl_3).

Gradifloracin (16)

Oxidation of salicyl alcohol benzoate (50.0 mg, 0.17 mmol) according to the reported procedure¹¹ (1.2 equiv DIEA, 1.5 equiv $[(\text{-})\text{-sparteine}]_2\text{Cu}_2\text{O}_2(\text{PF}_6)_2$, 3 Å MS, O₂, CH₂Cl₂, -78 °C, 40 h) generated dimer **16** as a grey solid (15.0 mg, 28%, 61% brsm) after silica gel chromatography. The ¹H, ¹³C NMR, IR, and HRMS of **16** were identical to literature reported data.^{24,39} $[\alpha]^{22}_{\text{D}} = -5.1^\circ$ ($c = 0.57$, CHCl₃; lit.⁵⁵ $[\alpha]^{22}_{\text{D}} = -13.6^\circ$, $c = 0.728$, CHCl₃). The *ee* for **16** (24%) was determined based on chiral HPLC analysis of its derivative **23**.

Dimer (±)-17

Oxidation of 2,6-dimethylphenol (20.0 mg, 0.164 mmol) according to the reported procedure¹¹ [1.0 equiv LiOH·H₂O, 1.2 equiv (*N,N'*-di-*tert*-butylethylene-diamine)₂Cu₂O₂(PF₆)₂, 3 Å MS, O₂, CH₂Cl₂, -78 °C, 16 h] generated dimer **17** as a light yellow solid (16.2 mg, 72%) after silica gel chromatography. ¹H, ¹³C NMR, IR, and HRMS of **17** were identical to literature reported data.²⁴

(1*R*,2*S*,6*R*,7*S*,9*R*)-11-*tert*-Butyl-9-hydroxy-9-methyltricyclo[5.2.2.0^{2,6}]-undeca-4,10-dien-8-one (19)

According to the general procedure, dimer (+)-**15** (30.0 mg, 0.083 mmol) and dicyclopentadiene **18** (54.9 mg, 0.415 mmol) were thermolyzed according to General Procedure C. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=15:1) to afford product **19** (38.0 mg, 93%) as a white solid. $R_f = 0.26$ (hexanes/EtOAc=7:1); mp 92–93 °C; IR (thin film): ν_{max} 3425, 2963, 2928, 1721, 1364 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 5.85 (dd, $J = 6.8, 2.0$ Hz, 1H), 5.62–5.59 (m, 1H), 5.34–5.32 (m, 1H), 3.25 (dd, $J = 2.0, 2.0$ Hz, 1H), 3.20–3.10 (m, 2H), 2.83 (dddd, $J = 9.2, 4.8, 2.4, 2.4$ Hz, 1H), 2.63 (bs, 1H), 2.46 (dddd, $J = 16.8, 9.2, 2.0, 2.0, 2.0$ Hz, 1H), 1.94 (dddd, $J = 16.8, 6.0, 2.0, 2.0, 2.0$ Hz, 1H), 1.21 (s, 3H), 0.92 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz): δ 214.9, 148.7, 132.8, 130.4, 122.2, 73.2, 52.6, 50.0, 47.5, 38.2, 34.2, 33.2, 28.2 (three carbons overlapping), 26.3; HRMS–ESI (m/z): $[\text{M}-\text{H}_2\text{O}+\text{H}]^+$ calcd for C₁₆H₂₁O, 229.1592; found, 229.1604. $[\alpha]^{22}_{\text{D}} = -90.0^\circ$ ($c = 0.65$, CHCl₃).

(1*R*,2*S*,6*R*,7*S*,9*S*)-11-*tert*-Butyl-9-hydroxy-9-methyl-3,5-dioxatricyclo-[5.2.2.0^{2,6}]undec-10-ene-4,8-dione (20)

Dimer (+)-**15** (30.0 mg, 0.083 mmol) and methyl vinyl ketone **9** (105.4 μL , 1.66 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=5:1 to 3:1) to afford product **20** (34.8 mg, 78%) as a colorless oil. $R_f = 0.13$ (hexanes/EtOAc=3:1); IR (thin film): ν_{max} 3423, 2965, 1786, 1731, 1367, 1162, 1058 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 5.99 (d, $J = 6.8, 1.2$ Hz, 1H), 5.36 (dd, $J = 8.0, 3.6$ Hz, 1H), 5.03 (dd, $J = 8.0, 3.2$ Hz, 1H), 3.81 (dd, $J = 3.6, 2.0$ Hz, 1H), 3.39 (dd, $J = 6.8, 3.2$ Hz, 1H), 2.56 (bs, 1H), 1.31 (s, 3H), 1.04 (s, 9H); ¹³C NMR (CDCl₃, 100 MHz): δ 205.8, 154.4, 147.9, 121.2, 74.5, 74.2, 71.8, 52.9, 45.6, 34.9, 27.8 (three carbons overlapping), 25.3; HRMS–ESI (m/z): $[\text{2M}+\text{H}]^+$ calcd for C₂₈H₃₇O₁₀, 533.2387; found, 533.2399. $[\alpha]^{22}_{\text{D}} = -176.9^\circ$ ($c = 1.09$, CHCl₃).

(12-Hydroxy-13-oxotetracyclo-[9.2.2.0^{2,10}.0^{3,8}]pentadeca-3,5,7,14-tetraen-12-yl)methyl benzoate (21)

Gradifloracin **16** (10.0 mg, 0.020 mmol) and indene **9b** (23.3 μL , 0.20 mmol) in mesitylene were irradiated in microwave reactor at 170 °C for 15 min. Following the general procedure the obtained crude mixture was purified by silica gel chromatography (hexanes/EtOAc=10:1 to 8:1) to afford product **21** (11.8 mg, 80%) as a light yellow oil. $R_f = 0.38$ (hexanes/EtOAc, 3:1). IR (thin film) ν_{max} : 3443, 2917, 2849, 1726, 1692, 1272 cm⁻¹; ¹H NMR (CDCl₃, 400

MHz): δ 8.08 (dd, $J = 8.4, 1.2$ Hz, 2H), 7.60–7.56 (m, 1H), 7.45 (dd, $J = 8.4, 8.4$ Hz, 2H), 7.19–7.13 (m, 3H), 7.12–7.09 (m, 1H), 6.35 (dd, $J = 7.2, 7.2$ Hz, 1H), 5.94 (dd, $J = 7.2, 7.2$ Hz, 1H), 4.45 (d, $J = 12.0$ Hz, 1H), 4.28 (d, $J = 12.0$ Hz, 1H), 3.83 (dd, $J = 9.2, 2.0$ Hz, 1H), 3.57 (ddd, $J = 6.4, 2.8, 1.2$ Hz, 1H), 3.50–3.43 (m, 1H), 3.29 (ddd, $J = 6.4, 2.8, 1.2$ Hz, 1H), 3.23 (dd, $J = 16.8, 10.4$ Hz, 1H), 3.13 (s, 1H), 2.69 (dd, $J = 16.8, 4.4$ Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 209.9, 166.6, 144.3, 142.4, 133.3, 133.2, 129.8 (two carbons overlap), 129.6, 128.6, 128.5 (two carbons overlap), 127.3, 126.7, 124.4, 124.0, 74.3, 68.4, 53.1, 50.1, 45.4, 37.7, 33.9; HRMS–ESI (m/z): $[\text{M} + \text{Na}]^+$ calcd for $\text{C}_{23}\text{H}_{20}\text{NaO}_4$, 383.1259; found, 383.1252. $[\alpha]_D^{22} = -22.3^\circ$ ($c = 0.24$, CHCl_3). The *ee* for **21** (24%) was determined using a Waters Breeze HPLC System (*ChiralPak AD-H*, 150×4.6 mm, 10% isopropanol in hexane, 1.0 mL/min, retention time 12.3 min (minor enantiomer) and 14.6 min (major enantiomer)) using UV detection at 254 nm.

9-Hydroxy-7,9-dimethyl-4-phenyl-4-azatricyclo[5.2.2.0^{2,6}]-undec-10-ene-3,5,8-trione (**22**)

Dimer **17** (20.0 mg, 0.072 mmol) and *N*-phenylmaleimide **6** (37.4 mg, 0.216 mmol) were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=2:1) to afford product **2** (44.6 mg, 99%) as a light yellow solid. $R_f = 0.08$ (hexanes/EtOAc=3:1); mp 161–163 °C; IR (thin film): ν_{max} 3444, 2977, 1703, 1698, 1385, 1186 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 7.45–7.40 (m, 2H), 7.39–7.34 (m, 1H), 7.17 (dd, $J = 7.6, 1.6$ Hz, 2H), 6.42 (dd, $J = 8.4, 6.4$ Hz, 1H), 5.95 (dd, $J = 8.4, 1.2$ Hz, 1H), 3.85 (dd, $J = 8.4, 3.6$ Hz, 1H), 3.54 (ddd, $J = 6.4, 3.2, 1.6$ Hz, 1H), 2.96 (d, $J = 8.0$ Hz, 1H), 2.95 (s, 1H), 1.57 (s, 3H), 1.32 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 210.3, 177.1, 174.4, 133.9, 132.9, 131.6, 129.1 (two carbons overlapping), 128.7, 126.4 (two carbons overlapping), 70.8, 50.5, 45.4, 44.9, 40.9, 25.6, 14.8; HRMS–ESI (m/z): $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{18}\text{NO}_4$, 312.1236; found, 312.1249.

12-Hydroxy-12-methyl-14-(propan-2-yl)-tetracyclo[9.2.2.0^{2,10}.0^{3,8}]-pentadeca-3,5,7,14-tetraene-13-one (**23**)

Dimer **17** (20.0 mg, 0.072 mmol) and indene **11** (83.9 μL , 0.72 mmol) were thermolyzed following the general procedure. The crude mixture was purified by preparative TLC (hexanes/EtOAc=3:1) to afford product **23** (28.7 mg, 78%) as a light yellow solid. The ^1H , ^{13}C NMR, IR, and HRMS of **16** were identical to literature reported data.^{1a}

8-hydroxy-1,8-dimethyl-3-oxatricyclo[5.2.2.0^{2,6}]-undec-10-en-9-one (**24**)

Dimer **17** (20.0 mg, 0.072 mmol) and 2,3-dihydrofuran **9d** (111.9 μL , 1.48 mmol) were thermolyzed following the general procedure. The concentrated mixture was dissolved in 2 mL of CH_2Cl_2 followed by addition of MP-TsOH resin (100 mg), and the reaction was shaken for 30 min. The MP-TsOH resin was filtered away and the filtrate was concentrated. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=8:1 to 7:1) to afford product **24** (25.6 mg, 85%) as a white solid. $R_f = 0.27$ (hexanes/EtOAc=2:1); mp 117–119 °C; IR (thin film): ν_{max} 3429, 2973, 2933, 1726, 1451, 1366, 1080 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 6.33 (dd, $J = 7.2, 7.2$ Hz, 1H), 5.75 (dd, $J = 8.4, 1.6$ Hz, 1H), 3.89–3.82 (m, 2H), 3.53 (ddd, $J = 8.4, 8.4, 8.4$ Hz, 1H), 3.18 (ddd, $J = 8.4, 8.4, 8.4$ Hz, 1H), 2.92–2.90 (m, 1H), 2.50 (s, 1H), 2.10–2.00 (m, 1H), 1.64–1.55 (m, 1H), 1.32 (s, 3H), 1.24 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 213.4, 134.1, 132.6, 84.5, 71.9, 69.1, 55.1, 46.4, 39.0, 30.9, 26.5, 14.5; LRMS m/z (% relative intensity): 231.1 $[(\text{M} + \text{Na})^+]$, 31, 191.1 (100), 163.1 (70), 141.1 (34).

9-hydroxy-11-methyl-4-phenyl-9-(propan-2-yl)-4-azatricyclo[5.2.2.0^{2,6}]undec-10-ene-3,5,8-trione (**25**)

Dimer **18** (30.0 mg, 0.090 mmol) and *N*-phenylmaleimide **6** (46.9 mg, 0.271 mmol) in mesitylene were thermolyzed following the general procedure. The crude mixture was purified by silica gel chromatography (hexanes/EtOAc=6:1 to 5:1) to afford product **25** (49.1 mg, 80%) as a white solid. $R_f = 0.31$ (hexanes/EtOAc=2:1); mp 169–170 °C; IR (thin film): $\nu_{\max}$ 3480, 2966, 2924, 1708, 1500, 1383, 1187 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 7.45~7.41 (m, 2H), 7.38~7.35 (m, 1H), 7.14 (dd, $J = 7.2, 1.2$ Hz, 2H), 6.03 (ddd, $J = 6.4, 1.6, 1.6$ Hz, 1H), 3.68 (dd, $J = 8.0, 3.2$ Hz, 1H), 3.59 (dd, $J = 3.2, 1.6$ Hz, 1H), 3.56 (dd, $J = 6.4, 3.2$ Hz, 1H), 3.36 (dd, $J = 8.0, 3.2$ Hz, 1H), 2.46 (s, 1H), 1.87 (sept., $J = 6.8$ Hz, 1H), 1.84 (d, $J = 1.6$ Hz, 3H), 0.98 (d, $J = 6.8$ Hz, 3H), 0.82 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz): δ 209.4, 177.5, 175.1, 136.0, 131.6, 129.1 (two carbons overlapping), 128.8, 126.3 (two carbons overlapping), 125.0, 75.9, 53.5, 42.9, 42.2, 39.8, 33.2, 20.7, 17.8, 16.2; HRMS–ESI (m/z): $[\text{M}+\text{H}]^+$ calcd for $\text{C}_{20}\text{H}_{22}\text{NO}_4$, 340.1549; found, 340.1553.

Computational Methods

All calculations were carried out with Gaussian 03 Revision E.01,⁴⁰ Spartan 06 or Spartan 08.⁴¹ Structures were optimized and characterized by frequency analysis at the B3LYP/6-31G(d) level of theory with all transition states showing a single imaginary vibrational mode corresponding to the expected reaction.⁴² Unscaled zero point vibrational energy (ZPVE) corrections have been applied to total energies. Energies and geometries for stationary points are summarized in the Supporting Information. Solvation was not included in these calculations because it is expected that our nonpolar solvents would not significantly affect reaction energetics.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was generously supported by the NIGMS CMLD Initiative (P50 GM067041 J.A.P., Jr.), the National Science Foundation (CHE-0910826 R.P.J.), and the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research. This project has been funded in part with Federal funds from the National Cancer Institute, National Institutes of Health under contract HHSN261200800001E. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products or organizations imply endorsement by the US Government. We thank Dr. Emil Lobkovsky (Cornell University) for X-ray crystal structure analysis and Anton Paar Corporation, CEM Corporation, and Waters Corporation for assistance with instrumentation.

References

1. For use of ortho-quinol dimers in retro-[4+2]/[4+2] cascades, see: (a) Singh VK, Deota PT, Bedekar AV. *J Chem Soc Perkin Trans 1*. 1992:903.(b) Singh V, Samanta B. *Tetrahedron Lett*. 1999; 40:1807.
2. Dong S, Hamel E, Bai R, Covell DG, Beutler JA, Porco JA Jr. *Angew Chem, Int Ed*. 2009; 48:1494.
3. For use of MOB-derived dimers, see: (a) Liao CC, Peddinti RK. *Acc Chem Res*. 2002; 35:856. [PubMed: 12379138] (b) Chittimalla SK, Shiao HY, Liao CC. *Org Biomol Chem*. 2006; 4:2267. [PubMed: 16729136] c) Snyder SA, Kontes F. *J Am Chem Soc*. 2009; 131:1745. [PubMed: 19191699]
4. For reviews on ortho-quinol and MOB-derived [4+2] cyclodimers, see: (a) Magdziak D, Meek SJ, Pettus TRR. *Chem Rev*. 2004; 104:1383. [PubMed: 15008626] (b) Quideau S, Pouysegu L, Deffieux D. *Synlett*. 2008:467.
5. Kappe CO. *Chem Soc Rev*. 2008; 37:1127. [PubMed: 18497926]

6. For a recent review on use of graphite in microwave reactions, see: (a) Besson T, Thiery V, Dubac J. *Microwaves in Organic Synthesis* (2). Wiley-VCH Weinheim, Germany 2006; 1:416. For a recent example employing graphite, see: (b) Cho HY, Ajaz A, Himali D, Waske PA, Johnson RP. *J Org Chem*. 2009; 74:4137. [PubMed: 19432412]
7. (a) Kreamsner JM, Kappe CO. *J Org Chem*. 2006; 71:4651. [PubMed: 16749800] (b) Razzaq T, Kreamsner JM, Kappe CO. *J Org Chem*. 2008; 73:6321. [PubMed: 18613726] For silicon carbide, see: (c) Harris GL. *Properties of Silicon Carbide*. Institute of Electrical Engineers London 1995 (d) Choyke WJ, Matsunami H, Pensl G. *Silicon Carbide: Recent Major Advances*. Springer Berlin 2004 (e) Sadow SE, Agarwal A. *Advances in Silicon Carbide Processing and Applications*. Artech House Inc Norwood, MA 2004
8. Damm M, Kappe CO. *Mol Diversity*. 2009; 13:529.
9. Kreamsner JM, Stadler A, Kappe CO. *J Comb Chem*. 2007; 9:285. [PubMed: 17348734] (b) Damm M, Kappe CO. *J Comb Chem*. 2009; 11:460. [PubMed: 19275226]
10. Beeler AB, Su S, Singleton CA, Porco JA Jr. *J Am Chem Soc*. 2007; 129:1413. [PubMed: 17263426]
11. (a) Li C, Johnson RP, Porco JA Jr. *J Am Chem Soc*. 2003; 125:5095–5106. [PubMed: 12708860] (b) Lei X, Johnson RP, Porco JA Jr. *Angew Chem, Int Ed*. 2003; 42:3913.
12. Dong S, Zhu J, Porco JA Jr. *J Am Chem Soc*. 2008; 130:2738. [PubMed: 18266375]
13. See Supporting Information for complete experimental details.
14. For UPLC, see: (a) Mazzeo JR, Neue UD, Kele M, Plumb RS. *Anal Chem*. 2005; 77:460 A. (b) Nováková L, Matysová L, Solich P. *Talanta*. 2006; 68:908. [PubMed: 18970409]
15. For a recent review on the dienone-phenol rearrangement, see: Whiting DA. Trost BM, Fleming I. *Comp Org Synth*. Pergamon Press Oxford 1991; 3:803–821.
16. Carman RM, Van Dongen JMAM. *Aust J Chem*. 1984; 37:2607.
17. Wannere CS, Paul A, Herges R, Houk KN, Schaefer HF III, Schleyer PvR. *J Comput Chem*. 2007; 28:344. [PubMed: 17109435]
18. Snider BB, Chen Z. *Org Prep Proced Int*. 1999; 31:537.
19. Kuo YH, Chen CH, Huang SL. *Chem Pharm Bull*. 1999; 46:181.
20. Competition between Cope rearrangements and retro-Diels-Alder reactions in similar bicyclic structures has been reported: (a) Su KJ, Mieusset JL, Arion VB, Brecker L, Brinker UH. *Org Lett*. 2007; 9:113. [PubMed: 17192098] (b) Jeong JP, Lee OS, Yang K. *Bull Korean Chem Soc*. 2002; 23:829. (c) Singh V, Sharma U, Prasanna V, Porinchu M. *Tetrahedron*. 1995; 51:6015.
21. For [3,3]-sigmatropic rearrangement of bridged 1,5-dienes, see: Bultman MS, Ma J, Gin DY. *Angew Chem, Int Ed*. 2008; 47:6821. and references cited therein.
22. Paquette LA, Hofferberth JE. *Org React*. 2003; 62:477.
23. (a) Nagahama S, Tazaki M. *Bull Chem Soc Jpn*. 1987; 60:4175. (b) Shimizu T, Hiranuma S, Qian ZH, Yoshioka H. *Synlett*. 1991:725.
24. Liao YH, Xu LZ, Yang SL, Dai J, Zhen YS, Zhu M, Sun NJ. *Phytochemistry*. 1997; 45:729.
25. Gagnepain J, Castet F, Quideau S. *Angew Chem, Int Ed*. 2007; 46:1533.
26. Dory YL, Roy AL, Soucy P, Deslongchamps P. *Org Lett*. 2009; 11:1197. [PubMed: 19228005]
27. Gagnepain J, Mereau R, Dejuguac D, Leger JM, Castet F, Deffieux D, Pouysegu L, Quideau S. *Tetrahedron*. 2007; 63:6493.
28. Details of computational methods and results are presented in the Supporting Information.
29. (a) Cieplak AS. *J Am Chem Soc*. 1981; 103:4540. (b) Macaulay JB, Fallis AG. *J Am Chem Soc*. 1990; 112:1136.
30. Hou HF, Peddinti RK, Liao CC. *Org Lett*. 2002; 4:2477. [PubMed: 12123355]
31. In each case, we find that the transition state DFT wavefunction is stable relative to becoming open shell. Thus, these reactions may be described as asynchronous but concerted. Further examples of transition state structures are shown in the Supporting Information.
32. Miller B. *Acc Chem Res*. 1975; 8:245. and references therein.
33. Ruocco KM, Goncharova EI, Young MR, Colburn NH, McMahon JB, Henrich CJ. *J Biomol Screen*. 2007; 12:133. [PubMed: 17175522]

34. Booth NL, Sayers TJ, Brooks AD, Thomas CF, Jacobsen K, Goncharova EI, McMahon JB, Henrich CJ. *Cancer Immunol Immunother.* 2009; 58:1229. [PubMed: 19089423]
35. Woldemichael GM, Vasselli JR, Gardella RS, McKee TC, Linehan WM, McMahon JB. *J Biomol Screen.* 2006; 11:678. [PubMed: 16858007]
36. Young MR, Li J, Rincon M, Flavell RA, Sathyanarayana BK, Hunziker R. *Proc Natl Acad Sci USA.* 1999; 96:9827. [PubMed: 10449779]
37. Rodriguez AM, Prieto P, de la Hoz A, Díaz-Ortiz A. *Org Biomol Chem.* 2011; 9:2371. [PubMed: 21321770]
38. Lebel H, Paquet V. *J Am Chem Soc.* 2004; 126:320. [PubMed: 14709098]
39. Liao Y, Xu L, Yang S, Dai J, Zhen Y, Zhu M, Sun N. *Phytochemistry.* 1997; 45:729.
40. Frisch, MJ.; Trucks, GW.; Schlegel, HB.; Scuseria, GE.; Robb, MA.; Cheeseman, JR.; Montgomery, JA., Jr; Vreven, T.; Kudin, KN.; Burant, JC.; Millam, JM.; Iyengar, SS.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, GA.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, JE.; Hratchian, HP.; Cross, JB.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, RE.; Yazyev, O.; Austin, AJ.; Cammi, R.; Pomelli, C.; Ochterski, JW.; Ayala, PY.; Morokuma, K.; Voth, GA.; Salvador, P.; Dannenberg, JJ.; Zakrzewski, VG.; Dapprich, S.; Daniels, AD.; Strain, MC.; Farkas, O.; Malick, DK.; Rabuck, AD.; Raghavachari, K.; Foresman, JB.; Ortiz, JV.; Cui, Q.; Baboul, AG.; Clifford, S.; Cioslowski, J.; Stefanov, BB.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, RL.; Fox, DJ.; Keith, T.; Al-Laham, MA.; Peng, CY.; Nanayakkara, A.; Challacombe, M.; Gill, PMW.; Johnson, B.; Chen, W.; Wong, MW.; Gonzalez, C.; Pople, JA. *Gaussian 0.3*, Revision E.01. Gaussian Inc; Wallingford, CT: 2004.
41. Spartan 06 and Spartan 08. Wavefunction Inc; Irvine, CA:
42. (a) Kohn W, Becke AD, Parr RG. *J Phys Chem.* 1996; 100:12974. (b) Becke AD. *J Chem Phys.* 1993; 98:5648. (c) Lee C, Yang Y, Parr RG. *Phys Rev B.* 1988; 37:785.

Figure 1.
Reaction Partners in Retro-[4+2]/[4+2] Cascade.

Figure 2.
Cope rearrangement vs. a retro-[4+2]/[4+2] process

Figure 3.
Examples of transition state structures.

Figure 4.
DFT results for cycloaddition and Cope rearrangement.

Figure 5.
(-)-31 shows higher selectivity for AP-1-dependent activity than other transcription factors.

Scheme 1.

Synthesis of (+)-chamaecypanone C utilizing a retro-[4+2]/[4+2] cascade

Scheme 2.
Dienone-phenol rearrangement of *o*-quinol **4**

Scheme 3.
Trapping experiment

Scheme 4.
DFT results for α -ketol rearrangement pathways

Scheme 5.
DFT results for dienone-phenol rearrangement

Table 1

Optimization of the Microwave-mediated retro-[4+2]/[4+2] Cascade

entry	solvent	conditions ^a	time (min)	conversion ^b (%)
1	DMF	A	30	>99 (90) ^{c,d}
2	DMA	A	30	>99 (92) ^d
3	mesitylene	A	30	>99 (98)
4 ^e	mesitylene	A	15	>99 (98)
5 ^f	mesitylene	B	90	>99 (97)

^a Reaction conditions: A, 3.0 equiv **6**, SiC chip, μ W, 188 °C (IR temperature); B, 3.0 equiv *N*-phenylmaleimide (**6**), 150 °C;

^b Conversion based on UPLC analysis.

^c Isolated yield of product **7** in parentheses.

^d Aqueous workup required.

^e Reaction conducted at 180 °C (IR temperature) under microwave irradiation.

^f Reaction reported in Reference 2.

Table 2Retro-[4+2]/[4+2] reactions to afford bicyclo[2.2.2]-octenones^a

entry	dienophile (equiv)	cycloadduct ^b	yield ^c (%)
1	(5 equiv) 9a	10a	85
2	(10 equiv) 9b	10b	99
3	(10 equiv) 9c	10c^d	98
4	(20 equiv) 9d	10d	85
5	(20 equiv) 9e	10e^d	76
6	(10 equiv) 9f	10f	90

^a Reaction conditions: dimer (-)-2, dienophile, SiC chip, mesitylene, μW , 180 °C, 15 min;

^b Single diastereomer isolated unless otherwise noted;

^c Isolated yield after column chromatography;

^d Approximately 9% of an inseparable minor product observed by ¹H-NMR (*ca.* 10:1 isomeric ratio).

Table 3

Retro-[4+2]/[4+2] cycloadditions with dienes^a

entry	dienophile (equiv)	cycloadduct	yield ^b (%)
1	11a (10 equiv)	12aα	81
		12aβ	18
2	11b (10 equiv)	13b	57
3	11c (10 equiv)	13c ^C	94
4	11d (5 equiv)	12d	98
5	11e (20 equiv)	12e	57
		13e	25
		14	9

^a Reaction conditions: dimer (–)-2, diene, SiC chip, mesitylene, μW, 180 °C, 15 min;^b Isolated yield after column chromatography;

^c Approximately 5% of an inseparable minor product was observed by ¹H-NMR analysis (*ca.* 19:1 isomeric ratio).

Table 4Retro-[4+2]/[4+2] cascade using other *o*-quinol precursors^a

entry	dimer	dienophile (equiv)	cycloadduct ^b	yield ^c (%)
1	(⁺)-15	(5 equiv) 11d	19	93
2	(⁺)-15	(20 equiv) 9e	20	78
3	16	(10 equiv) 9b	21	86
4	([±])-17	(3 equiv) 6	22	99
5	([±])-17	(10 equiv) 9b	23	78
6	([±])-17	(20 equiv) 9d	24	85

entry	dimer	dienophile (equiv)	cycloadduct ^b	yield ^c (%)
7	(±)-18	 6 (3 equiv)	 25	80

^a Reaction conditions: dimer, dienophile, SiC chip, mesitylene, μ W, 170 °C, 15 min;

^b Single diastereomer isolated;

^c Isolated yield after silica gel chromatography.

Table 5

B3LYP/6-31G(d) Reaction Energetics

entry	reactant(s)	transition state	product(s)
1	$E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS1 $\longrightarrow$ 23.9	2 x 11.6
2	(6S)-4 + $E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS2 $\longrightarrow$ 13.8	10a -17.3
3	(6S)-4 + $E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS3 $\longrightarrow$ 17.0	10d -18.5
4	(6S)-4 + $E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS4 $\longrightarrow$ 17.8	10b -14.9
5	(6S)-4 + $E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS5a $\longrightarrow$ 17.5	-26.9
		TS5b $\longrightarrow$ 16.9	-13.7
6	(6S)-4 + $E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS6 $\longrightarrow$ 23.5	-13.4
7	(6S)-4 + $E_{rel} = 0.0 \text{ kcal/mol}$	$\xrightarrow{\Delta}$ TS7 $\longrightarrow$ 22.9	-10.3