Published in final edited form as:

Health Psychol. 2011 September ; 30(5): 570–578. doi:10.1037/a0024210.

Implicit measures of early-life family conditions: Relationships to psychosocial characteristics and cardiovascular disease risk in adulthood

Meanne Chan, Edith Chen, Anita S. Hibbert, Jennifer H. K. Wong, and Gregory E. Miller
University of British Columbia

Abstract

Objectives—An implicit measure of early-life family conditions was created to help address potential biases in responses to self-reported questionnaires of early-life family environments. We investigated whether a computerized affect attribution paradigm designed to capture implicit, affective responses (anger, fear, warmth) regarding early-life family environments was a) stable over time, b) associated with self-reports of childhood family environments, c) able to predict adult psychosocial profiles (perceived social support, heightened vigilance), and d) able to predict adult cardiovascular risk (blood pressure) either alone or in conjunction with a measure of early-life socioeconomic status.

Method—Two studies were conducted to examine reliability and validity of the affect attribution paradigm (Study 1, N = 94) and associated adult psychosocial outcomes and cardiovascular risk (Study 2, N = 122).

Results—Responses on the affect attribution paradigm showed significant correlations over a 6-month period, and were moderately associated with self-reports of childhood family environments. Greater attributed negative affect about early-life family conditions predicted lower levels of current perceived social support and heightened vigilance in adulthood. Attributed negative affect also interacted with early-life socioeconomic status to marginally predict resting systolic blood pressure, such that those individuals high in early-life SES but who had implicit negative affect attributed to early-life family conditions had SBP levels that were as high as individuals low in early-life SES.

Conclusions—Implicit measures of early-life family conditions are a useful approach for assessing the psychosocial nature of early-life environments and linking them to adult psychosocial and physiological health profiles.

Keywords

early-life socioeconomic status; family environment; implicit affect; psychosocial profile; cardiovascular risk

The notion that early-life experiences, such as with low socioeconomic status (SES) or family adversity, can influence health years later has been receiving increasing attention in recent years. In particular, the early-life family environment provides a background for development that has long-term implications for mental and physical outcomes in adulthood (Conger & Donnellan, 2007). However, one of the prevailing issues in this field is whether there are biases in how adults respond to questions about their early-life family environments. To overcome some of these potential biases, this paper introduces an implicit measure of early-life family conditions (affect attribution) and discusses its implications for health research.

Growing evidence suggests that low early-life SES is a determinant of susceptibility to chronic diseases later in life. Individuals from low early life SES backgrounds are vulnerable to numerous health problems in adulthood, including respiratory and cardiovascular diseases, arthritis, certain cancers, as well as elevated inflammation levels and clustering of metabolic risk markers (Blane, Bartley, & Davey-Smith, 1997; Cohen et al., 2004; Danese et al., 2009; Galobardes, Lynch, & Davey-Smith, 2004; Galobardes, Smith, & Lynch, 2006; Lawlor, Ronalds, Macintyre, Clark, & Leon, 2006; Poulton et al., 2002). These relationships are generally true irrespective of SES in adulthood (Kittleson, Meoni, Wang, Chu, Ford, & Klag, 2006; Kuh, Hardy, Langenberg, Richards, & Wadsworth, 2002; Poulton et al., 2002), suggesting that early-life SES is not merely serving as a proxy for current socioeconomic conditions. In particular, some studies suggest there may be a “sensitive period,” whereby SES during the first five years of life predicts later life health outcomes (Cohen et al., 2004; Miller & Chen, 2007).

Understanding the psychosocial conditions during the first years of life may help shed light on why low early-life SES affects adult health (Conger & Donnellan, 2007; Matthews & Gallo, 2011; Shonkoff, Boyce, & McEwen, 2009). In particular, low SES has been closely tied to an adverse family environment marked by higher levels of family conflict, harsh restrictive parenting styles, chaotic or neglectful parenting, as well as inadequate emotional nurturing, such as family relationships that are cold, neglectful, and lacking support (Dodge, Pettit, & Bates, 1994; Emery & Laumann-Billings, 1998; Lehman et al., 2005; McLoyd, 1998).

In turn, there is increasing evidence that individuals who reported adverse family conditions during childhood have increased rates of physical health problems in adulthood (Repetti et al., 2002). Exposure to abuse during childhood has been linked with a heightened risk of neurological problems, respiratory problems, ischemic heart disease, cancer, skeletal fracture, and liver disease (Dong et al., 2004; Felitti et al., 1998; Wegman & Stetler, 2009). Individuals who reported a ‘risky family environment’ during upbringing show elevated physiological responses to stress (e.g. autonomic and cortisol), compromised metabolic functioning (Lehman, Taylor, Kiefe, & Seeman, 2005; Luecken, Rodriguez, & Appelhans, 2005; Taylor, Lerner, Sage, Lehman, & Seeman, 2004), and have elevated inflammatory profiles (Danese et al., 2009; Taylor, Lehman, Kiefe, & Seeman, 2006; Miller & Chen, 2010).

In sum, diverse research literatures consistently point to the long-term impact of adverse early-life family conditions on health. Some birth cohort or longitudinal studies were able to obtain assessments during childhood that would not be subject to recall biases later in life (e.g. Danese et al., 2009). However, many studies assess health status in adulthood and are constrained to asking adults to retrospectively describe the nature of their family environments during upbringing (e.g. Dong et al., 2004; Felitti et al., 1998). In general, these approaches are particularly vulnerable to respondent biases, especially when recalling interpersonal information (Metts, Sprecher, & Cupach, 1991). A review of previous studies using retrospective accounts of childhood adversity revealed that measurement error and bias exist in these reports (Hardt & Rutter, 2004). Importantly, reporting on one’s feelings about family conditions can be a sensitive issue, and it may be difficult for some people to be completely open and honest. Thus, retrospective self-report measures may not represent the optimal approach for assessing early-life family environments.

One method in the social psychology literature that is sometimes used in lieu of self-report measures (particularly for sensitive issues such as prejudice) is implicit attitude assessments, which are paradigms designed to capture attitudes that are expressed when individuals are unable to overtly monitor and control the influence of their attitudes on judgments (Fazio &

Olson, 2003; Gawronski & Bodenhausen, 2006). Since monitoring and control is limited, implicit measures are extremely useful for investigating socially sensitive attitudes such as prejudice (Fazio, Jackson, Dunton, & William, 1995; Greenwald, McGhee, & Schwartz, 1998; Wittenbrink, Judd, & Park, 1997). However, to our knowledge implicit measures have yet to be adopted within health studies of physiological risk markers.

We developed a behavioral paradigm based on that by Payne, Cheng, Govorun, and Stewart (2005) to measure implicit affect elicited by early-life family conditions. The original Affect Misattribution Procedure (AMP) is a computer-based implicit attitude assessment tool that combines projective testing with computer-based quantitative measurements of evaluations. Affect-laden pictures are presented visually and then paired with ambiguous stimuli, which participants in turn rate on dimensions like attractiveness or pleasantness. Misattributions of affect onto the ambiguous stimuli indicate unintentional expression of attitudes. The AMP can predict outcomes such as intended voting behavior and attitudes toward political candidates, as well as actual voting behavior during presidential elections (Payne et al., 2005; Payne, Krosnick, Pasek, Lelkes, Akhtar, & Tompson, 2010).

In the present set of studies, we created a novel application of the AMP in which we used a modified version to probe participants' feelings about their early-life family conditions. Participants' own childhood family photographs were used as the affect-laden picture, and the evaluative judgments of ambiguous images in our paradigm were made on three affect dimensions – anger, fear, and warmth. These dimensions were chosen based on the Risky Families model that posits a central role for conflict and aggression in the home (generating anger and fear) and to cold, unsupportive, and neglectful environments (indicating a lack of parental warmth) in risky family environments (Taylor et al., 2004). We examined the stability and validity of our affect attribution paradigm across two studies. In the first study, we tested the stability of the paradigm across a six month period, and tested whether this implicit measure of early-life family conditions correlated with explicit measures of childhood family environments and parenting styles. In the second study, we examined whether this implicit measure of early-life family conditions could predict unfavorable adult psychosocial profiles and cardiovascular risk (resting blood pressure) in a healthy sample of adults.

Study 1

Method

Participants—Ninety-four participants aged 18 to 43 ($M = 22.21$, $SD = 5.069$) participated in a three-wave study. Participants were recruited at the University of British Columbia (UBC), and represented a variety of ethnic groups, socioeconomic classes, as well as both student and non-student populations. The sample included 79.8% female participants, with the majority of participants being of Chinese descent (53.2%), followed by European descent (23.4%), and other ethnicities (23.4%). Eighty-six (91%) participants completed the second visit 1 month later, and 85 (90%) participants completed all 3 time points over the course of 7 months.

Measures

Affect attribution—Developed by Payne, Cheng, Govorun, and Stewart (2005), the Affect Misattribution Procedure (AMP) is a computer-based implicit attitude assessment tool that combines projective testing with computer-based quantitative measurements. The 'attitude object' image is presented for 75 ms, then a blank screen for 125 ms, then the ambiguous pictograph for 100 ms. Following this, the participant is asked to make an affective judgment of the ambiguous pictograph. They are instructed that the 'attitude object' image is

a warning signal for the ambiguous pictograph and that they should not respond to it. Of note, Payne et al. (2005) highlighted that across all studies, the task was unaffected by direct warnings to avoid influence of the affect-laden pictures, suggesting that it is resistant to correction attempts. In Payne's work, the AMP has proven to have excellent psychometric characteristics, with attitude ratings showing high levels of internal consistency ($\alpha = .88$) and reliable correlations with criterion variables like self-reported political attitudes and intended voting behavior.

Our study involved a modified version of the AMP. First, our 'attitude object' of interest consisted of two early childhood family photographs meant to elicit implicit affect toward one's early-life family conditions. Participants were asked to bring their own personal photographs for inclusion in the experiment, with the stipulations that each picture had to contain an image of themselves age 0–5, and include at least one primary caregiver. This age range was selected based on previous research pointing to a "sensitive period" that impacts health later in life (Cohen et al., 2004; Miller & Chen, 2007). Second, participants were asked to evaluate ambiguous images of Jackson Pollock-style splatter paintings, rather than the Chinese pictographs used in the original AMP studies, due to the large number of Chinese descent individuals (for whom the Chinese pictographs would have had meaning, rather than being ambiguous) in the Greater Vancouver community. Finally, we asked participants to rate each splatter painting on three affect dimensions – anger, fear and warmth – rather than the ratings of pleasantness made in the original AMP studies because of the relevance of these emotions to childhood family psychosocial environments.

In an initial study to validate the splatter paintings, eighty unique splatter paintings were generated. These splatter paintings were then evaluated by 10 psychology-trained raters on a 9-point scale ranging from unpleasant to pleasant. The 40 splatter paintings with the most 'neutral' ratings (mean ratings of 4.2 to 5.8) were selected for inclusion in the task. An additional 7 paintings were excluded because of high variance in their ratings. The remaining 33 splatter paintings were then reversed and rotated to create 6 images each, which led to a total of 198 splatter paintings that were included in the paradigm.

Participants completed 72 trials in our paradigm, which took on average five minutes to complete. Each trial involved a family photograph or a grey-colored control image followed by a splatter painting that had been randomly selected from the pool. The participant was then asked to indicate whether the painting did vs. did not convey a target affect (anger, fear, warmth). The trials varied in terms of the prime that was used (3 possibilities: 1 of 2 family photographs or a grey control image) and the affect rating that was made (3 possibilities: anger, fear, or warmth). There were a total of 8 trials for each combination of prime and rating. Ratings were averaged within each category to form separate attribution scores for the active and control primes. Values could range from 0 – 1.

Family environment—The 13-item Risky Families measure (RF) assesses early-life family environment (Taylor et al., 2004). Participants are asked to think about their family life between the ages of 5 and 15, and answer questions on a scale ranging from 1 ('Not at All') to 5 ('Very Often' or 'Very Much'). Question items include: "How often did a parent or other adult in the household swear at you, insult you, put you down, or act in a way that made you feel threatened?", and "How often would you say you were neglected while you were growing up, that is, left on your own to fend for yourself?". The RF measure has been shown to have high agreement and reliability with clinical interviews conducted and coded by trained clinical interviews (Taylor et al., 2004).

Parenting—Paternal and maternal warmth in the first 16 years of the participants' lives were measured using the 25-item Parental Bonding Instrument (Parker, Tupling, & Brown,

1979). Participants filled out the questionnaire for maternal and paternal bonding separately and had to rate how true the statements were on a 5-point Likert-type scale. Scores for the warmth subscale (12 items; e.g. “Spoke to me in a warm and friendly voice”) were computed for each parent and higher scores indicated higher levels of parental warmth. This questionnaire has been shown to have good test-retest reliability, and has been validated with actual parental behavior recorded by family observers and expert judges (Wilhelm & Parker, 1990).

Procedure—After arriving at the laboratory and providing informed written consent, participants were taken to a private room and seated at a computer. Participants were asked to bring in two childhood family photographs, and these were inserted into the affect attribution computer program. Participants completed the affect attribution task and questionnaires described above via MediaLab software, version 2008.1.22 (Jarvis, 2008). Participants were invited back to the lab one month (Visit 2) and six months (Visit 3) after their first visit. The same family photographs were used and the same study procedure was repeated. To thank them for their time, each participant received \$10 for each of three visits, with a \$10 bonus for completing all three visits. The study was approved by the Behavioral Research Ethics Board of the University of British Columbia.

Results

Stability across Visits

Overall, correlations of affect attribution tendencies over all three visits during the six month study period were highly significant. Pearson correlations revealed that attribution of anger in the first and second visits (one month apart), and the first and last visits (six months apart) were significant respectively, $r = .458, p < .001$ and $r = .522, p < .001$. Similarly, attribution of fear in the first and second visits, and the first and last visits were significantly correlated, $r = .670, p < .001$ and $r = .463, p < .001$. Lastly, attribution of warmth in the first and second visits, and the first and last visits were significantly correlated, $r = .524, p < .001$ and $r = .467, p < .001$. These correlations were somewhat weaker than the correlation over the same time period of a questionnaire, self-report measure of family environment, $r's > .891, p's < .001$. However, the stability estimates of our implicit paradigm are comparable to those of other implicit tasks (Cunningham, Preacher, & Banaji, 2001). Affect attribution responses were subsequently aggregated across the three visits to create a more reliable indicator.

Affect Attribution Analyses

Attribution of affect was computed by summing up the number of times participants attributed the different emotions towards the splatter paintings following a family photograph, and as well, following the grey colored control image. A percentage score was calculated (number of times participants endorsed an emotion divided by the number of trials probing that emotion) per each emotion for family photographs and for the control image. Attribution of affect following the two different family photographs were averaged to obtain one family affect attribution score per each affect dimension. In all analyses examining associations between affect attribution and other measures, affect attribution after the control image were subtracted out from affect attribution after a family photograph. This controlled for individual differences in the tendency to endorse the various affect descriptors.

After their own early childhood family photograph, participants attributed significantly less anger and fear in comparison to the control image, $M_{diff} > 3.089, t's > 2.021, p's < .04$. Also, participants attributed significantly more warmth after a family photograph, $M_{diff} = -14.259, t = -7.821, p < .001$.

Relationships among the three affect dimensions—Attributions of anger correlated with attributions of fear, $r = .659$, $p < .001$, but neither correlated with attribution of warmth, $r's < .002$, $p's > .380$.

Affect Attribution and the Family

Family environment—Pearson correlation analyses were conducted to examine the association between affect attribution and explicit measures of childhood family environment (Risky Family (RF) scores). Participants who reported more negative childhood family environments on the Risky Family measure also significantly attributed more anger after a family photograph, $r = .374$, $p < .001$. Similarly, these participants also significantly attributed more fear after a family photograph, $r = .266$, $p = .012$. However, higher Risky Family scores were not significantly correlated with attribution of warmth, $r = -.157$, $p = .143$.

Parenting—Paternal warmth during childhood was significantly correlated with less attribution of anger, indicating that participants who had warmer fathers during childhood also attributed significantly less anger after a family photograph, $r = -.319$, $p = .002$. Paternal warmth was also significantly correlated with less attribution of fear, $r = -.276$, $p = .009$. Furthermore, paternal warmth was significantly associated with greater attribution of warmth, $r = .223$, $p = .036$. Maternal warmth was correlated with less attribution of anger, $r = -.249$, $p = .019$, but not attribution of fear or attribution of warmth, $r's < -.165$, $p's > .123$.

In sum, these findings indicate that a new measure of implicit attributed affect about childhood family conditions was associated with explicit self-report measures of childhood family environments and parenting styles during upbringing, and showed consistency over a 6 month period. A second study was then conducted with a healthy sample of adults to examine the validity of this measure by testing whether this implicit early life measure could predict adult psychosocial and physiological profiles. In particular, individuals who grow up in adverse early-life environments have been found to display more negative affect, to perceive greater threat, and to have less social support as adults (Luecken, Appelhans, Kraft, & Brown, 2006; Repetti et al., 2002). In addition, adverse early life environments have been associated with greater cardiovascular risk profiles in adulthood (Repetti et al., 2002; Wegman & Stetler, 2009). As such, we tested associations of our implicit childhood family environment measure with adult depressive symptoms, vigilance for threat, social support, and resting blood pressure. In addition to the main effects of implicit affect on these adult psychosocial and physiological outcomes, we also tested the hypothesis that the quality of early childhood family environments might interact with broader social environment characteristics, such as socioeconomic status (SES) to impact outcomes. Previous research has documented that childhood family relationship quality can moderate or buffer the effects of socioeconomic status (SES) on cardiovascular and inflammatory profiles (Chen, Miller, Kobor, & Cole, in press; Evans, Kim, Ting, Tesher, & Shannis, 2007). Hence we tested whether implicit attributions about early childhood environments might also interact with early-life SES in predicting adult characteristics.

Study 2

Method

Participants—Participants were recruited from Vancouver, BC, Canada, through postings in local media and public transit as part of a larger study examining the biological embedding of early-life socioeconomic status. To be eligible, they had to be 1) between 15–55 years of age, 2) North American born, 3) fluent in English, 4) in good health, defined as

being free of acute infectious disease the past 2 weeks, and without a history of chronic disease, and 5) from high or low socioeconomic categories. Socioeconomic status (SES) was obtained using the National Statistics Socio-economic Classification guide (NS-SEC, 2005) to code current occupation (9-point scale), as well as early-life (age 0 to 5) occupation of parents. Those in midrange SES categories (3 to 4 on 9-point scale) were excluded from the larger study to better contrast low and high SES. One hundred and twenty two individuals ($M_{\text{age}} = 32.46$, $SD = 10.92$) participated in this study, with 52.5% of individuals in the low early-life SES category. The sample consisted of 54.1% female participants, with the majority of participants being of European descent (68.0%), followed by Chinese descent (13.1%), and other ethnicities (18.9%).

Measures

Attribution of affect—The same modified affect attribution paradigm from Study 1 was used. We replaced the grey colored control image in Study 1 with an image of a neutral face obtained from a face database created in the Karolinska Institute in Sweden (Lundqvist, Flykt, & Ohman, 1998; Oosterhof & Todorov, 2008), so that both the image of interest (early childhood family photograph) and the control image would contain faces. This face database contains previously established images of different emotional expressions and we used an image drawn from their neutral expression database.

Social support—An abbreviated version of the Interpersonal Support Inventory List (ISEL) developed by Cohen, Mermelstein, Kmack and Hoberman (1985) was used. The 12-item version of ISEL provides a global measure of perceived social support, with higher scores indicating greater support. The ISEL has demonstrated validity with moderate correlations to other social support measures, such as the Moos Family Environment Scale (FES) (Moos & Moos, 1981). Adequate internal and test-retest reliabilities have been found for the total ISEL scale and the subscales in several samples (Cohen et al., 1985).

Depression—Depressive symptoms were assessed using the Center for Epidemiological Studies Depression (CES-D) Scale (Radloff, 1977), which has been widely used in clinical trials in both general and psychiatric populations and has excellent psychometric characteristics with a Cronbach's alpha of .85. The 10-item version included items assessing the frequency of experiencing depressive cognitions, affect, and behaviors during the past one week (e.g., feeling depressed and lonely, having poor appetite and sleep). Responses range from 0 = none of the time or rarely (less than 1 day) to 3 = most or all of the time (5–7 days). Scores are summed, with higher scores reflecting greater depressive symptoms.

Vigilance of threat—To assess vigilance for threat, we administered a slightly modified version of Payne's weapon identification procedure (Payne et al., 2001). The original paradigm measures the tendency to misidentify neutral objects (e.g., tools) as threatening (e.g., guns) after presentation of African-American vs. European-American faces. Participants are asked to decide whether the second image is a gun or a tool. Incorrect responses, or in other words false-positives, occur when participants misidentify a tool as a gun (Payne et al., 2001).

In our modified version of this paradigm, we probed the tendency to perceive threat after either neutral (i.e. a parrot) or threatening (i.e. snake) images which were selected from the International Affective Picture System, and which were presented prior to images of a gun or a tool (Lang, Bradley, & Cuthbert, 2008). False-positives occur when participants misidentify a tool as a gun. In this study, we focused on the percentage of false-positives that occur after the neutral image, which provides a general measure of automatic heightened vigilance tendencies.

Blood pressure—Blood pressure was recorded with a standard occluding cuff on the participant's non-dominant arm. The VSM-100 BpTRU automatic blood pressure monitor is a reliable non-invasive device. Validation studies indicate that its measurements are within 5 mm Hg of a gold standard auscultatory mercury sphygmomanometer measurements 89.2% of the time, and within 10 mm Hg 96.4% of the time (Mattu, Heran, & Wright, 2004). Following a 5-minute period where participants acclimated to the device, blood pressure readings were taken every two minutes over a 6-minute period, totaling three readings that were averaged.

Procedure—Participants were seated in a private room to rest for five minutes. Blood pressure readings were then taken by placing the occluding cuff on the upper aspect of the participant's non-dominant arm with the microphone placed above an area where the brachial artery could be palpated. Participants were then seated at a computer and completed the same Study 1 affect attribution task and questionnaires described above via MediaLab software, version 2008.1.22 (Jarvis, 2008). This study was approved by the Clinical Research Ethics Board of the University of British Columbia.

Results

Affect Attribution Analyses

Attributions of anger and fear after a family photograph were significantly correlated with each other, $r = .565, p < .001$. However, attribution of warmth after a family photograph was not associated with attribution of anger and fear, r 's $< .064, p$'s $> .487$, similar to Study 1. Hence anger and fear attributions were aggregated to create a more reliable indicator of negative affect attribution. Warmth was retained as a separate indicator of positive affect attribution. See Table 1 for descriptive information.

Psychosocial Analyses

Multiple regression analyses were conducted to test whether attribution of negative affect towards early-life family conditions could predict adult psychosocial profiles. In all analyses, affect attribution after the control image was subtracted out from affect attribution after a family photograph. Covariates of age, gender, ethnicity, early-life and current SES were included.

Social support—Greater attribution of negative affect significantly predicted lower levels of current perceived social support, such that individuals who attributed more anger and fear after their own childhood family photographs reported lower levels of current perceived social support on the ISEL, $\beta = -.191, p = .028$.

Depression—Attribution of negative affect did not significantly predict levels of depressive symptoms as measured by the CES-D Scale, $\beta = .097, p = .287$.

Vigilance of threat—Greater attribution of negative affect significantly predicted higher vigilance scores, such that individuals who attributed more anger and fear after their own childhood family photograph exhibited greater tendencies to automatically overinterpret cues as threatening as adults, $\beta = .217, p < .018$.

In addition to main effects, we also tested for interactions between affect attribution and early-life SES, using the procedures recommended by Aiken and West (1991). No significant interactions were found for adult psychosocial outcomes, β 's $< |.118|, p$'s $> .349$. To further test the specificity of the main effects, we conducted subsequent analyses in which we included explicit childhood family environment measures (the Risky Family

questionnaire) as a simultaneous predictor of adult psychosocial profiles. Implicit childhood negative affect attributions continued to predict lower levels of adult perceived social support and a greater tendency toward heightened vigilance, above and beyond explicit childhood family environments, β 's $> .148$, p 's $< .074$.

Cardiovascular Risk

Multiple regression analyses were conducted to test whether resting blood pressure could be predicted from implicit negative affect attributions. Covariates of age, gender, ethnicity, and current SES were included in each analysis. Systolic and diastolic blood pressure were not predicted by affect attribution, β 's $< .085$, p 's $> .354$. However, the interaction between negative affect attributions and early-life SES was marginally significant in predicting systolic blood pressure (SBP), $\beta = .210$, $p = .089$. This interaction indicates that SBP levels are lowest for those individuals who are both from a high early-life SES background and who attributed low levels of negative affect after a family photograph. On the other hand, among those from a high early-life SES background, as attributed negative affect increases, SBP also increases, such that those individuals who are from a high early-life SES background but who attributed high levels of negative affect after a family photograph actually have SBP levels similar to individuals from low early-life SES backgrounds (see Figure 1). Subsequent analyses revealed that when entered simultaneously with explicit childhood family measures, the interaction between implicit negative affect toward early life family conditions and SES in predicting SBP remained marginally significant, $\beta = .218$, $p = .079$. In contrast, the interaction between negative affect attribution and early-life SES did not predict diastolic blood pressure (DBP), $\beta = .119$, $p = .332$.

Discussion

Across two study samples, we demonstrated reliability and validity for a novel measure of early-life family environments, an implicit affect attribution paradigm which tapped implicit affect about early-life family conditions in adults. Implicit affective judgments of anger, fear, and warmth were significantly correlated over a six month period, and this test-retest estimate was comparable to the consistency of personality traits over a 1 year period (Roberts & DelVecchio, 2000). Our paradigm showed construct validity in being associated with explicit self-report measures of childhood family environments as well as parenting styles during childhood. Our measure showed predictive validity in implicit negative affect toward one's childhood family being associated with unfavorable adult psychosocial profiles, including lower levels of current perceived social support and a greater tendency toward heightened vigilance (overinterpreting threat). Lastly, this measure of implicit affect marginally predicted an indicator of cardiovascular risk (e.g. elevated resting SBP) in conjunction with a measure of early-life family SES.

The finding that greater attributed negative affect after one's own early childhood family photograph was associated with lower levels of current perceived social support and a greater tendency toward heightened vigilance is consistent with a large body of literature documenting damaging psychosocial consequences for individuals from an adverse childhood family environment. For instance, previous research has demonstrated that children reared in a family environment characterized by overt conflict, aggression, and inadequate emotional nurturance, are at an increased risk of a wide array of current and future emotional or behavioral problems, such as anxiety, depression, conduct disorder, antisocial behavior, and suicide (Emery & Laumann-Billings, 1998; Repetti et al., 2002). Moreover, our findings document that implicit measures of affect regarding one's childhood family environment predict adult psychosocial profiles after controlling for explicit measures of childhood family environment, suggesting that implicit measures of childhood environments add unique predictive power beyond what explicit measures can explain.

We did not observe a direct association between our measure of implicit affect towards early-life family conditions and blood pressure. However, there was evidence that implicit affect interacted with early-life SES in systolic blood pressure (SBP) levels. Levels of SBP were lowest for those individuals who were both from a high early-life SES background and who implicitly attributed low levels of negative affect after a family photograph. Among those high in early-life SES, as attributed negative affect increased, SBP also increased. Thus, those individuals from a high early-life SES background who attributed negative affect to childhood family photographs actually had SBP levels similar to individuals low in early-life SES. This suggests that implicit affect toward early-life family conditions can at times override the effects of early-life SES on resting blood pressure, and suggests the importance of considering both implicit and explicit assessments when exploring the influences of early-life factors on later life health. All that said, because the findings were only marginally significant, they will need to be replicated in other samples before any definitive conclusions about the value of implicit measures in predicting health-relevant outcomes can be reached.

This finding is in line with a large body of evidence documenting damaging physical health outcomes for individuals who explicitly report adverse early-life family circumstances. For instance, individuals who reported a childhood risky family environment consistently showed increased rates of physical health issues in adulthood, such as elevated stress responses, major physical disorders, and a compromised metabolic functioning (Felitti et al., 1998; Lehman et al., 2005; Luecken et al., 2005; Taylor et al., 2004). We add to this literature by showing that indirectly measured implicit affect toward early-life family conditions have implications for cardiovascular functioning in adulthood. We unmasked a subgroup of individuals who may be at risk – those who were from a high early-life SES background but had implicit negative affect toward early-life family conditions. Thus, this suggests that our implicit measure may have incremental utility in health research, as this subgroup would not have been identified on the basis of explicit measures alone.

Limitations

There are several limitations in our two studies. First, we explored a limited selection of criterion variables. Future studies should conduct a broader assessment to determine the scope of our paradigm's associations with psychosocial and psychobiological outcomes relevant to health. Second, our validation study findings were cross-sectional and longitudinal studies are needed to determine whether implicit negative affect truly precede change in health-relevant processes. Third, our two samples were not identical in composition (and consisted primarily of Chinese- and European-descent adults with no chronic health conditions). As has been done in almost every validation study of implicit attitude measures (Nosek & Smyth, 2007; Payne, Cheng, Govorun, & Stewart, 2005), participants in Study 1 were a sample recruited from the university setting (both students and non-students). In contrast, Study 2 drew on data from a larger, ongoing community study, and hence the demographics of the two studies are different. Future prospective studies should examine our paradigm's temporal stability and associations with health processes in broader, population-based samples. Fourth, because participants self-selected their family photographs, participants from adverse childhood environments may have fewer photographs from which to choose. We are also unable to verify that participants' responses are specific to early childhood. Lastly, although automatic affective responses are less explicitly biased than self-report measures that require participants to recall detailed information from the past, emotional recall may also reflect some bias. Other measures that capture the early-life environment are still warranted.

Conclusion

Across two study samples, our affect attribution paradigm designed to capture implicit affect toward early-life family conditions showed moderate associations with explicit measures of childhood family environment, predicted unfavorable adult psychosocial processes, and interacted with a measure of SES to predict cardiovascular risk. Importantly, our results indicate that implicit characteristics can potentially override the effects of early-life characteristics on adult BP. These findings highlight the potential importance of incorporating implicit measures together with explicit measures when assessing early-life factors, especially given the relative ease with which these measures could be incorporated into future research studies (approximately 5 minutes to administer), the advantage that this approach brings of eliminating self-presentation biases, and the fact that these implicit measures have predictive power over and above explicit measures of early life family environments. Implicit measures may be useful when exploring socially sensitive situations, such as the early-life family environment, because it can circumvent social desirability pressures. This new affect attribution paradigm as an indirect measure of early-life factors may help address some of the prevailing issues surrounding retrospective assessments of early-life circumstances, thus providing a practical method for further exploring the implications of early-life social environments on later life health.

Acknowledgments

This research was supported by a grant from the National Institute of Child Health and Human Development (058502) to G. E. Miller. The authors thank Keith Payne for his assistance in designing the task, and Tara Martin, Kharah Ross, Sherylann Chan, Alvin Lim, and Sarah Liu for their assistance with data collection.

References

- Aiken, L.; West, S. Multiple Regression: Testing and Interpreting Interaction Effects. Thousand Oaks, CA: Sage Publications; 1991.
- Blane D, Bartley M, Smith GD. Disease aetiology and materialist explanations of socioeconomic mortality differentials. *The European Journal of Public Health*. 1997; 7(4):385–391.10.1093/eurpub/7.4.385
- Chen E, Miller GE, Kobor MS, Cole S. Maternal warmth buffers the effects of low early-life socioeconomic status on pro-inflammatory signalling in adulthood. *Molecular Psychiatry*. (in press).
- Cohen S, Doyle WJ, Turner RB, Alper CM, Skoner DP. Childhood socioeconomic status and host resistance to infectious illness in adulthood. *Psychosomatic Medicine*. 2004; 66(4):553–558.10.1097/01.psy.0000126200.05189.d366/4/553 [PubMed: 15272102]
- Cohen, S.; Mermelstein, R.; Kamarck, T.; Hoberman, HM. Measuring the functional components of social support. In: Sarason, IG.; Sarason, BR., editors. *Social support: Theory, research and applications*. Dordrecht: Martinus Nijhoff; 1985. p. 73-94.
- Conger RD, Donnellan MB. An interactionist perspective on the socioeconomic context of human development. *Annual Review of Psychology*. 2007; 58:175–199.10.1146/annurev.psych.58.110405.085551
- Cunningham WA, Preacher KJ, Banaji MR. Implicit attitude measures: Consistency, stability, and convergent validity. *Psychological Science*. 2001; 12(2):163–170.10.1111/1467-9280.00328 [PubMed: 11340927]
- Danese, A.; Moffitt, TE.; Harrington, H.; Milne, BJ.; Polanczyk, G.; Pariante, CM.; Caspi, A. Adverse childhood experiences and adult risk factors for age-related disease; *Archives of Pediatrics and Adolescent Medicine*. 2009. p. 1135-114. Retrieved from <http://archpedi.ama-assn.org/cgi/content/abstract/163/12/1135>
- Dodge KA, Pettit GS, Bates JE. Socialization mediators of the relation between socioeconomic status and child conduct problems. *Child Development*. 1994; 65:649–665.10.1111/j.1467-8624.1994.tb00774.x [PubMed: 8013245]

- Dong M, Giles WH, Felitti VJ, Dube SR, Williams JE, Chapman DP, Anda RF. Insights into causal pathways for ischemic heart disease: Adverse childhood experiences study. *Circulation*. 2004; 110:1761–1766.10.1161/01.CIR.0000143074.54995.7F [PubMed: 15381652]
- Dovidio, JF.; Fazio, RH. New technologies for the direct and indirect assessment of attitudes. In: Tanur, J., editor. *Questions about survey questions: Meaning, memory, attitudes, and social interaction*. New York: Russell Sage Foundation; 1992. p. 204–237.
- Emery RE, Laumann-Billings L. An overview of the nature, causes, and consequences of abusive family relationships: Toward differentiating maltreatment and violence. *American Psychologist*. 1998; 53(2):121–135.10.1037/0003-066x.53.2.121 [PubMed: 9491743]
- Evans GW, Kim P. Childhood poverty and health: Cumulative risk exposure and stress dysregulation. *Psychological Science*. 2007; 18(11):953–957.10.1111/j.1467-9280.2007.02008.x [PubMed: 17958708]
- Evans GW, Kim P, Ting AH, Tesher HB, Shannis D. Cumulative risk, maternal responsiveness, and allostatic load among young adolescents. *Developmental Psychology*. 2007; 43:341–351. [PubMed: 17352543]
- Fazio RH, Jackson JR, Dunton BC, Williams CJ. Variability in automatic activation as an unobtrusive measure of racial attitudes: A bona fide pipeline? *Journal of Personality and Social Psychology*. 1995; 69(6):1013–1027.10.1037/0022-3514.69.6.1013 [PubMed: 8531054]
- Fazio RH, Olson MA. Implicit measures in social cognition research: Their meaning and use. *Annual Review of Psychology*. 2003; 54(1):297–327.10.1146/annurev.psych.54.101601.145225
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, Marks JS. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) study. *American Journal of Preventive Medicine*. 1998; 14(4):245–258.10.1016/S0749-3797(98)00017-8 [PubMed: 9635069]
- Galobardes B, Lynch JW, Davey Smith G. Childhood socioeconomic circumstances and cause-specific mortality in adulthood: Systematic review and interpretation. *Epidemiologic Reviews*. 2004; 26(1):7–21.10.1093/epirev/mxh008 [PubMed: 15234944]
- Galobardes B, Smith GD, Lynch JW. Systematic review of the influence of childhood socioeconomic circumstances on risk for cardiovascular disease in adulthood. *Annals of Epidemiology*. 2006; 16(2):91–104.10.1016/j.annepidem.2005.06.053 [PubMed: 16257232]
- Gawronski B, Bodenhausen GV. Associative and propositional processes in evaluation: An integrative review of implicit and explicit attitude change. *Psychological Bulletin*. 2006; 132(5):692–731.10.1037/0033-2909.132.5.692 [PubMed: 16910748]
- Greenwald AG, McGhee DE, Schwartz JLK. Measuring individual differences in implicit cognition: The implicit association test. *Journal of Personality and Social Psychology*. 1998; 74(6):1464–1480.10.1037/0022-3514.74.6.1464 [PubMed: 9654756]
- Hardt J, Rutter M. Validity of adult retrospective reports of adverse childhood experiences: review of the evidence. *Journal of Child Psychology and Psychiatry*. 2004; 45(2):260–273.10.1111/j.1469-7610.2004.00218.x [PubMed: 14982240]
- Jarvis, BG. MediaLab (Version 2008.1.33) [Computer Software]. Empirisoft Corporation; New York, NY: 2008.
- Kittleson MM, Meoni LA, Wang NY, Chu AY, Ford DE, Klag MJ. Association of childhood socioeconomic status with subsequent coronary heart disease in physicians. *Archives of Internal Medicine*. 2006; 166(21):2356–2361.10.1001/archinte.166.21.2356 [PubMed: 17130389]
- Kuh D, Hardy R, Langenberg C, Richards M, Wadsworth MEJ. Mortality in adults aged 26–54 years related to socioeconomic conditions in childhood and adulthood: post war birth cohort study. *British Medical Journal (Clinical Research Ed)*. 2002; 325(7372):1076–1080.10.1136/bmj.325.7372.1076
- Lang, PJ.; Bradley, MM.; Cuthbert, BN. Technical Report A-8. University of Florida; Gainesville, FL: 2008. International affective picture system (IAPS): Affective ratings of pictures and instruction manual.
- Lawlor DA, Ronalds G, Leon DA, Macintyre S, Clark H. Family socioeconomic position at birth and future cardiovascular disease risk: Findings from the Aberdeen children of the 1950s cohort study.

- American Journal of Public Health. 2006; 96(7):1271–1277.10.2105/ajph.2005.066290 [PubMed: 16735637]
- Lehman BJ, Taylor SE, Kiefe CI, Seeman TE. Relation of childhood socioeconomic status and family environment to adult metabolic functioning in the CARDIA study. *Psychosomatic Medicine*. 2005; 67(6):846–854.10.1097/01.psy.0000188443.48405.eb [PubMed: 16314588]
- Leucken LJ, Appelhans BM, Kraft A, Brown A. Never far from home: A cognitive-affective model of the impact of early-life family relationships on physiological stress responses in adulthood. *Journal of Social and Personal Relationships*. 2006; 23(2):189–203.10.1177/0265407506062466
- Luecken LJ, Rodriguez AP, Appelhans BM. Cardiovascular stress responses in young adulthood associated with family-of-origin relationship experiences. *Psychosomatic Medicine*. 2005; 67(4): 514–521.10.1097/01.psy.0000160466.10397.18 [PubMed: 16046362]
- Lundqvist, D.; Flykt, A.; Ohman, A. Karolinska Directed Emotional Faces [Database of standardized facial images]. Psychology Section, Department of Clinical Neuroscience, Karolinska Hospital; Stockholm, Sweden: 1998. p. S-171-76.
- Matthews KA, Gallo LC. Psychological perspectives on pathways linking socioeconomic status and physical health. *Annual Review of Psychology*. 2011; 62:501–530.10.1146/annurev.psych.031809.130711
- Mattu GS, Heran BS, Wright JM. Comparison of the automated non-invasive oscillometric blood pressure monitor (BpTRU(TM)) with the auscultatory mercury sphygmomanometer in a paediatric population. *Blood Pressure Monitoring February*. 2004; 9(1):39–45.10.1097/01.mbp.0000122483.45242.f1
- McLoyd VC. Socioeconomic disadvantage and child development. *American Psychologist*. 1998; 53(2):185–204.10.1037/0003-066x.53.2.185 [PubMed: 9491747]
- Metts, S.; Sprecher, S.; Cupach, WR. Retrospective self-reports. In: Montgomery, BM.; Duck, S., editors. *Studying interpersonal interaction*. New York: The Guilford Press; 1991. p. 162-178.
- Miller GE, Chen E. Unfavorable socioeconomic conditions in early life presage expression of pro-inflammatory phenotype in adolescence. *Psychosomatic Medicine*. 2007; 69:402–409.10.1097/PSY.0b013e318068fcf9 [PubMed: 17556642]
- Miller GE, Chen E. Harsh family climate in early life presages the emergence of a proinflammatory phenotype in adolescence. *Psychological Science*. 2010; 21:848–856.10.1177/0956797610370161 [PubMed: 20431047]
- Moos, RH.; Moos, BS. *Manual for the Family Environment Scale*. Palo Alto: Consulting Psychology Press; 1981.
- Nosek BA, Smyth FL. A multitrait-multimethod validation of the Implicit Association Test: Implicit and explicit attitudes are related but distinct constructs. *Experimental Psychology*. 2007; 54(1):14–29.10.1027/1618-3169.54.1.14 [PubMed: 17341011]
- Office for National Statistics. *The National Statistics Socio-economic Classification User Manual*. 2005. Retrieved from http://www.statistics.gov.uk/methods_quality/ns_sec/
- Oosterhof NN, Todorov A. The functional basis of face evaluation. *Proceedings of the National Academy of Sciences of the USA*. 2008; 105:11087–11092. [PubMed: 18685089]
- Parker G, Tupling H, Brown LB. A parenting bonding instrument. *British Journal of Medical Psychology*. 1979; 52:1–10.
- Payne BK. Prejudice and perception: The role of automatic and controlled processes in misperceiving a weapon. *Journal of Personality and Social Psychology*. 2001; 81(2):181–192.10.1037/0022-3514.81.2.181 [PubMed: 11519925]
- Payne BK, Cheng CM, Govorun O, Stewart BD. An inkblot for attitudes: Affect misattribution as implicit measurement. *Journal of Personality and Social Psychology*. 2005; 89(3):277–293.10.1037/0022-3514.89.3.277 [PubMed: 16248714]
- Payne BK, Krosnick JA, Pasek J, Lelkes Y, Akhtar O, Tompson T. Implicit and explicit prejudice in the 2008 American presidential election. *Journal of Experimental Social Psychology*. 2010; 46(2): 367–374.10.1016/j.jesp.2009.11.001
- Poulton R, Caspi A, Milne BJ, Thomson WM, Taylor A, Sears MR, Moffitt TE. Association between children's experience of socioeconomic disadvantage and adult health: A life-course study. *The Lancet*. 2002; 360(9346):1640–1645.10.1016/S0140-6736(02)11602-3

- Radloff LS. The CES-D Scale. *Applied Psychological Measurement*. 1977; 1(3):385–401.10.1177/014662167700100306
- Repetti RL, Taylor SE, Seeman TE. Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*. 2002; 128(2):330–366.10.1037/0033-2909.128.2.330 [PubMed: 11931522]
- Roberts BW, DelVecchio WF. The rank-order consistency of personality traits from childhood to old age: A quantitative review of longitudinal studies. *Psychological Bulletin*. 2000; 126(1):3–25.10.1037/0033-2909.126.1.3 [PubMed: 10668348]
- Shonkoff JP, Boyce TW, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities. *Journal of American Medical Association*. 2009; 301(21):2252–2259.10.1001/jama.2009.754
- Taylor SE, Lehman BJ, Kiefe CI, Seeman TE. Relationship of early life stress and psychological functioning to adult C-reactive protein in the Coronary Artery Risk Development In young Adults study. *Biological Psychiatry*. 2006; 60(8):819–824.10.1016/j.biopsych.2006.03.016 [PubMed: 16712805]
- Taylor SE, Lerner JS, Sage RM, Lehman BJ, Seeman TE. Early environment, emotions, responses to stress, and health. *Journal of Personality*. 2004; 72(6):1365–1394.10.1111/j.1467-6494.2004.00300.x [PubMed: 15509286]
- Wegman HL, Stetler C. A meta-analytic review of the effects of childhood abuse on medical outcomes in adulthood. *Psychosomatic Medicine*. 2009; 71:805–812.10.1097/PSY.0b013e3181bb2b46 [PubMed: 19779142]
- Wilhelm K, Parker G. Reliability of the Parental Bonding Instrument and Intimate Bond Measure scales. *Australian and New Zealand Journal of Psychiatry*. 1990; 24(2):199–202.10.3109/00048679009077683 [PubMed: 2396963]
- Wittenbrink B, Judd CM, Park B. Evidence for racial prejudice at the implicit level and its relationship with questionnaire measures. *Journal of Personality and Social Psychology*. 1997; 72(2):262–274.10.1037/0022-3514.72.2.262 [PubMed: 9107001]

Figure 1.

The interaction between negative affect attribution tendencies and early-life (EL) SES predicting resting SBP. SBP levels are lowest for those individuals who are both from a high early-life SES background and have low levels of attributed negative affect after a family photograph. On the other hand, those individuals who are from a high early-life SES background but have high levels of attributed negative affect after a family photograph have blood pressure levels similar to individuals from low early-life SES backgrounds.

Table 1

Study 1 and 2 descriptive information including affect attribution tendencies (percent), resting blood pressure (mmHg) and heart rate (bpm), self-reported depressive symptoms and current perceived support, and vigilance tendencies (percent).

	Min.	Max.	Mean	Std. Deviation
Study 1				
Anger attributed to family photo (%)	0	87.50	24.81	22.84
Fear attributed to family photo (%)	0	87.50	24.48	22.58
Warmth attribute to family photo (%)	0	100.00	45.58	27.64
Anger attributed to control image (%)	0	100.00	27.90	25.01
Fear attributed to control image (%)	0	100.00	29.42	26.60
Warmth attributed to control image (%)	0	100.00	31.32	25.40
Study 2				
Negative affect attributed to family photo (%)	0	78.13	17.62	20.12
Warmth attributed to family photo (%)	0	100	49.49	32.42
Negative affect attributed to control image (%)	0	100	26.79	27.71
Warmth attributed to control image (%)	0	100	30.94	29.69
Systolic blood pressure (mmHg)	84	150	107.49	12.35
Diastolic blood pressure (mmHg)	49	120	70	11.18
Heart rate (bpm)	42	93	68.64	9.90
Depressive symptoms	0	26	7.85	5.00
Current perceived social support	20	48	39.95	6.34
Vigilance of threat (%)	0	100	18.56	19.03