Published in final edited form as:

Dev Psychopathol. 2011 August ; 23(3): 897–906. doi:10.1017/S0954579411000381.

Social encounters in daily life and two-year changes in metabolic risk factors in young women

Kharah Ross, B.Sc., Tara Martin, B.A., Edith Chen, Ph.D., and Gregory E. Miller, Ph.D.
Department of Psychology, University of British Columbia

Abstract

Research shows that poor social ties increase risks of morbidity and mortality from cardiovascular disease (CVD). However, little is known about the nature of everyday social encounters that give rise to this association, or when in the course of development they begin to shape disease-relevant biological processes. In this study, 122 adolescent females recorded the qualities of their everyday social interactions using electronic diaries. At the same time we measured components of the metabolic syndrome, a precursor to CVD that includes central adiposity, high blood pressure, insulin resistance, and lipid dysregulation. Metabolic symptoms were reassessed 12 and 24 months later. Hierarchical linear modeling revealed an association between negative social interactions and metabolic symptom trajectories. To the extent that participants had more intense negative social encounters in daily life, they showed increasing scores on a composite indicator of metabolic risk over two years. This association was independent of a variety of potential confounders, and persisted when symptoms of depression and broader personality traits were controlled. There was no association between positive social encounters and metabolic risk trajectories. These findings suggest that even in otherwise healthy adolescents, abrasive social encounters may accelerate the progression of CVD's early stages.

Keywords

Social interactions; adolescence; metabolic syndrome; cardiovascular disease; atherosclerosis

Introduction

Extensive research suggests that people with poor social ties are at risk for morbidity and mortality from cardiovascular disease (CVD; Barth, Schneider, & Von Kanel, 2010). A number of different constructs have been studied in this literature, including isolation, network diversity, social support, and marital status and quality. In general, individuals who are isolated and have limited access to and/or low quality of social support, and who have smaller or less diverse networks are at increased risk of CVD morbidity and mortality, independent of traditional cardiovascular (CV) risk factors (e.g., Berkman, Leo-Summers, & Horwitz, 1992; Eng, Rimm, Fitzmaurice, & Kawachi, 2002; Frasure-Smith et al., 2000; Gorkin et al., 1993; Kawachi et al., 1996; Knox, & Uvnas-Moberg, 1998; Kuper, Marmot, & Hemingway, 2002; Lett et al., 2005; Seeman, & Syme, 1987; Vogt, Mullooly, Ernst, Pope, & Hollis, 1992). There is also mounting evidence that social factors are associated with risk for the metabolic syndrome, a collection of symptoms that includes central adiposity, hypertension, insulin resistance and dyslipidaemia, and is known to be a precursor of and contributor to the pathogenesis of CVD (Grundy et al., 2005). The strongest evidence of this

phenomenon comes from long-term prospective studies that have shown that social constructs, i.e. childhood social isolation, poor marital quality, and the loss of a spouse through death or divorce, are associated with increased risk of metabolic syndrome onset or clustering of the symptoms that comprise it (e.g. Caspi, Harrington, Moffitt, Milne, & Poulton, 2006; Troxel, Matthews, Gallo, & Kuller, 2005; for an exception, see Ikeda, Kawachi, Iso, Inoue, & Tsugane, 2010).

To understand the mechanisms through which social factors contribute to CVD outcome, researchers have begun studying the biological consequences of social interactions. Research in this domain has focused on how both supportive and conflictual interactions influence biological processes involved in CVD pathogenesis. Much of this work has been done in laboratory settings. For example, studies have shown that support from others can ameliorate people's cardiac and vascular responses to brief lab stress (e.g., Kamarck, Manuch, & Jennings, 1990; Lepore, Mata Allen, & Evans, 1993; see review by Uchino, Carlisle, Birmingham, & Vaughn, 2010), which may be important because stress reactivity itself is a CVD risk factor in both healthy and clinical populations (e.g., Alderman, Ooi, Madhavan, & Cohen, 1990; Keys et al., 1971; Krantz et al., 1999; Manuck, Olsson, Hjemdahl, & Rehnqvist, 1992; Sheps et al., 2002; see review by Treiber et al., 2003). Lab studies have also evaluated the biological sequelae of social conflicts; particularly in married couples (see review by Kiecolt-Glaser, & Newton, 2001). These studies indicate that conflictual interactions raise blood pressure, trigger activity of the sympathetic nervous system (SNS) and hypothalamic-pituitary-adrenal (HPA) axis, and increase the production of inflammatory mediators (Baker et al., 2000; Graham, Christian, & Kiecolt-Glaser, 2007; Heffner, Kiecolt-Glaser, Loving, Glaser, & Malarkey, 2004; Kiecolt-Glaser et al., 1993; Kiecolt-Glaser et al., 1996; Kiecolt-Glaser et al., 1997; Kiecolt-Glaser et al., 2005; Smith et al., 2009). Each of these processes has been implicated in various stages of the development and manifestation of CVD (Kop, 1999).

More recent studies have explored the biological consequences of social interactions that occur in real world settings. Kamarck and colleagues (2002) followed 340 older adults over a 3-day period using electronic daily diaries and ambulatory blood pressure (ABP) monitors. They observed that episodes of social conflict were associated with greater systolic blood pressure (SBP) and diastolic blood pressure (DBP) activity, above and beyond potential confounds like posture and activity at the time of measurement. A similar pattern has been observed in adolescents: One study that used electronic diaries and ABP monitoring found that anxious- and avoidant-attachment boys had augmented ambulatory heart rate (HR) and blood pressure (BP) during social interactions (Gallo, & Matthews, 2006). Another found that among adolescents who have a tendency to read threat into ambiguous situations, ABP was higher during social interactions than it was when they were alone. This pattern was reversed in adolescents who tend not to read threat into ambiguous situations (Chen, Matthews, & Zhou, 2007). Day-to-day social interactions have also been related to indicators of atherosclerosis, the pathogenic condition that underlies much CVD. In particular, to the extent that men reported frequent interactions with their spouses, they showed less progression of atherosclerosis in the carotid artery over 3 years, provided they also had high marital quality (Janicki, Kamarck, Shiffman, Sutton-Tyrell, & Gwaltney, 2005). These associations were not observed among women.

Collectively, these findings suggest that social interactions have biological consequences of potential relevance to understanding CVD pathogenesis. However, a number of important questions in this literature have not yet been answered. The first has to do with the nature of the association between social interactions and disease processes. Do positive and negative encounters have separable and independent associations with the relevant outcomes? Or do they represent different ends of the same continuum? To what extent do they have

synergistic influences on disease processes, such that people who have more negative and fewer positive encounters are worst off? And can positive interactions serve as a buffer, offsetting any health effects of negative ones?

Another unanswered question in this literature has to do with what role broader personality characteristics play in driving these associations. Personality traits like extraversion, neuroticism/emotional stability, and agreeableness have strong influences on the quantity and quality of a person's social encounters (Doeven-Eggens, De Fruyt, Hendriks, Bosker, & van der Werf, 2008; Scarr, & McCartney, 1983; Swickert, Hittner, & Foster, 2010). These traits are also associated with risk for CVD morbidity and mortality (e.g., Shipley, Weiss, Der, Taylor, & Deary, 2007; Smith et al., 2006; Terracciano, Lockenhoff, Zonderman, Ferrucci, & Costa, 2008). Thus, it is important to determine the extent to which they are contributing to associations between social interactions and disease processes.

Finally, little is known about the role of social interactions and disease processes in a developmental context (Matthews, 2005). Researchers are increasingly emphasizing the importance of viewing pathogenesis of CVD from a life-course perspective. Autopsy studies indicate that atherosclerosis begins in the early decades of life (Kavey et al., 2003; McGill et al., 2000; Strong, Malcom, & McMahan, 1999) and CVD risk factors established in childhood and adolescence are generally stable across time (Morrison, Friedman, & Gray-McGuire, 2007). Adolescence is also of special interest psychologically, given the major socioemotional changes it often entails. Indeed, adolescence is a time when many youth struggle for independence from their parents, have their first romances, assemble their own friendship networks, and navigate the interpersonal complexities of their school environments. These struggles seem to have important consequences for mental health; indeed, the risk of depression increases markedly in the teenage years, particularly in females (get a cite here). These mood problems often exacerbate young women's social difficulties (Allen 2006, Lee 2010, Schrepferman 2006), creating a downward spiral that might also have implications for CVD. These considerations make adolescence a rich time of life to observe everyday social interactions and their biological consequences.

The goal of this study was to build on previous research by addressing these limitations. A sample of female adolescents recorded impressions of their day-to-day social encounters using electronic diaries. Symptoms of the metabolic symptom were assessed at the same time and annually thereafter for 2 years. We predicted that, to the extent that they had abrasive social interactions in day-to-day life, participants would display increased metabolic risk over the follow-up. We also expected positive interactions to be associated with better metabolic outcomes, and to attenuate any adverse health effects arising from negative ones. Finally, we expected these associations to be independent of personality characteristics, like neuroticism, agreeableness, and extraversion, which are associated with both social behavior and CVD outcomes.

Methods

Participants

Data were collected as part of a larger project on depression and atherosclerosis in young women at risk for affective disorders. The participants were recruited from the Vancouver, BC community through advertisements. Eligibility criteria included being (1) female, 15–19 years old, and fluent in English, (2) free of acute illness in the past 2 weeks, and (3) without a history of chronic medical or psychiatric disorders. A total of 157 young women were enrolled in the study; our analyses included only the 147 participants who were recruited to be at high risk for developing an initial episode of depression. High-risk was defined as having a first-degree relative with a history of affective disorder, and/or scoring in the top

quartile of the population on cognitive vulnerability to depression. (The other 10 participants, who were excluded from these analyses, were enrolled as low-risk controls.) Over the course of the 2 year follow-up, 25 of the 147 admissible participants were lost to attrition (17.0%): 6 of them moved away, we lost contact with 4, and 15 withdrew. Thus, the final sample consisted of 122 young, healthy female participants. All procedures and methods were approved by the University of British Columbia Research Ethics Board. Written consent was obtained from all participants prior to participation. For participants younger than 18 years of age, a parent or guardian consent was also obtained.

Procedure

Participants visited the research centre on six occasions over a 2.5 year period. The analyses in this article focus on the data gathered at study entry, 12 and 24 months later (referred to hereafter as Visits 1, 2, and 3, respectively.) These were the only study visits in which daily social interactions and metabolic symptoms were assessed. At each visit, participants arrived between 8 and 11 AM, following an overnight fasting period. After obtaining informed consent, participants completed a battery of interviews and questionnaires, described in more detail below, which were used to evaluate alternative explanations such as mood, personality characteristics, and health practices. Blood pressure values, waist circumference and a blood sample for assessment of high density lipoproteins (HDL), triglycerides, and glucose were obtained (see below for details). Following the testing sessions at Visits 1 and 2, participants were given a Palm Pilot which prompted them to rate the quality of social encounters in daily life. The diary was completed twice daily over a two-day period using a format similar to the Rochester Interaction Record (see below). In addition, participants were also asked questions about daily physical activity practices.

Measures

Perception of social encounters—On each day of monitoring, participants were queried twice about their social interactions (4 and 14 hours after waking). The first item asked if they had interacted with their closest friend, romantic partner, parent/guardian, and other friends since the last diary entry. For each category of interaction answered ‘yes,’ the Palm Pilot prompted them to rate the encounter(s) on 8 negative (conflict, criticism, disappointment, exclusion, sense of inferiority, anger, shame or embarrassment, prying) and 7 positive (pleasantness, intimacy, helpful advice, helpful assistance, care, sense of confidence, value or respect) dimensions. These items were drawn from the Rochester Interaction Record (Reis, & Wheeler, 1991) and the Diary of Ambulatory Behavioral States (Kamarck et al., 1998). Ratings were made on a 0 (not at all) to 4 (very much/a lot) scale. For each encounter, composite positive and negative scales were calculated by averaging ratings across the appropriate dimensions. Both of the scales were internally consistent, with Cronbach’s $\alpha = 0.75$ for negative perception of social encounters and $\alpha = 0.82$ for positive. Scores on the negative scale could range from 0 to 32, whereas those on the positive scale could range from 0 to 28. Scores on the two scales were inversely correlated in a modest fashion, $r = -0.17$, $p = 0.05$. Participants’ ratings of their interactions were fairly stable over time: over the year that elapsed between diary ratings at Visits 1 and 2, the stability of scores on the negative scale was $r = 0.36$ ($p < 0.001$) and on the positive scale was $r = 0.60$ ($p < 0.001$). Thus, ratings were collapsed across these periods to form more trait-like indicators of positive and negative interaction tendencies. Each of these indicators was comprised of 8 separate diary entries (2 entries per day, for 2 days, at both Visits 1 and 2), which could have included ratings of up to 4 separate social encounters each.

Metabolic risk—Following recommendations as issued in a joint statement by the American Heart Association and National Heart, Lung and Blood Institute, metabolic risk was assessed by considering levels of high-density lipoproteins (HDL) and triglycerides, as

well as fasting glucose, SBP and DBP, and waist circumference (Grundy et al., 2005). To measure resting SBP and DBP, participants were seated in a chair with an occluding cuff on one arm. Following a 5-minute rest period, 4 BP readings, spaced 2 minutes apart, were collected using an automatic, calibrated, oscillometric BP monitor (BPM-100, VSM MedTech, Coquitlam, BC). Appropriately sized BP cuffs were selected according to the diameter of the participant's arm. Average SBP and DBP were calculated by averaging the last 3 measures (the first reading is excluded as it tends to be elevated due to the novelty of the procedure). This device and protocol has been validated in pediatric populations and yield BP readings that meet the standards of the British Hypertension Society for accuracy and reliability (Mattu, Heran, & Wright, 2004).

Waist circumference was measured from the side at the midpoint between the upper iliac crest and lower costal margin at the mid-axillary line using a standard measuring tape. Measurements were taken at least twice, and repeated until a consistent reading was obtained.

Overnight fasting blood samples were collected at each visit to quantify other metabolic parameters. A 10 mL sample of blood was collected into a serum separator Vacutainer tube (Becton-Dickinson, Oakville ON) through antecubital venipuncture. The sample was spun for 12 minutes at $1200 \times g$. After the serum had been aspirated, it was frozen at -30°C until assayed in batch. The analyses were performed in the Clinical Chemistry laboratory of St Paul's Hospital in Vancouver. Triglycerides were determined enzymatically on a Hitachi 747 (Kyowa Medex, Japan) after hydrolysis to glycerol. This method has an inter-assay coefficient of variation of 1.1%. High-density lipoprotein was measured using standard enzymatic techniques with cholesterol esterase and cholesterol oxidase, after low-density, intermediate density, and very-low density lipoproteins had been precipitated through centrifugation, on a Hitachi 911 instrument (Kyowa Medex, Japan). This method has an inter-assay coefficient of variation of 5.1%. Glucose was measured with an enzymatic technique that uses hexokinase and glucose-6-phosphate dehydrogenase enzymes on an ADVIA 1650 Chemistry System (Bayer Diagnostics, Tarrytown, NY) with an inter-assay coefficient of variation of 1.2%.

Because our sample was recruited to be young and healthy, none of the participants met criteria for diagnosis with the metabolic syndrome either at study entry or during the follow-up period. Therefore, in order to model the evolution of metabolic risk during follow-up, and based on previous research that has focused on these processes in healthy youth (e.g., Eisenmann, 2008; Ekelund et al., 2005), we computed a composite score comprised of values on the six components of the metabolic syndrome: HDL, triglycerides, fasting blood glucose, waist circumference, SBP and DBP. Scores on each component were converted to a z-score; HDL was reverse coded to match risk direction of the others. Composite metabolic risk was then calculated for each visit by averaging the z scores across components. Scores were fairly stable across time, with correlations of 0.47 ($p < 0.001$) and 0.44 ($p < 0.001$) across one and two years of follow-up, respectively.

Alternative explanations—Several other variables were assessed as potential alternative explanations for any association observed between social interaction tendencies and metabolic risk trajectories. These included demographic characteristics (age, ethnicity, socioeconomic status (SES)), oral contraceptive (OC) use, depressed mood (as measured by the BDI, see below), daily physical activity, and neuroticism, agreeableness and extraversion (as measured by the Big Five Inventory (BFI), see below). Socioeconomic status was indexed by the highest years of education completed by either the participant's mother or father.

Depressed mood: Depression has profound influences on the quantity and quality of people's social interactions, and is also independently associated with increased risk for the metabolic syndrome and CVD more generally (Kinder et al., 2004; Rugulies 2002). Coupled with the fact that this sample was recruited specifically to be at risk for affective disorders, these considerations led us to consider whether depression was underlying any observed associations between social interactions and metabolic outcomes. Thus, we had participants complete the BDI (Beck, Ward, Mendelson, Mock, & Erbaugh, 1961) at Visits 1, 2 and 3. The BDI is a widely used measure of the intensity of depressive symptoms. It consists of 21 items assessing symptoms and attitudes of depression experienced during the past week, with possible scores for each question ranging from 0 to 3. The BDI was internally consistent in our sample, with an average Cronbach's α of 0.93 across visits.

Neuroticism, agreeableness, extraversion: Personality was assessed at Visit 1 using the BFI (1991), a 44-item inventory that provides assessment of the Big Five dimensions of personality. Participants are asked to rate the degree to which they agree with a characteristic presented in a statement, with responses ranging from 1 (strongly disagree) to 5 (strongly agree). For the purposes of this article, we considered 3 dimensions closely linked to social interactions and health outcomes in past research: neuroticism, which is characterized by traits like tenseness, moodiness, and anxiety (8 items) (Cronbach's $\alpha = 0.76$); extraversion, which encompasses traits like talkative, energetic, and assertive (8 items) (Cronbach's $\alpha = 0.84$); and agreeableness, which includes traits like sympathetic, kind and affectionate (9 items) (Cronbach's $\alpha = 0.75$).

Physical activity: Low levels of physical activity have been associated general negative mood (e.g., Dunn, Trivedi, & O'Neal, 2001; Janisse, Nedd, Escamilla, & Nies, 2004; Penedo, & Dahn, 2005; Scully, Kremer, Meade, Graham, & Dudgeon, 1998) and increased risk of the metabolic syndrome (e.g., Ekelund et al., 2005; Laaksonen et al., 2002; Lakka et al., 2003). To rule out physical activity as a possible confound, we chose to include it as a control variable in analyses. At the end of each day of monitoring participants were asked about physical activity using an item from the Paffenbarger Activity Scale (Paffenbarger, Blair, Lee, & Hyde, 1993). They were asked "Did you engage in any activity today akin to brisk walking, jogging, bicycling, etc., long enough to work up a sweat?" If they responded affirmatively, they were asked for how long. The average per-day minutes of exercise was then averaged over the monitoring periods following Visits 1 and 2. This approach to assessing physical fitness has been validated in previous research, which shows that self-reported brisk activity is related to VO2 max scores during treadmill tests (Paffenbarger et al., 1993).

Results

The sample consisted of healthy adolescent females, as can be seen from the information presented in Table 1. Participants were primarily of Asian and European descent, and whose parents had, on average, a university education. Table 2 contains descriptive statistics for individual components of the metabolic index. Over the two-year follow-up period the sample as a whole displayed increases in HDL, $F(1.8, 169) = 8.384, p = 0.001$, and glucose, $F(2, 184) = 4.951, p = 0.008$. There was also a trend towards increasing SBP over follow-up, $F(2, 184) = 2.452, p = 0.081$. The sample as a whole did not show changes in triglycerides, $F(1.5, 134) = 1.19, p = 0.296$, DBP, $F(2, 184) = 0.346, p = 0.708$ or waist circumference, $F(1.8, 167) = 0.682, p = 0.494$, over time. The sample as a whole did not show changes in the metabolic composite over the follow-up period, $p = .75$. However, further analyses revealed that there was marked within- and between-person variability in metabolic trajectories. The intra-class correlation (ICC) for the metabolic composite was 0.54, indicating that about half of the total variance in this construct over time was between

participants. The other half was due to within-person changes and person by time interactions. These figures suggest that there is sufficient variability in metabolic trajectories in the sample to justify exploring the potential influence of participants' social encounters.

Statistical Approach

To determine whether perceived quality of social encounters predicted changes in metabolic risk over time, we estimated a series of multi-level models with HLM 6.08 (Raudenbush, Bryk, & Congdon, 2006). The within-person (level 1) models included a variable reflecting time since study entry (in months). The between-person (level 2) models included the relevant indices reflecting social interactions, as well as 5 standard control variables: age, ethnicity (European descent vs. non-European descent), OC use, parental education, and average minutes of physical activity. In later models we added depressive symptoms and personality characteristics as covariates, to evaluate their role in shaping the observed associations. In all cases we estimated random slope models, in which Level 2 error terms were allowed to vary freely. Models were estimated with robust standard errors.

Preliminary Results

To estimate how much the control variables explained metabolic change over time, we ran an initial model that included time from study entry at level-1, and age, ethnicity, oral contraceptive use, and parental education at level-2. The results are presented in Table 3. Age was associated with change in metabolic composite over time, $b = 3.85 \times 10^{-3}$, $SE = 1.68 \times 10^{-3}$, $p = 0.024$, such that participants who were older at study had larger increases in metabolic composite scores over the follow-up. None of the other standard control variables was associated with the metabolic composite, either at study entry or over time.

Positive and Negative Social Encounters

Next we evaluated our principal hypotheses by including variables reflecting the quality of social interactions in the level-2 models. As Table 4 shows, older subjects, $b = 4.15 \times 10^{-3}$, $SE = 1.67 \times 10^{-3}$, $p = 0.015$, continued to show larger increases in metabolic symptoms over the follow-up. Over and above the contribution of the control variables, negative social interactions were associated with metabolic composite trajectories. As Figure 1 shows this was a positive association, such that participants who had more intense negative social encounters in day-to-day life showed increasing scores on the metabolic composite over the 2-year follow-up, $b = 3.79 \times 10^{-3}$, $SE = 1.91 \times 10^{-3}$, $p = 0.049$. Conversely, participants with less intense negative encounters showed a decline in metabolic composite scores over time.

Further analyses revealed that there was no association between the intensity of positive encounters and the metabolic composite at baseline, $b = -0.10$, $SE = 6.2 \times 10^{-2}$, $p = 0.10$, or over follow-up, $b = 3.34 \times 10^{-3}$, $SE = 1.9 \times 10^{-3}$, $p = 0.083$. There also was no evidence of a statistical interaction between the two dimensions of social encounters, $b = 3.5 \times 10^{-4}$, $SE = 2.0 \times 10^{-3}$, $p = 0.86$. Thus, positive encounters did not offset the apparent metabolic consequences of negative ones.

Alternative Explanations

The final models evaluated alternative explanations. We began by testing whether symptoms of depression might be driving the observed associations. BDI scores from Visits 1, 2 and 3 were added to the level-1 models. However, the intensity of negative social encounters continued to predicted changes in the metabolic composite under these conditions, $b = 5.4 \times 10^{-3}$, $SE = 2.1 \times 10^{-3}$, $p = 0.013$, suggesting this association was not being driven by underlying depressive symptoms.

We next evaluated the contribution of the personality traits extraversion, neuroticism, and agreeableness. Variables reflecting these traits were added to the level-2 equations in separate models. None of the variables were associated with scores on the metabolic composite at study entry. Only extraversion was associated with changes in the metabolic composite, $b = 5.8 \times 10^{-4}$, $SE = 2.9 \times 10^{-4}$, $p = 0.05$. Moreover, the intensity of negative encounters continued to predict larger increases in metabolic composite scores over time when extraversion, $b = 4.0 \times 10^{-3}$, $SE = 1.9 \times 10^{-3}$, $p = 0.037$, neuroticism, $b = 3.9 \times 10^{-3}$, $SE = 1.9 \times 10^{-3}$, $p = 0.045$, and agreeableness, $b = 4.3 \times 10^{-3}$, $SE = 1.9 \times 10^{-3}$, $p = 0.027$ were in the model, suggesting these traits were not driving the observed associations.

Discussion

With mounting evidence that poor social ties increase the risk of morbidity and mortality from CVD, greater focus has been placed on exploring the biological consequences of everyday social interactions. The goal of this study was to determine whether the quality of these interactions presaged changes in symptoms of the metabolic syndrome in healthy young women at risk for depression. We found evidence that the intensity of negative, but not positive, social encounters was associated with trajectories of metabolic risk over a two-year period. To the extent that they had more intense negative encounters in day-to-day life, participants showed increasing scores on the metabolic composite over follow-up. Conversely, participants with less intense negative interactions showed a decline in metabolic composite scores over the same timeframe. This observation persisted after adjustment for age, ethnicity, SES, OC use, daily physical activity, depressed mood, and personality traits (agreeableness, neuroticism, and extraversion). Positive encounters were not associated with metabolic trajectories, and did not offset the apparent influence of negative ones. These results further our understanding of how everyday social interactions affect the development of metabolic processes involved in CVD. In doing so they may help to explain why broader constructs, like perceived social support or social network diversity, relate to CVD outcomes in such a robust fashion.

We predicted that negative social interactions would be associated with worsening metabolic trajectories. The results were consistent with this hypothesis. To the degree that they experienced more intense negative encounters, participants showed increasing scores on the metabolic composite over time. This association was fairly small in its magnitude, with each standard deviation (SD) increment in negative interactions associated with a 3.8×10^{-3} SD greater increase in metabolic composite trajectories. Differences of this size are unlikely to have any immediate clinical significance, especially in this population of healthy adolescents. However, if this dynamic was sustained through adolescence and adulthood it would presumably contribute to the development of frank metabolic syndrome. Apart from their potential clinical implications, these findings are noteworthy in two more theoretical respects: First, they suggest the presence of a graded association between social interactions and metabolic symptoms, whereby even small variations in people's tendencies to have negative exchanges seem to have biological repercussions. Second, they suggest that this dynamic begins early in life, when people are still in good overall metabolic health. The implication is that aversive social encounters, often viewed as a normative development experience for adolescents, may set off biological trajectories that, if sustained, ultimately contribute to the development of CVD.

We also expected that positive social interactions would relate to better metabolic outcomes, or at least offset the consequences of negative encounters. However, neither of these hypotheses was borne out. These findings indicate that, at least in our sample, there are no metabolic benefits associated with having regular positive social interactions. When considered alongside the roughly linear effects that emerged for negative encounters, these

results suggest that young women are likely to accrue more metabolic benefits by reducing the intensity of negative social interactions than increasing the intensity of positive ones. In general, this pattern of findings is consistent with literature on well-being, which suggests that negative events tend to be more impactful than positive ones (Baumeister, Bratslavsky, Finkenauer, & Vohs, 2001; Finch, Okun, Barrera, Zautra, & Reich, 1989; Finch, Okun, Pool, & Ruehlman, 1999; Newsom, Rook, Nishishiba, Sorkin, & Mahan, 2005; Rook, 2001; Schuster, Kessler, & Aseltine, 1990). It is also consistent with an early study of BP responses to marital conflict, which concluded that “not being nasty matters more than being nice” (Ewart, Taylor, Kraemer, & Agras, 1991, p. 155). Of course, it is also possible that the null findings herein stem from problems with the study’s research design. The diary sampling period was fairly brief. If influential positive encounters occur infrequently, our sampling period could have missed them altogether. This also could have posed a problem for testing the offset hypothesis, particularly if positive encounters act as buffers against very intense (but infrequent) negative exchanges. It is also possible that our diary did not tap the dimensions of positive social interactions that are most important for health.

Though these data suggest that negative interactions worsen metabolic symptoms over time, the underlying mechanism through which this process occurs is not yet understood. Our analyses help to eliminate some of the most plausible explanations, such as confounding by age, ethnicity, SES, physical activity, and depressed mood. The observed patterns also do not seem to be a reflection of the influence of personality characteristics like neuroticism, agreeableness, and extraversion, which shape the quality of people’s social encounters and also have been related to CVD outcomes. Of course, it is possible that more narrow facets of personality not captured in the Big 5’s major dimensions could play a role in the observed associations.

Assuming the observed associations reflect a process that is causal in nature, they could be driven by the direct biological consequences of the social encounters themselves. As we noted earlier in the paper, acute episodes of social conflict raise blood pressure (Gallo, & Matthews, 2006; Kamarck et al., 2002), which is a component of the metabolic syndrome. Conflictual interactions also trigger activity of the SNS and the HPA axis (Graham et al., 2007; Heffner et al., 2004; Kiecolt-Glaser et al., 1996; Kiecolt-Glaser et al., 1997; Kiecolt-Glaser et al., 2005), whose hormonal products are thought to increase fat accumulation and impair glucose control, among other things (Bjorntorp, & Rosmond, 1999; Brunner et al., 2002). And in young women, more chronic exposure to negative social encounters has been prospectively linked to upregulation of inflammatory processes (Miller, Rohleder, & Cole, 2009), which are known to play a key role in the pathogenesis of most components of the metabolic syndrome (Hotamisligil, 2006). It will be important for the next wave of studies in this area to evaluate the role these proposed mechanisms play in the link between social encounters and metabolic trajectories. Meanwhile, there could be value in targeting social processes as a way to intervene and slow early disease processes. If our findings do indeed reflect a causal process, interventions that reduce the frequency and/or severity of negative interactions could have consequences for metabolic trajectories. Indeed, there is evidence that family systems therapy, which aims to improve parent-child relationships, can improve blood glucose control in teenagers with diabetes (see papers by Wysocki, especially *Diabetes Care*. 2007 Mar;30(3):555-60). Such interventions might also have benefits for psychopathology and comorbidity of mental and physical illnesses.

Future research is also needed to explain the temporal dimension of our findings. The metabolic disparities associated with abrasive encounters were not apparent at baseline, but seemed to gradually accumulate as the study’s follow-up period progressed. The reasons for this pattern are unclear. During adolescence there are profound changes in body size and shape, as well as blood pressure, insulin resistance, and other metabolic processes related to

CVD (e.g., see Moran et al., 2008). At this stage of life there are also considerable within-person symptom fluctuations, with nearly 50% of youth classified as having metabolic syndrome losing their diagnosis within 3 years (Goodman et al., 2007). It may be the influence of social encounters only becomes apparent in the later stages of adolescence, when the frequency of these fluctuations has subsided and more stable risk trajectories becomes established.

This study has several limitations that should be considered. First, because it focused on young women who were at risk for affective disorders, it is unclear how broadly the results can be generalized to other populations. The women studied herein may be especially sensitive to negative social encounters, and some may have attributional tendencies that would exaggerate responsibility for, and consequences of, abrasive interactions that occurred with friends, family, or even strangers. Thus, replication of these findings in other populations is necessary before they can be safely generalized. Second, our daily diary sampling was limited to two 48-hour periods. Although this duration was adequate to capture the individual differences in negative interactions that associated with metabolic trajectories, it may have been too short to tap positive encounters that are infrequent but highly impactful. In future research, more extended periods of sampling would help to alleviate this concern, and allow for a more robust test of the hypotheses we considered. Third, as is the case with any study that has an observational design, there are potential alternative explanations for our findings that involve unmeasured confounds. Particularly worrisome in this regard are genetic liabilities that predispose individuals to both abrasive social encounters and symptoms of metabolic syndrome. One especially rigorous way to evaluate whether our findings are causal in nature would be to assess metabolic endpoints in people who received (or did not receive) an intervention to improve the quality of their social interactions (e.g., couples therapy).

In summary, this study's findings suggest that, in adolescent females, there is a graded association between the negativity of social encounters and trajectories of metabolic symptoms over time. Positive interactions do not seem to have an influence on metabolic profiles. They also do not appear to function as a buffer against the metabolic consequences of negative interactions. Collectively, these findings shed light on some of the mechanisms that might underlie the broader epidemiologic linkage between social relationships and CVD morbidity and mortality (e.g., Barth et al., 2010; Berkman et al., 1992; Eng et al., 2002; Frasure-Smith et al., 2000; Gorkin et al., 1993; Kawachi et al., 1996; Knox, & Uvnas-Moberg, 1998; Kuper et al., 2002; Lett et al., 2005; Seeman, & Syme, 1987; Vogt et al., 1992). They also highlight the important role that adolescence experiences may play in early stages of the disease process.

Acknowledgments

This research was supported by grants from the Canadian Institutes of Health Research (67191), the National Alliance for Research on Schizophrenia and Depression, and the Heart and Stroke Foundation of Canada to Gregory E. Miller.

References

- Alderman MH, Ooi WL, Madhavan S, Cohen H. Blood pressure reactivity predicts myocardial infarction among treated hypertensive patients. *Journal of Clinical Epidemiology*. 1990; 43:859–866. [PubMed: 2213075]
- Baker B, Paquette M, Szalai JP, Driver H, Perger T, Helmers K, et al. The influence of marital adjustment on 3-year left ventricular mass and ambulatory blood pressure in mild hypertension. *Archives of Internal Medicine*. 2000; 160:3453–3458. [PubMed: 11112239]

- Barth J, Schneider S, von Kanel R. Lack of social support in the etiology and prognosis of coronary heart disease: A systematic review and meta-analysis. *Psychosomatic Medicine*. 2010; 72:229–238. [PubMed: 20223926]
- Baumeister RF, Bratslavsky E, Finkenauer C, Vohs KD. Bad is stronger than good. *Review of General Psychology*. 2001; 5:323–370.
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Archives of General Psychiatry*. 1961; 4:561–571. [PubMed: 13688369]
- Berkman LF, Leo-Summers L, Horwitz RI. Emotional support and survival after myocardial infarction: A prospective, population-based study of the elderly. *Annals of Internal Medicine*. 1992; 117:1003–1009. [PubMed: 1443968]
- Bjorntorp P, Rosmond R. Hypothalamic origin of the metabolic syndrome X. *Annals of the New York Academy of Sciences*. 1999; 892:297–307. [PubMed: 10842670]
- Brunner EJ, Hemingway H, Walker BR, Page M, Clarke P, Juneja M, et al. Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome. *Circulation*. 2002; 106:2659–2665. [PubMed: 12438290]
- Caspi A, Harrington H, Moffitt TE, Milne BJ, Poulton R. Socially isolated children 20 years later. *Archives of Pediatric and Adolescent Medicine*. 2006; 160:805–811.
- Chen E, Matthews KA, Zhou F. Interpretations of ambiguous social situations and cardiovascular responses in adolescents. *Annals of Behavioral Medicine*. 2007; 34:26–36. [PubMed: 17688394]
- Doeven-Eggens L, De Fruyt F, Hendriks AAJ, Bosker RJ, Van der Werf MPC. Personality and personal network type. *Personality and Individual Differences*. 2008; 45:689–693.
- Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Medicine & Science in Sports & Exercise*. 2001; 33:S587–S597. [PubMed: 11427783]
- Eisenmann JC. On the use of a continuous metabolic syndrome score in pediatric research. *Cardiovascular Diabetology*. 2008; 7
- Ekelund U, Brage S, Franks PW, Hennings S, Emms S, Wareham NJ. Physical activity energy expenditure predicts progression toward the metabolic syndrome independently of aerobic fitness in middle-aged health Caucasians. *Diabetes Care*. 2005; 28:1195–1200. [PubMed: 15855588]
- Eng PM, Rimm EB, Fitzmaurice G, Kawachi I. Social ties and change in social ties in relation to subsequent total and cause-specific mortality and coronary heart disease incidence in men. *American Journal of Epidemiology*. 2002; 155:700–709. [PubMed: 11943687]
- Ewart CK, Taylor CB, Kraemer HC, Agras WS. High blood pressure and marital discord: not being nasty matters more than being nice. *Health Psychology*. 1991; 10:155–163. [PubMed: 1879387]
- Finch JF, Okun MA, Barrera MJ, Zautra AJ, Reich JW. Positive and negative social ties among older adults: Measurement models and the prediction of psychological distress and well being. *American Journal of Community Psychology*. 1989; 17:585–605. [PubMed: 2534023]
- Finch JF, Okun MA, Pool GJ, Ruehlman LS. A comparison of the influence of conflictual and supportive social interactions on psychological distress. *Journal of Personality*. 1999; 67:581–621. [PubMed: 10444852]
- Frasure-Smith N, Lesperance F, Gravel G, Masson A, Juneau M, Talajic M, et al. Social support, depression, and mortality during the first year after myocardial infarction. *Circulation*. 2000; 101:1919–1924. [PubMed: 10779457]
- Gallo LC, Matthews KA. Adolescents' attachment orientation influences ambulatory blood pressure responses to everyday social interactions. *Psychosomatic Medicine*. 2006; 68:253–261. [PubMed: 16554391]
- Goodman E, Daniels SR, Meigs JB, Dolan LM. Instability in the diagnosis of metabolic syndrome in adolescents. *Circulation*. 2007; 115:2316–2322. [PubMed: 17420347]
- Gorkin L, Schron EB, Brooks MM, Wiklund I, Kellen J, Verter J, et al. Psychosocial predictors of mortality in the Cardiac Arrhythmia Suppression Trial-1 (CAST-1). *American Journal of Cardiology*. 1993; 71:263–267. [PubMed: 8427165]
- Graham, JE.; Christian, LM.; Kiecolt-Glaser, JK. Close relationships and immunity. In: Ader, R., editor. *Psychoneuroimmunology*. 4. San Diego: Elsevier; 2007. p. 781-798.

- Grundey SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of metabolic syndrome: An American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Circulation*. 2005; 112:2735–2752. [PubMed: 16157765]
- Heffner KL, Kiecolt-Glaser JK, Loving TJ, Glaser R, Malarkey WB. Spousal support satisfaction as a modifier of physiological responses to marital conflict in younger and older couples. *Journal of Behavioral Medicine*. 2004; 27:233–254. [PubMed: 15259454]
- Hotamisligil GS. Inflammation and metabolic disorders. *Nature*. 2006; 444:860–867. [PubMed: 17167474]
- Ikedo A, Kawachi I, Iso H, Inoue M, Tsugane S. Gender difference in the association between social support and metabolic syndrome in Japan: The ‘enkai’ effect? *Journal of Epidemiology and Community Health*. 2010
- Janicki DL, Kamarck TW, Shiffman S, Sutton-Tyrell K, Gwaltney CJ. Frequency of spousal interaction and 3-year progression of carotid artery intima medial thickness: The Pittsburgh Healthy Heart Project. *Psychosomatic Medicine*. 2005; 67:889–896. [PubMed: 16314593]
- Janisse HC, Nedd D, Escamilla S, Nies MA. Physical activity, social support, and family structure as determinants of mood among European-American and African-American women. *Women & Health*. 2004; 39:101–116.
- Kamarck TW, Janicki DL, Shiffman S, Polk DE, Muldoon MF, Liebenauer LL, et al. Psychosocial demands and ambulatory blood pressure: A field assessment approach. *Physiology and Behavior*. 2002; 77:699–704. [PubMed: 12527022]
- Kamarck TW, Manuck SB, Jennings JR. Social support reduces cardiovascular reactivity to psychological challenge: A laboratory model. *Psychosomatic Medicine*. 1990; 52:42–58. [PubMed: 2305022]
- Kamarck TW, Shiffman S, Smithline L, Goodie JL, Paty JA, Gnys M, et al. Effects of task strain, social conflict, and emotional activation on ambulatory cardiovascular activity: Daily life consequences of recurring stress in a multiethnic adult sample. *Health Psychology*. 1998; 17:17–29. [PubMed: 9459066]
- Kavey RW, Daniels SR, Lauer RM, Atkins DL, Haymann LL, Taubert K. American Heart Association guidelines for primary prevention of atherosclerotic cardiovascular disease beginning in childhood. *Circulation*. 2003; 107:1562–1566. [PubMed: 12654618]
- Kawachi I, Colditz GA, Ascherio A, Rimm EB, Giovannucci E, Stampfer MJ, et al. A prospective study of social networks in relation to total mortality and cardiovascular disease in men in the USA. *Journal of Epidemiology and Community Health*. 1996; 50:245–251. [PubMed: 8935453]
- Keys A, Taylor HL, Blackburn H, Brozek J, Anderson JT, Simonson E. Mortality and coronary heart disease among men studied for 23 years. *Archives of Internal Medicine*. 1971; 128:201–214. [PubMed: 5564199]
- Kiecolt-Glaser JK, Glaser R, Cacioppo JT, MacCallum RC, Snyder-Smith M, Kim C, et al. Marital conflict in older adults: Endocrinological and immunological correlates. *Psychosomatic Medicine*. 1997; 59:339–349. [PubMed: 9251151]
- Kiecolt-Glaser JK, Loving TJ, Stowell JR, Malarkey WB, Lemeshow S, Dickinson SL, et al. Hostile marital interactions, pro-inflammatory cytokine production, and wound healing. *Archives of General Psychiatry*. 2005; 62:1377–1384. [PubMed: 16330726]
- Kiecolt-Glaser JK, Malarkey WB, Chee M, Newton T, Cacioppo JT, Mao H, et al. Negative behavior during marital conflict is associated with immunological down-regulation. *Psychosomatic Medicine*. 1993; 55:395–409. [PubMed: 8265740]
- Kiecolt-Glaser JK, Newton TL. Marriage and health: His and hers. *Psychological Bulletin*. 2001; 127:472–503. [PubMed: 11439708]
- Kiecolt-Glaser JK, Newton T, Cacioppo JT, MacCallum RC, Glaser R, Malarkey WB. Marital conflict and endocrine function: Are men really more physiologically affected than women? *Journal of Consulting and Clinical Psychology*. 1996; 64:324–332. [PubMed: 8871417]
- Kinder LS, Carnethon MR, Palaniappan LP, King AC, Fortmann SP. Depression and the metabolic syndrome in young adults: Findings from the third National Health and Nutrition Examination Survey. *Psychosomatic Medicine*. 2004; 66:316–322. [PubMed: 15184689]

- Knox SS, Uvnas-Moberg K. Social isolation and cardiovascular disease: An atherosclerotic pathway? *Psychoneuroendocrinology*. 1998; 23:877–890. [PubMed: 9924742]
- Kop WJ. Chronic and acute psychological risk factors for clinical manifestations of coronary artery disease. *Psychosomatic Medicine*. 1999; 61:476–487. [PubMed: 10443756]
- Krantz DS, Santiago HT, Kop WJ, Merz CNB, Rozanski A, Gottdiener JS. Prognostic value of mental stress testing in coronary artery disease. *American Journal of Cardiology*. 1999; 84:1292–1297. [PubMed: 10614793]
- Kuper H, Marmot M, Hemingway H. Systematic review of prospective cohort studies of psychosocial factors in the etiology and prognosis of coronary heart disease. *Seminars in Vascular Medicine*. 2002; 2:267–314. [PubMed: 16222620]
- Laaksonen DE, Lakka H, Salonen JT, Niskanen LK, Rauramaa R, Lakka TA. Low levels of leisure-time physical activity and cardiorespiratory fitness predict development of the metabolic syndrome. *Diabetes Care*. 2002; 25:1612–1618. [PubMed: 12196436]
- Lakka TA, Laaksonen DE, Lakka H, Mannikko N, Niskanen LK, Rauramaa R, et al. Sedentary lifestyle, poor cardiorespiratory fitness, and the metabolic syndrome. *Medicine & Science in Sports & Exercise*. 2003; 35:1279–1286. [PubMed: 12900679]
- Lepore SJ, Mata Allen KA, Evans GW. Social support lowers cardiovascular reactivity to an acute stressor. *Psychosomatic Medicine*. 1993; 55:518–524. [PubMed: 8310112]
- Lett HS, Blumenthal JA, Babyak MA, Strauman TJ, Robins C, Sherwood A. Social support and coronary heart disease: Epidemiologic evidence and implications for treatment. *Psychosomatic Medicine*. 2005; 67:869–878. [PubMed: 16314591]
- Manuck SB, Olsson G, Hjemdahl P, Rehnqvist N. Does cardiovascular reactivity to mental stress have prognostic value in post-infarction patients? A pilot study. *Psychosomatic Medicine*. 1992; 54:102–108. [PubMed: 1553397]
- Matthews KA. Psychological perspectives on the development of coronary heart disease. *American Psychologist*, November. 2005:783–796.
- Mattu GS, Heran BS, Wright JM. Overall accuracy of the BpTRU - an automated electronic blood pressure device. *Blood Pressure Monitoring*. 2004; 9:47–52. [PubMed: 15021078]
- McGill HCJ, McMahan CA, Herderick EE, Malcom GT, Tracy RE, Strong JP. Origin of atherosclerosis in childhood and adolescence. *American Journal of Clinical Nutrition*. 2000; 72:1307S–1315S. [PubMed: 11063473]
- Miller GE, Rohleder N, Cole SW. Chronic interpersonal stress predicts activation of pro- and anti-inflammatory signaling pathways 6 months later. *Psychosomatic Medicine*. 2009; 71:57–62. [PubMed: 19073750]
- Moran A, Jacobs DRJ, Steinberger J, Steffen LM, Pankow JS, Hong CP, Sinaiko AR. Changes in insulin resistance and cardiovascular risk during adolescence: establishment of differential risk in males and females. *Circulation*. 2008; 117:2361–2368. [PubMed: 18427135]
- Morrison JA, Friedman LA, Gray-McGuire C. Metabolic syndrome in childhood predicts adult cardiovascular disease 25 years later: The Princeton Lipid Research Clinics Follow-up Study. *Pediatrics*. 2007; 120:340–345. [PubMed: 17671060]
- Newsom JT, Rook KS, Nishishiba M, Sorkin DH, Mahan TL. Understanding the relative importance of positive and negative social exchanges: Examining specific domains and appraisals. *Journal of Gerontology*. 2005; 60B:P304–P312.
- Paffenbarger RSJ, Blair SN, Lee I, Hyde RT. Measurement of physical activity to assess health effects in free-living populations. *Medicine & Science in Sports & Exercise*. 1993; 25:60–70. [PubMed: 8423758]
- Penedo FJ, Dahn JR. Exercise and well-being: A review of mental and physical health benefits associated with physical activity. *Current Opinion in Psychiatry*. 2005; 18:189–193. [PubMed: 16639173]
- Raudenbush, SW.; Bryk, AS.; Congdon, RT. HLM Version 6.03. Chicago, IL: Scientific Software International; 2006.
- Reis, HT.; Wheeler, L. Studying social interaction with the Rochester Interaction Record. In: Zanna, MP., editor. *Advances in Experimental Social Psychology*. San Diego: Academic Press, Inc; 1991. p. 269-317.

- Rook KS. Emotional health and positive versus negative social exchanges: A daily diary analysis. *Applied Developmental Science*. 2001; 5:86–97.
- Rugulies R. Depression as a predictor for coronary heart disease: A review and meta-analysis. *American Journal of Preventive Medicine*. 2002; 23:51–61. [PubMed: 12093424]
- Scarr S, McCartney K. How people make their own environments: A theory of genotype → environmental effects. *Child Development*. 1983; 54:424–435. [PubMed: 6683622]
- Schuster TL, Kessler RC, Aseltine RHJ. Supportive interactions, negative interactions, and depressed mood. *American Journal of Community Psychology*. 1990; 18:423–438. [PubMed: 2264558]
- Scully D, Kremer J, Meade MM, Graham R, Dudgeon K. Physical exercise and psychological well being: A critical review. *British Journal of Sports Medicine*. 1998; 32:111–120.10.1136/bjism.32.2.111 [PubMed: 9631216]
- Seeman TE, Syme SL. Social networks and coronary artery disease: A comparison of the structure and function of social relations as predictors of disease. *Psychosomatic Medicine*. 1987; 49:341–354. [PubMed: 3615763]
- Sheps DS, McMahon RP, Becker L, Carney RM, Freedland KE, Cohen JD, et al. Mental stress-induced ischemia and all-cause mortality in patients with coronary artery disease: Results from the Psychophysiological Investigations of Myocardial Ischemia study. *Circulation*. 2002; 105:1780–1784. [PubMed: 11956119]
- Shipley BA, Weiss A, Der G, Taylor MD, Deary IJ. Neuroticism, extraversion, and mortality in the UK Health and Lifestyle Survey: A 21-year prospective cohort study. *Psychosomatic Medicine*. 2007; 69:923–931. [PubMed: 17991814]
- Smith TW, MacKenzie J. Personality and risk of physical illness. *Annual Review of Clinical Psychology*. 2006; 2:435–467.
- Smith TW, Uchino BN, Berg CA, Florsheim P, Pearce G, Hawkins M, et al. Conflict and collaboration in middle-aged and older couples: II. Cardiovascular reactivity during marital interaction. *Psychology and Aging*. 2009; 24:274–286. [PubMed: 19485647]
- Strong JP, Malcom GT, McMahan CA, Tracy RE, Newman WP, Herdrick EE, et al. Prevalence and extent of atherosclerosis in adolescents and young adults: Implications for prevention from the Pathobiological Determinants of Atherosclerosis in Youth Study. *JAMA*. 1999; 281:727–735. [PubMed: 10052443]
- Swickert RJ, Hittner JB, Foster A. Big Five traits interact to predict perceived social support. *Personality and Individual Differences*. 2010; 48:736–741.
- Terracciano A, Lockenhoff CE, Zonderman AB, Ferrucci L, Costa PT. Personality predictors of longevity: Activity, emotional stability, and conscientiousness. *Psychosomatic Medicine*. 2008; 70:621–627. [PubMed: 18596250]
- Treiber FA, Kamarck TW, Schneiderman N, Sheffield D, Kapuku G, Taylor T. Cardiovascular reactivity and development of preclinical and clinical disease states. *Psychosomatic Medicine*. 2003; 65:46–62. [PubMed: 12554815]
- Troxel WM, Matthews KA, Gallo LC, Kuller LH. Marital quality and occurrence of the metabolic syndrome in women. *Archives of Internal Medicine*. 2005; 165:1022–1027. [PubMed: 15883241]
- Uchino BN, Carlisle M, Birmingham W, Vaughn AA. Social support and the reactivity hypothesis: Conceptual issues in examining the efficacy of received support during acute psychological stress. *Biological Psychology*. 2010
- Vogt TM, Mullooly JP, Ernst D, Pope CR, Hollis JF. Social networks as predictors of ischemic heart disease, cancer, stroke and hypertension: Incidence, survival and mortality. *Journal of Clinical Epidemiology*. 1992; 45:659–666. [PubMed: 1607905]

Figure 1.

Negative social encounters and metabolic composite scores over two years. Participants who had more negative social encounters in day-to-day life showed increasing scores on the metabolic composite over follow-up, $b = 3.79 \times 10^{-3}$, $SE = 1.91 \times 10^{-3}$, $p = 0.049$. Conversely, participants with fewer negative social encounters showed a decline in metabolic composite scores over the same timeframe.

Table 1

Characteristics of the Sample (N = 122)

Variable	Mn +/- SD or Number (%)
Ethnicity	
European descent	67 (48.6)
Non-European descent	71 (51.4)
Age (years)	17.0 +/- 1.34
SES (highest parental years of education)	16.1 +/- 2.96
Using oral contraceptives	27 (19.9)
Intensity of social encounters	
Average negative encounter (0–32)	3.13 +/- 2.45
Average positive encounter (0–28)	11.6 +/- 3.57
Average daily brisk physical activity (minutes)	23.4 +/- 27.1
Agreeableness	33.6 +/- 5.19
Neuroticism	24.6 +/- 5.52
Extraversion	26.5 +/- 5.80
Depressed mood from BDI	
Visit 1	7.18 +/- 5.91
Visit 2	6.77 +/- 5.69
Visit 3	6.00 +/- 5.18

Table 2

Descriptive statistics for metabolic parameters over the course of study.

Variable	Mn +/- SD or Number (%)
HDL (mmol/L)	
Visit 1	1.48 +/- 0.282
Visit 2	1.52 +/- 0.297
Visit 3	1.58 +/- 0.306
Triglycerides (mmol/L)	
Visit 1	0.865 +/- 0.342
Visit 2	0.964 +/- 0.747
Visit 3	0.961 +/- 0.417
Fasting blood glucose (mmol/L)	
Visit 1	4.52 +/- 0.356
Visit 2	4.53 +/- 0.483
Visit 3	4.65 +/- 0.385
Systolic BP (mmHg)	
Visit 1	105 +/- 8.43
Visit 2	103 +/- 8.61
Visit 3	103 +/- 9.91
Diastolic BP (mmHg)	
Visit 1	66.7 +/- 7.77
Visit 2	66.6 +/- 8.28
Visit 3	67.4 +/- 8.81
Waist circumference (cm)	
Visit 1	72.0 +/- 7.33
Visit 2	71.3 +/- 6.37
Visit 3	71.8 +/- 7.91

Table 3

Results of HLM model predicting metabolic profiles over time from control variables.

Variable	B	SE	P
Intercept (Baseline)			
Ethnicity	-0.105	0.080	0.193
Age	-0.033	0.030	0.281
SES	-0.015	0.014	0.309
Physical Activity	1.73×10^{-4}	1.61×10^{-3}	0.915
OC	0.046	0.097	0.632
Slope (Trajectory)			
Ethnicity	-1.66×10^{-3}	3.91×10^{-3}	0.672
Age	3.85×10^{-3}	1.68×10^{-3}	0.024
SES	5.49×10^{-4}	7.78×10^{-4}	0.482
Physical Activity	7.40×10^{-5}	8.10×10^{-5}	0.363
OC	-2.22×10^{-3}	5.17×10^{-3}	0.668

Table 4

Results of HLM model predicting metabolic profiles over time from control variables and negative social interactions.

Variable	B	SE	P
Intercept (Baseline)			
Ethnicity	-0.103	0.082	0.211
Age	-0.034	0.030	0.265
SES	-0.015	0.015	0.322
Physical Activity	1.67×10^{-4}	1.61×10^{-4}	0.918
OC	0.045	0.098	0.644
Negative Encounters	-2.52×10^{-3}	0.048	0.959
Slope (Trajectory)			
Ethnicity	-2.60×10^{-3}	3.79×10^{-3}	0.495
Age	4.15×10^{-3}	1.67×10^{-3}	0.015
SES	3.85×10^{-4}	7.49×10^{-4}	0.607
Physical Activity	7.00×10^{-5}	7.80×10^{-5}	0.369
OC	-3.61×10^{-3}	5.13×10^{-3}	0.483
Negative Encounters	3.79×10^{-3}	1.91×10^{-3}	0.049