Published in final edited form as:

Urology. 2010 September 1; 76(3): 722–. doi:10.1016/j.urology.2010.01.030.

Long Term-Survival after Radical Prostatectomy for men with High Gleason Sum

Phillip M. Pierorazio, MD, Thomas J. Guzzo, MD, Misop Han, MD, Trinity J. Bivalacqua, MD, PhD, Jonathan I. Epstein, MD, Edward M. Schaeffer, MD, PhD, Mark Schoenberg, MD, Patrick C. Walsh, MD, and Alan W. Partin, MD, PhD

Keywords

Prostate Cancer; High-Risk; Gleason sum

RESPONSE TO EDITORIAL COMMENTS REGARDING MANUSCRIPT URL-D-10-00014

We would like to thank the authors of both editorials for their insightful commentary regarding our manuscript. We agree that the ultimate goal should be to preoperatively recognize these select patients with high Gleason prostate cancer and favorable surgical outcomes. Further studies into clinical parameters, molecular markers and imaging technologies are needed to accurately identify these men.

Phillip Pierorazio, MD

Corresponding Author: Phillip M. Pierorazio, James Buchanan Brady Urological Institute, Johns Hopkins Medical Institutions, 600 N. Wolfe Street, Marburg 134, Baltimore, Maryland 21287, philpierorazio@jhmi.edu, (410) 955-0351 phone, (410) 614-3695 fax.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.