Open Access

Anomalous Origin of the Left Circumflex Coronary Artery From the First Diagonal Branch Presented as Acute Myocardial Infarction

Jung Hyun Kim, MD¹, Geun Jin Ha, MD², Myung Jun Seong, MD², Jin Wook Jung, MD³,
So Yeon Kim, MD⁴, Sung Hee Moon, MD⁵, and Young Soo Lee, MD²

¹Division of Cardiology, Gimcheon Jeil Hospital, Gimcheon,

²Divisions of Cardiology and ⁵Radiology, Daegu Catholic University Medical Center, Daegu,

³Pohang St. Mary Hospital, Pohang, ⁴Division of Cardiology, Dongguk University Medical Center, Gyeongju, Korea

ABSTRACT

Coronary artery anomalies are diagnosed in 0.6 to 1.5% of patients who undergo coronary angiography (CAG). They may present with life threatening conditions but are generally asymptomatic. Recognition and adequate visualization of the anomaly is essential for correct management of the condition. However, in some cases the exact orifice and course of an anomalous coronary vessel cannot be selectively identified by CAG. In this report, a 54-year-old man was admitted to the hospital with acute inferior myocardial infarction and had an anomalous origin of the left circumflex coronary artery (LCX) from the first diagonal branch (D1). In CAG, the right CAG showed no significant stenosis and fortunately we found an anomalous origin of the LCX from the D1. The course of LCX was precisely established by 64-slice multi-detector computed tomography. (**Korean Circ J 2011;41:612-614**)

KEY WORD: Coronary vessel anomalies.

Introduction

The incidence of coronary artery anomalies in a routine coronary angiography (CAG) series is between 0.6% and 1.5%.¹⁾ The ostium of an aberrant vessel may be difficult to identify in the diagnostic procedure, but recognition and adequate visualization of the anomaly is essential for appropriate patient management, especially in patients undergoing evaluation for percutaneous coronary intervention (PCI) and cardiac surgery.²⁾

This report describes the anomalous origin of the left cir-

cumflex coronary artery (LCX) from the first diagonal branch in a 54-year-old man, which presented as an acute inferior myocardial infarction.

Case

A 54-year-old man with hypertension and 30 pack-years smoking history presented with chest pain. The resting electrocardiogram was demonstrating ST-segment elevation in leads II, III and aVF. CAG showed no critical stenosis in the right coronary artery (RCA) and no visualization of the LCX. Initially, we considered total occlusion of ostium of the LCX but the guide wire could not bypass the site which usually exists in an ostium of the LCX. We then identified the small artery originating from the first diagonal branch with thrombolysis in the myocardial infarction flow grade 1 (Fig. 1A and B). After balloon angioplasty was accomplished for that artery and D1, its course was similar to that of the LCX (Fig. 1C and D). To trace the exact anatomical course of coronary arteries, 64-slice multi-detector computed tomography (MDCT) was performed. MDCT demonstrated an anomalous origin of the LCX coursed to the left atrioventricular groove

Received: May 31, 2010

Revision Received: July 19, 2010

Accepted: November 1, 2010

Correspondence: Young Soo Lee, MD, Division of Cardiology, Daegu Catholic University Medical Center, 3056-6 Daemyeong 4-dong, Nam-gu, Daegu 705-718, Korea
Tel: 82-53-650-3010, Fax: 82-53-621-3166
E-mail: mdlees@cu.ac.kr

• The authors have no financial conflicts of interest.

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Fig. 1. Coronary angiographic finding. A and B: coronary angiography shows the small artery (arrow) with TIMI flow grade 1 originating from the D1 (arrowhead). C and D: after percutaneous coronary intervention for the small artery, the coronary angiography reveals an anomalous origin of the LCX from the D1. A and C: left anterior oblique cranial view. B and D: left anterior oblique caudal views. TIMI: thrombolysis in myocardial infarction, LAD: left anterior descending coronary artery, LCX: left circumflex coronary artery, D1: first diagonal branch.

from the first diagonal branch (Fig. 2).

Discussion

Coronary artery anomalies have been identified in 0.6 to 1.5% of coronary angiograms.¹⁾ Previous studies reported that LCX anomalies were the third most common coronary artery anomaly diagnosed by using CAG.³⁾ A total of 71% of patients with LCX anomaly had significant stenosis of the proximal LCX and 11% had severe atherosclerosis in this vessel alone.⁴⁾ The origin of the LCX from the right sinus of Valsalva or RCA is a relatively common anatomical variation¹⁾⁵⁾ and some cases reported the origin of the LCX from the pulmonary artery.⁶⁾ However, an anomalous origin of the LCX from the first diagonal branch is a very rare congenital anomaly.

Since the variable clinical presentation and prognosis of coronary anomalies depends on the proximal course of the

anomalous arising coronary artery in relation to the great vessels, early detection and exact delineation of their proximal course are crucial.²⁾ Conventional CAG has traditionally been used to diagnose coronary anomalies but in some cases the exact origin of anomalous coronary vessels cannot be selectively identified by this technique.⁶⁾⁷⁾ The three dimensional acquisitions of either MDCT or cardiovascular magnetic resonance allow unambiguous interpretation of the locations of coronary origins.⁶⁾⁷⁾ For our patient, initially we recognized an anomalous origin of the LCX from the first diagonal branch by conventional CAG and finally the origin of the LCX could be precisely established by MDCT.

REFERENCES

- 1) Yamanaka O, Hobbs RE. *Coronary artery anomalies in 126,595 patients undergoing coronary arteriography.* *Cathet Cardiovasc Diagn* 1990;21:28-40.
- 2) Gavrielatos G, Letsas KP, Pappas LK, Antonellis I, Kardaras F. *Anom-*

Fig. 2. 64-slice multi-detector computed tomography demonstrates an anomalous origin of the LCX coursed to the left atrioventricular (AV) groove from the D1. LAD: left anterior descending coronary artery, LCX: left circumflex coronary artery, D1: first diagonal branch, LA: left atrium, LV: left ventricle.

alous origin of the entire coronary system with separate ostia within the right sinus of valsalva: a rare congenital anomaly and a review of the literature. *Cardiology* 2007;107:209-12.

- 3) Aydinlar A, Cicek D, Senturk T, et al. Primary congenital anomalies of the coronary arteries: a coronary arteriographic study in Western Turkey. *Int Heart J* 2005;46:97-103.
- 4) Wilkins CE, Betancourt B, Mathur VS, et al. Coronary artery anomalies: review of over 10,000 patients from the Clayton Cardiovascular Laboratories. *Tex Heart Inst J* 1988;15:166-73.
- 5) Lee YS, Lee JB, Kim KS. Anomalous origin of the left circumflex coro-

nary artery from the right sinus of valsalva identified by imaging with multidetector computed tomography. *Korean Circ J* 2006;36:823-5.

- 6) Korosoglou G, Ringwald G, Giannitsis E, Katus HA. Anomalous origin of the left circumflex coronary artery from the pulmonary artery: a very rare congenital anomaly in an adult patient diagnosed by cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2008;10:4.
- 7) Post JC, van Rossum AC, Bronzwaer JG, et al. Magnetic resonance angiography of anomalous coronary arteries: a new gold standard for delineating the proximal course. *Circulation* 1995;92:3163-71.