Published in final edited form as:

J Am Chem Soc. 2009 December 30; 131(51): 18206–18207. doi:10.1021/ja907271a.

Direct Observation of an Enamine Intermediate in Amine Catalysis

Xueyong Zhu, Fujie Tanaka, Richard A. Lerner*, Carlos F. Barbas III*, and Ian A. Wilson*

Departments of Chemistry and Molecular Biology and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

Abstract

An enamine intermediate is believed to be the central feature of biological catalysts, such as aldolases and small molecule amine organocatalysts. Despite decades of investigation of naturally occurring aldolase enzymes and recent studies on designed aldolase antibodies and organocatalysts, direct structural observation of an enamine intermediate has proven to be rare. Herein, we report observation of a stable enamine intermediate in the crystal structure of an aldolase antibody 33F12 in complex with a 1,3-diketone derivative. This enamine complex structure provides strong evidence that fewer residues are essential for amine catalysis within the hydrophobic environments of this catalytic antibody than speculated for natural aldolase enzymes, and should serve to guide future studies aimed at the rational design of these types of catalysts, as well as organocatalysts. Indeed, enamine catalysis in proteins might be more simplistic than previously imagined.

The advent of aldolase catalytic antibodies¹ and the translation of this chemistry to small organic molecules has led to the burgeoning field of organocatalysis.² The centerpiece of amine-catalyzed reactions is presumed to be an enamine, the nascent nucleophile in enantioselective C-C and C-X bond-forming reactions.^{1,2} In order to more fully understand amine catalysis via an enamine mechanism, we have sought to observe enamine intermediates of aldolase antibodies directly. Enamines, however, are typically unstable in water; equilibrium between formation and hydrolysis of an enamine makes direct observation of an enamine within protein crystals difficult.^{1,2} Aldolase antibodies were initially raised against a 1,3-diketone hapten under the assumption that its reaction with a nucleophilic lysine in the binding pocket would yield an enaminone or vinylogous amide, a stabilized enamine (Fig. 1A). Spectroscopic analysis of reactions of these aldolase antibodies with 1,3-diketones suggested that the enaminone, was, indeed, formed.^{1a-c} We now report direct observation of the enamine by crystallographic analysis of the adduct formed when aldolase antibody 33F12 reacts with a 1,3-diketone derivative **1** (Fig. 1A).

wilson@scripps.edu; foleyral@scripps.edu; carlos@scripps.edu.

Supporting Information Available: Experimental procedures, crystal structure determination, and structure of the Fab complex (Table S1; Figure S1). This material is available free of charge via the Internet at <http://pubs.acs.org>.

To obtain crystals of enzymes containing unstable enamine and related intermediates (i.e., carbinolamine and iminium), special conditions and procedures, such as soaking of crystals in a substrate solution and flash freezing, are typically required.³ To isolate unstable intermediates within crystals, additional interactions between substrate functional groups and amino acid residues of the catalyst (such as charge interactions between substrate phosphate groups and acidic residues) may be needed.³ However, such interactions involving substrates often complicate analysis of the essential features required for catalysis. In the analysis of enamine complexes with natural enzymes, the assignment of functional roles to amino acid residues involved in catalysis or in substrate binding has been difficult.³ Amino acid residues that interact with particular substrate functional groups may play less of a role in the catalytic machinery than in substrate binding. To understand the essentials of enamine catalysis, it may be necessary to investigate catalysts that have more limited interactions with their substrates. Due to its mechanism of elicitation, aldolase antibody 33F12 (and its sequence-related variant 38C2) acquired a promiscuous active site and, as a consequence, is able to catalyze aldol, retro-aldol, and enamine/iminium-based^{1b,d,h} transformations of a very wide range of substrates.¹ It was originally proposed that substrate interactions with the binding site of the catalyst would be based on hydrophobic interactions and covalent imine and subsequent enamine formation.¹ Therefore, aldolase antibodies such as 33F12 should serve as a simplified model system to observe the requirements for an effective aminocatalyst.

We co-crystallized 33F12 Fab' with 1,3-diketone **1** and determined its structure by molecular replacement to 1.9 Å resolution (Table S1). The Fab' complex resembles that of the native Fab'^{1c} with an overall r.m.s.d. of 1.0 Å (C $_{\alpha}$ atoms), and only 0.5 Å for the Fv domain (Fig. S1). The binding site pocket is a narrow, elongated cleft more than 11 Å deep with the reactive catalytic lysine, LysH93, at its base (Fig. 1B and 1C). LysH93 is surrounded by mostly hydrophobic residues that substantially lower its pK_a, a mechanism that appears now to be shared with the natural enzyme acetoacetate decarboxylase.^{1c,4} The LysH93-diketone covalent adduct is clearly present in the complex structure (Fig. 1B, 1C and S1), and conclusively demonstrates enamine formation. The enamione portion is well ordered in the electron density maps, but the rest of the 1,3-diketone is less well defined, consistent with the experimental observations that the antibody accepts a wide range of substrate through non-specific hydrophobic interactions.^{1c}

The ϵ -amine of LysH93 is covalently linked to the C2 atom of the 1,3-diketone and the C1, C2-N ϵ , C3, C4=O, and C5 atoms of the enamione moiety are all in the same plane (Fig. 1). Upon covalent adduct formation, no significant conformational changes occur within the antibody-binding site, except for a few side-chain rotamers, such for AsnL34 (Fig. 1B). The enamine moiety makes van der Waals interactions with TyrL36, SerL89, PheL98, TyrH95, and TrpH103, while the phenylamide substituent interacts with AsnL34, GlyL91, TrpH33, TyrH95, PheH96 and TyrH97 (Fig. 1B). Importantly, hydrophilic residues (AsnL34, SerL89, SerH100 and TyrL36) are in close proximity to the enamine moiety (Fig. 1B, 1D). The distances between the phenolic oxygen of TyrL36 and the enamine N ϵ (from LysH93) and C3 (from the diketone) are 4.2 and 4.0 Å, respectively, while the SerH100 hydroxyl is located 4.2 and 5.1 Å from the enamine N ϵ and C3, respectively. Additionally, the SerL89 hydroxyl is within 3.9 Å of the enamine C3.

Based on these structural observations, we hypothesize that TyrL36 acts as the general acid and base directly (Scheme 1), or indirectly through a water molecule, with assistance of SerH100, SerL89 and/or AsnL34 for the formation and hydrolysis of enamine and imine intermediates. Water-assisted catalysis has been proposed for aldolases and designed retro-aldol catalysts.⁵ For the aldol reaction of acetone and an aldehyde, as in Scheme 1, LysH93 first makes a nucleophilic attack on the carbonyl carbon of acetone to form a carbinolamine;

the phenolic OH of TyrL36 may act as an acid to activate the carbonyl. Next, water elimination from the carbinolamine results in formation of an iminium ion. TyrL36 may assist in the elimination of water by providing a charged environment. The iminium intermediate is then converted to an enamine, in which Tyr L36 could act as the base to abstract the α -proton of the iminium. For the C–C bond formation step, TyrL36 may then activate the carbonyl group of the aldehyde substrate through a hydrogen bond or proton transfer. Hydrolysis of the iminium affords the aldol product, where TyrL36 could again act as a base to form OH^- by abstraction of a proton from a water molecule in the hydrolysis step. Water molecules may also be involved in the catalysis depending on the actual distances between TyrL36 and the lysine ϵ -amino group during catalysis.^{5,6} The role of TyrL36 was further supported by the TyrL36Phe mutant of aldolase antibody 38C2 which did not bind to diketone **1** derivatives (unpublished).

Once the enamionone is formed, no additional interactions are required to retain the enamionone. TyrL36 does not need to provide direct interactions with the enamionone formed in the crystal structure. The distances between the TyrL36 and the enamionone moiety suggest its role in general acid/base catalysis either directly or through water. No amino acid residues other than TyrL36, SerL89, SerH100, and AsnL34 are observed in close enough proximity that could potentially provide charged interactions or hydrogen bonds during catalysis. Thus, this enamine complex structure provides strong evidence that fewer residues are essential for amine catalysis within the hydrophobic environments of this catalytic antibody than speculated for natural aldolase enzymes,³ and this structure should serve to guide future studies aimed at the rational design of these types of catalysts⁵ as well as organocatalysts². Indeed, enamine catalysis in proteins might be more simplistic than previously imagined.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the staff of the ALS beamline 821 for data collection and the support by NIH AI042266 (I.A.W.) and CA27489 (C.F.B., R.A.L. and I.A.W.) and the Skaggs Institute for Chemical Biology. Coordinates and structure factors for the 33F12 Fab' hapten complex have been deposited in the RCSB Protein Data Bank with accession code 3FO9.

References

1. (a) Wagner J, Lerner RA, Barbas CF III. *Science*. 1995; 270:179.(b) Bjornestedt R, Zhong G, Lerner RA, Barbas CF III. *J Am Chem Soc*. 1996; 118:11720.(c) Barbas CF III, Heine A, Zhong G, Hoffmann T, Gramatikova S, Bjornestedt R, List B, Anderson J, Stura EA, Wilson IA, Lerner RA. *Science*. 1997; 278:2085. [PubMed: 9405338] (d) Zhong G, Hoffmann T, Lerner RA, Danishefsky S, Barbas CF III. *J Am Chem Soc*. 1997; 119:8131.(e) Zhong G, Shabat D, List B, Anderson J, Sinha SC, Lerner RA, Barbas CF III. *Angew Chem, Int Ed*. 1998; 37:2481.(f) Sinha SC, Barbas CF III, Lerner RA. *Proc Natl Acad Sci USA*. 1998; 95:14603. [PubMed: 9843936] (g) Zhong G, Lerner RA, Barbas CF III. *Angew Chem Int Ed*. 1999; 38:3738.(h) Shabat D, Lode HN, Pertl U, Reisfeld RA, Rader C, Lerner RA, Barbas CF III. *Proc Natl Acad Sci, USA*. 2001; 98:7528. [PubMed: 11404472] (i) Tanaka F, Barbas CF III. *J Immunol Methods*. 2002; 269:67. [PubMed: 12379353] (j) Tanaka, F.; Barbas, CF, III. Antibody-catalyzed aldol reactions. In: Mahrwald, R., editor. *Modern Aldol Reactions*. Vol. 1. Wiley-VCH; 2004. p. 273-310.(k) Zhu X, Tanaka F, Hu Y, Heine A, Fuller R, Zhong G, Olson AJ, Lerner RA, Barbas CF III, Wilson IA. *J Mol Biol*. 2004; 343:1269. [PubMed: 15491612]
2. (a) Notz W, Tanaka F, Barbas CF III. *Acc Chem Res*. 2004; 37:580. [PubMed: 15311957] (b) Melchiorre P, Marigo M, Carlone A, Bartoli G. *Angew Chem Int Ed Engl*. 2008; 47:6138.

- [PubMed: 18666089] (c) Dondoni A, Massi A. *Angew Chem, Int Ed.* 2008; 47:4638.(d) Bertelsen S, Jørgensen KA. *Chem Soc Rev.* 2009; 38:2178. [PubMed: 19623342] (e) Barbas CF III. *Angew Chem, Int Ed.* 2008; 47:42.
3. (a) Heine A, DeSantis G, Luz JG, Mitchell M, Wong CH, Wilson IA. *Science.* 2001; 294:369. [PubMed: 11598300] (b) Allard J, Grochlski P, Sygusch J. *Proc Natl Acad Sci USA.* 2001; 98:3679. [PubMed: 11274385] (c) St-Jean M, Sygusch J. *J Biol Chem.* 2007; 282:31028–31037. [PubMed: 17728250]
4. Ho MC, Ménétret JF, Tsuruta H, Allen KN. *Nature.* 2009; 459:393. [PubMed: 19458715]
5. Jiang L, Althoff EA, Clemente FR, Doyle L, Rothlisberger D, Zanghellini A, Gallaher JL, Betker JL, Tanaka F, Barbas CF III, Hilvert D, Houk KN, Stoddard BL, Baker D. *Science.* 2008; 319:1387. [PubMed: 18323453]
6. Fullerton SW, Griffiths JS, Merkel AB, Cheriyan M, Wymer NJ, Hutchins MJ, Fierke CA, Toone EJ, Naismith JH. *Bioorg Med Chem.* 2006; 14:3002. [PubMed: 16403639]

Figure 1.

Enamine-forming reaction and the aldolase antibody binding site. (A) Reaction of antibody 33F12 with 1,3-diketone hapten **1**. A reactive lysine in 33F12 attacks one of the carbonyl groups of **1** to form carbinolamine **2** that subsequently collapses to iminium ion **3**. A stable covalent adduct, enaminone **4** is formed when the imine tautomerizes. (B) Binding site of the 33F12-hapten complex. The light chain and heavy chain backbone is represented as a tube in light and dark grey, respectively, with side chains of residues discussed in the text shown in stick form and labeled. (C) Shape complementarity of the enamine lysine-diketone adduct. A slice is shown through the antibody molecular surface and colored by electrostatic potential with 1.4 Å sphere radius. (D) 33F12-enamine interactions with some important binding site residues. Atom contact distances are given in Angstroms.

Scheme 1.
Plausible role of TyrL36 in general acid/base catalysis.