Findings that shed new light on the possible pathogenesis of a disease or an adverse effect

Fusobacterium necrophorum – beyond Lemierre's syndrome

Lokesh Shahani,¹ Nancy Khardori²

¹Department of Internal Medicine, Southern Illinois University School of Medicine, Springfield, Illinois, United States;

²Division of Infectious Disease, Department of Internal Medicine, SIU School of Medicine, Springfield, Illinois, United States

Correspondence to Dr Lokesh Shahani, lokesh83@hotmail.com

Summary

Fusobacterium necrophorum is a non-sporulating anaerobic gram negative bacillus and has traditionally been associated with Lemierre's syndrome. The authors report a 34-year-old male who presented to the emergency department with a week's history of dull epigastric pain. Significant medical history included chronic pancreatitis secondary to alcohol use. The patient had radiological evidence of acute on chronic pancreatitis with thrombosis of the portal vein and multiple intrahepatic abscesses. CT-guided drainage of left upper quadrant revealed fluid collection in the pancreatic bed. The fluid culture grew *F necrophorum* and the patient was treated with tigecycline for 4 weeks. The patient improved symptomatically and his follow-up computerised axial tomography scan 2 months later showed resolving liver abscess, cavernous transformation of the portal vein and stable findings of chronic pancreatitis. This could represent an infection of the peripancreatic tissue with *F necrophorum* further leading to pylephlebitis.

BACKGROUND

Fusobacterium necrophorum is an anaerobic gram negative bacillus forming a part of the family Bacteroidaceae. These bacteria are a predominant part of the oral flora. It has traditionally been associated with Lemierre's syndrome. Lemierre syndrome, also known as postanginal septicemia or necrobacillosis, is an uncommon but potentially life-threatening complication of acute pharyngo-tonsillitis. It is characterised by acute primary infection of the oropharynx causing secondary thrombophlebitis of the internal jugular vein and associated with subsequent septicemia and septic embolisation.¹ When pharyngitis due to *Fusobacterium* species occurs, the physical proximity of the vessels in the lateral pharyngeal space permits extension from the peritonsillar space to the internal jugular vein. This usually occurs in less than a week from the development of pharyngitis. Once invasion of the internal jugular vein is achieved, the resultant bacteremia triggers platelet aggregation and thrombus formation.² Thrombus formation and rapid bacterial growth result in a nidus for metastatic septic embolisation, commonly to the lungs and less commonly to the large joints.³

We report a patient with radiological evidence of acute on chronic pancreatitis with thrombosis of the portal vein and multiple intrahepatic abscesses. *F necrophorum* was isolated from the peripancreatic and perihepatic fluid. This could represent an infection of the pancreatic and the peripancreatic tissue with *F necrophorum* further leading to thrombophlebitis of the portal vein and subsequent septic embolisation. This could reflect pathology similar to the clinical features seen in Lemierre's syndrome.

CASE PRESENTATION

A 34-year-old Caucasian male presented to the emergency department with a week's history of dull epigastric pain. The pain had been increasing in intensity and was

discomforting to the patient. No aggravating or relieving factors were noted and there was no radiation of the pain. The patient had history of excessive alcohol use until 1 year prior to current presentation. He was diagnosed with chronic pancreatitis one year back and has had abdominal pain on and off since then. He also had steatorrhea suggestive of pancreatic exocrine dysfunction; however previous investigation had not shown any clinical significant vitamin deficiency. The current presentation was described as an exacerbation of his previous symptoms. There were no other associated symptoms and the patient was able to tolerate a liquid diet.

The patient was afebrile and haemodynamically stable. He was cachectic and pallor was noted. Respiratory and cardiovascular examinations were unremarkable. Epigastric tenderness was elicited and moderate abdominal distention without any fluid thrill was noted. Neurological examination including cranial nerve examination was unremarkable.

INVESTIGATIONS

Routine laboratory investigations showed low haemoglobin of 12.8 g/dl, elevated white blood cell count of 26 100 cells/mm³ with absolute neutrophil count of 25 300 cells/mm³. Platelet count was in the normal range as the electrolytes, renal function and liver function tests. Computerised axial tomography (CAT) scan of abdomen and pelvis showed acute pancreatitis superimposed upon underlying chronic calcified pancreatitis with fluid collection in the lesser sac, porta hepatis, perisplenic region and left subphrenic region consistent with pseudocysts. There were low density lesions in both lobes of the liver consistent with intrahepatic abscesses (figure 1). There was acute thrombosis of left and right portal vein branches. The spleen was enlarged with lesions consistent with splenic abscesses and there was occlusion of the splenic vein and

Figure 1 Computerised axial tomography scan of abdomen showing thrombosis of the portal vein along with hepatic and splenic abscesses.

superior mesenteric vein thrombosis with numerous collaterals throughout the upper abdomen. There was however no evidence of hepatic parenchymal changes suggestive of hepatic cirrhosis.

TREATMENT

The patient was initially started on intravenous vancomycin and meropenem. The infectious disease service was consulted and antibiotics were changed to intravenous tigecycline and meropenem. The blood cultures remained negative after a week of incubation. CT-guided drainage of left upper quadrant revealed fluid collection in the pancreatic bed. This was initially haemorrhagic and later purulent. The fluid culture grew *F necrophorum* on the third day. Considering contiguous infection from the pancreatic and peripancreatic tissue causing septic thrombophlebitis of the portal vein, other causes such as hypercoagulable state were not considered.

CAT scan of the neck and chest was specifically ordered to evaluate for the classical features of Lemierre's syndrome associated with *F necrophorum*. There was no evidence of the internal jugular vein. The patient further failed to give any preceding history suggestive of any oropharyngeal or respiratory infection. Lungs are the most common site of metastasis³; however the patient did not have any radiological evidence of any pulmonary involvement. Features of lateral pharyngeal space involvement leading

to Horner's syndrome or trapezius muscle involvement were not present.

OUTCOME AND FOLLOW-UP

Meropenem was discontinued and the patient was continued on tigecycline for 4 weeks. The patient was followed up in the clinic. He improved symptomatically and his follow-up CAT scan 2 months later showed resolving liver abscess, cavernous transformation of the portal vein and stable findings of chronic pancreatitis. Cavernous transformation of the portal vein is the development of periportal collaterals around the recanalising or occluded main portal vein which occurs secondary to chronic portal vein thrombosis. Numerous collateral vessels that vary in character from small to markedly enlarged and serpiginous end up functionally 'replacing' the portal vein.

DISCUSSION

Septic thrombophlebitis of the portal vein also known as pylephlebitis is usually a complication of any intra-abdominal or pelvic infection that occurs in the region drained by the portal venous system, especially diverticulitis or appendicitis. Contiguous spread from nearby intra-abdominal infection has also been described. The diagnosis of septic thrombophlebitis is made on culture data in light of radiological diagnosis of acute thrombosis of any vessel.⁴ Enteric gram negative bacilli and anaerobes, especially *Bacteroides*

fragilis are the most common organisms associated with pylephlebitis.⁴ There have been some isolated cases of pylephlebitis associated with *Fusobacterium* species.^{5–9} Most of these patients had an identified gastrointestinal source of their portal vein infection. Liver abscesses and bowel ischemia are acute complications of pylephlebitis. Portal hypertension could be a long-term complication.

We report a patient with radiological evidence of acute on chronic pancreatitis with thrombosis of the portal vein and multiple intrahepatic abscesses. *F necrophorum* was the only organism isolated from the peripancreatic and perihepatic fluid. This could represent an infection of the pancreatic and the peripancreatic tissue with *F necrophorum* and contagious spread leading to pylephlebitis. Hepatic and splenic abscess could represent metastatic septic emboli from the infection.

Learning points

- ▶ *F necrophorum* is an anaerobic gram negative bacillus forming a part of the family Bacteroidaceae. These bacteria are a predominant part of the oral flora.
- ▶ Lemierre syndrome is characterised by acute primary infection of the oropharynx causing secondary thrombophlebitis of the internal jugular vein and associated with subsequent septicemia and septic embolisation.
- ▶ When pharyngitis due to *Fusobacterium* species occurs, the physical proximity of the vessels in the lateral pharyngeal space permits extension from the peritonsillar space to the internal jugular vein; and the resultant bacteremia triggers platelet aggregation and thrombus formation.
- ▶ Septic thrombophlebitis of the portal vein also known as pylephlebitis is usually a complication of any intra-abdominal or pelvic infection that occurs in the region drained by the portal venous system, however contiguous spread from nearby intra-abdominal infection has also been described.
- ▶ Liver abscesses and bowel ischemia are acute complications of pylephlebitis. Portal hypertension could be a long-term complication.

Competing interests None.

Patient consent Obtained.

REFERENCES

1. **Lemierre A.** On certain septicemias due to anaerobic organisms. *Lancet* 1936;**1**:701–3.
2. **Forrester LJ,** Campbell BJ, Berg JN, *et al.* Aggregation of platelets by *Fusobacterium necrophorum*. *J Clin Microbiol* 1985;**22**:245–9.
3. **Chirinos JA,** Lichtstein DM, Garcia J, *et al.* The evolution of Lemierre syndrome: report of 2 cases and review of the literature. *Medicine (Baltimore)* 2002;**81**:458–65.
4. **Plemmons RM,** Dooley DP, Longfield RN. Septic thrombophlebitis of the portal vein (pylephlebitis): diagnosis and management in the modern era. *Clin Infect Dis* 1995;**21**:1114–20.
5. **El Braks R,** Harnois F, Boutros N, *et al.* Mesenteric adenitis and portal vein thrombosis due to *Fusobacterium nucleatum*. *Eur J Gastroenterol Hepatol* 2004;**16**:1063–6.
6. **Etienne M,** Gueit I, Abboud P, *et al.* *Fusobacterium nucleatum* hepatic abscess with pylephlebitis associated with idiopathic CD4(+) T lymphocytopenia. *Clin Infect Dis* 2001;**32**:326–8.
7. **Bultink IE,** Dorigo-Zetsma JW, Koopman MG, *et al.* *Fusobacterium nucleatum* septicemia and portal vein thrombosis. *Clin Infect Dis* 1999;**28**:1325–6.
8. **Soo R,** Gosbell I, Gallo J, *et al.* Septic portal vein thrombosis due to *Fusobacterium necrophorum*. *Aust N Z J Med* 1999;**29**:569–70.
9. **Verna EC,** Larghi A, Faddoul SG, *et al.* Portal vein thrombosis associated with *Fusobacterium nucleatum* septicemia in a patient with ulcerative colitis. *J Clin Gastroenterol* 2004;**38**:611–12.

This pdf has been created automatically from the final edited text and images.

Copyright 2011 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Please cite this article as follows (you will need to access the article online to obtain the date of publication).

Shahani L, Khadori N. *Fusobacterium necrophorum* – beyond Lemierres syndrome. *BMJ Case Reports* 2011;**10**:1136/bcr.07.2011.4527, Published XXX

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow