Published in final edited form as:

Arch Med Res. 2011 August ; 42(6): 523–531. doi:10.1016/j.arcmed.2011.10.010.

Association Study of Candidate Gene Polymorphisms and Obesity in a Young Mexican-American Population from South Texas

Jorge Duran-Gonzalez^a, Ixiu Ortiz^a, Enrique Gonzales^a, Nicole Ruiz^a, Manti Ortiz^a, Arthur Gonzalez^a, Edna K. Sanchez^a, Eugenia Curet^d, Susan Fisher-Hoch^c, Anne Rentfro^b, Huiqi Qu^c, and Saraswathy Nair^a

^aDepartment of Biological Sciences and Center of Biomedical Studies, University of Texas School of Public Health, Brownsville, Texas

^bSchool of Nursing and Center of Biomedical Studies, University of Texas School of Public Health, Brownsville, Texas

^cDivision of Epidemiology, University of Texas School of Public Health, Brownsville, Texas

^dStudent Health Services, University of Texas at Brownsville, Brownsville, Texas

Abstract

Background and Aims—Obesity is increasingly a health problem and a risk factor for diabetes in young Mexican-American populations. Genetic association studies in older, mostly non-Hispanic populations have reported that polymorphisms in the candidate genes *HSD11B1*, *CRP*, *ADIPOQ*, *PPARG*, *ANKK1*, *ABCC8* and *SERPINF1* are associated with obesity or diabetes. We analyzed the polymorphisms rs846910, rs1205, rs1501299, rs1801282, rs1800497, rs757110 and rs1136287 in these candidate genes, for association with obesity and metabolic traits in a young Mexican-American population from south Texas.

Methods—Genotyping of the seven common SNPs were performed by allelic discrimination assays in 448 unrelated Mexican Americans (median age = 16 years) from south Texas. χ^2 tests and regression analyses using additive models were used for genetic association analyses adjusting for covariates; *p* values were corrected for multiple testing by permutation analyses.

Results—rs1800497 (*ANKK1*) shows association with waist circumference (*p* = 0.009) and retains the association (*p* = 0.03) after permutation testing. Analysis of metabolic quantitative traits shows that rs846910 (*HSD11B1*) was associated with HOMA-IR (*p* = 0.04) and triglycerides (*p* = 0.03), and rs1205 (*CRP*) with HOMA-IR (*p* = 0.03) and fasting glucose levels (*p* = 0.007). However, the quantitative traits associations are not maintained after permutation analysis. None of the other SNPs in this study showed associations with obesity or metabolic traits in this young Mexican-American population.

© 2011 Instituto Mexicano del Seguro Social (IMSS). Published by Elsevier Inc. All rights reserved.

Address reprint requests to: Saraswathy Nair, 80 Fort Brown, Brownsville, TX, 78520: Phone: 956-882-5108; FAX 956-882-5043; Saraswathy.nair@utb.edu.

The authors state that there are no conflicts of interest to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusions—We report a potential association between rs1800497 (linked to changes in brain dopamine receptor levels) and central obesity in a young Mexican-American population.

Keywords

Obesity; Diabetes; Polymorphism; Mexican-American; Genetic association

Introduction

Obesity and overweight represent health problems that are risk factors for the development of other chronic conditions such as diabetes, cancer and cardiovascular disease (CVD) (1). According to the information provided by the National Health and Nutrition Examination Survey (NHANES) in the U.S., the prevalence of obesity in persons >20 years is 33.9%. However, this number increases to 39.3% in the Mexican-American population (2). It has also been suggested that Mexican immigrants have a 64% higher chance to develop the obese phenotype than US-born whites (3). In the southernmost region of Texas, in the Rio Grande Valley (RGV), a recent socioeconomic study shows that prevalence of obesity in an older Mexican-American population ranges from 55.5–57%, the highest rate observed nationwide (4). In younger Mexican Americans, the increasing rates of obesity, metabolic syndrome (MetS) or type 2 diabetes (T2D) are disproportionately higher than in other populations (5,6). In a study of adolescent Mexican-Americans in RGV, 50% are either overweight or obese, 41% have central obesity and 27% exhibited insulin resistance (7). Environmental factors such as reduced physical activity, intake of energy-dense food and/or cultural and socioeconomic factors play an important role in the development of obesity (8). However, evidence from recent studies suggests that genetic factors account for 40–90% of the body mass index (BMI) variations among populations (9). The future of medicine, whether it is prevention, diagnosis, and therapeutic care for human diseases includes personalized genomics. This involves the identification of genetic factors that confer risk of susceptibility to obesity and related metabolic abnormalities in individuals. More than 127 biologically relevant candidate genes with DNA variations associated with obesity-related phenotypes have been identified (10). In the present study we analyzed the association of seven single nucleotide polymorphisms (SNPs), rs846910, rs1205, rs1501299, rs1801282, rs1800497, rs757110 and rs1136287, with obesity and related metabolic traits in a young Mexican-American cohort of 448 non-related individuals. These seven SNPs were chosen because they have been shown to be associated with obesity or obesity-related phenotypes in previous studies or are located in well-established obesity candidate genes (Table 1). The chosen candidate genes are *HSD11B1*, *CRP*, *ADIPOQ*, *PPARG*, *ANKK1*, *ABCC8* and *SERPINF1*. Individuals from this cohort include previously recruited RGV Mexican-American adolescents (7) as well as newly recruited Mexican-American college students. None were from the older Mexican-American Cameron County Hispanic Cohort of RGV (4).

Subjects and Methods

Subjects and Clinical Measurements

A total of 448 volunteers from the Brownsville population (southern Texas) consented to participate. All subjects (and parents for subjects <18 years) provided informed consent prior to participation in the study. Consent was specifically obtained to save identified DNA samples for genetic studies. All studies were approved by the IRBs (UTHSC-Houston and UTB). This study was conducted under the basic principles of the Declaration of Helsinki. Inclusion criteria include Mexican-American etiology. First-degree familiars were excluded from statistical analysis.

Fasting blood samples were drawn from all subjects into EDTA-coated tubes and red blood cells separated by centrifugation. The supernatant comprising plasma and the buffy coat layer including white blood cells were aspirated separately and frozen immediately in aliquots at -70°C .

Anthropometric measurements including height, weight, and waist circumference (WC) were done according to standardized protocols. WC cut-offs for central obesity were calculated using the parameters proposed by Taylor (11) for individuals <20 years. For subjects 21 years and older, the values were >88 cm for women and >102 cm for men (12). BMI was calculated according to the formula: weight in kilograms (kg)/height in meters (m^2). For subjects <20 years of age, calculations were made converting to age- and gender-standardized percentiles based on the U.S. Centers for Disease Control and Prevention 2000 growth charts (<http://www.cdc.gov/growthcharts/>), which are not race specific. Results were classified as normal weight if BMI was <85 th percentile, overweight if BMI was ≥ 85 th and <95 th percentile, or obese if BMI was ≥ 95 th percentile. For subjects 20 years of age and older, BMI $25\text{--}29.9$ kg/m^2 was classified as overweight and BMI ≥ 30 kg/m^2 as obese (12).

Metabolic measurements include fasting blood glucose, blood pressure, homeostasis model of insulin resistance (HOMA-IR) and triglycerides. HOMA-IR was calculated by the formula (fasting insulin [$\mu\text{U}/\text{ml}$] $\times$ fasting glucose [mmol/l]/22.5) and classified as insulin resistant if the result was ≥ 3.16 (13).

DNA Extraction and Genotyping

DNA was extracted from 1 ml of blood sample using a QIAmp DNA Mini Kit (Qiagen, Valencia, CA) according to the manufacturer's protocol. DNA quantification was carried out using a Nanodrop spectrophotometer (Thermo Scientific, Waltham, MA) and the amount of recovered DNA was determined using 96-well plates prepared with 50 ng per well of DNA. These plates were left to dry overnight and then stored at -20°C until used. TaqMan SNP Genotyping Assays were used for genotyping in the Step One Plus Real-Time PCR system (Applied Biosystems, Carlsbad, CA). Five- μl PCR reactions were performed using 1 μl DNA (50 ng/ μl) or ddH₂O for negative controls, 0.125 μl probe (20 $\mu\text{mol}/\text{L}$), and 2.25 μl Master Mix (20 $\mu\text{mol}/\text{l}$) (Applied Biosystems). The program consisted of an initial denaturation step at 95°C for 10 min followed by 40 cycles of 95°C for 15 sec, 60°C for 1 min and a final extension step of 60°C for 30 sec.

Statistical Analysis

Data are presented as mean and $\pm$ standard deviation (SD). Genotype distributions of the seven SNPs were analyzed for deviation from Hardy-Weinberg equilibrium using χ^2 test. Logistic regression analysis assuming an additive model was performed to determine the association between SNPs and BMI as an obesity measure and WC as central obesity value. The associations were adjusted for gender and age as covariates. Odds ratios (ORs) with 95% confidence intervals (CIs) are presented in the tables with respect to the risk alleles; p value of 0.05 was considered significant and empirical significance levels were computed by permutation test. Linear regression analysis was used for quantitative traits under an additive genetic model adjusted for age, gender and BMI. All statistical analysis were performed using PLINK v.1.07 (<http://pngu.mgh.harvard.edu/~purcell/plink>) (14), and SPSS v.17.0 (SPSS, Chicago, IL).

Quanto Software (v.1.2.4) (<http://hydra.usc.edu/gxe/>) was used to calculate the power of the sample. A 39% prevalence of obesity among Mexican-Americans (2), additive genetic model for any SNP and a MAF of at least 30% shows that the sample size of the present study had an 80% power with an $\alpha = 0.05$ to detect an effect size of 1.47.

Results

Clinical and anthropometric characteristics are shown in Table 2. The median age of the cohort was 16 years and was composed predominately of female subjects (65%). Based on median value of BMI, the male population in our study is slightly overweight (BMI 25.4). According to BMI values, 28.3% of the sample population is obese, and 21.3% are overweight. However, a total of 53.1% of the subjects have WC measurements correlated with central obesity and considered a risk factor for MetS.

Distributions of the genotype and allele frequencies of the seven SNPs were in accordance with Hardy-Weinberg equilibrium using χ^2 test (Table 3). The *ANKK1* SNP rs1800497 (G/A) shows a higher frequency of the (A) allele in subjects with obese BMI values than in those with non-obese BMI values ($p = 0.04$) as also when comparing subjects with central obesity vs. subjects with no central obesity ($p = 0.009$) and after adjusting for age and gender (Table 4). After using permutation test to empirically derive significance corrected for multiple testing, only association with WC values of central obesity remained significant ($p = 0.03$). There were no trends of association found for the remaining six SNPs with BMI or WC.

Analysis of the quantitative metabolic traits shows statistical significance for the association between the *HSD11B1* SNP rs846910 with HOMA-IR and triglyceride values ($p = 0.04$ and $p = 0.03$, respectively). For the *CRP* SNP rs1205, a statistical significance was also observed with HOMA-IR and fasting plasma glucose levels ($p = 0.03$ and $p = 0.007$, respectively) (Table 5). Nevertheless, after permutation testing, none of the values of associations with quantitative traits remained statistically significant. No associations were observed for the other SNPs.

Discussion and Conclusions

Although 21.3% of the young Mexican-American population evaluated in this study is overweight, only 28.3% are obese as defined by BMI. However, when WC is considered, 53% of the subjects are classified as being centrally obese. These results, as well as results from an earlier study, suggest that WC measurement may be a more effective measurement in young Mexican-Americans to assess obesity and metabolic risk than BMI (7). Rankinen et al. (2005) summarized obesity and/or obesity-related phenotype associations of various SNPs in the candidate genes in this study, in multiple populations (15). We summarized in Table 1 (16–30) previously described associations between obesity and obesity-related phenotypes and the selected SNPs in this study. To date, limited or no information exists in the literature regarding the associations of rs846910, rs1205, rs757110 and rs1136287 located in *HSD11B1*, *CRP*, *ABCC8* and *SERPINF1*, respectively, with obesity in the Mexican-American population.

The *TaqIA* polymorphism (A1 allele) originally identified in the dopamine receptor (*DRD2*) has been associated with lower levels of dopamine receptors (31–35). The *TaqI* A1 allele is associated with obesity as well as eating disorders and addictive behavior in heterogeneous analyzed groups (27–30,32,36,37). It has been reported that dysfunction in the dopamine signaling pathway (involved in food reward) in the brain may contribute to obesity (32). Recently, it was reported that *TaqIA* (A1 and A2 alleles) polymorphism, now referred to as rs18000497, is located 9489 bp downstream from the 3' end of *DRD2* in the last exon of the adjacent gene ankyrin repeat and kinase domain containing 1 (*ANKK1*) transcribed in the opposite direction (38). Although this SNP has been shown to affect *DRD2* expression, we thus refer to rs1800497 as a SNP in *ANKK1*. The rs1800497 (A) allele corresponds to the *TaqI* (A1) allele and the rs18700497 (G) allele corresponds to the *TaqI* (A2) allele.

In this study, an association was observed with the (A) allele of rs1800497 and central obesity in a young Mexican-American population. Considering the genetic admixture of this Mexican population, further structured association studies will be needed to validate this potentially novel genetic association. From a biological perspective, the dopamine neurotransmitter is important in food intake regulation via the mesolimbic reward circuit of the brain (32) and higher BMIs are associated with a lesser number of brain dopamine receptors (37). However, despite the plausible biological explanation, association between obesity and the rs1800497 SNP is not universal, and these results have not been replicated in Pima Indian, Nauruan and Australian populations (39,40). The association of this polymorphism in obese subjects and binge eaters with high sensitivity to food reward in a mixed Canadian population is also not fully understood (41). Among risk factors, glucocorticoid excess is a well-known cause of obesity (42). Recent evidence suggests that even if circulating levels of glucocorticoids are not elevated, excessive production of glucocorticoids in peripheral tissues is associated with obesity and insulin resistance (43,44). The enzyme 11 beta hydroxysteroid dehydrogenase Type 1 (*HSD11B1*), which converts the inactive cortisone into the active cortisol, is expressed in insulin-sensitive tissues such as adipose tissue, muscle, liver and central nervous system (45). Elevated levels of *HSD11B1* expression in adipose tissue have been shown to be associated with obesity, T2D, and changes in insulin sensitivity in several populations (43,46). Allelic variants and mRNA levels of *HSD11B1* have been previously associated with obesity, insulin sensitivity, hypertension and T2D (26,47,48).

In the present study we observed a trend towards association between the (A) allele of rs846910 (*HSD11B1*) and HOMA-IR and triglycerides values, which are predictive measures for MetS. In Pima Indians, the (A) allele is associated with increased insulin resistance as well as T2D. Because Mexican-Americans have Native American heritage, the observed trend of association suggests the possibility of shared genetic susceptibility to diabetes through variations in *HSD11B1*.

C-reactive protein (CRP) is an important inflammatory marker, and circulating levels of this protein may have a prognostic value in identifying subjects with increased risk of developing T2D, obesity and or insulin resistance in Caucasian, African-American and Peruvian populations (49,50). It has also been shown that CRP levels may be modified by mutations in the *CRP* gene (51). Although we observed an association between the (T) allele of rs1205 with HOMA-IR and fasting glucose, the statistical significance did not remain after correction for multiple testing.

One of the most studied obesity and obesity-related phenotype candidate genes is the peroxisome proliferator-activated receptor- γ 2 (*PPARG*) because it plays a role as the master transcriptional regulator for adipocyte differentiation and metabolism genes. The rs1801282 SNP (also known as Pro12Ala) is commonly associated with obesity and diabetes in several populations, ethnic groups and meta-analysis studies (15,52). Previous reports of association between the polymorphism rs1801282 (Pro12Ala) of the *PPARG* gene and obesity in the Mexican-American (17,19) or Mexican-Mestizo populations (16) could not be replicated in this study. However, the minor allele (G) frequencies were similar between this cohort and other Caucasian and Hispanic cohorts (10–13.5%) (19,53–55). The most important differences from the previous reports (Table 1) and this study are the ages of the subjects. In this cohort the median age was 16 years, a relatively young and healthy population when compared to the other reports with median age values from 38.7–58.6 years. Nevertheless, a future study in this Mexican-American population including older participants could verify the previously reported associations of *PPARG* SNPs with obesity. Furthermore, other studies show only association between the *PPARG* SNP and obesity in a gender-specific manner in Mediterranean (Greek and Tunisian) populations (56,57), which we did not

observe in this study. A novel HuGE meta-analysis shows that *PPAR*rs1801282 was associated with a reduction in T2D risk; however, they also report a moderate level of heterogeneity, which can lead to variations among ethnic groups (52). A few studies show associations not with T2D, but with metabolic changes associated with MetS (58,59). However, we did not observe associations with metabolic traits in this study.

The adiponectin gene (*ADIPOQ*) encodes a major adipokine secreted from white adipose tissue and plays an important role in energy metabolism and immune response. Associations of polymorphisms in this gene with obesity and insulin resistance have been investigated in Caucasian and Hispanic populations (21,60,61). The pancreatic β -cell K_{ATP} channel is composed of two essential subunits: the Kir6.2 and the sulfonylurea receptor 1 (SUR1) and are encoded by the two adjacent genes, the potassium inwardly-rectifying channel, subfamily J, member 11 (*KCNJ11*) and the ATP-binding cassette, subfamily C member 8 (*ABCC8*), respectively (62). This channel plays a central role in glucose-stimulated insulin secretion from pancreatic beta cells, and meta-analysis studies show that polymorphisms in *KCNJ11* have been associated with T2D (63). Polymorphisms in *ABCC8* have been correlated with hyperinsulinemic hypoglycemia and T2D in Japanese and Caucasian populations (64,65). The serpin peptidase inhibitor, clade F, member 1 (*SERPINF1*), also known as anti-plasmin pigment epithelium derived factor (PEDF) is an adipokine secreted by adipocyte cells with levels comparable to that of adiponectin. Although, at first, it was isolated as a neurotrophic factor capable of converting retinoblastoma tumor cells into differentiated nonproliferative neurons, novel studies showed that expression of *SERPINF1* in adipocyte cells correlates with obesity and insulin resistance in murine models (66,67). It is also highly expressed in hepatocytes, and accumulation of triglycerides in PEDF null mouse hepatocytes has been reported and postulated as one of the factors underlying fatty liver disease in MetS and T2D (67). Serum PEDF levels have been positively associated with adiposity in diabetic Japanese and obese Caucasian subjects (68,69). The *ADIPOQ* SNP rs1501299 has been analyzed previously in Mexican-American population by Richardson (21), but no association was found with obesity, T2D and HOMA-IR. Our study confirms this finding. rs757110 (Ala1369Ser) in *ABCC8* is associated with T2D in Caucasian populations (65). However, other SNPs in *ABCC8* in large-scale association studies in Caucasian subjects from the UK show no association with diabetes (63). Thus, the lack of associations with obesity and metabolic traits in this study and previous reports for rs1501299 (*ADIPOQ*) (21,70) and rs757110 (*ABCC8*) are consistent. Replication in a larger cohort size may be required to rule out any small-to-moderate effects of these variants on obesity and T2D risk in the Mexican-American population. Although we hypothesized that the potential candidate gene *SERPINF1* (PEDF) may have genetic associations with obesity or metabolic traits based on biological function, we did not observe any such associations in this cohort.

Limitations in this study are the relatively small cohort size and lack of gold standard measurements of obesity and insulin resistance such as body fat measurements and euglycemic-hyperinsulinemic clamp measurements. The age of the subjects may possibly be another limitation of this study because we sampled a significantly younger, healthier population, whereas the onset of T2D is typically at older ages. On the other hand, such a cohort may be useful to identify inherent genetic risks for obesity or diabetes before environmental influences play a significant role in the onset of these diseases. In summary, we report the lack of genetic association between rs846910 (*HSD11B1*), rs1205 (*CRP*), rs1501299 (*ADIPOQ*), rs1801282 (*PPARG*), rs757110 (*ABCC8*), and rs1136287 (*SERPINF1*) and obesity and metabolic quantitative traits in a young Mexican American population from south Texas. Lack of associations with these SNPs does not necessarily rule out the six genes as potential obesity candidate genes. We report, for the first time, a possible association between rs1800497 and central obesity in this population.

Acknowledgments

This work was supported by the intramural Borderplex Council Diabetes/Obesity Research funds awarded to S. Nair. The undergraduate students I. Ortiz and E.K. Sanchez were supported as MBRS RISE scholars by the National Institutes of Health (NIH) 5R25 GM065925-7. Dr. A. Rentfro's work on the collection of the adolescent clinical data was supported by NIH National Center for Minority Health Disparities P20 MD000170-05.

References

1. Kopelman P. Health risks associated with overweight and obesity. *Obesity Rev.* 2007; 8:13–17.
2. Flegal KM, Carroll MD, Ogden CL, et al. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA.* 2010; 303:235–241. [PubMed: 20071471]
3. Singh GK, Kogan MD, van Dyck PC. Changes in state-specific childhood obesity and overweight prevalence in the United States from 2003 to 2007. *Arch Pediatr Adol Med.* 2010; 164:598–607.
4. Fisher-Hoch SP, Rentfro AR, Salinas JJ, et al. Socioeconomic status and prevalence of obesity and diabetes in a Mexican American community, Cameron County, Texas, 2004–2007. *Prev Chronic Dis.* 2010; 7:A53. [PubMed: 20394692]
5. Lawrence JM, Mayer-Davis EJ, Reynolds K, et al. Diabetes in Hispanic American youth: prevalence, incidence, demographics, and clinical characteristics: the SEARCH for Diabetes in Youth Study. *Diabetes Care.* 2009; 32(suppl 2):S123–S132. [PubMed: 19246577]
6. Duncan GE. Prevalence of diabetes and impaired fasting glucose levels among US adolescents: National Health and Nutrition Examination Survey, 1999–2002. *Arch Pediatr Adolesc Med.* 2006; 160:523–528. [PubMed: 16651496]
7. Rentfro AR, Nino JC, Pones RM, et al. Adiposity, biological markers of disease, and insulin resistance in Mexican American adolescents, 2004–2005. *Prev Chronic Dis.* 2011; 8:A40. [PubMed: 21324254]
8. Bayles B. Perceptions of childhood obesity on the Texas-Mexico border. *Public Health Nursing.* 2010; 27:320–328. [PubMed: 20626832]
9. Fawcett KA, Barroso I. The genetics of obesity: FTO leads the way. *Trends Genet.* 2010; 26:266–274. [PubMed: 20381893]
10. Yang WJ, Kelly T, He J. Genetic epidemiology of obesity. *Epidemiol Rev.* 2007; 29:49–61. [PubMed: 17566051]
11. Taylor RW, Jones IE, Williams SM, et al. Evaluation of waist circumference, waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by dual-energy X-ray absorptiometry, in children aged 3–19 y. *Am J Clin Nutr.* 2000; 72:490–495. [PubMed: 10919946]
12. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: executive summary. Expert Panel on the Identification, Evaluation, and Treatment of Overweight in Adults. *Am J Clin Nutr.* 1998; 68:899–917. [PubMed: 9771869]
13. Keskin M, Kurtoglu S, Kendirci M, et al. Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics.* 2005; 115:e500–e503. [PubMed: 15741351]
14. Purcell S, Cherny SS, Sham PC. Genetic power calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics.* 2003; 19:149–150. [PubMed: 12499305]
15. Rankinen T, Zuberi A, Chagnon YC, et al. The human obesity gene map: the 2005 update. *Obesity.* 2006; 14:529–644. [PubMed: 16741264]
16. Canizales-Quinteros S, Aguilar-Salinas CA, Ortiz-Lopez MG, et al. Association of PPARG2 Pro12Ala variant with larger body mass index in Mestizo and Amerindian populations of Mexico. *Human Biol.* 2007; 79:111–119. [PubMed: 17985660]
17. Cole SA, Mitchell BD, Hsueh WC, et al. The Pro12Ala variant of peroxisome proliferator-activated receptor-gamma 2 (PPAR-gamma 2) is associated with measures of obesity in Mexican Americans. *Int J Obesity.* 2000; 24:522–524.

18. Cruz C, Camarena B, Mejia JM, et al. The dopamine D2 receptor gene TaqI A1 polymorphism and alcoholism in a Mexican population. *Arch Med Res.* 1995; 26:421–426. [PubMed: 8555738]
19. Hsueh WC, Cole SA, Shuldiner AR, et al. Interactions between variants in the beta(3)-adrenergic receptor and peroxisome proliferator-activated receptor-gamma 2 genes and obesity. *Diabetes Care.* 2001; 24:672–677. [PubMed: 11315829]
20. Konishi T, Calvillo M, Leng AS, et al. Polymorphisms of the dopamine D-2 receptor, serotonin transporter, and GABA(A) receptor beta(3) subunit genes and alcoholism in Mexican-Americans. *Alcohol.* 2004; 32:45–52. [PubMed: 15066703]
21. Richardson DK, Schneider J, Fourcaudot MJ, et al. Association between variants in the genes for adiponectin and its receptors with insulin resistance syndrome (IRS)-related phenotypes in Mexican Americans. *Diabetologia.* 2006; 49:2317–2328. [PubMed: 16955209]
22. Iizuka H, Awata T, Osaki M, et al. Promoter polymorphisms of the pigment epithelium-derived factor gene are associated with diabetic retinopathy. *Biochem Biophys Res Commun.* 2007; 361:421–426. [PubMed: 17658465]
23. Siitonen N, Pulkkinen L, Lindstrom J, et al. Association of ADIPOQ gene variants with body weight, type 2 diabetes and serum adiponectin concentrations: the Finnish Diabetes Prevention Study. *BMC Med Genet.* 2011; 12:5. [PubMed: 21219602]
24. Eiriksdottir G, Smith AV, Aspelund T, et al. The interaction of adiposity with the CRP gene affects CRP levels: age, gene/environment susceptibility—Reykjavik study. *Int J Obes (Lond).* 2009; 33:267–272. [PubMed: 19139754]
25. Elbein SC, Sun J, Scroggin E, et al. Role of common sequence variants in insulin secretion in familial type 2 diabetic kindreds: the sulfonylurea receptor, glucokinase, and hepatocyte nuclear factor 1alpha genes. *Diabetes Care.* 2001; 24:472–478. [PubMed: 11289470]
26. Nair S, Lee YH, Lindsay RS, et al. 11 beta-Hydroxysteroid dehydrogenase Type 1: genetic polymorphisms are associated with type 2 diabetes in Pima Indians independently of obesity and expression in adipocyte and muscle. *Diabetologia.* 2004; 47:1088–1095. [PubMed: 15156315]
27. Blum K, Sheridan PJ, Wood RC, et al. Dopamine D2 receptor gene variants: association and linkage studies in impulsive-addictive-compulsive behaviour. *Pharmacogenetics.* 1995; 5:121–141. [PubMed: 7550364]
28. Comings DE, Flanagan SD, Dietz G, et al. The dopamine D2 receptor (DRD2) as a major gene in obesity and height. *Biochem Med Metab Biol.* 1993; 50:176–185. [PubMed: 8260195]
29. Noble EP, Noble RE, Ritchie T, et al. D2 dopamine receptor gene and obesity. *Int J Eat Disord.* 1994; 15:205–217. [PubMed: 8199600]
30. Thomas GN, Critchley JA, Tomlinson B, et al. Relationships between the taqI polymorphism of the dopamine D2 receptor and blood pressure in hyperglycaemic and normoglycaemic Chinese subjects. *Clin Endocrinol (Oxf).* 2001; 55:605–611. [PubMed: 11894971]
31. Lucht M, Samochowiec A, Samochowiec J, et al. Influence of DRD2 and ANKK1 genotypes on apomorphine-induced growth hormone (GH) response in alcohol-dependent patients. *Prog Neuropsychopharmacol Biol Psychiatry.* 2010; 34:45–49. [PubMed: 19796663]
32. Stice E, Spoor S, Bohon C, et al. Relation between obesity and blunted striatal response to food is moderated by TaqIA A1 allele. *Science.* 2008; 322:449–452. [PubMed: 18927395]
33. Ritchie T, Noble EP. Association of seven polymorphisms of the D2 dopamine receptor gene with brain receptor-binding characteristics. *Neurochem Res.* 2003; 28:73–82. [PubMed: 12587665]
34. Jonsson EG, Nothen MM, Grunhage F, et al. Polymorphisms in the dopamine D2 receptor gene and their relationships to striatal dopamine receptor density of healthy volunteers. *Mol Psychiatry.* 1999; 4:290–296. [PubMed: 10395223]
35. Blum K, Braverman ER, Wood RC, et al. Increased prevalence of the Taq I A1 allele of the dopamine receptor gene (DRD2) in obesity with comorbid substance use disorder: a preliminary report. *Pharmacogenetics.* 1996; 6:297–305. [PubMed: 8873216]
36. Davis C, Levitan RD, Kaplan AS, et al. Dopamine transporter gene (DAT1) associated with appetite suppression to methylphenidate in a case-control study of binge eating disorder. *Neuropsychopharmacology.* 2007; 32:2199–2206. [PubMed: 17314918]
37. Wang GJ, Volkow ND, Logan J, et al. Brain dopamine and obesity. *Lancet.* 2001; 357:354–357. [PubMed: 11210998]

38. Neville MJ, Johnstone EC, Walton RT. Identification and characterization of ANKK1: a novel kinase gene closely linked to DRD2 on chromosome band 11q23.1. *Human Mutat.* 2004; 23:540–545.
39. Jenkinson CP, Hanson R, Cray K, et al. Association of dopamine D2 receptor polymorphisms Ser311Cys and TaqIA with obesity or type 2 diabetes mellitus in Pima Indians. *Int J Obes Relat Metab Disord.* 2000; 24:1233–1238. [PubMed: 11093282]
40. Southon A, Walder K, Sanigorski AM, et al. The Taq IA and Ser311 Cys polymorphisms in the dopamine D2 receptor gene and obesity. *Diabetes Nutr Metab.* 2003; 16:72–76. [PubMed: 12848308]
41. Davis C, Levitan RD, Kaplan AS, et al. Reward sensitivity and the D2 dopamine receptor gene: a case-control study of binge eating disorder. *Prog Neuropsychopharmacol Biol Psychiatry.* 2008; 32:620–628. [PubMed: 18262320]
42. Feldman BJ. Glucocorticoids influence on mesenchymal stem cells and implications for metabolic disease. *Pediatr Res.* 2009; 65:249–251. [PubMed: 19262295]
43. Lindsay RS, Wake DJ, Nair S, et al. Subcutaneous adipose 11 beta-hydroxysteroid dehydrogenase type 1 activity and messenger ribonucleic acid levels are associated with adiposity and insulinemia in Pima Indians and Caucasians. *J Clin Endocrinol Metab.* 2003; 88:2738–2744. [PubMed: 12788882]
44. Staab CA, Maser E. 11 beta-Hydroxysteroid dehydrogenase type 1 is an important regulator at the interface of obesity and inflammation. *J Steroid Biochem Mol Biol.* 2010; 119:56–72. [PubMed: 20045052]
45. Morgan SA, Sherlock M, Gathercole LL, et al. 11beta-hydroxysteroid dehydrogenase type 1 regulates glucocorticoid-induced insulin resistance in skeletal muscle. *Diabetes.* 2009; 58:2506–2515. [PubMed: 19675138]
46. Karlsson C, Jernas M, Olsson B, et al. Differences in associations between HSD11B1 gene expression and metabolic parameters in subjects with and without impaired glucose homeostasis. *Diabetes Res Clin Practice.* 2010; 88:252–258.
47. Franks PW, Knowler WC, Nair S, et al. Interaction between an 11betaHSD1 gene variant and birth era modifies the risk of hypertension in Pima Indians. *Hypertension.* 2004; 44:681–688. [PubMed: 15452033]
48. Koska J, de Court, Wake DJ, et al. 11beta-hydroxysteroid dehydrogenase type 1 in adipose tissue and prospective changes in body weight and insulin resistance. *Obesity.* 2006; 14:1515–1522. [PubMed: 17030962]
49. Meng YX, Ford ES, Li CY, et al. Association of C-reactive protein with surrogate measures of insulin resistance among nondiabetic US adults: findings from National Health and Nutrition Examination Survey 1999–2002. *Clin Chem.* 2007; 53:2152–2159. [PubMed: 17951292]
50. Gelaye B, Revilla L, Lopez T, et al. Association between insulin resistance and C-reactive protein among Peruvian adults. *Diabetol Metab Syndr.* 2010; 2:30. [PubMed: 20482756]
51. Marioni RE, Deary IJ, Murray GD, et al. Genetic variants associated with altered plasma levels of C-reactive protein are not associated with late-life cognitive ability in four Scottish samples. *Behavior Genet.* 2010; 40:3–11.
52. Gouda HN, Sagoo GS, Harding AH, et al. The association between the peroxisome proliferator-activated receptor-gamma 2 (PPARG2) Pro12Ala gene variant and type 2 diabetes mellitus: a HuGE review and meta-analysis. *Am J Epidemiol.* 2010; 171:645–655. [PubMed: 20179158]
53. Fornage M, Jacobs DR, Steffes MW, et al. Inverse effects of the PPAR(gamma)2 Pro12Ala polymorphism on measures of adiposity over 15 years in African Americans and whites. The CARDIA study. *Metabolism.* 2005; 54:910–917. [PubMed: 15988700]
54. Nelson TL, Fingerlin TE, Moss L, et al. The PPARGgamma Pro12Ala polymorphism is not associated with body mass index or waist circumference among Hispanics from Colorado. *Ann Nutr Metab.* 2007; 51:252–257. [PubMed: 17587797]
55. Yen CJ, Beamer BA, Negri C, et al. Molecular scanning of the human peroxisome proliferator activated receptor gamma (hPPAR gamma) gene in diabetic Caucasians: identification of a Pro12Ala PPAR gamma 2 missense mutation. *Biochem Biophys Res Commun.* 1997; 241:270–274. [PubMed: 9425261]

56. Ben Ali S, Ben Yahia F, Sediri Y, et al. Gender-specific effect of Pro12A1a polymorphism in peroxisome proliferator-activated receptor gamma-2 gene on obesity risk and leptin levels in a Tunisian population. *Clin Biochem.* 2009; 42:1642–1647. [PubMed: 19733160]
57. Dedoussis GV, Vidra N, Butler J, et al. Peroxisome proliferator-activated receptor-gamma (PPAR gamma) Pro12Ala polymorphism and risk for pediatric obesity. *Clin Chem Lab Med.* 2009; 47:1047–1050. [PubMed: 19634985]
58. Ghossaini M, Meyre D, Lobbens S, et al. Implication of the Pro 12Ala polymorphism of the PPAR-gamma 2 gene in type 2 diabetes and obesity in the French population. *BMC Med Genet.* 2005; 22:6.
59. Johansson LE, Danielsson P, Norgren S, et al. Interaction between PPARG Pro12Ala and ADIPOQ G276T concerning cholesterol levels in childhood obesity. *Int J Pediatr Obesity.* 2009; 4:119–125.
60. Fumeron F, Aubert R, Siddiq A, et al. Adiponectin gene polymorphisms and adiponectin levels are independently associated with the development of hyperglycemia during a 3-year period—The Epidemiologic Data on the Insulin Resistance Syndrome prospective study. *Diabetes.* 2004; 53:1150–1157. [PubMed: 15047634]
61. Gu HF, Abulaiti A, Ostenson CG, et al. Single nucleotide polymorphisms in the proximal promoter region of the adiponectin (APM1) gene are associated with type 2 diabetes in Swedish Caucasians. *Diabetes.* 2004; 53:S31–S35. [PubMed: 14749263]
62. Koster JC, Permutt MA, Nichols CG. Diabetes and insulin secretion—the ATP-sensitive K⁺ channel (K-ATP) connection. *Diabetes.* 2005; 54:3065–3072. [PubMed: 16249427]
63. Gloyn AL, Weedon MN, Owen KR, et al. Large-scale association studies of variants in genes encoding the pancreatic beta-cell KATP channel subunits Kir6.2 (KCNJ11) and SUR1 (ABCC8) confirm that the KCNJ11 E23K variant is associated with type 2 diabetes. *Diabetes.* 2003; 52:568–572. [PubMed: 12540637]
64. Sakamoto Y, Inoue H, Keshavarz P, et al. SNPs in the KCNJ11-ABCC8 gene locus are associated with type 2 diabetes and blood pressure levels in the Japanese population. *J Hum Genet.* 2007; 52:781–793. [PubMed: 17823772]
65. Florez JC, Burtt N, de Bakker PIW, et al. Haplotype structure and genotype-phenotype correlations of the sulfonylurea receptor and the islet ATP-sensitive potassium channel gene region. *Diabetes.* 2004; 53:1360–1368. [PubMed: 15111507]
66. Famulla S, Lamers D, Hartwig S, et al. Pigment epithelium-derived factor (PEDF) is one of the most abundant proteins secreted by human adipocytes and induces insulin resistance and inflammatory signaling in muscle and fat cells. *Diabetes.* 2010; 59:A98.
67. Chung C, Doll JA, Gattu AK, et al. Anti-angiogenic pigment epithelium-derived factor regulates hepatocyte triglyceride content through adipose triglyceride lipase (ATGL). *J Hepatol.* 2008; 48:471–478. [PubMed: 18191271]
68. Nakamura K, Yamagishi S, Adachi H, et al. Serum levels of pigment epithelium-derived factor (PEDF) are positively associated with visceral adiposity in Japanese patients with type 2 diabetes. *Diabetes Metab Res Rev.* 2009; 25:52–56. [PubMed: 18677713]
69. Wang P, Smit E, Brouwers MC, et al. Plasma pigment epithelium-derived factor is positively associated with obesity in Caucasian subjects, in particular with the visceral fat depot. *Eur J Endocrinol.* 2008; 159:713–718. [PubMed: 18787046]
70. Sutton BS, Weinert S, Langefeld CD, et al. Genetic analysis of adiponectin and obesity in Hispanic families: the IRAS Family Study. *Hum Genet.* 2005; 117:107–118. [PubMed: 15843989]
71. Spitz MR, Detry MA, Pillow P, et al. Variant alleles of the D2 dopamine receptor gene and obesity. *Nutr Res.* 2000; 20:371–380.

Table 1

Examples of previous association studies of the investigated candidate gene SNPs, with obesity and/or obesity-related phenotypes

Gene/SNP	First author/Year/Reference	Ethnic group	n	Phenotype	p
<i>HSD11B1</i> rs846910 *	Nair 2004 (26)	Pima Indian	706	T2D	0.01
<i>CRP</i> rs1205 *	Eiriksdottir 2009 (24)	Pima Indian	127	Insulin sensitivity	0.03
<i>ADIPOQ</i> rs1501299	Sutton 2005 (70)	Caucasian (Icelandic)	2130	BMI	0.045
	Sitonen 2011 (23)	Hispanic	811	BMI, WC	NS
	Richardson 2006 (21)	Caucasian (Finnish)	507	Body weight	0.02
<i>PPARG</i> rs1801282	Fornage 2005 (53)	Mexican-American	439	Obesity, T2D, HOMA-IR	NS
	Cole 2000 (17)	Caucasian	1954	BMI, WC	0.01, 0.01
	Hsueh 2001 (19)	Mexican-American	921	BMI, WC	0.015, 0.028
	Canizales-Quinteros 2007 (16)	Mexican-American	453	Obesity	0.03
<i>ABCC8</i> rs757110 *	Florez 2004 (65)	Mexican-Mestizo	131	Obesity	0.007
<i>SERPINF1</i> rs1136287 *	Hiroiyuki 2007 (22)	Caucasians, Asians, African Americans	>3400	T2D	0.02
<i>ANKK1</i> rs1800497	Noble 1994 (29)	Japanese	416	Diabetic retinopathy	NS
	Spitz 2000 (71)	Non-Hispanic and Hispanic	73	Obesity	<0.002
	Thomas 2001 (30)	Caucasian	176	Obesity BMI	0.002
		Chinese	990	Obesity BMI, WC	0.02

n = subjects included in the study.

NS, not significant; BMI, body mass index; WC, waist circumference; T2D, type 2 diabetes; HOMA-IR, homeostasis model assessment of insulin resistance.

* SNPs not investigated in Hispanic cohorts to date.

Table 2

Anthropometric and clinical characteristics of the study group

Characteristics	Median	±SD	Interquartile	Min	Max
M/F	155/293				
Age (Years)	16	±3.53	15–19	14	30
BMI (kg/m ²)					
General	24.69	±6.28	21.86–29.06	15.67	52.21
Females	24.25	±6.29	21.86–29.05	17.17	52.21
Males	25.4	±6.25	21.98–29.46	15.67	47.65
WC (cm)					
General	82	±16.26	74.92–94	30	157
Women	81	±14.79	74–92	40.5	128.73
Men	83.35	±18.62	76–97	30	157
SBP (mmHg)	110	±11.25	100–118	85	156
DBP (mmHg)	69	±8.89	62–74	43	97
Fasting plasma glucose (mmol/l)	85 ^a	±17.76	79.65–93	43	266
LogTG (mmol/l)	1.81 ^b	±0.22	1.6627–1.9530	1.2552	2.6053
HOMA-IR	2.13 ^c	±2.08	1.3252–3.3296	0.1377	19.7481

BMI, body mass index; WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure; HOMA-IR, homeostasis model assessment of insulin resistance; Log TG, triglycerides logarithmically transformed.

Number of subjects :

^a 239,

^b 302,

^c 297.

Table 3
Genotype and allele frequencies of the seven analyzed SNPs and Hardy-Weinberg calculations

SNPs (genes)	Major/minor allele	Major allele homozygote (%)	Heterozygote (%)	Minor allele homozygote (%)	MAF	HWE <i>p</i>
rs846910 (<i>HSD11B1</i>)	G/A	230 (74.3)	80 (23.2)	7 (2.5)	0.14	1.0
rs1205 (<i>CRP</i>)	C/T	134 (40.4)	155 (49.5)	28 (10.1)	0.34	0.07826
rs1501299 (<i>ADIPOQ</i>)	G/T	207 (64.4)	93 (29.1)	17 (6.4)	0.21	0.1585
rs1801282 (<i>PPARG</i>)	C/G	236 (75.3)	73 (22.2)	8 (2.5)	0.13	0.3604
rs1800497 (<i>ANKK1</i>)	G/A	120 (38.3)	148 (46.5)	49 (15.2)	0.38	0.8129
rs757110 (<i>ABCC8</i>)	A/C	127 (39.5)	153 (49)	37 (11.5)	0.36	0.4631
rs1136287 (<i>SERPINF1</i>)	T/C	94 (29.8)	165 (53.5)	58 (16.7)	0.43	0.3638

HWE *p*, Hardy Weinberg calculation; MAF, minor allele frequency.

Table 4
Association of SNPs with obesity (BMI) on top and central obesity (WC) on bottom, adjusted for age and gender

SNP (Gene)	Major/Minor allele	<i>n</i>		MAF			<i>p</i>	Permuted <i>p</i>
		Non-obese/Obese	Non-obese (BMI)	Obese (BMI)	OR (95%CI)			
rs846910 (<i>HSD11B1</i>)	G/A	179/98	0.13	0.18	0.8524 (0.3804–1.91)		0.6	0.9
rs1205 (<i>CRP</i>)	C/T	186/101	0.33	0.32	0.9546 (0.611–1.491)		0.8	1.0
rs1501299 (<i>ADIPOQ</i>)	G/T	181/99	0.20	0.25	1.274 (0.7844–2.07)		0.3	0.9
rs1801282 (<i>PPARG</i>)	C/G	184/101	0.13	0.14	1.713 (0.8212–3.573)		0.1	0.6
rs1800497 (<i>ANKK1</i>)	G/A	181/106	0.34	0.42	1.47 (1.005–2.15)		0.04	0.2
rs757110 (<i>ABCC8</i>)	A/C	179/105	0.37	0.34	0.8108 (0.5356–1.227)		0.3	0.9
rs1136287 (<i>SERPINF1</i>)	T/C	178/103	0.44	0.43	0.9356 (0.6428–1.362)		0.7	0.9
SNP (gene)	Major/Minor allele	Non-obese/Centrally obese	Non-obese (WC)	Centrally obese (WC)	OR (95%CI)		<i>p</i>	Permuted <i>p</i>
rs846910 (<i>HSD11B1</i>)	G/A	167/185	0.11	0.16	1.035 (0.527–2.031)		0.9	1.0
rs1205 (<i>CRP</i>)	C/T	174/190	0.34	0.34	1.026 (0.7084–1.487)		0.8	1.0
rs1501299 (<i>ADIPOQ</i>)	G/T	170/185	0.21	0.20	1.144 (0.7299–1.793)		0.5	0.9
rs1801282 (<i>PPARG</i>)	C/G	173/190	0.12	0.14	1.872 (0.8402–4.172)		0.1	0.5
rs1800497 (<i>ANKK1</i>)	G/A	169/197	0.31	0.44	1.54 (1.114–2.129)		0.009	0.03
rs757110 (<i>ABCC8</i>)	A/C	168/195	0.37	0.34	0.8375 (0.591–1.187)		0.3	0.9
rs1136287 (<i>SERPINF1</i>)	T/C	166/191	0.40	0.45	1.174 (0.851–1.618)		0.3	0.9

n = number of subjects genotyped.

OR, odds ratio; MAF, minor allele frequency.

Association of SNPs with quantitative traits adjusted for gender, age and BMI

Table 5

SNP (Gene)	Genotype	HOMA-IR		Fasting plasma glucose (mmol/l)		Log TG (mmol/l)		SBP (mmHg)		DBP (mmHg)	
		Median (SD)	p	Permuted p	p	Permuted p	p	Median (SD)	Permuted p	Median (SD)	Permuted p
rs846910 (<i>HSD11B1</i>)	GG	1.682 (2.242)	0.04	0.2	0.7	0.9	0.03	109.5 (11.09)	0.8	67.98 (8.765)	0.3
	GA	1.987 (1.917)			44.17 (45.08)		1.308 (0.8685)	110.7 (11.18)		68.82 (9.098)	
	AA	2.579 (1.691)			64.16 (37.23)		1.783 (0.112)	106.0 (6.585)		68.33 (10.92)	
rs1205 (<i>CRP</i>)	CC	1.811 (2.574)			49.25 (44.34)		1.245 (0.8702)	109.7 (11.42)	0.4	68.07 (8.756)	0.9
	CT	1.619 (1.742)	0.03	0.2	50.87 (44.85)	0.007	0.1	109.5 (10.69)	0.5	68.01 (9.139)	0.1
	TT	1.944 (2.062)			50.88 (58.3)		1.288 (0.8744)	109.8 (12.38)		69.53 (9.367)	
rs1501299 (<i>ADIPOQ</i>)	GG	1.688 (1.818)	0.6	0.9	47.18 (44.24)	0.9	0.5	109.4 (10.41)	0.2	67.97 (9.005)	0.9
	GT	1.889 (2.877)			60.27 (48.63)		1.135 (0.8971)	109.8 (11.8)		68.62 (8.68)	
	TT	1.29 (1.501)			53.67 (44.93)		1.053 (0.949)	114.0 (15.44)		69.18 (9.43)	
rs1801282 (<i>PPARG</i>)	CC	1.804 (2.263)			52.84 (45.17)		1.256 (0.8747)	109.5 (10.79)		68.22 (9.197)	
	CG	1.537 (1.74)	0.1	0.7	43.36 (47.51)	0.1	0.6	108.6 (12.02)	0.07	67.74 (8.488)	0.8
	GG	1.361 (1.873)			36.0 (54.06)		1.027 (0.9871)	118.6 (11.72)		71.11 (7.849)	
rs1800497 (<i>ANKK1</i>)	GG	1.461 (1.701)			52.49 (44.02)		1.146 (0.8863)	109.1 (10.75)		67.04 (7.687)	
	GA	1.903 (2.155)	0.4	0.9	50.4 (47.13)	0.7	0.8	110.2 (11.85)	0.8	69.54 (9.395)	0.8
	AA	2.216 (3.038)			50.41 (47.71)		1.364 (0.8503)	109.9 (10.83)		67.12 (9.648)	
rs757110 (<i>ABCC8</i>)	AA	1.688 (1.858)			51.76 (44.09)		1.206 (0.8752)	111.0 (11.1)		68.56 (8.328)	
	AC	1.841 (2.444)	0.7	0.9	53.48 (47.61)	0.7	0.5	109.6 (11.91)	0.1	68.22 (9.482)	0.5
	CC	1.799 (2.021)			37.33 (44.28)		1.234 (0.8992)	106.6 (8.136)		68.24 (8.804)	
rs1136287 (<i>SERPINF1</i>)	TT	1.773 (1.784)	0.7	1.0	53.23 (48.54)	0.1	0.4	110.0 (10.33)	0.4	68.78 (7.997)	0.2
	TC	1.873 (2.459)			52.96 (45.22)		1.216 (0.8829)	109.9 (11.34)		68.53 (9.308)	
	CC	1.462 (1.888)			48.46 (43.41)		1.201 (0.87120)	108.8 (11.980)		67.05 (8.376)	

SBP, systolic blood pressure; DBP, diastolic blood pressure; HOMA-IR, homeostasis model assessment of insulin resistance; Log TG, triglycerides logarithmically transformed.