Published in final edited form as:

Spine (Phila Pa 1976). 2011 December 1; 36(25 Suppl): S218–S225. doi:10.1097/BRS.0b013e3182387f96.

How can animal models inform on the transition to chronic symptoms in whiplash?

Beth A. Winkelstein, PhD

Departments of Bioengineering & Neurosurgery, University of Pennsylvania, 240 Skirkanich, 210 S. 33rd St., Philadelphia, PA 19104-6321, Phone: (215) 573-4589, Fax: (215) 573-2071

Beth A. Winkelstein: winkelst@seas.upenn.edu

Abstract

Study Design—A non-systematic review of the literature.

Objective—The objective was to present general schema for mechanisms of whiplash pain and review the role of animal models in understanding the development of chronic pain from whiplash injury.

Summary of Background Data—Extensive biomechanical and clinical studies of whiplash have been performed to understand the injury mechanisms and symptoms of whiplash injury. However, only recently have animal models of this painful disorder been developed based on other pain models in the literature.

Methods—A non-systematic review was performed and findings were integrated to formulate a generalized picture of mechanisms by chronic whiplash pain develops from mechanical tissue injuries.

Results—The development of chronic pain from tissue injuries in the neck due to whiplash involves complex interactions between the injured tissue and spinal neuroimmune circuits. A variety of animal models are beginning to define these mechanisms.

Conclusion—Continued work is needed in developing appropriate animal models to investigate chronic pain from whiplash injuries and care must be taken to determine whether such models aim to model the injury event or the pain symptom.

Keywords

spine; whiplash; pain; neck; animal model

This review describes the current understanding of how chronic pain develops, with a focus on whiplash-related pathomechanisms. Background on pain modeling is introduced with relevance to symptoms and sensitivity. Several working hypotheses describing mechanisms of pain symptom development and maintenance are presented, including concepts of the neurophysiology of pain initiation from injury and longer-term signal processing and the central nervous system's (CNS) neuroimmunologic involvement. A schema is presented highlighting the cellular and molecular processes for acute pain, as well as for the transition to a persistent pain state. Specific findings in a variety of animal model systems are also summarized that inform on the pain mechanisms originating from tissues relevant to whiplash injury. Lastly, potential considerations for animal modeling and interpretation of findings for a human disease state are presented.

Animal Modeling of Pain Symptoms

In vivo models of injury have provided the major platform to define relationships between the time course and extent of activation of the nociceptive cascade and development and maintenance of pain symptoms. A broad range of animal models has been developed to study several pain states, including neuropathic, inflammatory, peripherally-mediated and centrally-induced. Behavioral assessments enable pain measures in those models, covering different modalities, anatomic regions and paradigms of testing (Figure 1). Behavioral metrics are defined in the context of the applied stimulus and are described in terms related to the clinical symptoms. *Hyperalgesia* is an increased response to a stimulus which is normally painful, and includes all conditions of increased pain sensitivity [1]. *Allodynia* is a particular case of hyperalgesia in which a stimulus that is usually not painful becomes noxious, reflecting that the stimulus and responses differ [1]. Both of these responses are gauges of hypersensitivity in both humans and animals, and so, serve as important and useful quantitative outcomes for modeling of pain symptoms [2]. Various animal models of pain have been developed to mimic the symptoms and the syndromes of acute and chronic pain with an equally broad range of inciting pathologies. Indeed, that body of work is quite extensive and is beyond the scope of this review; several other reviews cover this topic in more detail [2–4]

Pain Mechanisms – Determining the Transition from Acute to Chronic States

An inciting event that stimulates nerve fibers initiates local responses and can lead to central signaling events that include neuronal and immune responses (Figure 2). In acute or nociceptive pain, these cascades resolve; however, when chronic pain develops, CNS modifications lead to hypersensitivity and central sensitization in which the spinal neural circuits respond with a heightened response to stimuli [2,5]. The CNS also mounts a widespread and sustained neuroimmune response further contributing to neuronal plasticity and pain maintenance (Figure 2).

Acute pain results from noxious stimuli that activate small-diameter unmyelinated C-fibers or thinly myelinated A δ sensory neurons in the periphery [6]. Many different neurotransmitters communicate from the peripheral afferents to spinal cord neurons. Nociceptive pain protects against tissue injury and continues *only* in the presence of sustained noxious stimuli [5]. After a tissue injury, local nociceptors become sensitized and exhibit both lower thresholds for firing and increased firing rates when exposed to a previously non-noxious stimulus [5,7,8]. This peripheral sensitization is due largely to the production and peripheral release of neurotransmitters and/or other factors from neurons and non-neuronal cells that can infiltrate the injured area (Figure 2) [7,9]. Multiple mediators, including bradykinin, amines, prostanoids, NGF, and protons, act rapidly via receptors on nociceptor terminals to sensitize these neurons [9]. This generalized nociceptive pain response encompasses the typical cascade of an acutely-painful episode, where the balance of injury, repair and healing is achieved and the collection of electrophysiological and chemical events resolves. However, for *persistent* pain the local, spinal, and supraspinal cascades become pathologically and permanently altered [7].

The contribution of the local and widespread inflammatory response plays a large role in the transition from acute to chronic pain. Persistent pain is due, in large part, to the initiation and maintenance of spinal sensitization via plasticity in the neuronal circuits. Although the exact mechanism by which the spinal cord transitions to a hyperexcitable state remains somewhat unknown, many hypotheses have emerged. These theories are highlighted briefly here; more extensive discussions can be found elsewhere [5,9–11]. Simultaneously, different processes

occur that induce this central sensitization. The low-threshold A β afferents, which normally do not transmit nociceptive signals, are recruited to transmit spontaneous and movement-induced pain [10]. The central hyperexcitability is further exacerbated by a “wind-up” response of repetitive C-fiber stimulation, expanding receptive fields, and spinal neurons taking on properties of wide dynamic range neurons [12,13]. Ultimately, A β -fibers stimulate the postsynaptic neurons, whereas these A β -fibers previously transmitted only to non-painful innocuous stimuli. Amplification of noxious signals from afferents at the synapse in the spinal cord can occur and this long-term potentiation can exacerbate pain [11].

Although the neuronal responses in the nervous system contribute to the onset of pain, neuroimmune responses throughout the nervous system also potentiate persistent pain (Figure 2) [14–17]. Tissue injury leads to the release of many chemical mediators that enhance neuronal activity and also initiate a neuroimmune cascade in which many peripheral non-neuronal cells are activated (Figure 2) [7,14,15]. These resident cells, including mast and Schwann cells, release mediators such as histamine, prostaglandins, cytokines and chemokines, that recruit other infiltrating immune cells (Figure 2). The cytokine cascade is responsible for the infiltration of immune cells to the site of injury [8,14,15]. Proinflammatory cytokines trigger the release of other inflammatory mediators that sensitizes nociceptors and further maintains neuronal excitability and sensitization, which leads to pain [8,14] (Figure 2). However, some cytokines, such as TNF- α and IL-1 β , can also directly sensitize nociceptors during inflammation since sensory neurons express their receptors [8,18].

Neurotrophins and growth factors (NGF, in particular) are important mediators and modulators of pain [19–22]. Although neurotrophic factors are necessary for the long-term survival of neurons, there is emerging evidence for their roles as crucial mediators of persistent pain. NGF is upregulated in many chronic pain conditions and modulates pain through BDNF which is released from the spinal terminals of a subset of nociceptors [21,23]. BDNF acts through the trkB receptor and leads to central sensitization by potentiating glutamate neurotransmission in the dorsal horn and promoting the synaptic rearrangement of the A β -fibers [14,21,24]. Neurotrophins also have indirect actions on neurotransmission since they are synthesized and released by several types of immune cells [14], which themselves interface with neurons and can modulate synaptic activity.

Activation of peripheral nociceptors also activates glial cells in the CNS. Microglia are the resident macrophages in the CNS and astrocytes help with maintaining homeostasis at neuronal synapses. Both types of glial cells respond to injury by changing their morphology, proliferating, and releasing inflammatory mediators [17,25–27]. Spinal microglia are activated early after peripheral injury by some neuromodulators, including excitatory amino acids, substance P and others [27,28]; when activated they release several proinflammatory cytokines (IL-1 β , TNF- α , IL-6), as well as nitric oxide, prostaglandins and NGF [16,25,29]. In addition, BDNF released from stimulated microglia modulate neuronal signaling in the superficial dorsal horn [30]. These mediators, in turn, induce the exaggerated release of neurotransmitters from presynaptic neurons, sensitize the post-synaptic membrane, activate neighboring astrocytes and enhance microglial activity [16,25]. This positive feedback sustains the continued further release of pain mediators, facilitating the development of neuronal hypersensitivity and leading to persistent pain.

Relevance to Whiplash – Considerations for Animal Modeling

There is growing physiologic evidence that various structures in the neck may generate pain and models have been developed for their injury [2,18]. Although the exact pain mechanisms for whiplash remain open to debate, there is consensus on the potential sites of

injury and how pain can originate from them. Recent research for cervical spine pain has focused particularly on understanding mechanisms of acute and chronic whiplash injury. These findings are discussed with regard to injury biomechanics and the nociceptive mechanisms related to pain generation and maintenance.

The Facet Joint

Many of the hypotheses of facet-mediated whiplash pain imply that “abnormal” motion patterns develop in the spine, that induce local injury or dysfunction to the specific anatomical components of the facet joint. During these injuries, the lower cervical spine undergoes extension while the upper spine flexes, modifying the local biomechanical environment of the facet joint by inducing: (1) compression of the facet joint, and/or (2) excessive capsular ligament strain [31–35].

The potential for tissue loading and pain generation is primarily relevant to the ligamentous capsule that encloses the facet joint [18,35]. Many engineering studies demonstrate the facet capsule to be stretched more during whiplash than in normal physiologic motions [37–43]. However, frank rupture of the cervical facet capsule has not been reported for any whiplash kinematic [38,39,44]. Even transient tensile loading of the cervical facet capsule can activate both short and long-lasting pain pathways [45–48]. For example, short-duration capsule loading activates and saturates mechanoreceptive and nociceptive afferents [45], suggesting an immediate mechanism to initiate pain. Interestingly, in a rat model of single exposure of the facet joint to tension loading that results in similar distortions in the capsule that saturate nerve afferents also produce persistent sensitivity [47,49–53]. The capsular biomechanics associated with those responses are very similar to those reported for the human capsule during whiplash simulations [34,38,39,54]. Using animal model systems, it has been further determined that those same transient painful joint loading scenarios are also associated with the production of collagen fiber disorganization [55,56], and axonal swelling and altered morphology [48], which provide a potential means to initiate pain maintenance.

Transient facet capsule loading that produces persistent pain also leads to a complicated cellular cascade in the affected afferents in the dorsal root ganglia (DRG) and spinal cord. Both neurons and accessory cells in the DRG and spinal cord exhibit modified phenotypic responses to painful joint loading (Figures 2 & 3) [47,53,57]. In particular, painful facet capsule tension increases expression of a marker of the activated stress response in DRG neurons in order to attempt to re-establish protein homeostasis that can be altered by neuronal damage [53,58]. Gene transcription of substance P in afferents is also induced [52] suggesting that painful loading may induce axonal dysfunction that subsequently disrupts normal protein production. Those afferent modifications are complemented by increased spinal activity of the glutamatergic system and neuronal hyperexcitability, that are sustained in association with persistent pain [47,57]. In contrast, spinal substance P is detected to decrease early and may be a consequence of increased protein utilization in the spinal cord and/or injury-induced axonal dysfunction that can affect protein transport [52]. Substance P can regulate the spinal glial responses [28] which has an important impact on pain because spinal glia are critical in the onset and maintenance of *chronic* pain [17,25–27] and are associated with persistent sensitivity in that same model [49,59]. Cytokines are also modulated in both the DRG and spinal cord after loading of the facet joint [50]. Cytokine expression parallels spinal glial activation and both of these responses increase after painful subfailure capsule loading; however, neither may specifically drive *chronic* pain because joint loading that produces greater injury does not independently produce more behavioral sensitivity [50,59]. This apparent disconnect between the central immune response and facet joint loading indicates there may be other cellular processes that maintain facet-mediated persistent pain.

The Nerve Root & DRG

The potential for nerve root and DRG trauma exists due to the flexibility of the cervical spine, coupled with the neck's potential to undergo loading due to motions of the head and torso. The cervical nerve roots can be at particular risk for injury due to increased hydrostatic pressure that can be established during rapid head and neck motions [60–63]. Although there is growing research in these areas, most of the current understanding of chronic pain developing from nerve root injury is from a large body of work in the lumbar spine. However, a unique porcine model was developed to study the potential mechanisms of whiplash-induced pain [60,61].

Aldman's group hypothesized that pressure gradients in the CNS from blood flow resistance can be generated during rapid spinal motions such as whiplash [60,61]. The spinal vasculature regulates blood volume in order to accommodate any changes in spinal canal size. Yet, during rapid head/neck motions, resistance to blood flow can generate pressure gradients in and outside the spinal canal [60–63]. These gradients can directly apply pressure to the nerve roots and spinal ganglia that can induce cellular injury [61,63]. Coupling these hypothesized pressure-wave effects with the direct mechanical compression and tethered stretching of the neural tissues that can occur by abnormal rapid spinal motions, several painful injury mechanisms emerge [60,62]. In fact, plasma membrane breakdown of spinal ganglia cells has been induced in an *in vivo* porcine model of rapid head-neck extension [60–63]. Although it suggested a potential mode of injury to neuron cell bodies, that work did not provide any direct measure of pain.

Emerging research in persistent radiculopathic pain has highlighted the influence of injury on the neurophysiologic responses that lead to the establishment of chronic pain. The injury profile at the time of nerve root injury modulates neuropeptide expression in the DRG and the spinal cord, associated with development of persistent pain [64–66]. Cervical root compression that produces persistent sensitivity shows greater allodynia at later time points after more severe dorsal root compression [64,66]. There is a corresponding depletion of substance P in the DRG that also depends on the injury severity, suggesting its synaptic release and utilization may be increased in persistently painful injuries [67,68]. Spinal expression of CGRP also decreases significantly with increasing nerve root loading and pain [64,66]. Further, both the extent of axonal damage and Wallerian degeneration in the compressed nerve root are directly related to the likelihood of developing persistent allodynia [66,68].

Much of what is understood about the development of persistently painful radiculopathy comes from models of lumbar radicular pain [69–73]. It is clear from studies of lumbar and cervical nerve root compression that direct nerve root trauma (even transient) can induce long-lasting pain and a constellation of physiologic responses throughout the nervous system [64,65,68,69,74–76]. The onset of behavioral hypersensitivity is immediate following trauma and parallels a host of widespread neuronal and neuroinflammatory responses, including endoneurial edema, axonal membrane leakage and Wallerian degeneration [73,77–79]. In response to axonal damage, macrophages infiltrate the injury site, phagocytose myelin debris, and induce the release of IL-1 and TNF- α [67]. By blocking the TNF- α pathway using a soluble TNF receptor at the time of injury, behavioral sensitivity is reduced after injury along with decreases in both TNF- α and IL-1 β mRNA [29,80,81]. Axonal degeneration and inflammation can also increase electrical activity in adjacent intact axons [72,82], leading to spinal sensitization and the immune sequelae that further exacerbate sensitization (Figure 2). Spinal glial cells become activated and modulate other immunologic changes leading to persistent pain, via cytokine and growth factor production [15,80,83,84]. Spinal glial activation is also sensitive to the severity of loading and the presence of chemical irritation in radicular pain [74,76,85], with greater degrees of nerve

root compression inciting greater activation of spinal astrocytes [71,74]. The differentiation in activation is apparent early and is sustained in parallel with persistent pain symptoms [76], suggesting that astrocytes may directly respond to the changes in the dorsal horn that are induced by damaged primary afferents [86,87]. Spinal microglial activation is also induced and appears to be directly related to the *onset* of pain [76]. However, spinal microglia proliferation induced early after cervical nerve root compression may only indicate tissue trauma and not be due to nociceptive signaling and pain [88].

Lessons from Animal Models of Pain & Implications

Given the extreme complexity of the mechanisms of neck injuries that *do and do not result in chronic pain*, it must be recognized that there are a variety of factors that have important roles. For example, many biomechanical factors have been shown to alter neuronal function: rate, loading duration, and load magnitude, among others [47,51,52,57,59,64,74,75,89,90]. The factors which confound the injuries themselves, the physiologic milieu at the time of injury, can also potentially directly modulate the severity and extent of pain [70,76,81,85,88,91]. For example, it can be hypothesized that injury following an already preexisting *minor* tissue injury may produce a more severe nociceptive response than in the absence of any preexisting damage. It is not unlikely that an initial injury produces the local inflammatory changes discussed earlier, which in turn may lower that tissue's threshold for mechanical injury that can lead to symptoms (Figure 2). In fact, the mechanical insult required to produce behavioral hypersensitivity in the presence of an inflammatory insult is less than that required in its absence, despite producing the same degree of behavioral sensitivity [70,76,81,85,88], corroborating clinical studies in which patients with preexisting spinal degeneration experience more severe and longer lasting neck pain symptoms [92,93]. It is possible that degeneration can contribute to inflammatory changes in the facet joint that increase its susceptibility to whiplash injury so when its tissues undergo loading that may not normally be painful, the nerve fibers may be pre-sensitized and have lower thresholds for response than those previously required to initiate nociception or that may induce a more widespread central response and chronic pain. Nonetheless, an integrated approach is required in order to fully understand the complicated disease of chronic pain, especially in the context of whiplash injury.

Although animal models are integral and necessary in order to define mechanisms of painful injury and those factors that drive the transition to chronic pain, there are several important caveats about using such platforms, in particular to understand a mechanically-based injury such as whiplash. Anatomic geometry and scaling are paramount among these considerations. For example, most animal models of pain have been developed in quadrupeds that do not have the same mechanical demands, anatomic relationships and locomotion demands as the bipedal human. This must not be ignored particularly for spine-based pathologies in which the tissue loading can be substantially different among species. However, the neurologic function, tissue responses and relationships between them have been well-documented for a number of species. Nonetheless, care must be taken when developing such models and interpreting results.

It is also important to consider whether an animal model is serving as a proxy to model the pathology/injury or the symptoms that develop. Within that context it is also important to note that the current animal models of whiplash pain measure only evoked pain and do not capture the spontaneous pain that is reported clinically. In this sense, animal models could be enhanced by the development of assessments for behavioral outcomes that measure the pain in the referred pain areas observed clinically. Certainly, the behavioral and physiological evidence that is already available strongly suggests spontaneous pain may be induced in these models, but additional work is needed to develop relevant and sensitive

assays to detect it. In fact, animal models of whiplash injury and chronic pain offer great promise since they represent a “reduced” model system in which many of the confounding factors that are present in the clinical problem – such as litigation status, compensation issues and psychosocial factors – can be controlled or avoided altogether. Conversely, the complexity of the clinical presentation, which includes a variety of physical function and stress responses [94], also is not fully-captured in the laboratory model.

The complexity of the clinical syndrome of chronic whiplash pain is further extended by the recent reports that genetics play a key role in pain persistence, accounting for many discrepancies observed for seemingly similar injuries [95]. The utility of animal models is both highlighted and hampered by this. While the clinical picture of persistent pain in humans is indeed intricate, similar genetic specificity has been defined for different strains of rats exhibited different responses to a standardized painful tissue injury [96]. Moreover, such diversity could be exploited in a rat model to determine if and which genetic variants may be more predictive of chronic pain from whiplash. Such studies are needed and would undoubtedly provide additional valuable insight into the clinical management of this syndrome. Indeed, it may be possible to induce behavioral outcomes that model the clinical picture but it is imperative to consider then the pathology, tissue origin and pathways being mapped – the question to consider becomes “is it a model of the injury or the symptoms?”

Acknowledgments

The author gratefully acknowledges the National Institutes of Health (NIAMS; Grant #AR056288) and the Catharine Sharpe Foundation for support.

The manuscript submitted does not contain information about medical device(s)/drug(s). Institutional and Foundation funds were received to support this work. No benefits in any form have been or will be received from a commercial party related directly or indirectly to the subject of this manuscript.

References

1. Loeser JD, Treede RD. The Kyoto protocol of IASP Basic Pain Terminology. *Pain*. 2008; 137:473–7. [PubMed: 18583048]
2. DeLeo JA, Winkelstein BA. Physiology of chronic spinal pain syndromes: from animal models to biomechanics. *Spine*. 2002; 27:2526–37. [PubMed: 12435988]
3. Le Bars D, Gozariu M, Cadden SW. Animal models of nociception. *Pharmacol Rev*. 2001; 53:597–652. [PubMed: 11734620]
4. Winkelstein BA. Mechanisms of central sensitization, neuroimmunology & injury biomechanics in persistent pain: implications for musculoskeletal disorders. *J Electromyogr Kinesiol*. 2004; 14:87–93. [PubMed: 14759754]
5. Costigan M, Scholz J, Woolf CJ. Neuropathic Pain: A maladaptive response of the nervous system to damage. *Annu Rev Neurosci*. 2009; 32:1–32. [PubMed: 19400724]
6. Cavanaugh JM. Neural mechanisms of lumbar pain. *Spine*. 1995; 20:1804–9. [PubMed: 7502138]
7. Julius D, Basbaum AI. Molecular mechanisms of nociception. *Nature*. 2001; 413:203–10. [PubMed: 11557989]
8. Verri WA Jr, Cunha TM, Parada CA, et al. Hypernociceptive role of cytokines and chemokines: targets for analgesic drug development? *Pharmacol Ther*. 2006; 112:116–38. [PubMed: 16730375]
9. Hucho T, Levine JD. Signaling pathways in sensitization: toward a nociceptor cell biology. *Neuron*. 2007; 55:365–76. [PubMed: 17678851]
10. Djouhri L, Lawson SN. Abeta-fiber nociceptive primary afferent neurons: a review of incidence and properties in relation to other afferent A-fiber neurons in mammals. *Brain Res Brain Res Rev*. 2004; 46:131–45. [PubMed: 15464202]
11. Ruscheweyh R, Wilder-Smith O, Drdla R, et al. Long-term potentiation in spinal nociceptive pathways as a novel target for pain therapy. *Mol Pain*. 2011; 7:20. [PubMed: 21443797]

12. Cook AJ, Woolf CJ, Wall PD, et al. Dynamic receptive-field plasticity in rat spinal-cord dorsal horn following C-primary afferent input. *Nature*. 1987; 325:151–3. [PubMed: 3808072]
13. Ji RR, Kohno T, Moore KA, et al. Central sensitization and LTP: do pain and memory share similar mechanisms? *Trends Neurosci*. 2003; 26:696–705. [PubMed: 14624855]
14. Moalem G, Tracey DJ. Immune and inflammatory mechanisms in neuropathic pain. *Brain Research Reviews*. 2006; 51:240–64. [PubMed: 16388853]
15. DeLeo JA, Yezierski RP. The role of neuroinflammation and neuroimmune activation in persistent pain. *Pain*. 2001; 90:1–6. [PubMed: 11166964]
16. Inoue K. The function of microglia through purinergic receptors: Neuropathic pain and cytokine release. *Pharmacology & Therapeutics*. 2006; 109:210–26. [PubMed: 16169595]
17. Saab CY, Waxman SG, Hains BC. Alarm or curse? The pain of neuroinflammation. *Brain Research Reviews*. 2008; 58:226–35. [PubMed: 18486228]
18. Siegmund GP, Winkelstein BA, Ivancic PC, et al. The anatomy and biomechanics of acute and chronic whiplash injury. *Traffic Inj Prev*. 2009; 10:101–12. [PubMed: 19333822]
19. Boucher TJ, McMahon SB. Neurotrophic factors and neuropathic pain. *Curr Opin Pharmacol*. 2001; 1:66–72. [PubMed: 11712538]
20. Millan MJ. The induction of pain: an integrative review. *Prog Neurobiol*. 1999; 57:1–164. [PubMed: 9987804]
21. Pezet S, McMahon SB. Neurotrophins: mediators and modulators of pain. *Annu Rev Neurosci*. 2006; 29:507–38. [PubMed: 16776595]
22. Winkelstein BA, Kras JV. Is there an antinociceptive role for peripheral brain-derived neurotrophic factor? *Spine J*. 2010; 10:733–5. [PubMed: 20650413]
23. Gould HJ 3rd, Gould TN, England JD, et al. A possible role for nerve growth factor in the augmentation of sodium channels in models of chronic pain. *Brain Res*. 2000; 854:19–29. [PubMed: 10784102]
24. Matayoshi S, Jiang N, Katafuchi T, et al. Actions of brain-derived neurotrophic factor on spinal nociceptive transmission during inflammation in the rat. *J Physiol*. 2005; 569:685–95. [PubMed: 16210356]
25. DeLeo JA, Tanga FY, Tawfik VL. Neuroimmune activation and neuroinflammation in chronic pain and opioid tolerance/hyperalgesia. *Neuroscientist*. 2004; 10:40–52. [PubMed: 14987447]
26. Suter MR, Wen YR, Decosterd I, et al. Do glial cells control pain? *Neuron Glia Biology*. 2007; 3:255–68. [PubMed: 18504511]
27. Cao H, Zhang YQ. Spinal glial activation contributes to pathological pain states. *Neurosci Biobehav Rev*. 2008; 32:972–83. [PubMed: 18471878]
28. Marriott I. The role of tachykinins in central nervous system inflammatory responses. *Front Biosci*. 2004; 9:2153–65. [PubMed: 15353277]
29. Rothman SM, Winkelstein BA. Cytokine antagonism reduces pain and modulates spinal astrocytic reactivity after cervical nerve root compression. *Ann Biomed Eng*. 2010; 38:2563–76. [PubMed: 20309734]
30. Coull JA, Beggs S, Boudreau D, et al. BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature*. 2005; 438:1017–21. [PubMed: 16355225]
31. Panjabi MM, Cholewicki J, Nibu K, et al. Simulation of whiplash trauma using whole cervical spine specimens. *Spine*. 1998; 23:17–24. [PubMed: 9460147]
32. Panjabi MM, Cholewicki J, Nibu K, et al. Capsular ligament stretches during in vitro whiplash simulations. *J Spinal Disord*. 1998; 11:227–32. [PubMed: 9657548]
33. Stemper BD, Yoganandan N, Gennarelli TA, et al. Localized cervical facet joint kinematics under physiological and whiplash loading. *J Neurosurg Spine*. 2005; 3:471–76. [PubMed: 16381210]
34. Pearson AM, Ivancic PC, Ito S, et al. Facet joint kinematics and injury mechanisms during simulated whiplash. *Spine*. 2004; 29:390–97. [PubMed: 15094535]
35. Ivancic PC, Pearson AM, Tominaga Y, et al. Biomechanics of Cervical Facet Dislocation. *Traffic Inj Prev*. 2008; 9:606–11. [PubMed: 19058109]

36. Kallakuri S, Singh A, Chen C, et al. Demonstration of substance P, calcitonin gene-related peptide, and protein gene product 9.5 containing nerve fibers in human cervical facet joint capsules. *Spine*. 2004; 29:1182–6. [PubMed: 15167655]
37. Yang KH, King AI. Mechanism of facet load transmission as a hypothesis for low-back pain. *Spine*. 1984; 9:557–65. [PubMed: 6238423]
38. Winkelstein BA, Nightingale RW, Richardson WJ, et al. The cervical facet capsule and its role in whiplash injury: a biomechanical investigation. *Spine*. 2000; 25:1238–46. [PubMed: 10806500]
39. Siegmund GP, Myers BS, Davis MB, et al. Mechanical evidence of cervical facet capsule injury during whiplash: a cadaveric study using combined shear, compression, and extension loading. *Spine*. 2001; 26:2095–101. [PubMed: 11698885]
40. Yoganandan N, Pintar FA. Biomechanics of human head-neck in rear impacts. *Int J of Vehicle Design*. 2003; 32:53–67.
41. Luan F, Yang KH, Deng B, et al. Qualitative analysis of neck kinematics during low-speed rear-end impact. *Clin Biomech*. 2000; 15:649–57.
42. Siegmund GP, Davis MB, Quinn KP, et al. Head-turned postures increase the risk of cervical facet capsule injury during whiplash. *Spine*. 2008; 33:1643–9. [PubMed: 18594456]
43. Yoganandan N, Pintar FA, Cusick JF. Biomechanical analyses of whiplash injuries using an experimental model. *Accid Anal Prev*. 2002; 34:663–71. [PubMed: 12214961]
44. Quinn KP, Winkelstein BA. Detection of altered collagen fiber alignment in the cervical facet capsule after whiplash-like joint retraction. *Ann Biomed Eng*. 2011; 39:2163–73. [PubMed: 21538155]
45. Lu Y, Chen C, Kallakuri S, et al. Neural response of cervical facet joint capsule to stretch: a study of whiplash pain mechanism. *Stapp Car Crash J*. 2005; 49:49–65. [PubMed: 17096268]
46. Lu Y, Chen C, Kallakuri S, et al. Development of an in vivo method to investigate biomechanical and neurophysiological properties of spine facet joint capsules. *Eur Spine J*. 2005; 14:565–72. [PubMed: 15690211]
47. Quinn KP, Dong L, Golder FJ, et al. Neuronal hyperexcitability in the dorsal horn after painful facet joint injury. *Pain*. 2010; 151:414–21. [PubMed: 20739124]
48. Kallakuri S, Singh A, Lu Y, et al. Tensile stretching of cervical facet joint capsule and related axonal changes. *Eur Spine J*. 2008; 17:556–63. [PubMed: 18080147]
49. Lee KE, Davis MB, Mejilla RM, et al. In vivo cervical facet capsule distraction: mechanical implications for whiplash and neck pain. *Stapp Car Crash J*. 2004; 48:373–95. [PubMed: 17230274]
50. Lee KE, Davis MB, Winkelstein BA. Capsular ligament involvement in the development of mechanical hyperalgesia after facet joint loading: behavioral and inflammatory outcomes in a rodent model of pain. *J Neurotrauma*. 2008; 25:1383–93. [PubMed: 19061382]
51. Lee KE, Thinnies JH, Gokhin DS, et al. A novel rodent neck pain model of facet-mediated behavioral hypersensitivity: implications for persistent pain and whiplash injury. *J Neurosci Methods*. 2004; 137:151–9. [PubMed: 15262055]
52. Lee KE, Winkelstein BA. Joint distraction magnitude is associated with different behavioral outcomes and substance P levels for cervical facet joint loading in the rat. *J Pain*. 2009; 10:436–45. [PubMed: 19327645]
53. Dong L, Odeleye AO, Jordan-Sciutto KL, et al. Painful facet joint injury induces neuronal stress activation in the DRG: implications for cellular mechanisms of pain. *Neurosci Lett*. 2008; 443:90–4. [PubMed: 18675314]
54. Sundararajan S, Prasad P, Demetropoulos CK, et al. Effect of head-neck position on cervical facet stretch of post mortem human subjects during low speed rear end impacts. *Stapp Car Crash J*. 2004; 48:331–72. [PubMed: 17230273]
55. Quinn KP, Lee KE, Ahaghotu CC, et al. Structural changes in the cervical facet capsular ligament: potential contributions to pain following subfailure loading. *Stapp Car Crash J*. 2007; 51:169–87. [PubMed: 18278597]
56. Quinn KP, Bauman JA, Crosby ND, et al. Anomalous fiber realignment during tensile loading of the rat facet capsular ligament identifies mechanically induced damage and physiological dysfunction. *J Biomech*. 2010; 43:1870–5. [PubMed: 20381048]

57. Dong L, Winkelstein BA. Simulated whiplash modulates expression of the glutamatergic system in the spinal cord suggesting spinal plasticity is associated with painful dynamic cervical facet loading. *J Neurotrauma*. 2010; 27:163–74. [PubMed: 19772459]
58. Markowitz AJ, White MG, Kolson DL, et al. Cellular interplay between neurons and glia: toward a comprehensive mechanism for excitotoxic neuronal loss in neurodegeneration. *Cellscience*. 2007; 4:111–46. [PubMed: 19122795]
59. Winkelstein BA, Santos DG. An intact facet capsular ligament modulates behavioral sensitivity and spinal glial activation produced by cervical facet joint tension. *Spine*. 2008; 33:856–62. [PubMed: 18404104]
60. Svensson MY, Aldman B, Bostrom O, et al. Transient pressure gradients in the pig spinal canal during experimental whiplash motion causing membrane dysfunction in spinal ganglion nerve cells. *Orthopade*. 1998; 27:820–6. [PubMed: 9894236]
61. Aldman, B. An analytical approach to the impact biomechanics of head and neck injury. *Proceedings of the 39th American Association for Automotive Medicine Conference*; Des Plaines, IA. 1986. p. 439-61.
62. Eichberger A, Darok M, Steffan H, et al. Pressure measurements in the spinal canal of post-mortem human subjects during rear-end impact and correlation of results to the neck injury criterion. *Accid Anal Prev*. 2000; 32:251–60. [PubMed: 10688481]
63. Bostrom O, Fredriksson R, Haland Y, et al. Comparison of car seats in low speed rear-end impacts using the BioRID dummy and the new neck injury criterion (NIC). *Accid Anal Prev*. 2000; 32:321–28. [PubMed: 10688488]
64. Hubbard RD, Chen Z, Winkelstein BA. Transient cervical nerve root compression modulates pain: Load thresholds for allodynia and sustained changes in spinal neuropeptide expression. *J Biomech*. 2008; 41:677–85. [PubMed: 17976629]
65. Rothman SM, Kreider RA, Winkelstein BA. Spinal neuropeptide responses in persistent and transient pain following cervical nerve root injury. *Spine*. 2005; 30:2491–6. [PubMed: 16284585]
66. Hubbard RD, Winkelstein BA. Dorsal root compression produces myelinated axonal degeneration near the biomechanical thresholds for mechanical behavioral hypersensitivity. *Exp Neurol*. 2008; 212:482–9. [PubMed: 18572166]
67. Kobayashi S, Baba H, Uchida K, et al. Effect of mechanical compression on the lumbar nerve root: Localization and changes of intradiscal inflammatory cytokines, nitric oxide, and cyclooxygenase. *Spine*. 2005; 30:1699–705. [PubMed: 16094269]
68. Hubbard RD, Quinn KP, Martinez JJ, et al. The role of graded nerve root compression on axonal damage, neuropeptide changes, and pain-related behaviors. *Stapp Car Crash J*. 2008; 52:33–58. [PubMed: 19085157]
69. Rutkowski MD, Winkelstein BA, Hickey WF, et al. Lumbar nerve root injury induces central nervous system neuroimmune activation and neuroinflammation in the rat: relationship to painful radiculopathy. *Spine*. 2002; 27:1604–13. [PubMed: 12163719]
70. Kawakami M, Hashizume H, Nishi H, et al. Comparison of neuropathic pain induced by the application of normal and mechanically compressed nucleus pulposus to lumbar nerve roots in the rat. *J Orthop Res*. 2003; 21:535–9. [PubMed: 12706028]
71. Winkelstein BA, DeLeo JA. Nerve root injury severity differentially modulates spinal glial activation in a rat lumbar radiculopathy model: considerations for persistent pain. *Brain Res*. 2002; 956:294–301. [PubMed: 12445698]
72. Obata K, Yamanaka H, Fukuoka T, et al. Contribution of injured and uninjured dorsal root ganglion neurons to pain behavior and the changes in gene expression following chronic constriction injury of the sciatic nerve in rats. *Pain*. 2003; 101:65–77. [PubMed: 12507701]
73. Sekiguchi M, Kikuchi S, Myers RR. Experimental spinal stenosis - Relationship between degree of cauda equina compression, neuropathology, and pain. *Spine*. 2004; 29:1105–11. [PubMed: 15131438]
74. Hubbard RD, Winkelstein BA. Transient cervical nerve root compression in the rat induces bilateral forepaw allodynia and spinal glial activation: Mechanical factors in painful neck injuries. *Spine*. 2005; 30:1924–32. [PubMed: 16135981]

75. Rothman SM, Nicholson KJ, Winkelstein BA. Time-dependent mechanics and measures of glial activation and behavioral sensitivity in a rodent model of radiculopathy. *J Neurotrauma*. 2010; 37:803–14. [PubMed: 20121422]
76. Rothman SM, Winkelstein BA. Chemical and mechanical nerve root insults induce differential behavioral sensitivity and glial activation that are enhanced in combination. *Brain Res*. 2007; 1181:30–43. [PubMed: 17920051]
77. Olmarker K, Rydevik B, Holm S. Edema formation in spinal nerve roots induced by experimental, graded compression - an experimental study on the pig cauda-equina with special reference to differences in effects between rapid and slow onset of compression. *Spine*. 1989; 14:569–73. [PubMed: 2546258]
78. Kobayashi S, Yoshizawa H. Effect of mechanical compression on the vascular permeability of the dorsal root ganglion. *J Orthop Res*. 2002; 20:730–9. [PubMed: 12168661]
79. Kobayashi S, Yoshizawa H, Hachiya Y, et al. Vasogenic edema induced by compression injury to the spinal nerve root - distribution of intravenously injected protein tracers and gadolinium-enhanced magnetic resonance imaging. *Spine*. 1993; 18:1410–24. [PubMed: 8235811]
80. Winkelstein BA, Rutkowski MD, Sweitzer SM, et al. Nerve injury proximal or distal to the DRG induces similar spinal glial activation and selective cytokine expression but differential behavioral responses to pharmacologic treatment. *J Comp Neurol*. 2001; 439:127–39. [PubMed: 11596043]
81. Rothman SM, Huang Z, Lee KE, et al. Cytokine mRNA expression in painful radiculopathy. *J Pain*. 2009; 10:90–9. [PubMed: 18848809]
82. Li YB, Dorsi MJ, Meyer RA, et al. Mechanical hyperalgesia after an L5 spinal nerve lesion in the rat is not dependent on input from injured nerve fibers. *Pain*. 2000; 85:493–502. [PubMed: 10781924]
83. Obata K, Yamanaka H, Dai Y, et al. Contribution of degeneration of motor and sensory fibers to pain behavior and the changes in neurotrophic factors in rat dorsal root ganglion. *Exp Neurol*. 2004; 188:149–60. [PubMed: 15191811]
84. Hashizume H, DeLeo JA, Colburn RW, et al. Spinal glial activation and cytokine expression after lumbar root injury in the rat. *Spine*. 2000; 25:1206–17. [PubMed: 10806496]
85. Winkelstein BA, DeLeo JA. Mechanical thresholds for initiation and persistence of pain following nerve root injury: mechanical and chemical contributions at injury. *J Biomech Eng*. 2004; 126:258–63. [PubMed: 15179857]
86. Sapunar D, Modric-Jednacak K, Grkovic I, et al. Effect of peripheral axotomy on pain-related behavior and dorsal root ganglion neurons excitability in NPY transgenic rats. *Brain Res*. 2005; 1063:48–58. [PubMed: 16259969]
87. Hogan QH. Role of decreased sensory neuron membrane calcium currents in the genesis of neuropathic pain. *Croat Med J*. 2007; 48:9–21. [PubMed: 17309135]
88. Rothman SM, Guarino BB, Winkelstein BA. Spinal microglial proliferation is evident in a rat model of painful disc herniation both in the presence of behavioral hypersensitivity and following minocycline treatment sufficient to attenuate allodynia. *J Neurosci Res*. 2009; 87:2709–17. [PubMed: 19382225]
89. Sturzenegger M, Radanov BP, DiStefano G. The effect of accident mechanisms and initial findings on the long-term course of whiplash injury. *J Neurol*. 1995; 242:443–9. [PubMed: 7595675]
90. Winkelstein BA, Weinstein JN, DeLeo JA. The role of mechanical deformation in lumbar radiculopathy: an in vivo model. *Spine*. 2002; 27:27–33. [PubMed: 11805632]
91. Ellenberg MR, Honet JC, Treanor WJ. Cervical radiculopathy. *Arch Phys Med Rehabil*. 1994; 75:342–52. [PubMed: 8129590]
92. Miles KA, Maimaris C, Finlay D, et al. The incidence and prognostic significance of radiological abnormalities in soft tissue injuries to the cervical spine. *Skeletal Radiol*. 1988; 17:493–6. [PubMed: 3201276]
93. Maimaris C, Barnes MR, Allen MJ. 'Whiplash injuries' of the neck: a retrospective study. *Injury*. 1988; 19:393–6. [PubMed: 3267643]
94. Sterling M, Jull G, Kenardy J. Physical and psychological factors maintain long-term predictive capacity post-whiplash injury. *Pain*. 2006; 122:102–8. [PubMed: 16527397]

95. Max MB, Stewart WF. The molecular epidemiology of pain: a new discipline for drug discovery. *Nat Rev Drug Discov.* 2008; 7:647–58. [PubMed: 18587382]
96. LaCroix-Fralish ML, Rutkowski MD, Weinstein JN, et al. The magnitude of mechanical allodynia in a rodent model of lumbar radiculopathy is dependent on strain and sex. *Spine.* 2005; 30:1821–7. [PubMed: 16103850]
97. Lord SM, Barnsley L, Wallis BJ, et al. Chronic cervical zygapophysial joint pain after whiplash. A placebo-controlled prevalence study. *Spine.* 1996; 21:1737–44. [PubMed: 8855458]

Key Points

1. Animal models provide utility for defining injury and pain mechanisms in whiplash.
2. Mechanisms by which pain becomes chronic involve the integration of peripheral and central neuronal and immune responses.
3. The development of chronic pain from whiplash injury depends on a variety of factors.
4. Animal modeling of painful injuries, like whiplash, must be interpreted carefully in the context of the goal of modeling the injury event or the resulting symptoms.

Figure 1.

Maps of the regions of referred pain in the human and evoked-pain in the rat enable comparisons between these two species and interpretation of common behavioral tests of sensitivity in the rodent in the context of clinical symptoms in the human. **Top Left.** The distribution of spontaneous whiplash pain in human patients, modified from Lord et al. [97].

Top Right. Distributions of evoked-pain regions are also observed in the rat indicating the similar “coathanger” distribution. **Bottom Right.** Dermatomes indicating spinal nerve innervations of the forepaw of the rat.

Figure 2.

Trauma in the periphery induces the release of chemical mediators that can activate local nociceptors, lead to spinal sensitization, and produce pain. Nociceptor activation leads to neurotransmitter and neurotrophin synthesis in the DRG. In addition, local inflammation involves the recruitment and stimulation of many non-neuronal cells to release inflammatory mediators that modulate neuronal excitability. Both of these neuronal and immune responses can induce central sensitization in the spinal cord that increases the excitability of neurons in the dorsal horn. Spinal glial cells are activated and release inflammatory mediators that further potentiate the neuronal and glial responses, contributing to persistent pain.

Figure 3.

(a) Representative extracellular voltage recordings and neuronal firing rate of spinal neurons during repeated stimulation of the forepaw with a 26g filament following a painful facet capsule injury (distraction) or sham. (b) Evoked spinal neuronal firing is significantly (*) elevated after painful injury. (c) Neuronal firing upon noxious paw stimulation is strongly correlated with the magnitude of stimulus to elicit paw withdrawal (a lower threshold is more sensitivity).