Published in final edited form as:

J Phys Chem B. 2011 December 15; 115(49): 14784–14788. doi:10.1021/jp206486z.

Changes in Apparent Molar Water Volume and DKP Solubility Yield Insights on the Hofmeister Effect

Alexander Y. Payumo, R. Michael Huijon, Deauna D. Mansfield, Laurel M. Belk, Annie K. Bui, Anne E. Knight, and Daryl K. Eggers*

Department of Chemistry, San José State University, San José, California 95192-0101 USA

Abstract

This study examines the properties of a 4×2 matrix of aqueous cations and anions at concentrations up to 8.0 M. The apparent molar water volume, as calculated by subtracting the mass and volume of the ions from the corresponding solution density, was found to exceed the molar volume of ice in many concentrated electrolyte solutions, underscoring the non-ideal behavior of these systems. The solvent properties of water were also analyzed by measuring the solubility of diketopiperazine (DKP) in 2.000 M salt solutions prepared from the same ion combinations. Solution rankings for DKP solubility were found to parallel the Hofmeister series for both cations and anions, whereas molar water volume concurred with the cation series only. The results are discussed within the framework of a desolvation energy model that attributes solute-specific changes in equilibria to solute-dependent changes in the free energy of bulk water.

Keywords

Non-ideal solution; apparent water density; electrolytes; backbone hydration; desolvation energy

1. INTRODUCTION

The interior of a living cell is as an extremely crowded environment where macromolecules occupy over 30% of the total solution volume.^{1,2} If one also considers the presence of smaller solutes and the fact that biomacromolecules are polyelectrolytes,³ it becomes evident that life's vital reactions take place in a highly non-ideal solution, in marked contrast to the dilute experimental conditions in which the properties of the individual reactants are typically characterized. In addition to excluded volume effects on reaction rates and equilibria due to macromolecular crowding,^{4,5} the innate properties of water may be altered in biological systems, although this issue is rarely addressed in the literature.⁶ New biophysical methods, such as in-cell NMR spectroscopy,^{7,8} have been developed for examining proteins in their natural environment, but such techniques may not be feasible or adequate for characterizing all biomolecules of interest, including the solvent.

To gain a better understanding of molecular activity in vivo, the properties of model compounds and model reactions may be determined in aqueous, non-ideal solutions of defined composition. In the current work, solutions of concentrated electrolytes are examined to quantify two properties of water: the apparent molar water volume and the

*Corresponding Author Phone: 408-924-4960. Fax: 408-924-4945. daryl.eggers@sjsu.edu..

ASSOCIATED CONTENT

Supporting Information. Solution densities with corresponding values of ϕV_l (Tables S1-S3), DKP solubility in 2 M solutions (Table S4), and a critical analysis of partial molar volumes from apparent molar volumes. This material is available free of charge via the Internet at <http://pubs.acs.org>.

maximum solubility of diketopiperazine. Diketopiperazine (DKP), also referred to as cyclic diglycine or glycine anhydride, is an amide containing compound with the same bond connectivity as that found in the backbone of a protein. Electrolyte solutions are interesting for these studies because specific ions have been shown to influence the properties of biomolecules in a recurring order, known to biochemists as the Hofmeister series.^{9,10} Although it has been hypothesized often that ion-specific effects are mediated through changes in the properties of water, the underlying basis for the Hofmeister series is still highly debated,¹¹ and a single theory may not capture all nuances of this complicated biophysical problem. In vitro studies with non-ideal solutions should facilitate a deeper understanding of water's role in cell biology, and a thoughtful selection of electrolytes for such studies may also yield insights on the forces that drive the rankings of the Hofmeister series.

2. EXPERIMENTAL METHODS

Stock solutions of each salt were made at room temperature from ultrapure water (Millipore, Milli-Q system) and the following reagents: LiCl (99%, Fisher), KCl (99%, Sigma), CsCl (99.99%, Acros), N(CH₃)₄Cl (97%, Aldrich), LiC₂H₃O₂ (dihydrate, 98%, Acros), KC₂H₃O₂ (99%, Fisher), CsC₂H₃O₂ (99.9%, Aldrich), N(CH₃)₄C₂H₃O₂ (hydrate, 95%, Acros), and LiClO₄ (trihydrate, 99%, Acros). For solubility experiments, solutions were supplemented with 0.010 M Tris hydrochloride prepared from a 1.00 M stock buffer solution of pH 7.4 (Sigma, T2663). The final pH value of each salt solution was not adjusted. Diketopiperazine was obtained from Sigma (G7251).

Stock solutions were diluted with ultrapure water to obtain final electrolyte concentrations in the range of 0.2500 to 8.000 M, depending on the specific salt and its maximum solubility. All solutions were analyzed at atmospheric pressure and at four temperatures, $T = 278.15$, 298.15, 310.15, and 323.15 K. Solutions were maintained in capped vials at the desired air temperature for several hours in a closed incubator prior to density measurements at the same temperature using a high-precision oscillating U-tube density meter (model DMA 5000, Anton Parr). This instrument has a rated uncertainty of $\pm 1 \times 10^{-6} \text{ g}\cdot\text{cm}^{-3}$ and a temperature uncertainty of ± 0.001 degrees. The density meter was calibrated with air and ultrapure water, following the manufacturer's protocol. The apparent molar water volume of each solution was calculated with the following equation:

$$\varphi V_1 = \frac{M_1 \left[1000 - C_2 (4/3) \pi (r_a^3 + r_b^3) N_A \right]}{1000\rho - M_2 C_2} \quad (1)$$

where φV_1 is the apparent molar water volume in units of $\text{cm}^3\cdot\text{mol}^{-1}$, r_a and r_b are the Pauling radii of the cation and anion in centimeters, N_A is Avogadro's number, C_2 is the salt concentration in molarity, ρ is the solution density in $\text{g}\cdot\text{cm}^{-3}$, M_1 is the molecular mass of water ($18.015 \text{ g}\cdot\text{mol}^{-1}$), and M_2 is the formula mass of the anhydrous salt in $\text{g}\cdot\text{mol}^{-1}$. Values for the ionic radii were taken from Marcus¹² as listed here in Ångstroms: $\text{Li}^+ = 0.60$; $\text{K}^+ = 1.33$; $\text{Cs}^+ = 1.69$; $\text{N}(\text{CH}_3)_4^+ = 2.80$; $\text{Cl}^- = 1.81$; $\text{ClO}_4^- = 2.36$; $\text{CH}_3\text{COO}^- = \text{OAc}^- = 2.32$.

Diketopiperazine solubilities in 2.000 M salt solutions were determined by density analysis, using the density meter described above. For a given experiment, 8-10 vials were prepared with 3.50 cm^3 of salt solution in each vial and increasing amounts of DKP. The mass range of DKP addition was selected such that three or more vials were below the saturation point and three or more vials were above the saturation point. The vials were placed in an incubator at the desired air temperature and stirred on a rotary mixing device at low speed (Barnstead International, Labquake model 400110). Samples were allowed to equilibrate for

36-48 hours before analysis. Each sample was loaded into the inlet port of the density meter through a syringe filter of 0.22 μ m pore size (Millipore, SLGP033RB). DKP solubility was determined by plotting solution density versus milligrams of DKP added for each salt solution. The pre-saturation points were fitted to a linear equation of positive slope, the densities of the post-saturation points were averaged, and the intersection of the two regimes was interpreted as the saturation point.^{13,14} The concentration of DKP at the saturation point in units of g DKP per 100 g of water may be converted from mg DKP added per cm³ of initial solution volume using the following expression:

$$C_i^\Phi = \frac{10 \cdot c^\Phi}{\rho_0 - (M_2C_2 + M_3C_3) / 1000} \quad (2)$$

where C_i^Φ is the solubility of DKP in solution i in units of g·(100g H₂O)⁻¹, c^Φ is the solubility of DKP in units of mg·cm⁻³, ρ_0 is the original density of the electrolyte solution before addition of DKP, M_2C_2 refers to the formula mass and concentration of the electrolyte, and M_3C_3 refers to the formula mass (157.6 g·mol⁻¹) and concentration (0.010 M) of the Tris buffer. The original density value is employed, as opposed to the density at saturation, because the total volume of solution used to calculate c^Φ and the precise concentration of electrolyte are known only before addition of DKP.

3. RESULTS AND DISCUSSION

In this study, a 4 × 2 matrix of salt combinations was examined; Li⁺, K⁺, Cs⁺, and N(CH₃)₄⁺ were paired with two anions, chloride and acetate (OAc⁻). In addition, lithium perchlorate was added to the solution list to provide a third anion for testing; this salt was deemed most appropriate for comparison of the anions because lithium is one of few cations that permits the preparation of perchlorate solutions above 1.0 M concentration. The ions were selected to encompass a broad range of effects based on their relative positions within the Hofmeister series. The densities of the nine electrolyte combinations were determined for a range of solute concentrations at atmospheric pressure and at four temperatures: $T = 278.15, 298.15, 310.15,$ and 323.15 K (see Tables S1-S3 of Supporting Information).

3.1 Apparent Molar Water Volume

The measured solution density values were converted to apparent molar water volumes, ${}^\phi V_1$, by subtracting the known volume and mass of each ion in each solution (Equation 1). This calculation, which treats the volume of the solute as a constant and allows the density of water to fluctuate, is a departure from the traditional approach that treats the density of water as a constant and attributes changes in solution volume to changes in the apparent molar volume of the solute, ${}^\phi V_2$.¹⁵ Apparent molar solute volumes and their use in calculation of partial molar volumes, $\bar{V}_1$ and $\bar{V}_2$, are examined and discussed in the Supporting Information.

A sampling of ${}^\phi V_1$ results for the 4 × 2 matrix is given in Figures 1 and 2. The apparent molar water volumes are presented as a function of salt concentration at one temperature, 310.15 K (Figure 1), and as a function of temperature at one concentration, 2.000 M (Figure 2). Interestingly, the apparent molar water volume of nearly all salt solutions in this study exceeds the molar volume of neat water at the same temperature, and, in several solutions of high salt concentration, the apparent molar water volume exceeds the molar volume of ice, 19.65 cm³·mol⁻¹ at 273.15 K (Figure 1).

With regard to the cations, a direct concordance was observed between the Hofmeister series and ${}^\phi V_1$. For each temperature, concentration, and anion pairing that was examined, the

apparent molar water volume increased in the order $\text{Li}^+ < \text{K}^+ < \text{Cs}^+ < \text{N}(\text{CH}_3)_4^+$. This ranking corresponds with an increase in radius and with a decrease in the effective surface charge density of each cation. For the anions, the results indicate that each chloride solution has a lower ${}^\phi V_I$ value than the corresponding acetate solution when paired with the same cation (compare Figure 1a, 1b). This outcome is consistent with the effective charge density of the two anions, *i.e.* the ion of highest charge density (Cl^-) resulted in the lowest value of ${}^\phi V_I$.

More insight on anion-specific effects is obtained by comparing the apparent molar water volume of solutions made from lithium salts of chloride, acetate, and perchlorate at 310.15 K (Figure 3). When the solutions are compared on a basis of equal ion numbers per unit volume, the solute-dependent value of ${}^\phi V_I$ increases in the order $\text{Cl}^- < \text{OAc}^- < \text{ClO}_4^-$ at all concentrations. It is important to note that this result is not consistent with the Hofmeister rankings of anions; acetate and perchlorate are found at opposite ends of the anion series with chloride positioned near the center. Thus, the anion-induced changes in water structure that lead to changes in apparent molar water volume do not correlate with the Hofmeister effect. Others have also observed a lack of correlation between water structure and the Hofmeister series using pressure perturbation calorimetry¹⁶ and nonlinear optical spectroscopy of Langmuir monolayers.¹⁷

Except for the specific case of LiCl at 0.500 M, the value of ${}^\phi V_I$ was found to be greater than the corresponding volume of pure water at the same temperature in all solutions. At first thought, this result might be alarming because many ions are known to cause electrostriction of the solvent, yielding a molar volume less than neat water. It should be noted, however, that the phenomenon of electrostriction is applicable only to dilute solutions. As pointed out by Marcus, the average center-to-center ion spacing of a homogeneously-dispersed solution of spherical ions may be approximated by $0.940c^{-1/3}$ nm, where c is the salt concentration in molarity.^{18,19} After subtracting the radii of the anion and cation, there may be space for only a few water molecules in solutions above 1.0 M concentration. For example, in a solution of 6.0 M CsCl, for which the ionic diameters are 0.169 nm for Cs^+ and 0.181 nm for Cl^- , the available hydration space is $0.517 - (0.169 + 0.181)/2 = 0.342$ nm, corresponding to slightly more than one water molecule between ions. Thus, for most of the non-ideal solutions reported here, water molecules are under the constant influence of one or more ions at any given instant in time. Under these conditions, the structural properties of water are expected to differ drastically from that observed in neat solution. Presumably, the tug-of-war between ions for waters of solvation contributes to the non-ideal behavior of concentrated electrolyte solutions, as exemplified by the observation that ${}^\phi V_I$ may exceed the value of ice (Figure 1).

Apparent molar water volumes are not commonly reported in the literature. Dougherty has calculated a reciprocally-related parameter, the apparent water density, by subtracting the mass and volume of the electrolyte from the solution density.²⁰ The Dougherty study includes CsCl in a graphical presentation of the data, from which one may estimate an apparent molar water density of $0.963 \text{ g}\cdot\text{cm}^{-3}$ for 2.0 M CsCl at 298 K. Taking the reciprocal of this value and multiplying by the molecular mass of water yields an apparent molar water volume of $18.7 \text{ cm}^3\cdot\text{mol}^{-1}$, in close agreement with the value of $18.65 \text{ cm}^3\cdot\text{mol}^{-1}$ reported here (Table S1). The Dougherty analysis was limited to solutions up to 3.0 M concentration and employed ionic radii from crystal data that differ slightly from the radii used in the current work.

3.2 Diketopiperazine Solubility

Solubility experiments with DKP were performed for a control solution of low ionic strength (0.010 M Tris buffer) and for 2.000 M electrolyte solutions using density measurements to

ascertain the saturation point. Initial work was completed at the physiological temperature of $T = 310.15$ K for all salts in the 4×2 matrix. In general, DKP solubility followed the rankings of the Hofmeister series of cations; at $T = 310.15$ K, solubility decreased in the order $\text{Li}^+ > \text{K}^+ \approx \text{Cs}^+ > \text{N}(\text{CH}_3)_4^+$ for both the chloride and acetate solutions (Figure 4). It is often noted in the literature that the cation series of Hofmeister effects is harder to discern than the anion series, but the cations in this study yield clear and reproducible changes in the solubility of DKP. A similar ranking of cations, in reverse order relative to highest DKP solubility, was deduced for solutions that enhance the structure of apomyoglobin confined to the pores of a silica glass matrix.²¹

Solubility experiments were expanded to $T = 278.15$, 298.15 , and 323.15 K for the control solution and for 2.000 M solutions of lithium chloride, lithium acetate, and lithium perchlorate. The DKP solubility results for the 2.000 M lithium solutions are summarized in Figure 5. For a given temperature, the solubility of DKP decreases in accord with the Hofmeister series of anions such that $\text{ClO}_4^- \gg \text{Cl}^- > \text{water} > \text{OAc}^-$. The high solubility of DKP in lithium perchlorate solutions is consistent with the denaturing effect of perchlorate on protein structure. Because the amide units of the polypeptide backbone are normally buried in the core of a folded protein and not in contact with solvent, an increase in DKP solubility is analogous to the increase in exposure of the amide units of a protein in the unfolded state.

A solubility of 1.69 g per 100 g of H_2O , as reported in Table S4 for DKP in water at 298.15 K, is in excellent agreement with a value of 1.68 g per 100 g of solvent, as obtained by density measurements and reported independently by the groups of Bolen^{13,22} and Lee.^{14,23} Most previous studies on DKP solubility have focused on urea or osmolyte solutions. The Lee group has measured DKP solubility in 2.0 M electrolyte solutions of KBr, KCl, and KOAc at 298.15 K.²³ The DKP solubilities reported in the current work for KCl and KOAc solutions were carried out at 310.15 K, precluding a direct comparison, but it should be noted that the Lee group also found a correlation with the Hofmeister series of anions; DKP solubility decreased in the order $\text{Br}^- > \text{Cl}^- > \text{OAc}^-$ for all concentrations of potassium salts up to 4.0 M concentration.²³

The solubility approach was also employed in the classical works of Robinson and Jencks who tested acetyl tetraglycine ethyl ester as a model compound for the backbone of a protein.^{24,25} The rankings of salt effects on the solubility of the tetraglycine derivative are in agreement with the rankings reported here for DKP (Figures 4-5, Table S4).

3.3 Relationship between Solubility and Desolvation Energy

Recently, our laboratory proposed an equation for relating solubility of a model compound in two or more solutions to the corresponding desolvation energies:²⁶

$$RT \ln \left(\frac{C_A^\Phi}{C_B^\Phi} \right) = \hat{n} \left(C_A^\Phi \Delta \bar{G}_A^{H_2O} - C_B^\Phi \Delta \bar{G}_B^{H_2O} \right) \quad (3)$$

where C_i^Φ is the saturation concentration of the model compound (Φ) in moles per mole of water, subscripts A and B denote the two solutions, $\hat{n}$ is the moles of water that solvate one molecule of Φ , and $\Delta \bar{G}_i^{H_2O}$ is the molar desolvation energy for Φ in solution i . The desolvation energy term is defined by the following relationship:

$$\Delta \bar{G}_i^{H_2O} = \bar{G}_i^{bulk} - \bar{G}^{solv} \quad (4)$$

where $\bar{G}_i^{bulk}$ represents the average molar free energy of the bulk water in solution i and $\bar{G}^{solv}$ is the average molar free energy of the $\hat{n}$ water molecules in the solvation sphere of model compound Φ that are displaced upon precipitation. The value of $\bar{G}^{solv}$ is a function of the surface chemistry of the model compound, and the value of $\bar{G}_i^{bulk}$ is a function of all solutes in solution i , including secondary solutes that do not participate directly in the reaction of interest.²⁶ Because water molecules are in a dynamic equilibrium between multiple subpopulations of differing energy, as defined by the chemical boundaries of all solutes and surfaces in contact with the solvent, it seems reasonable to treat the thermodynamics of bulk water as a number-weighted average of all of the subpopulations. With this thermodynamic framework in mind, the Hofmeister effect is expected to correlate with changes in the average free energy of the bulk aqueous phase, and it is not surprising that the measurement of a specific structure-related property of water does not always follow the Hofmeister series of ions. Reaction equilibria are dictated by changes in free energy, and, therefore, it seems logical that the effects of electrolytes, and secondary solutes in general, are realized through their effects on the free energy of bulk water.

In the context of the current work, an electrolyte that increases the free energy of bulk water will result in a more-positive desolvation energy, increase the solubility of DKP, and destabilize protein structure. Conversely, an electrolyte that decreases the free energy of bulk water will correspond to a more-negative desolvation energy, decrease DKP solubility, and enhance protein structure. Thus, based on differences in DKP solubility, $N(CH_3)_4Cl$ and all acetate solutions in this study should be stabilizers, $LiCl$ should be a weak destabilizer, $LiClO_4$ should be a strong destabilizer, and both KCl and $CsCl$ should have small or negligible effects on protein structure at 310.15 K (Figures 4-5). Further experiments are required to obtain the solute-specific free energy terms given in Equations 3 and 4.

CONCLUSIONS

The apparent molar water volume was calculated for solutions of four chloride salts, four acetate salts, and one perchlorate salt at high molar concentrations. The results indicate that φV_l increases with increasing solute concentration and is greater in magnitude than pure water at the same temperature for all solutions above 1 M concentration. At the highest electrolyte concentrations, φV_l exceeds the molar volume of ice for several ion pairings, underscoring the non-ideal behavior of these systems. The solution rankings for apparent molar water volume were consistent with the cation series of Hofmeister ions but not with the anion series, supporting the idea that the Hofmeister effect is not a function of any specific structural feature of water but, rather, follows a thermodynamic quantity, such as the change in the Gibbs free energy of bulk water, that accompanies all changes in water structure.

The solvent properties of the electrolyte solutions in this study were compared using DKP as a model compound. The highest solubility was found in lithium perchlorate solutions, consistent with the protein-denaturing property of perchlorate. For all cations and anions examined, the DKP solubility rankings followed the Hofmeister effect. This result is expected if changes in solubility are governed by changes in the free energy of bulk water and if changes in the free energy of bulk water follow the Hofmeister series of ions. Further characterization of water in non-ideal solutions may lead to a better understanding of hydration effects on reaction equilibria in biological systems, including binding equilibria and protein folding.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was funded by the National Institutes of Health (SC3 GM089591), as initiated under a previous NIH subproject award to D.K.E. (S06 GM008192). A.Y.P. was supported by an NIH MARC scholarship for undergraduate research (T34 GM008253).

REFERENCES

1. Zimmerman SB, Trach SO. *J. Mol. Biol.* 1991; 222:599–620. [PubMed: 1748995]
2. Cayley S, Lewis BA, Guttman HJ, Record MT Jr. *J. Mol. Biol.* 1991; 222:281–300. [PubMed: 1960728]
3. Record MT, Courtenay ES, Cayley S, Guttman HJ. *Trends Biochem. Sci.* 1998; 23:190–194. [PubMed: 9612084]
4. Zimmerman SB, Minton AP. *Annu. Rev. Biophys. Biomol. Struct.* 1993; 22:27–65. [PubMed: 7688609]
5. Zhou H-X, Rivas G, Minton AP. *Annu. Rev. Biophys.* 2008; 37:375–397. [PubMed: 18573087]
6. Ball P. *ChemPhysChem.* 2008; 9:2677–2685. [PubMed: 18855966]
7. Crowley PB, Chow E, Papkovskaia T. *ChemBioChem.* 2011; 12:1043–1048. [PubMed: 21448871]
8. Schlesinger AP, Wang Y, Tadeo X, Millet O, Pielak GJ. *J. Am. Chem. Soc.* 2011; 133:8082–8085. [PubMed: 21534566]
9. Collins KD, Washabaugh MW. *Quarterly Rev. Biophys.* 1985; 18:323–422.
10. Collins KD. *Biophys. J.* 1997; 72:65–76. [PubMed: 8994593]
11. Tobias DJ, Hemminger JC. *Science.* 2008; 319:1197–1198. [PubMed: 18309069]
12. Marcus, Y. *Ion Properties.* Marcel Dekker, Inc; New York: 1997.
13. Liu Y, Bolen DW. *Biochemistry.* 1995; 34:12884–12891. [PubMed: 7548045]
14. Venkatesu P, Lee M-J, Lin H. *J. Phys. Chem. B.* 2007; 111:9045–9056. [PubMed: 17602514]
15. Høiland, H. *Chp. 2 of Thermodynamic Data for Biochemistry and Biotechnology.* Hinz, H-J., editor. Springer-Verlag; New York: 1986.
16. Batchelor JD, Olteanu A, Tripathy A, Pielak GJ. *J. Am. Chem. Soc.* 2004; 126:1958–1961. [PubMed: 14971928]
17. Gurai MC, Lim S-M, Castellana ET, Albertorio F, Kataoka S, Cremer PS. *J. Am. Chem. Soc.* 2004; 126:10522–10523. [PubMed: 15327293]
18. Marcus Y. *Chem. Rev.* 2009; 109:1346–1370. [PubMed: 19236019]
19. Marcus Y. *J. Solution Chem.* 2009; 38:513–516.
20. Dougherty RC. *J. Phys. Chem. B.* 2001; 105:4514–4519.
21. Eggers DK, Valentine JS. *J. Mol. Biol.* 2001; 314:911–922. [PubMed: 11734007]
22. Auton M, Bolen DW. *Biochemistry.* 2004; 43:1329–1342. [PubMed: 14756570]
23. Venkatesu P, Lee M-J, Lin H-M. *Biochem. Eng. J.* 2006; 32:157–170.
24. Robinson DR, Jencks WP. *J. Am. Chem. Soc.* 1965; 87:2470–2479. [PubMed: 14327155]
25. Nandi PK, Robinson DR. *J. Am. Chem. Soc.* 1972; 94:1299–1308. [PubMed: 5060273]
26. Eggers DK. *Biochemistry.* 2011; 50:2004–2012. [PubMed: 21284393]

Figure 1. Apparent molar water volume at 310.15 K as a function of electrolyte concentration for (a) the chloride salts, and (b) the acetate salts. The solid horizontal line of each panel denotes the molar volume of neat water at 310.15 K, and the dashed horizontal line denotes the molar volume of ice at 273.15 K.

Figure 2. Apparent molar water volume in 2.000 M salt solutions as a function of temperature for (a) the chloride salts, and (b) the acetate salts. The bottom curve of each panel denotes the molar volume of neat water at the corresponding temperature, as calculated from a standard water density table.

Figure 3.

Apparent molar water volume of lithium solutions at 310.15 K as a function of electrolyte concentration. Note that, for a given salt concentration, the apparent molar water volume increases in the order $\text{Cl}^- < \text{OAc}^- < \text{ClO}_4^-$, which deviates from the Hofmeister series of anions.

Figure 4. Diketopiperazine solubility in salt solutions relative to solubility in water at 310.15 K. For all solubility values in units of $\text{mg}\cdot\text{cm}^{-3}$ solution or $\text{g}\cdot(100\text{ g H}_2\text{O})^{-1}$, see Table S4.

Figure 5. Diketopiperazine solubility in water and in 2.000 M lithium salt solutions as a function of temperature.