Published in final edited form as:

Psychol Med. 2012 April ; 42(4): 717–726. doi:10.1017/S0033291711001814.

Longitudinal predictors of past-year non-suicidal self-injury and motives among college students

Holly C. Wilcox, PhD^{1,2}, Amelia M. Arria, PhD^{3,4}, Kimberly M. Caldeira, MS³, Kathryn B. Vincent, MA³, Gillian M. Pinchevsky, MA⁵, and Kevin E. O'Grady, PhD⁶

¹Department of Psychiatry and Behavioral Sciences, Johns Hopkins University, Baltimore, MD, USA

³Center on Young Adult Health and Development (CYAHD), University of Maryland School of Public Health, Department of Family Science, College Park, MD, USA

⁴Treatment Research Institute, Philadelphia, PA, USA

⁵Department of Criminology and Criminal Justice, University of South Carolina, Columbia, SC, USA

⁶Department of Psychology, University of Maryland, College Park, MD, USA

Abstract

Background—Non-suicidal self-injury (NSSI) is the deliberate and direct injuring of body tissue without suicidal intent for purposes not socially sanctioned. Few studies have examined the correlates of NSSI among young adults. This paper aimed to identify predictors of lifetime and past-year NSSI; describe motives for NSSI and disclosure of NSSI to others.

Method—Interviews were conducted annually with 1,081 students enrolled in the College Life Study, a prospective longitudinal study conducted at one large public mid-Atlantic university. NSSI characteristics were assessed at Year 4. Demographic and predictor variables were assessed during Years 1 through 4. Multivariate logistic regression models were used to identify correlates of lifetime NSSI and predictors of past-year NSSI.

Results—The prevalence of past-year and lifetime NSSI were 2% and 7%, respectively (>70% were female for both lifetime and past-year NSSI). Seven percent of NSSI cases self-injured once, whereas almost half self-injured six or more times. Independent predictors of past-year NSSI were maternal depression, non-heterosexual orientation, affective dysregulation, and depression. Independent predictors of lifetime NSSI were depression, non-heterosexual orientation, paternal depression, and female sex. One in six participants with NSSI attempted suicide by young adulthood. The three most commonly reported motives for NSSI were mental distress, coping, and situational stressors. Two-thirds (89%) told someone about their NSSI, most commonly a friend (68%).

Conclusions—This study identified unique predictors of NSSI, which should help elucidate its etiology and has implications for early identification and interventions.

²Corresponding Author: Holly C. Wilcox, The Johns Hopkins Hospital, 550 North Broadway, Room 206A, Baltimore, MD 21205, USA. Phone: 410-502-0629; Fax: 410-955-8691; hwilcox1@jhmi.edu.

Location of work: Center on Young Adult Health and Development, University of Maryland School of Public Health, 8400 Baltimore Avenue, Suite 100, College Park, MD 20740, USA.

Declaration of Interest

The investigators acknowledge funding from the National Institute on Drug Abuse (R01DA14845, Dr. Arria, PI) and the American Foundation for Suicide Prevention. All authors report no competing interests.

Keywords

college students; deliberate self-harm; NSSI; self-injurious behavior; young adults

Introduction

Non-suicidal self-injury (NSSI) is the direct and intentional destruction of one's body tissue without suicidal intent (Nock & Prinstein, 2004; Nock *et al.*, 2006; Shaffer & Jacobson, 2010). These behaviors range from self-cutting, scratching, and burning to implanting objects under the skin. While no national estimates of NSSI exist, prevalence estimates from individual community studies show that NSSI appears to be more common in adolescents [~15% (Ross & Heath, 2002; Muehlenkamp & Gutierrez, 2004)] and young adults [17% (Whitlock *et al.*, 2006)] than adult populations [4% (Briere & Gil, 1998) and 6% (Klonsky, 2011)].

The existing knowledge base regarding NSSI is limited in part because of methodological limitations of existing studies. For example, the majority of studies have used cross-sectional designs and clinical samples. The few prospective studies of NSSI with community samples did not sort out the temporality between constructs, were exploratory due to small sample size (Hankin & Abela, 2011), or focused on one developmental pathway (e.g., parental criticism on NSSI, peer influences on NSSI) without assessing a broad range of predictors (Yates *et al.*, 2008; Prinstein *et al.*, 2010). Studies of college students have utilized convenience samples from psychology courses without appropriate control groups and typically have low response rates. Additionally, the assessment of NSSI has varied widely between studies; many include wound-picking, which inflates the prevalence of NSSI. Lastly, most studies have not followed their sample past the peak period of risk for NSSI.

Moreover, studies have investigated a narrow range of potential correlates for NSSI based on models of adolescent suicidal behaviors, despite evidence that those who engage in NSSI without prior suicide attempts have different characteristics and psychiatric profiles from suicide attempters (Muehlenkamp & Gutierrez, 2004). Jacobson *et al.* (2008) found that the participants who engaged in NSSI without prior suicide attempts were more similar to their non-self-harming peers than their peers who had attempted suicide.

Several studies have examined individual-level characteristics that might place an individual at risk for NSSI in the face of adverse environmental or familial stressors. Stressful experiences such as family conflict and sexual or physical abuse are the most commonly cited environmental risk factors for NSSI (Weierich & Nock, 2008; Bureau *et al.*, 2010). Linehan (1993) proposed that the interaction of biologically-based vulnerability to intense emotion and early environments characterized by adversity and the stifling of emotional expression are associated with emotional dysregulation which can lead to maladaptive behaviors such as NSSI. Other important environmental risk factors for NSSI include interpersonal problems with family and peers, such as more conflict, less cohesion, and less support (Adrian *et al.*, 2011); alienation from parents; and parental criticism (Yates *et al.*, 2008). With respect to personal factors, Crowell *et al.* (2008) found more opposition/defiance and less positive affect among self-injuring adolescents as compared with adolescents who do not self-injure. The link between NSSI and depression is well established (Garrison *et al.*, 1993; Jacobson & Gould, 2007) but has not been adequately studied in prospective studies of community-residing young adults. In a small scale study, Hankin and Abela (2011) found that the onset of maternal depression predicted NSSI.

Albeit limited, studies of small clinical samples have suggested genetic or physiological mechanisms of NSSI linked to maladaptive stress responses (e.g., Kaess *et al.*, 2011). Deliberto and Nock (2008) found that those with NSSI had more family members with impulsivity-related psychopathology. Herpertz *et al.* (1997) found that NSSI was associated with impaired serotonergic function, a condition associated with impulsivity and aggression. Crowell *et al.* (2008) also found that adolescents with NSSI had lower levels of peripheral serotonin (5-HT); however, in adolescents with higher levels of 5-HT, parent-child negativity was associated with NSSI.

The motives for engaging in NSSI are critical for guiding treatment decisions and designing preventive interventions (Lloyd-Richardson *et al.*, 2007; Peterson *et al.*, 2008). The most commonly reported motive for NSSI in community samples is related to regulating negative affect states (Laye-Gindhu & Schonert-Reichl, 2005; Klonsky, 2009). This temporary relief from distress may reinforce NSSI and make repetition likely. Other motives for NSSI are to stop disassociation and to draw parental/peer attention (Nock & Prinstein, 2004), or as a means of self-punishment or eliciting care (Peterson *et al.*, 2008).

The purpose of the present study was to investigate a wide range of potential individual and familial unique predictors of NSSI among a large cohort of young adults originally sampled as part of the College Life Study, a longitudinal study of health-risk behaviors. Qualitative data were used to examine motives for NSSI and whether NSSI was disclosed to others.

Method

Participants and procedures

Sample selection took place in two stages. First, a screening survey was administered to 3,401 (response rate=89%) incoming first-time, first-year students ages 17 to 19, during new-student orientation in 2004 at one large, public university in the mid-Atlantic region of the U.S. Next, purposive sampling strategies were employed to oversample students who had used an illicit drug or nonmedically used a prescription drug at least once prior to college. This stratified random sample of screener participants was selected to participate in a series of annual follow-up assessments ($n=1,253$, response rate=87%). The sample was representative of the first-year class with respect to race, gender, and socioeconomic status (Arria *et al.*, 2008a). Sampling weights were computed to produce prevalence estimates that represent the general population of screened students; however, the results did not differ appreciably, so unweighted results are presented herein. Additional details regarding recruitment and sampling can be found elsewhere (Arria *et al.*, 2008a; Vincent *et al.*, under second review).

The baseline assessment consisted of self-administered questionnaires and an interview administered by a trained interviewer during participant's first year of college ("Year 1"). Similar follow-up assessments were administered annually thereafter. After a complete description of the study was given to the participants, written informed consent was obtained following IRB-approved protocols. A federal Certificate of Confidentiality was obtained. Participants received cash incentives for each assessment. Annual follow-up rates were 91% ($n=1,142/1,253$), 88% ($n=1,101$) and 88% ($n=1,097$). The current sample consisted of 1,081 participants who provided complete NSSI data, and were enrolled in college for at least one semester during the study period (2004–2008). Most (94.4%) were still enrolled in college in Year 4. Sixteen participants were excluded due to missing data on NSSI.

Measures

Non-suicidal self-injury—In Year 4, a self-administered module to measure NSSI was adapted from Whitlock et al. (2006). The following question assessed lifetime occurrence of NSSI: “Sometimes people do things to hurt themselves on purpose, like cutting, scratching, burning, or injuring themselves in other ways. Have you ever done something like that with the intention of hurting yourself? (Please do not count suicide attempts).” Answer choices were “No”, “Yes”, or “Don’t know”. NSSI frequency and past-year NSSI were assessed by asking, “How often in your life have you done these kinds of things with the intention of hurting yourself (but without the intention of committing suicide)?” and “When was the last time you did something like this with the intention of hurting yourself (but without the intention of committing suicide)?”, respectively. Participants were asked about NSSI motives (“What were your reasons for doing these things to hurt yourself?”) in an open-ended way; later, responses were coded using the method reported by Polk and Liss (2009). Participants were also asked “Have you ever told anyone else that you did things to hurt yourself on purpose? If so, whom did you tell?” Multiple responses were permitted and options included “physician”, “other health care professional”, “therapist, counselor, or other mental health professional”, “parent”, “other family member”, “clergy”, “friend”, “boyfriend, girlfriend, or significant other”, “someone else” (and were asked to specify), and “no one”.

Demographics—Sex was coded as observed during Year 1. Data on race was self-reported, and later dichotomized as White vs. non-White. Socioeconomic status (SES) was estimated by the mean Adjusted Gross Income of participants’ home ZIP codes in the last year before college (MelissaDATA, 2003). Sexual orientation was self-reported annually, with response options of “heterosexual”, “homosexual”, “bisexual”, or “unsure”, and later dichotomized into heterosexual vs. other.

Suicide ideation and attempt—Self-administered questions on suicide ideation and attempt were adapted from the Composite International Diagnostic Interview (Robins *et al.*, 1988) in Year 4: “When was the last time you seriously thought about committing suicide?” and “When was the last time you attempted suicide?” Responses were later dichotomized as “Never” vs. “Less than 24 hours ago” to “More than a year ago”.

Depression, victimization, and exposure to domestic violence—In Years 3 and 4, participants were asked, “Have you ever been diagnosed with depression?” and, if yes, they were asked their age at diagnosis. Victimization, exposure to domestic violence, and age at first occurrence were assessed during Years 2 through 4 as part of a self-administered life events questionnaire based on two widely used life events scales (Sarason *et al.*, 1978; Compas *et al.*, 1987). For the lifetime NSSI analyses, data for these variables were collapsed into three levels: “never”, “prior to college”, and “during college”. For the prospective analyses on past-year NSSI at Year 4, data from Years 2 and 3 were collapsed into “never” and “by Year 3”.

Affective dysregulation—Affective dysregulation was assessed during Year 1 using the Dysregulation Inventory (Mezzich *et al.*, 2001). Participants were asked to indicate how often each statement is true in describing their behavior. Examples include “It is very difficult for you not to think about your fears and worries”, “You slam the door when you are mad”, and “When you are emotionally upset, it lasts for one or two hours even if the problem is gone”. Responses are scored from 0 for “never true” to 3 for “always true.” Items were summed for the 28 items comprising the affective dysregulation subscale (Cronbach’s $\alpha=0.884$, range 0 to 84). Higher scores indicate higher self-reported emotional reactivity and

low control over one's emotional state. A binary variable was created representing the highest quartile compared to the other quartiles.

Parental history of depression—During Year 4, a self-administered family tree questionnaire assessed family history of depression (Mann *et al.*, 1985). Participants were asked to categorize each biological parent regarding the presence or absence of depression, e.g., 1) *Definitely No*: This person definitely does not have depression; 2) *Maybe Yes*: this person could possibly have depression; 3) *Definitely Yes*: this person has been diagnosed with depression. A definite or possible diagnosis of depression was coded as present, “definitely no” was coded as absent, and “Don’t know” and “Don’t remember” were coded as missing.

Perceived social support—In Year 3, participants completed the Social Support Appraisals Scale, a 23-item self-administered assessment measuring subjective perceptions of social support currently received, such as feeling loved and esteemed, and feeling involved with family, friends, and others (Vaux, 1986). Items are scored on a four-point Likert scale and summed (after reverse-coding five items) to derive a total score (Cronbach's $\alpha=0.916$). Scores range from 23 to 92, with lower scores indicating stronger perceived social support.

Conduct problems and impulsive sensation-seeking—Early conduct problems were assessed in Year 1 with the Conduct Disorder Screener (Johnson *et al.*, 1995; Nurco *et al.*, 1999), which asks about the frequency of 18 conduct problems that may have occurred before age 18, and which correspond to the DSM-IV Conduct Disorder symptoms (American Psychiatric Association, 1994), with the sole exception of forgery. An index of the number of conduct problems experienced was computed with a maximum possible score of 18. Participants self-administered the Zuckerman-Kuhlman Personality Questionnaire-Short Form (Zuckerman, 2002) in Year 1, which includes a seven-item subscale measuring impulsive sensation-seeking (Zuckerman, 2002; Arria *et al.*, 2008b).

Cannabis use disorder—During Years 1 through 4, participants who used cannabis five or more times in the past year were assessed for cannabis use disorder (CUD), using questions based in part on the National Survey on Drug Use and Health (NSDUH) interview (Substance Abuse and Mental Health Services Administration, 2003). Items in this series correspond to DSM-IV criteria (American Psychiatric Association, 1994) for abuse and dependence. Students who used cannabis less than five times in the past year skipped out of this series and were automatically coded for the absence of CUD, similar to procedures used in the NSDUH.

Alcohol use disorder—Alcohol use disorder (AUD) was assessed using the same procedures as CUD with the exception that withdrawal symptoms were also assessed for alcohol dependence.

Statistical analyses

Analyses were conducted in three steps. First, bivariate logistic regression models were conducted of lifetime NSSI with the four demographic and thirteen independent variables (see Table 1). Second, a multivariate logistic regression model (and negative binomial regression of NSSI frequency) was used to estimate cross-sectional associations between eleven independent variables (those listed in Table 1 except suicide ideation and attempt) and lifetime NSSI, adjusted for demographic variables. Third, a multivariate logistic regression model was used to predict prospective associations between the eleven independent variables assessed by Year 3 and past-year NSSI at Year 4, adjusting for

demographic variables and excluding individuals whose most recent NSSI occurred more than a year ago. Lifetime suicide ideation and attempt were not included in the multivariate models because temporality could not be resolved for the prospective model, and we wanted the cross-sectional and prospective models to include the same constructs. For all multivariate models, we retained only those variables that were significantly associated with NSSI in bivariate analyses. To obtain a more parsimonious ‘final’ model, we eliminated all non-significant variables, and then re-entered them one by one to retain any that might achieve statistical significance. Only the ‘final models’ are presented in the tables. Statistical analyses were performed using STATA version 10.0 (StataCorp, 2007).

Results

Seventy-five individuals (7%) reported NSSI and 24 individuals (2%) reported past-year NSSI. Of these, 26 (35%) had their most recent episode of NSSI during college; 34 (45%) prior to college; and for 15 (20%) the timing could not be determined. Five (7%) self-injured once and 51% self-injured six or more times (data not shown in table).

Overall, 54% of participants were female, 73% White, and 8% non-heterosexual (Table 1). In bivariate models, lifetime NSSI was significantly associated with being female, non-heterosexual, victimized, exposed to domestic violence during college, and having high affective dysregulation, low social support, and a depression diagnosis. Individuals with NSSI were also more likely to have experienced suicide ideation, a suicide attempt, and maternal and/or paternal depression. Of the 75 individuals with lifetime NSSI, 12 (16%) ever made a suicide attempt, 31 (41%) reported lifetime suicide ideation, and 38 (51%) were ever diagnosed with depression. Lifetime NSSI was not associated with AUD or CUD.

In the multivariate logistic regression analyses for lifetime NSSI (Table 2), female sex ($AOR=1.8$; $95\% CI=1.0-3.1$; $p=0.046$), non-heterosexual orientation ($AOR=3.8$; $95\% CI=1.9-7.4$; $p<0.001$), paternal depression ($AOR=1.9$; $95\% CI=1.1-3.3$; $p=0.030$), and depression diagnosis both prior to college ($AOR=7.3$; $95\% CI=3.9-13.5$; $p<0.001$) and during college ($AOR=2.6$; $95\% CI=1.1-5.9$; $p=0.026$) were all independently associated with lifetime NSSI. These variables maintained statistical significance after adjustment for lifetime suicide attempt.

Prospective analyses for past-year NSSI (Table 3) showed that the following variables were independent predictors of past-year NSSI: non-heterosexual orientation ($AOR=6.2$; $95\% CI=1.8-22.1$; $p=0.005$), maternal depression ($AOR=5.3$; $95\% CI=1.7-16.0$; $p=0.003$), affective dysregulation ($AOR=2.6$; $95\% CI=1.1-6.4$; $p=0.038$), and depression diagnosis by Year 3 ($AOR=2.9$; $95\% CI=1.0-8.2$; $p=0.045$). These same four variables were independently associated with lifetime NSSI frequency; 67% of those with past-year NSSI self-injured 10 or more times, compared with 44% of those whose most recent NSSI was prior to college. All variables maintained statistical significance after adjustment for lifetime suicide attempt, except depression diagnosis by Year 3 ($AOR=2.7$; $95\% CI=0.9-7.6$; $p=0.058$).

Table 4 lists the motives for NSSI. Free text was recorded by interviewers on 107 motives from the 75 participants with NSSI. The most frequently reported motive was mental distress (60%; e.g., “*anger, anxiety, sadness*”; “*mental illness*”; “*I was unmedicated with Bipolar Disorder*”), followed by coping (28%; e.g., “*physical pain blocks emotional pain*”; “*release mental pain*”; “*felt like it would make me feel better*”; “*calmed me when I was upset*”; “*distraction, expulsion of nervous energy*”; “*to try and centralize my pain*”; “*makes me focus*”; “*very overwhelmed and needed a release*”), situational stressors (25%; e.g., relationship, parental divorce, school), and attention seeking (8%); 16% did not know or cite

a motive, or refused to answer. Alcohol problems and sensation-seeking were the least frequently reported motives (3% each). Motives were endorsed with similar frequency with regard to past-year NSSI.

A minority (11%) told no one about their NSSI (Table 5). Most told a friend and/or significant other (68% and 64%, respectively), followed by therapist (40%), parent (37%), other family member (13%), physician (13%), another health care professional (8%), and clergy member (1%).

Discussion

In this study, 7% of the sample engaged in NSSI at least once in their lifetime and 2% in the past year (>70% were female for lifetime and past-year NSSI). Five (7%) with lifetime NSSI self-injured once and 51% self-injured six or more times. Although suicide attempt was independently associated with lifetime NSSI, most individuals with NSSI (84%) never attempted suicide, consistent with the literature from community samples of adolescents and college students (Muehlenkamp & Gutierrez, 2004; Whitlock & Knox, 2007).

Non-heterosexual orientation and a diagnosis of depression predicted past-year NSSI and had independent associations with lifetime NSSI. Maternal depression and affective dysregulation predicted past-year NSSI whereas lifetime NSSI was associated with female sex and paternal depression. Those with past-year NSSI were more likely to have self-injured 10 or more times than those whose last episode with NSSI prior to college (67% vs. 44%, respectively); the same independent predictors of recent NSSI were independently associated with lifetime NSSI frequency. Jacobson and colleagues (2008), in a chart review of outpatient adolescents attending a depression and suicide clinic, noted the importance of examining the frequency and recency of deliberate self-harm behaviors (including NSSI) as it is associated with more severe psychopathology. The three most commonly reported motives for NSSI were mental distress, as a means of coping, and situational stressors; consistent in other studies. This study provides some evidence that NSSI in college is likely to be repetitive and persistent with motives involving emotional regulation in response to situational stress and mental distress.

Several study limitations merit attention. Although the entire sample size of the cohort was large, the subsets of individuals with lifetime ($n=75$) and past-year NSSI ($n=24$) were fairly small as is reflected in the broad confidence intervals for the past-year analyses. These data are based on self-report and thus subject to bias. We do not know when or why participants told others about their NSSI. The NSSI and maternal and paternal depression variables were collected during Year 4 and the exact timing of onset could not be sorted out. It is possible that individuals with depression were more likely to be aware of a parental history of depression than those who were not depressed. Because our sample was limited to individuals from a single public university, results may not be generalizable to students in other areas of the country or at smaller, private settings.

Despite such limitations, this study has a number of counterbalanced strengths. The overall design is prospective, longitudinal, the sample was large, and the assessment battery was extensive. The cohort had passed through the peak period of risk for NSSI (Jacobson & Gould, 2007). Our emphasis on prediction of past-year NSSI limited biases associated with retrospective recall. Instead of constraining response choices regarding motives, we allowed free expression of their reasons [similar to Polk and Liss (2009)]. The sample was recruited as a cohort so the comparison group comprised individuals in the cohort without NSSI, which is an improvement over many existing studies.

Lifetime prevalence of NSSI in our sample was 7%, much lower than in other college samples, such as 17% found by Whitlock et al. (2006) and 38% found by Gratz et al. (2002), but similar to the 5.9% ($n=26$) reported by Klonsky (2011) on a sample of 439 adults in the U.S. recruited by random-digit dialing. This discrepancy might be explained by differences in sampling, differing NSSI definitions (inclusion of wound-picking) and varying assessment time frames (past year, lifetime). Our lower estimate cannot be explained by bias from interviewer-administered format since we used self-administered reports, but could be related to characteristics of our sample (older age) or a cohort effect. Moreover, we cannot rule out the possibility of recall bias (i.e., participants forgetting NSSI episodes).

The link between non-heterosexual orientation and NSSI is in agreement with several other studies (Whitlock *et al.*, 2006; Deliberto & Nock, 2008). Our results suggest that NSSI that persists into young adulthood may be different than adolescent-limited NSSI, given that it has different correlates (e.g., affective dysregulation and maternal depression) and greater frequency. Mental distress, coping, and situational stressors were the main motives for NSSI, suggesting there is a mechanism driving NSSI that persists into young adulthood. Stress reactivity may be impaired in those with frequent or persistent NSSI, with some studies showing increased physiological arousal while completing a distressing task compared with those without prior NSSI (Nock & Mendes, 2008). Additionally, preliminary evidence suggests that NSSI might occur in response to hyperactivity of the neuroendocrine systems (Sachsse *et al.*, 2002). Thus, biological vulnerabilities could be important risk factors for frequent or persistent NSSI as well as stressful environmental factors that could trigger physiological vulnerabilities.

Similar to Heath et al. [86%; (2009)], 89% of individuals in our sample with NSSI told others about it, most commonly their friends (68%). A recent longitudinal study by Prinstein et al. (2010) suggests that peers might influence some individuals' engagement in self-injury among younger females. Nock and Prinstein (2004) found that internal and social motives are the most commonly endorsed. Community studies of adolescents have reported social factors to be influential [e.g., "to try to get a reaction from someone" or attention seeking (Lloyd-Richardson *et al.*, 2007)], but these motives were endorsed by only 8% in our sample. Our data provide strong support for internal rather than social motives underlying NSSI. It could be that motives for NSSI change over the course of development (Lloyd-Richardson *et al.*, 2007) or that those with NSSI that persists into adulthood have different motives.

Although drug and alcohol abuse has been observed to co-exist with NSSI among clinical populations (Nock *et al.*, 2006), AUD and CUD were not observed to be significant risk factors for NSSI in our sample. Our findings comport with Jacobson et al. (2008), who found, among outpatient adolescents, that the prevalence of substance use disorder was not significantly elevated among individuals with NSSI as compared to individuals without NSSI.

These findings have important implications for accurate identification, prevention, and treatment of NSSI. Campus-based mental health professionals who treat students with NSSI may not be aware of clinical management strategies. Some may view NSSI as a manipulative act and disregard its clinical relevance. NSSI has been proposed to appear in the Diagnostic and Statistical Manual of Mental Disorders-5 (American Psychiatric Association, 2010). Currently, NSSI is part of the criteria for Borderline Personality Disorder, but frequently those with NSSI do not meet criteria for Borderline Personality Disorder; thus clinical management and treatment recommendations are often obscured (American Psychiatric Association, 1994). Despite the prevalence and significant consequences of NSSI, few intervention approaches for NSSI have been empirically

supported (Prinstein, 2008). Other than Dialectical Behavior Therapy, an intensive treatment program for self-injurers with Borderline Personality Disorder (Kliem *et al.*, 2010), no empirically supported treatment for NSSI exists. Evidence supporting Dialectical Behavior Therapy's efficacy in adolescents and college students is scarce. If mental distress is indeed a primary motive for NSSI, coping strategies for tolerating and regulating intense emotions could be an intervention target for NSSI prevention.

Longitudinal research following larger community samples through the peak period of risk for NSSI is needed to sort out the course and temporal sequencing of NSSI and other suspected risk and protective factors which could be targets for intervention. This study identified predictors of NSSI, which might provide clues to its etiology and have implications for intervention.

Acknowledgments

Special thanks are given to Laura Garnier-Dykstra, Lauren Stern, Emily Winick, the interviewing team, and the participants.

References

- Adrian M, Zeman J, Erdley C, Lisa L, Sim L. Emotional dysregulation and interpersonal difficulties as risk factors for nonsuicidal self-injury in adolescent girls. *Journal of Abnormal Child Psychology*. 2011; 39:389–400. [PubMed: 20953828]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV*. American Psychiatric Press; Washington, DC: 1994.
- American Psychiatric Association. [Accessed October 1 2010] *DSM-5 Development*. 2010. (<http://www.dsm5.org/>)
- Arria AM, Caldeira KM, O'Grady KE, Vincent KB, Fitzelle DB, Johnson EP, Wish ED. Drug exposure opportunities and use patterns among college students: Results of a longitudinal prospective cohort study. *Substance Abuse*. 2008a; 29:19–38. [PubMed: 19042196]
- Arria AM, Caldeira KM, Vincent KB, O'Grady KE, Wish ED. Perceived harmfulness predicts nonmedical use of prescription drugs among college students: Interactions with sensation-seeking. *Prevention Science*. 2008b; 9:191–201. [PubMed: 18633709]
- Briere J, Gil E. Self-mutilation in clinical and general population samples: Prevalence, correlates, and functions. *American Journal of Orthopsychiatry*. 1998; 68:609–620. [PubMed: 9809120]
- Bureau JF, Martin J, Freynet N, Poirier AA, Lafontaine MF, Cloutier P. Perceived dimensions of parenting and non-suicidal self-injury in young adults. *Journal of Youth and Adolescence*. 2010; 39:484–494. [PubMed: 19882378]
- Compas BE, Davis GE, Forsythe CJ, Wagner BM. Assessment of major and daily stressful events during adolescence: The Adolescent Perceived Events Scale. *Journal of Consulting and Clinical Psychology*. 1987; 55:534–541. [PubMed: 3624609]
- Crowell SE, Beauchaine TP, McCauley E, Smith CJ, Vasilev CA, Stevens AL. Parent-child interactions, peripheral serotonin, and self-inflicted injury in adolescents. *Journal of Consulting and Clinical Psychology*. 2008; 76:15–21. [PubMed: 18229978]
- Deliberto TL, Nock MK. An exploratory study of correlates, onset, and offset of non-suicidal self-injury. *Archives of Suicide Research*. 2008; 12:219–231. [PubMed: 18576203]
- Garrison CZ, Addy CL, McKeown RE, Cuffe SP, Jackson KL, Waller JL. Nonsuicidal physically self-damaging acts in adolescents. *Journal of Child and Family Studies*. 1993; 3:339–352.
- Gratz KL, Conrad SD, Roemer L. Risk factors for deliberate self-harm among college students. *American Journal of Orthopsychiatry*. 2002; 72:128–140. [PubMed: 14964602]
- Hankin BL, Abela JRZ. Nonsuicidal self-injury in adolescence: Prospective rates and risk factors in a 2 year longitudinal study. *Psychiatry Research*. 2011; 186:65–70. [PubMed: 20807667]

- Heath NL, Ross S, Toste JR, Charlebois A, Nedecheva T. Retrospective analysis of social factors and nonsuicidal self-injury among young adults. *Canadian Journal of Behavioural Science*. 2009; 41:180–186.
- Herpertz S, Sass H, Favazza A. Impulsivity in self-mutilative behavior: Psychometric and biological findings. *Journal of Psychiatric Research*. 1997; 31:451–465. [PubMed: 9352472]
- Jacobson CM, Gould M. The epidemiology and phenomenology of non-suicidal self-injurious behavior among adolescents: A critical review of the literature. *Archives of Suicide Research*. 2007; 11:129–147. [PubMed: 17453692]
- Jacobson CM, Muehlenkamp JJ, Miller AL, Turner JB. Psychiatric impairment among adolescents engaging in different types of deliberate self-harm. *Journal of Clinical Child and Adolescent Psychology*. 2008; 37:363–375. [PubMed: 18470773]
- Johnson EO, Arria AM, Borges G, Ialongo N, Anthony JC. The growth of conduct problem behaviors from middle childhood to early adolescence: Sex differences and the suspected influence of early alcohol use. *Journal of Studies on Alcohol*. 1995; 56:661–671. [PubMed: 8558898]
- Kaess M, Hille M, Parzer P, Maser-Gluth C, Resch F, Brunner R. Alterations in the neuroendocrinological stress response to acute psychosocial stress in adolescents engaging in nonsuicidal self-injury. *Psychoneuroendocrinology*. 2011 e-pub ahead of print, June 13, 2011.
- Kliem S, Kroger C, Kosfelder J. Dialectical behavior therapy for borderline personality disorder: A meta-analysis using mixed-effects modeling. *Journal of Consulting and Clinical Psychology*. 2010; 78:936–951. [PubMed: 21114345]
- Klonsky ED. The functions of self-injury in young adults who cut themselves: Clarifying the evidence for affect-regulation. *Psychiatry Research*. 2009; 166:260–268. [PubMed: 19275962]
- Klonsky ED. Non-suicidal self-injury in United States adults: Prevalence, sociodemographics, topography, and functions. *Psychological Medicine*. 2011 e-pub ahead of print, January 5, 2011.
- Laye-Gindhu A, Schonert-Reichl KA. Nonsuicidal self-harm among community adolescents: Understanding the “whats” and “whys” of self-harm. *Journal of Youth and Adolescence*. 2005; 34:447–457.
- Linehan, MM. Cognitive-behavioral treatment of borderline personality disorder. The Guilford Press; New York: 1993.
- Lloyd-Richardson E, Perrine N, Dierker L, Kelley ML. Characteristics and functions of non-suicidal self-injury in a community sample of adolescents. *Psychological Medicine*. 2007; 37:1183–1192. [PubMed: 17349105]
- Mann RE, Sobell LC, Sobell MB, Pavan D. Reliability of a family tree questionnaire for assessing family history of alcohol problems. *Drug and Alcohol Dependence*. 1985; 15:61–67. [PubMed: 4017879]
- MelissaDATA. [Accessed May 28 2008] Income tax statistics lookup. 2003. (<http://www.melissadata.com/lookups/taxzip.asp>)
- Mezzich AC, Tarter RE, Giancola PR, Kirisci L. The dysregulation inventory: A new scale to assess the risk for substance use disorder. *Journal of Child and Adolescent Substance Abuse*. 2001; 10:35–43.
- Muehlenkamp J, Gutierrez PM. An investigation of differences between self-injurious behavior and suicide attempts in a sample of adolescents. *Suicide and Life-Threatening Behavior*. 2004; 34:12–23. [PubMed: 15106884]
- Nock MK, Joiner TE, Gordon KH, Lloyd-Richardson E, Prinstein MJ. Non-suicidal self-injury among adolescents: Diagnostic correlates and relation to suicide attempts. *Psychiatry Research*. 2006; 144:65–72. [PubMed: 16887199]
- Nock MK, Mendes WB. Physiological arousal, distress tolerance, and social problem-solving deficits among adolescent self-injurers. *Journal of Consulting and Clinical Psychology*. 2008; 76:28–38. [PubMed: 18229980]
- Nock MK, Prinstein MJ. A functional approach to the assessment of self-mutilative behavior. *Journal of Consulting and Clinical Psychology*. 2004; 72:885–890. [PubMed: 15482046]
- Nurco DN, Blatchley RJ, Hanlon TE, O’Grady KE. Early deviance and related risk factors in the children of narcotic addicts. *The American Journal of Drug and Alcohol Abuse*. 1999; 25:25–45. [PubMed: 10078976]

- Peterson J, Freedenthal S, Sheldon C, Andersen R. Nonsuicidal self injury in adolescents. *Psychiatry*. 2008; 5:20–26. [PubMed: 19724714]
- Polk E, Liss M. Exploring the motivations behind self-injury. *Counselling Psychology Quarterly*. 2009; 22:233–241.
- Prinstein MJ. Introduction to the special section on suicide and nonsuicidal self-injury: A review of unique challenges and important directions for self-injury science. *Journal of Consulting and Clinical Psychology*. 2008; 76:1–8. [PubMed: 18229976]
- Prinstein MJ, Heilbron N, Guerry JD, Franklin JC, Rancourt D, Simon V, Spirito A. Peer influence and nonsuicidal self injury: Longitudinal results in community and clinically-referred adolescent samples. *Journal of Abnormal Child Psychology*. 2010; 38:669–682. [PubMed: 20437255]
- Robins LN, Wing J, Wittchen HU, Helzer JE, Babor TF, Burke J, Farmer A, Jablenski A, Pickens R, Regier DA, Sartorius N, Towle LH. The Composite International Diagnostic Interview: An epidemiologic instrument suitable for use in conjunction with different diagnostic systems and in different cultures. *Archives of General Psychiatry*. 1988; 45:1069–1077. [PubMed: 2848472]
- Ross S, Heath N. A study of the frequency of self-mutilation in a community sample of adolescents. *Journal of Youth and Adolescence*. 2002; 31:61–77.
- Sachsse U, Von der Heyde S, Huether G. Stress regulation and self-mutilation. *The American Journal of Psychiatry*. 2002; 159:672–672. [PubMed: 11925313]
- Sarason IG, Johnson JH, Siegel JM. Assessing the impact of life changes: Development of the Life Experiences Survey. *Journal of Consulting and Clinical Psychology*. 1978; 46:932–946. [PubMed: 701572]
- Shaffer, D.; Jacobson, C. Proposal to the DSM-V childhood disorder and mood disorder work groups to include 9 non-suicidal self-injury (NSSI) as a DSM-V disorder. *American Psychiatric Association*; 2010. p. 1-21.
- StataCorp. *Stata Statistical Software: Release 10*. StataCorp LP; College Station, TX: 2007.
- Substance Abuse and Mental Health Services Administration. 2002 National Survey on Drug Use and Health Questionnaire. Office of Applied Studies; Rockville, MD: 2003.
- Vaux A. The Social Support Appraisals (SS-A) Scale: Studies of reliability and validity. *American Journal of Community Psychology*. 1986; 14:195–219.
- Vincent KB, Kasperski SJ, Caldeira KM, Garnier-Dykstra LM, Pinchevsky GM, O'Grady KE, Arria AM. Maintaining superior response rates in a longitudinal study: Experiences from the College Life Study. *International Journal of Multiple Research Approaches*. (under second review).
- Weierich MR, Nock MK. Posttraumatic stress symptoms mediate the relation between childhood sexual abuse and nonsuicidal self-injury. *Journal of Consulting and Clinical Psychology*. 2008; 76:39–44. [PubMed: 18229981]
- Whitlock J, Eckenrode J, Silverman D. Self-injurious behaviors in a college population. *Pediatrics*. 2006; 117:1939–1948. [PubMed: 16740834]
- Whitlock J, Knox KL. The relationship between self-injurious behavior and suicide in a young adult population. *Archives of Pediatrics and Adolescent Medicine*. 2007; 161:634–640. [PubMed: 17606825]
- Yates TM, Tracy AJ, Luthar SS. Nonsuicidal self-injury among “privileged” youths: Longitudinal and cross-sectional approaches to developmental process. *Journal of Consulting and Clinical Psychology*. 2008; 76:52–62. [PubMed: 18229983]
- Zuckerman, M. Zuckerman-Kuhlman Personality Questionnaire (ZKPQ): An alternative five-factorial model. In: de Raad, B.; Perugini, M., editors. *Big Five Assessment*. Hogrefe & Huber; Seattle: 2002. p. 377-396.

Table 1

Characteristics of the sample

Demographic Variables	Overall (n=1081)				NSSI (n=75)				p
	n	%	n	%	n	%	n	%	
Sex (female)	582	54	527	52	55	73	<0.001		
Race (non-White)	295	27	276	27	19	25	0.693		
SES (highest quartile) ^b	266	25	249	25	17	23	0.662		
Sexual orientation (homosexual/bisexual/not sure)	84	8	68	7	16	21	<0.001		
Psychosocial Variables									
Victimization prior to college ^c	77	7	68	7	9	12	0.021		
Victimization during college ^c	64	6	54	5	10	13	0.001		
Exposure to domestic violence prior to college ^d	79	7	70	7	7	9	0.267		
Exposure to domestic violence during college ^d	31	3	26	2	5	7	0.018		
Top quartile, affective dysregulation	238	23	211	21	27	36	0.001		
Diagnosis of depression prior to college ^e	89	8	62	6	27	36	<0.001		
Diagnosis of depression during college ^e	64	6	53	5	11	15	<0.001		
Lifetime suicide ideation	135	12	104	10	31	41	<0.001		
Lifetime suicide attempt	29	3	17	2	12	16	<0.001		
Possible/definite maternal depression	240	22	201	20	39	52	<0.001		
Possible/definite paternal depression	230	23	199	20	31	41	<0.001		
Lifetime alcohol use disorder ^f	609	56	568	56	41	55	0.986		
Lifetime cannabis use disorder ^f	287	27	265	26	22	29	0.446		
Conduct problems	mean	SD	mean	SD	mean	SD			
Impulsivity	4.6	2.7	4.6	2.7	4.9	2.8	0.567		
Social support	33.8	8.1	33.5	7.9	36.5	9.0	0.003		

^aNSSI: Non-suicidal self-injury

^b The mean adjusted gross income reported by the Internal Revenue Service for each participant's home ZIP code during their last year in high school.

^c Victimization by Year 3 was reported by 120/1028(12%) individuals overall [104/962(11%) without NSSI; 16/66(24%) with NSSI, $p=0.003$].

^d Exposure to domestic violence by Year 3 was reported by 95/1028 (9%) individuals overall [86/963 (9%) without NSSI; 9/65(14%) with NSSI, $p=0.396$].

^e Depression diagnosis by Year 3 was reported by 122/1040 (12%) individuals overall [88/966(9%) without NSSI; 34/74 (46%) with NSSI, $p<0.001$].

^f Alcohol use disorders by Year 3 was reported by 521/967(54%) individuals overall [484/902(54%) without NSSI; 37/65(57%) with NSSI, $p=0.610$]. Cannabis use disorders by Year 3 was reported by 267/967(28%) individuals overall [247/902(27%) without NSSI; 20/65(31%) with NSSI, $p=0.555$].

Table 2

Multivariate analyses for lifetime non-suicidal self-injury, as compared to those without lifetime non-suicidal self-injury (NSSI)

Demographic Variables	Lifetime NSSI		
	AOR^a	95%CI	p
Sex (female)	1.8	1.0–3.1	0.046
Sexual orientation (homosexual/bisexual/not sure)	3.8	1.9–7.4	<0.001
Psychosocial Variables			
Diagnosis of depression prior to college ^b	7.3	3.9–13.5	<0.001
Diagnosis of depression during college ^b	2.6	1.1–5.9	0.026
Possible/definite paternal depression	1.9	1.1–3.3	0.030

^aEstimates were adjusted for all independent variables shown.

^bReference group was individuals who were never diagnosed with depression.

Table 3

Prospective multivariate model for past-year non-suicidal self-injury, as compared to those without lifetime non-suicidal self-injury (NSSI)

Demographic Variables	Past-Year NSSI		
	<i>AOR^a</i>	<i>95%CI</i>	<i>p</i>
Sexual orientation (homosexual/bisexual/not sure)	6.2	1.8–22.1	0.005
Psychosocial Variables			
Diagnosis of depression by Year 3	2.9	1.0–8.2	0.045
Top quartile of affective dysregulation	2.6	1.1–6.4	0.038
Possible/definite maternal depression	5.3	1.7–16.0	0.003

^aEstimates were adjusted for all independent variables shown.

Table 4Motives for non-suicidal self-injury (NSSI)^a

	<i>n</i>	%
Mental distress	45	60
Coping mechanism	21	28
Situational stressors	19	25
Attention seeking	6	8
Alcohol problems	2	3
Sensation-seeking	2	3
No reason listed, don't know, refused to answer	12	16

^aResponse options were not provided, free text was recorded by the interviewer, participants often reported more than one motive for NSSI; 107 motives were provided by 75 participants.

Table 5To whom participants disclosed their non-suicidal self-injury (NSSI)^a

	<i>n</i>	%
Friend	51	68
Girl-, Boyfriend, significant other	48	64
Therapist, Counselor	30	40
Parent	28	37
Other Family Member	10	13
Physician	10	13
No one	8	11
Other Health Care Professional	6	8
Other	3	4
Clergy	1	1

^aMultiple responses were permitted