Unicompartmental Versus Total Knee Arthroplasty Database Analysis

Is There a Winner?

Matthew C. Lyons MBBS, FRACS, Steven J. MacDonald MD, FRCSC,
Lyndsay E. Somerville MSc, Douglas D. Naudie MD, FRCSC,
Richard W. McCalden MD, FRCSC

Published online: 27 October 2011
© The Association of Bone and Joint Surgeons® 2011

Abstract

Background TKA and unicompartmental knee arthroplasty (UKA) are both utilized to treat unicompartmental knee arthrosis. While some surgeons assume UKA provides better function than TKA, this assumption is based on greater final outcome scores rather than on change in scores and many patients with UKA have higher preoperative scores.

Questions/purposes We therefore asked whether TKA would demonstrate (1) better change in clinical outcome scores from preoperative to postoperative states and (2) better survivorship than UKA.

Methods We evaluated 4087 patients with 5606 TKAs and 179 patients with 279 UKAs performed between 1978 and 2009. Patients with TKA were older and heavier than patients with UKA (mean age, 68 versus 66 years; mean BMI, 32 versus 29). We compared preoperative, latest

postoperative, and change in Knee Society Clinical Rating System (KSCRS), SF-12, and WOMAC scores. Minimum followup was 2 years (UKA: mean, 7 years; range, 2.0–23 years; TKA: mean, 6.5 years; range, 2.0–33 years). Preoperative outcome measure scores (WOMAC, SF-12, KSCRS) were higher in the UKA group.

Results Patients with UKA had higher postoperative KSCRS and SF-12 mental scores. Changes in score for all WOMAC domains were similar between groups. Total KSCRS changes in score were similar between groups, although patients with TKA had higher knee scores (49 versus 43) but lower function scores than UKA (21 versus 26). Cumulative revision rate was higher for UKA than for TKA (13% versus 7%). Kaplan-Meier survivorship at 5 and 10 years was 95% and 90%, respectively, for UKA and 98% and 95%, respectively, for TKA.

Conclusions While patients with UKA had higher pre- and postoperative scores than patients with TKA, the changes in scores were similar in both groups and survival appeared higher in patients with TKA.

Level of Evidence Level III, therapeutic study. See the Guidelines for Authors for a complete description of levels of evidence.

One or more of the authors act as consultants for Smith & Nephew, Inc (Memphis, TN, USA) (RWM, DDN) and DePuy Orthopaedics, Inc (Warsaw, IN, USA) (SJM) although no payments were received related to this work. The institution of the authors has received funding from DePuy, Smith & Nephew, and Stryker Orthopaedics (Mahwah, NJ, USA).

Clinical Orthopaedics and Related Research neither advocates nor endorses the use of any treatment, drug, or device. Readers are encouraged to always seek additional information, including FDA approval status, of any drug or device before clinical use. Each author certifies that his or her institution has approved the human protocol for this investigation, that all investigations were conducted in conformity with ethical principles of research, and that informed consent for participation in the study was obtained.

M. C. Lyons, S. J. MacDonald (✉), L. E. Somerville,
D. D. Naudie, R. W. McCalden
Division of Orthopaedic Surgery, University of Western Ontario
& London Health Sciences Centre, University Campus,
339 Windermere Road, London, ON N6A 5A5, Canada
e-mail: steven.macdonald@lhsc.on.ca

Introduction

TKA and unicompartmental knee arthroplasty (UKA) are both utilized to treat unicompartmental knee arthrosis. TKA has long been considered the gold standard operative intervention for knee arthrosis due to demonstrated predictability, durability, and effectiveness in the treatment of pain and restoration of function [3, 9, 11, 16, 17, 53]. Advocates of UKA cite several advantages over conventional

TKA, including minimal access surgery, preservation of bone stock, and ease of revision [5, 32, 37, 41, 56], as well as superior knee ROM and kinematics, less blood loss, faster recovery, and a decreased hospital admission [4, 20–22, 25–27, 50, 62]. Satisfaction after either TKA or UKA is similar [54]. UKA has demonstrated survivorship of greater than 90% for more than 10 years after implantation in multiple studies [6, 12, 39, 40, 48, 49, 59]. The cumulative revision rate is higher for UKA than for conventional TKA, ranging from 82% to 98% at 5 to 22 years for UKA compared to 91% to 98.9% at 5 to 19 years for TKA (Table 1) [5, 6, 10, 12, 28–30, 33, 52].

Table 1. Comparison of survivorship of UKA and TKA in our study and other studies in the literature

Study	Intervention	Number	Survival (%)	Years
Argenson et al. [6]	UKA	160	94	10
Emerson and Higgins [13]	UKA	55	85	10
Gioe et al. [19]	UKA	516	88.6	10
Murray et al. [39]	UKA	143	98	10
Naudie et al. [40]	UKA	113	90	10
O'Rourke et al. [43]	UKA	136	84	20
Price et al. [48]	UKA	439	93	15
Price and Svard [49]	UKA	85	91	16
Squire et al. [57]	UKA	140	84	22
Steele et al. [58]	UKA	203	85.9	20
Svard and Price [59]	UKA	124	95	10
Tabor and Tabor [60]	UKA	67	91	5
Tabor and Tabor [60]	UKA	67	84	10
Vorlat et al. [61]	UKA	140	82	10
Whittaker et al. [64]	UKA	79	89	5
Australian Orthopaedic Association National Joint Replacement Registry [7]	UKA	31,884	86.6	9
Current study	UKA	219	94.6	5
Current study	UKA	219	90.4	10
Abdeen et al. [1]	TKA	100	92.4	19
Emmerson et al. [14]	TKA	109	95	10
Gill and Joshi [18]	TKA	404	92.6	17
Keating et al. [28]	TKA	4583	98.9	15
Parsch et al. [45]	TKA	141	91	14
Petrou et al. [47]	TKA	100	96.1	15
Ranawat et al. [52]	TKA	112	94.1	15
Australian Orthopaedic Association National Joint Replacement Registry [7]	TKA	231,409	94.9	9
Current study	TKA	5606	98.4	5
Current study	TKA	5606	94.9	10

UKA = unicompartmental knee arthroplasty.

The therapeutic effect of UKA and TKA is widely assessed with validated clinical outcome measures, with many studies describing better functional results in UKA [40–42]. Historically, authors have focused on absolute preoperative and postoperative scores; however, the change in score from preoperative to postoperative has been reported recently to represent the effect of the intervention under investigation [31, 35, 38, 51]. Since patients receiving UKA are typically younger with a higher level of preoperative function, the postoperative results tend to be higher.

We asked whether TKA would demonstrate (1) better change in clinical outcome scores from preoperative to postoperative states and (2) better survivorship than UKA.

Patients and Methods

We retrospectively identified from our database two cohorts of patients, those having a primary TKA and those having a UKA between 1978 and 2009. There were 4087 patients with 5606 primary TKAs and 179 patients with 219 UKAs performed at a single institution. For UKA to be undertaken, the patient must have satisfied the following criteria: isolated medial compartmental osteoarthritis; fixed flexion deformity of less than 5°; an active ROM of greater than 90°; and less than 15° of varus deformity. TKA was undertaken in patients who did not satisfy the criteria for UKA or who satisfied the criteria but elected a TKA.

Patients undergoing joint arthroplasty had demographic data collected prospectively, which was then entered into an institutional clinical database. This information included surgical information and preoperative and postoperative scores using WOMAC (pain, stiffness, function, total) [8], SF-12 (physical component summary [PCS] and mental component summary [MCS], Version 1) [63], and Knee Society Clinical Rating System (KSCRS) (function, knee, total) [24]. The minimum followup was 2 years for both groups (UKA: mean, 7.12 ± 4.45 years; range, 2.0–23.24 years; TKA: mean, 6.50 ± 3.96 years; range, 2.0–33.36 years). No patients were recalled specifically for this study; all data were obtained from medical records and no patients were lost to followup. Survivorship was defined as revision surgery for any aseptic cause. Institutional review board approval was obtained before initiation of this study.

A post hoc power analysis was performed and it was demonstrated our available sample size would result in 100% power to detect a significant change. The mean ($\pm$ SD) age was greater ($p < 0.001$) in the TKA group (67.65 ± 9.31 years) than in the UKA group (66.02 ± 8.18 years) at surgery (Table 2). The BMI was higher ($p < 0.001$) in the TKA group (31.8 ± 6.55) than in the UKA group (29.3 ± 4.69). The TKA cohort had a higher ($p < 0.001$) percentage of female patients than the UKA cohort (60.9% versus 47.3%).

The mean followup tended to be shorter ($p = 0.07$) in the TKA group (6.50 ± 3.96 years) than in the UKA group (7.12 ± 4.45 years). All preoperative outcome measure scores (WOMAC [Table 3], SF-12 [Table 4], KSCRS [Table 5]) were higher in the UKA group.

Table 2. Patient demographics by procedure

Variable	Procedure		p value
	TKA	UKA	
Sex (% female)	60.9	47.3	< 0.001
Followup time after surgery (years)*	6.50 ± 3.96	7.12 ± 4.45	0.07
Aseptic revisions (%)	6.40%	12.9%	< 0.001
Age (years)*	67.65 ± 9.31	66.02 ± 8.18	< 0.001
Height (cm)*	165.3 ± 10.39	167.2 ± 10.08	0.001
Weight (kg)*	86.9 ± 18.9	81.7 ± 15.8	< 0.001
BMI*	31.8 ± 6.55	29.3 ± 4.69	< 0.001

* Values are expressed as mean $\pm$ SD; UKA = unicompartmental knee arthroplasty.

Table 3. Preoperative, latest, and change in WOMAC scores

WOMAC domain	Procedure	Mean	SD	p value*
Preoperative				
Stiffness	TKA	40.67	19.98	0.06
	UKA	43.70	20.37	
Pain	TKA	46.15	17.77	0.06
	UKA	48.63	18.16	
Function	TKA	45.44	17.47	< 0.001
	UKA	51.42	17.68	
Total	TKA	44.76	16.14	0.002
	UKA	48.75	16.52	
Latest				
Stiffness	TKA	67.84	24.09	0.79
	UKA	68.45	23.41	
Pain	TKA	74.83	23.32	0.56
	UKA	76.40	22.09	
Function	TKA	69.25	23.13	0.002
	UKA	74.15	22.79	
Total	TKA	71.41	21.66	0.11
	UKA	73.97	21.05	
Change				
Stiffness	TKA	28.23	27.03	0.32
	UKA	25.41	23.78	
Pain	TKA	29.97	24.79	0.89
	UKA	30.04	22.96	
Function	TKA	25.64	23.66	0.97
	UKA	25.61	20.14	
Total	TKA	27.96	22.47	0.70
	UKA	27.25	19.63	

* Grouped by procedure; UKA = unicompartmental knee arthroplasty.

All TKAs were performed by a midline incision with a medial parapatellar approach and eversion of the patella. UKAs received a medial parapatellar approach with an

Table 4. Preoperative, latest, and change in SF-12 scores

SF-12 domain	Procedure	Mean	SD	p value*
Preoperative				
MCS	TKA	52.74	10.99	0.17
	UKA	54.24	10.18	
PCS	TKA	30.19	7.72	0.03
	UKA	32.04	8.74	
Latest				
MCS	TKA	52.22	10.18	0.41
	UKA	52.96	9.59	
PCS	TKA	36.97	10.98	0.005
	UKA	39.15	11.13	
Change				
MCS	TKA	−0.34	11.03	0.76
	UKA	−1.20	11.09	
PCS	TKA	7.64	11.67	0.03
	UKA	9.88	11.37	

* Grouped by procedure; MCS = mental component summary; PCS = physical component summary; UKA = unicompartmental knee arthroplasty.

Table 5. Preoperative, latest, and change in KSCRS scores

KSCRS domain	Procedure	Mean	SD	p value*
Preoperative				
Function	TKA	44.52	15.63	< 0.001
	UKA	53.21	13.18	
Knee	TKA	41.09	15.34	< 0.001
	UKA	47.82	15.92	
Total	TKA	85.73	24.45	< 0.001
	UKA	101.10	22.71	
Latest				
Function	TKA	65.74	27.06	< 0.001
	UKA	79.55	22.42	
Knee	TKA	89.72	13.48	0.33
	UKA	90.58	13.64	
Total	TKA	155.63	33.96	< 0.001
	UKA	170.87	29.88	
Change				
Function	TKA	21.36	26.22	< 0.001
	UKA	25.65	22.26	
Knee	TKA	49.24	19.45	0.001
	UKA	42.88	21.29	
Total	TKA	70.62	36.21	0.76
	UKA	69.56	32.80	

* Grouped by sex/procedure; KSCRS = Knee Society Clinical Rating System; UKA = unicompartmental knee arthroplasty.

incision from the superomedial border of the patellar to 1.5 cm distal to medial tibial plateau articular surface. The patella was subluxated for exposure and the ACL and the patellofemoral and lateral compartments were inspected critically to ensure a UKA was a suitable intervention. Both prostheses had all components cemented. The patellar was selectively resurfaced in the TKA cohort. We inserted a drain in all knees on closure and routinely removed the drain on the first postoperative day.

Patients were routinely mobilized fully weightbearing on the first postoperative day under the supervision of a physiotherapist. Walking aids were utilized as required. Ongoing outpatient physiotherapy was employed to ensure patients regained an adequate ROM and strength. This was continued to the satisfaction of the physiotherapist and surgeon, typically ceasing at the 6-week review.

The followup regime for all patients was a 2-week wound review with clinical assessments performed at 6 weeks, 3 months, 1 year, 2 years (± 2 months), and alternate years thereafter unless there was clinical concern for earlier review. At each review, a clinical and radiographic review was undertaken. ROM was assessed with a goniometer. Patients completed WOMAC, SF-12, and KSCRS questionnaires. All scores were entered in the hospital database at the time of review. All elements of the KSCRS (knee, function, total) were individually analyzed. Preoperative and latest postoperative scores were analyzed. Changes in score were also analyzed. Recent studies have suggested the change in clinical outcome scores represents the effect of the intervention under investigation [31, 35, 38, 51]. We believe this allows a valid comparison between treatment groups. We determined differences in demographics between the TKA and UKA groups. Differences in sex distribution was assessed using crosstabs with the chi square test. Differences in BMI, age, and followup time

between the TKA and UKA groups were assessed using a Mann-Whitney U-test for two-group comparison. All of our outcome measures (WOMAC, SF-12, KSCRS) were evaluated preoperatively and postoperatively by the Mann-Whitney U-test. Kaplan-Meier survivorship was undertaken to assess durability of the implants. We used SPSS[®] v.19 (SPSS Inc, Chicago, IL, USA) for all analyses.

Results

In all latest outcome measure scores, the UKA group had higher scores than the TKA group. The change in score (preoperatively to latest postoperatively) demonstrated no between-group differences for the WOMAC (Table 3) and SF-12 MCS (Table 4). The change in score for the SF-12 PCS was higher in the UKA sample. The latest KSCRS score was higher in UKAs for both total and function scores; however, the knee scores were not different. The change in score for KSCRS scores was not different between the groups for total scores but favored TKA over UKA for knee scores (49 versus 43; $p < 0.001$) and UKA over TKA for function scores (26 versus 21; $p < 0.001$) (Table 5).

The cumulative revision rate was higher ($p = 0.006$) for UKAs than for TKAs (37 revisions [13.3%] versus 399 revisions [7.1%]). Kaplan-Meier survivorship at 5 and 10 years was 98.4% (95% CI, 98.2%–98.6%) and 94.9% (95% CI, 94.6%–95.2%), respectively, for TKA and 94.6% (95% CI, 93.2%–96%) and 90.4% (95% CI, 88.4%–92.4%), respectively, for UKA (Fig. 1). Indications for revision were comparable between groups, with two of 37 UKAs (5.4%) and seven of 399 TKAs (1.8%) revised for pain of unknown origin (Table 6). There were some failure modes that were unique to TKAs (periprosthetic fracture and metal-backed patella failure being the most common).

Fig. 1 Kaplan-Meier survivorship at 5 and 10 years with aseptic revisions as the end point was 98.4% (95% CI, 98.2%–98.6%) and 94.9% (95% CI, 94.6%–95.2%), respectively, for TKA and 94.6% (95% CI, 93.2%–96%) and 90.4% (95% CI, 88.4%–92.4%), respectively, for UKA.

Table 6. Indications for revision arthroplasty

Reason for revision	TKA (n = 399 of 5606)	UKA (n = 37 of 279)
Overall all cause revision rate	7.1%	13.3%
Disease progression	47 (0.8%)	19 (6.8)
Polyethylene wear/osteolysis	148 (2.6%)	6 (2.2%)
Aseptic loosening	63 (1.1%)	5 (1.8%)
Instability	34 (0.6%)	1 (0.3%)
Pain unknown origin	7 (0.1%)	2 (0.7%)
Infection	40 (0.7%)	1 (0.3%)
Stiffness	7 (0.1%)	
Periprosthetic fracture	25 (0.4%)	
Implant failure (metal-backed patella)	16 (0.3%)	
Malalignment	5 (0.1%)	
Extensor mechanism failure	3 (0.1%)	
Other	4 (0.1%)	3 (1.1%)

TKA = total knee arthroplasty; UKA = unicompartmental knee arthroplasty.

Discussion

Treatment options for knee arthrosis include both UKA and TKA. Proposed advantages of UKA over TKA include both technical factors (less invasive surgery, preservation of bone stock, comparative ease of revision) [5, 32, 37, 41, 56] and improved clinical outcomes (superior knee kinematics, less blood loss, faster recovery, shorter inpatient stay) [4, 20–22, 25–27, 50, 62]. Clinical outcome scores have previously focused on absolute values at preoperative and postoperative review. Recent studies have identified the change in score (preoperative to postoperative) as a measure of the effect of the intervention [31, 35, 38, 51]. Survival analysis is an accepted method of evaluating the durability of a prosthesis and both prostheses have documented survival rates in excess of 90% at 10 years (Table 1). Nevertheless, TKA is more frequently performed due to the perception that TKA is a more durable operation [36]. We hypothesized TKA would demonstrate (1) better change in clinical outcome scores from preoperative to postoperative states and (2) better survivorship than UKA.

Limitations of this study include the following. First, the cohorts were nonrandomized and unevenly distributed, with UKA numbers representing 4.6% of the TKA population. The Australian Joint Registry reports UKA represents 7.5% of the TKA numbers [7]. It is possible surgeon proficiency could affect the outcome of UKA; however, surgeon-specific differences in outcome were not identified. We believe our patient cohorts reflect the

relative UKA/TKA usage in the general population having knee arthroplasty. Furthermore, changes in score were employed to allow comparisons between groups. Second, the data derived for this study span three decades. While this is almost certainly a confounding factor with respect to temporal variations for operative indications, technique, and implants, the same confounding variables are applied to both prostheses. The orthopaedic community's understanding of prosthetic design and surgical technique has evolved over that time period. Notwithstanding the narrower indications for UKA, TKA is utilized in all remaining cases of knee arthrosis requiring surgical intervention. The longevity of a database is the premise of its use and is an invaluable tool for observing large patient populations over a period of time. Third, UKA demonstrates a higher failure rate but tends to have a marginally better function. This effect can be ascribed to nonmatched patient cohorts as UKAs are performed in younger, leaner males with less severe disease confined to a single compartment, better ultimate function, and the capacity for disease progression in the lateral compartment. The perception that UKA is revised more frequently than TKA because it is easier to do so is not consistent with our study or that of the Australian Joint Registry [7]. Pain of unknown origin accounted for 6.2% and 6.0% in UKA and TKA revisions, respectively, in the Australian Joint Registry [7]. Only 0.7% of our UKA series were revised for pain of unknown origin compared with 0.1% for TKA. Most revisions were undertaken in both groups for definable indications.

Improvement in function is well documented in both TKA and UKA [2, 3, 5, 15, 40, 42, 55, 57, 64]. UKA advocates purport a marginally better functional outcome over TKA [5, 20, 62]. This study confirms previous reports of patients with UKA having higher absolute postoperative clinical outcome measures (SF-12, WOMAC, KSCRS). These patients, however, also had higher preoperative scores and it is the change in score (preoperative versus postoperative) that determines the effect of the intervention. The change in score for all WOMAC and SF-12 domains demonstrated no difference between the groups. The changes in KSCRS score were not different between the groups for total scores and favored TKA over UKA for knee scores and UKA over TKA for function scores. When changes in scores are considered, both interventions are equally effective in improving function. When comparing outcomes between these interventions, it is critical to evaluate change in scores and not raw postoperative scores alone. It is also important to recognize there is a ceiling effect to scoring systems currently in use when evaluating patients with knee arthroplasty postoperatively. Often the final scores are at, or close to, the maximum score. Therefore, subtle improvements, or advantages, of one implant or another would not be perceived due to this ceiling effect.

Early survival studies of UKA demonstrated revision rates of 15% to 28% [23, 34, 44]. More recent articles report substantial increases in survival, with midterm survivorship for UKA of 84% to 98% (Table 1) [10, 12, 14, 19, 39, 40, 42, 46, 48, 57–61]. O'Rourke et al. [43] followed by Price and Svard [49] documented long-term survival rates of UKA at 20 years of 84% and 91%, respectively. TKA has established survivorship of 92% to 100% in long-term studies [2, 3, 5, 15, 19, 28, 52, 53]. We demonstrated cumulative revision rates of 13.3% and 7.1% at a mean followup of 7.12 ± 4.45 years and 6.50 ± 3.96 years for UKA and TKA, respectively. Kaplan-Meier survivorship at 10 years was 94.9% for TKA and 90.4% for UKA (Fig. 1). This study is consistent with the most recent Australian Joint Registry cumulative revision rates at 9 years of 5.1% and 13.4% for TKA and UKA, respectively [7]. Our study and the body of literature confirm TKA is the more durable therapeutic intervention.

In summary, consistent with the literature, we found patients with UKA had higher absolute clinical outcome scores for function preoperatively and at followup than patient with TKA. Change in clinical outcome scores were similar in both groups. The durability of both prostheses was assessed by survival analysis, showing TKA was revised less frequently than UKA.

Acknowledgments The authors thank Drs Robert Bourne and Jim McAuley for their guidance, leadership, and words of wisdom.

References

1. Abdeen AR, Collen SB, Vince KG. Fifteen-year to 19-year follow-up of the Insall-Burstein-I total knee arthroplasty. *J Arthroplasty*. 2010;25:173–178.
2. Ackroyd CE, Whitehouse SL, Newman JH, Joslin CC. A comparative study of the medial Sled and Kinematic total knee arthroplasties: ten-year survivorship. *J Bone Joint Surg Br*. 2002;84:667–672.
3. Aglietti P, Buzzi R, De Felice R, Giron F. The Insall-Burstein total knee replacement in osteoarthritis: a 10-year minimum follow-up. *J Arthroplasty*. 1999;14:560–565.
4. Akizuki S, Mueller JK, Horiuchi H, Matsunaga D, Shibakawa A, Komistek RD. In vivo determination of kinematics for subjects having a Zimmer Unicompartamental High Flex Knee System. *J Arthroplasty*. 2009;24:963–971.
5. Amin AK, Patton JT, Cook RE, Gaston M, Brenkel IJ. Unicompartamental or total knee arthroplasty? Results from a matched study. *Clin Orthop Relat Res*. 2006;451:101–106.
6. Argenson JN, Chevrol-Benkedache Y, Aubaniac JM. Modern unicompartamental knee arthroplasty with cement: a three to ten-year follow-up study. *J Bone Joint Surg Am*. 2002;84:2235–2239.
7. Australian Orthopaedic Association National Joint Replacement Registry. Annual Report. Adelaide, Australia: AOA; 2010. Available at: www.dmac.adelaide.edu.au/aoanjrr/. Accessed March 11, 2011.
8. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol*. 1988;15:1833–1840.
9. Colizza WA, Insall JN, Scuderi GR. The posterior stabilized total knee prosthesis: assessment of polyethylene damage and osteolysis after a ten-year-minimum follow-up. *J Bone Joint Surg Am*. 1995;77:1713–1720.
10. Deshmukh RV, Scott RD. Unicompartamental knee arthroplasty: long-term results. *Clin Orthop Relat Res*. 2001;392:272–278.
11. Diduch DR, Insall JN, Scott WN, Scuderi GR, Font-Rodriguez D. Total knee replacement in young, active patients: long-term follow-up and functional outcome. *J Bone Joint Surg Am*. 1997;79:575–582.
12. Dudley TE, Gioe TJ, Sinner P, Mehle S. Registry outcomes of unicompartamental knee arthroplasty revisions. *Clin Orthop Relat Res*. 2008;466:1666–1670.
13. Emerson RH Jr, Higgins LL. Unicompartamental knee arthroplasty with the Oxford prosthesis in patients with medial compartment arthritis. *J Bone Joint Surg Am*. 2008;90:118–122.
14. Emmerson KP, Moran CG, Pinder IM. Survivorship analysis of the Kinematic Stabilizer total knee replacement: a 10- to 14-year follow-up. *J Bone Joint Surg Br*. 1996;78:441–445.
15. Epinette JA, Manley MT. Hydroxyapatite-coated total knee replacement: clinical experience at 10 to 15 years. *J Bone Joint Surg Br*. 2007;89:34–38.
16. Fetzer GB, Callaghan JJ, Templeton JE, Goetz DD, Sullivan PM, Kelley SS. Posterior cruciate-retaining modular total knee arthroplasty: a 9- to 12-year follow-up investigation. *J Arthroplasty*. 2002;17:961–966.
17. Font-Rodriguez DE, Scuderi GR, Insall JN. Survivorship of cemented total knee arthroplasty. *Clin Orthop Relat Res*. 1997;345:79–86.
18. Gill GS, Joshi AB. Long-term results of Kinematic Condylar knee replacement: an analysis of 404 knees. *J Bone Joint Surg Br*. 2001;88:355–358.
19. Gioe TJ, Killeen KK, Hoeffel DP, Bert JM, Comfort TK, Scheltens K, Mehle S, Grimm K. Analysis of unicompartamental knee arthroplasty in a community-based implant registry. *Clin Orthop Relat Res*. 2003;416:111–119.
20. Griffin T, Rowden N, Morgan D, Atkinson R, Woodruff P, Maddern G. Unicompartamental knee arthroplasty for the treatment of unicompartamental osteoarthritis: a systematic study. *ANZ J Surg*. 2007;77:214–221.
21. Hollinghurst D, Stoney J, Ward T, Gill HS, Newman JH, Murray DW, Beard DJ. No deterioration of kinematics and cruciate function 10 years after medial unicompartamental arthroplasty. *Knee*. 2006;13:440–444.
22. Hopper GP, Leach WJ. Participation in sporting activities following knee replacement: total versus unicompartamental. *Knee Surg Sports Traumatol Arthrosc*. 2008;16:973–979.
23. Insall J, Aglietti P. A five to seven-year follow-up of unicompartmental arthroplasty. *J Bone Joint Surg Am*. 1980;62:1329–1337.
24. Insall JN, Dorr LD, Scott RD, Scott WN. Rationale of the Knee Society Clinical Rating System. *Clin Orthop Relat Res*. 1989;248:13–14.
25. Isaac SM, Barker KL, Danial IN, Beard DJ, Dodd CA, Murray DW. Does arthroplasty type influence knee joint proprioception? A longitudinal prospective study comparing total and unicompartamental arthroplasty. *Knee*. 2007;14:212–217.
26. Jahromi I, Walton NP, Dobson PJ, Lewis PL, Campbell DG. Patient-perceived outcome measures following unicompartamental knee arthroplasty with mini-incision. *Int Orthop*. 2004;28:286–289.
27. Jeer PJ, Cossey AJ, Keene GC. Haemoglobin levels following unicompartamental knee arthroplasty: influence of transfusion practice and surgical approach. *Knee*. 2005;12:358–361.
28. Keating EM, Meding JB, Faris PM, Ritter MA. Long-term followup of nonmodular total knee replacements. *Clin Orthop Relat Res*. 2002;404:34–39.

29. Koskinen E, Eskelinen A, Paavolainen P, Pulkkinen P, Remes V. Comparison of survival and cost-effectiveness between unicompartmental arthroplasty and total knee arthroplasty in patients with primary osteoarthritis: a follow-up study of 50,493 knee replacements from the Finnish Arthroplasty Register. *Acta Orthop*. 2008;79:499–507.
30. Koskinen E, Paavolainen P, Eskelinen A, Pulkkinen P, Remes V. Unicompartmental knee replacement for primary osteoarthritis: a prospective follow-up study of 1,819 patients from the Finnish Arthroplasty Register. *Acta Orthop*. 2007;78:128–135.
31. Kostamo T, Bourne RB, Whittaker JP, McCalden RW, MacDonald SJ. No difference in gender-specific hip replacement outcomes. *Clin Orthop Relat Res*. 2009;467:135–140.
32. Levine WN, Ozuna RM, Scott RD, Thornhill TS. Conversion of failed modern unicompartmental arthroplasty to total knee arthroplasty. *J Arthroplasty*. 1996;11:797–801.
33. Lewold S, Robertsson O, Knutson K, Lidgren L. Revision of unicompartmental knee arthroplasty: outcome in 1,135 cases from the Swedish Knee Arthroplasty study. *Acta Orthop Scand*. 1998;69:469–474.
34. Lindstrand A, Stenstrom A, Lewold S. Multicenter study of unicompartmental knee revision: PCA, Marmor, and St Georg compared in 3,777 cases of arthrosis. *Acta Orthop Scand*. 1992;63:256–259.
35. MacDonald SJ, Charron KD, Bourne RB, Naudie DD, McCalden RW, Rorabeck CH. The John Insall Award. Gender-specific total knee replacement: prospectively collected clinical outcomes. *Clin Orthop Relat Res*. 2008;466:2612–2616.
36. McAllister CM. The role of unicompartmental knee arthroplasty versus total knee arthroplasty in providing maximal performance and satisfaction. *J Knee Surg*. 2008;21:286–292.
37. McAuley JP, Engh GA, Ammeen DJ. Revision of failed unicompartmental knee arthroplasty. *Clin Orthop Relat Res*. 2001;392:279–282.
38. McCalden RW, Charron KD, MacDonald SJ, Bourne RB, Naudie DD. Does morbid obesity affect the outcome of total hip replacement? An analysis of 3290 THRs. *J Bone Joint Surg Br*. 2011;93:321–325.
39. Murray DW, Goodfellow JW, O'Connor JJ. The Oxford medial unicompartmental knee arthroplasty: a ten-year survival study. *J Bone Joint Surg Br*. 1998;80:983–989.
40. Naudie D, Guerin J, Parker DA, Bourne RB, Rorabeck CH. Medial unicompartmental knee arthroplasty with the Miller-Gallante prosthesis. *J Bone Joint Surg Am*. 2004;86:1931–1935.
41. Newman J, Pydisetty RV, Ackroyd C. Unicompartmental or total knee replacement: the 15-year results of a prospective randomised controlled trial. *J Bone Joint Surg Br*. 2009;91:52–57.
42. Newman JH, Ackroyd CE, Shah NA. Unicompartmental or total knee replacement? Five-year results of a prospective randomised trial of 102 osteoarthritic knees with unicompartmental arthritis. *J Bone Joint Surg Br*. 1998;80:862–865.
43. O'Rourke MR, Gardner JJ, Callaghan JJ, Liu SS, Goetz DD, Vittetoe DA, Sullivan PM, Johnston RC. The John Insall Award. Unicompartmental knee replacement: a minimum twenty-one-year followup, end-result study. *Clin Orthop Relat Res*. 2005;440:27–37.
44. Palmer SH, Morrison PJ, Ross AC. Early catastrophic tibial component wear after unicompartmental knee arthroplasty. *Clin Orthop Relat Res*. 1998;350:143–148.
45. Parsch D, Krüger M, Moser MT, Geiger F. Follow-up of 11–16 years after modular fixed-bearing TKA. *Int Orthop*. 2009;33:431–435.
46. Perkins TR, Gunckle W. Unicompartmental knee arthroplasty: 3 to 10 year results in a community hospital setting. *J Arthroplasty*. 2002;17:293–297.
47. Petrou G, Petrou H, Tilkeridis C, Stavrakis T, Kapetsis T, Kremmidas N, Gavras M. Medium-term results with a primary cemented rotating-hinge total knee replacement: a 7- to 15-year follow-up. *J Bone Joint Surg Br*. 2004;86:813–817.
48. Price AJ, Short A, Kellett C, Beard D, Gill H, Pandit H, Dodd CA, Murray DW. Ten-year in vivo wear measurement of a fully congruent mobile bearing unicompartmental knee arthroplasty. *J Bone Joint Surg Br*. 2005;87:1493–1497.
49. Price AJ, Svard U. A second decade lifetable survival analysis of the Oxford unicompartmental knee arthroplasty. *Clin Orthop Relat Res*. 2011;469:174–179.
50. Price AJ, Webb J, Topf H, Dodd CA, Goodfellow JW, Murray DW. Rapid recovery after Oxford unicompartmental arthroplasty through a short incision. *J Arthroplasty*. 2001;16:970–976.
51. Rajgopal V, Bourne RB, Chesworth BM, MacDonald SJ, McCalden RW, Rorabeck CH. The impact of morbid obesity on patient outcomes after total knee arthroplasty. *J Arthroplasty*. 2008;23:795–800.
52. Ranawat CS, Flynn WF Jr, Saddler S, Hansraj KK, Maynard MJ. Long-term results of the total condylar knee arthroplasty: a 15-year survivorship study. *Clin Orthop Relat Res*. 1993;286:94–102.
53. Ritter MA, Berend ME, Meding JB, Keating EM, Faris PM, Crites BM. Long-term followup of anatomic graduated components posterior cruciate-retaining total knee replacement. *Clin Orthop Relat Res*. 2001;388:51–57.
54. Robertsson O, Borgquist L, Knutson K, Lewold S, Lidgren L. Use of unicompartmental instead of tricompartmental prostheses for unicompartmental arthrosis in the knee is a cost-effective alternative: 15,437 primary tricompartmental prostheses were compared with 10,624 primary medial or lateral unicompartmental prostheses. *Acta Orthop Scand*. 1999;70:170–175.
55. Rougraff BT, Heck DA, Gibson AE. A comparison of tricompartmental and unicompartmental arthroplasty for the treatment of gonarthrosis. *Clin Orthop Relat Res*. 1991;273:157–164.
56. Saldanha KA, Keys GW, Svard UC, White SH, Rao C. Revision of Oxford medial unicompartmental knee arthroplasty to total knee arthroplasty—results of a multicentre study. *Knee*. 2007;14:275–279.
57. Squire MW, Callaghan JJ, Goetz DD, Sullivan PM, Johnston RC. Unicompartmental knee replacement: a minimum 15 year followup study. *Clin Orthop Relat Res*. 1999;367:61–72.
58. Steele RG, Hutabarat S, Evans RL, Ackroyd CE, Newman JH. Survivorship of the St Georg Sled medial unicompartmental knee replacement beyond ten years. *J Bone Joint Surg Br*. 2006;88:1164–1168.
59. Svard UC, Price AJ. Oxford medial unicompartmental knee arthroplasty: a survival analysis of an independent series. *J Bone Joint Surg Br*. 2001;83:191–194.
60. Tabor OB Jr, Tabor OB. Unicompartmental arthroplasty: a long-term follow-up study. *J Arthroplasty*. 1998;13:373–379.
61. Vorlat P, Putzeys G, Cottenie D, Van Isacker T, Pouliart N, Handberg F, Casteleyn PP, Gheysen F, Verdonk R. The Oxford unicompartmental knee prosthesis: an independent 10-year survival analysis. *Knee Surg Sports Traumatol Arthrosc*. 2006;14:40–45.
62. Walton NP, Jahromi I, Lewis PL, Dobson PJ, Angel KR, Campbell DG. Patient-perceived outcomes and return to sport and work: TKA versus mini-incision unicompartmental knee arthroplasty. *J Knee Surg*. 2006;19:112–116.
63. Ware J Jr, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care*. 1996;34:220–233.
64. Whittaker JP, Naudie DD, McAuley JP, McCalden RW, MacDonald SJ, Bourne RB. Does bearing design influence midterm survivorship of unicompartmental arthroplasty? *Clin Orthop Relat Res*. 2010;468:73–81.