Esthetic Evaluation of the Treated Unilateral Cleft Lip Using Photographs and Image Analysis Software: A Retrospective Study

S. Sabitha · M. Veerabahu · B. Vikraman

Received: 30 October 2010 / Accepted: 30 April 2011 / Published online: 28 June 2011
© Association of Oral and Maxillofacial Surgeons of India 2011

Abstract

Aim The aim of this study was to objectively evaluate the surgical outcome of unilateral cleft lip patients treated by Dr. Adenwalla's modification of Millard's rotation advancement technique using image analysis software.

Method Of the unilateral cleft lip cases done by a single surgeon using Dr. Adenwalla's modification of Millard's rotation advancement technique, only 11 cases satisfied the photographic standardization criteria (frontal view with both ears visible to minimize rotation and least nostril show to minimize tilt) and were included in this study. These photographs were acquired into the imageJ Software and Farkas cleft-related landmarks were marked on the lip area. The values of treated cleft side and the normal side were used to work out a Cleft Lip Component Symmetry Index. A value of 100 would indicate perfect symmetry of the lip. A value of either less or more would mean a degree of asymmetry.

Result The horizontal lip length, vertical lip length from the top of the Cupid's bow to subnasale and nostril floor width were slightly increased on treated side of the lip and the vertical lip length from the top of the Cupid's bow to the alar base, the Cupid's bow width and vermillion length on the treated side were slightly decreased when compared to the normal side. Even though there were some differences, it was not statistically significant.

The horizontal and vertical lip lengths, the Cupid's bow width, vermillion length and nostril floor width were almost symmetrical.

Conclusion Computer assisted anthropometric analysis of photographs using imageJ software demonstrates that Dr. Adenwalla's modification of Millard's rotation and advancement technique produces esthetic lip symmetry consistently.

Keywords Esthetic evaluation · Cleft lip- photographs · Image analysis software

Introduction

Cleft lip is one of the most common congenital anomalies. Its surgical repair is the most challenging procedure encountered by surgeons. The goal in primary unilateral cleft lip surgery is to recreate normal lip architecture, which is esthetic and symmetrical to the normal side. The desire of every surgeon is the restoration of the Cupid's bow, the flawless continuity of the vermillion free border and the body of the upper lip.

Over the decades, in pursuit of esthetic primary correction of unilateral cleft lip, various techniques have been proposed (Le Mesurier 1949, Tennison 1952, Cronin 1966, Millard 1976, Noordhoff 1984) [1–3]. Each technique has its own theory supporting the procedure. The Millard's rotation advancement flap is the most widely used technique but has some disadvantages of residual notching and absence of muscle in the repaired region. Based on this Dr. Adenwalla proposed a technique which includes

1. Producing adequate rotation and advancement following Millard's technique.

S. Sabitha (✉)

Vinayaka Mission's Sankarachariyar Dental College
and Hospital, NH 47, Sankari Main Road, Ariyanoor,
Salem 636 308, Tamil Nadu, India
e-mail: surgsabi@gmail.com

M. Veerabahu · B. Vikraman

Department of Oral and Maxillofacial Surgery, Ragas Dental
College and Hospital, Chennai, Tamil Nadu, India

2. De-epithelializing the vermillion flaps and leaving a filler of muscles on both the edges.
3. Dissecting the vermillion muscle from skin and mucosa for 3–4 mm using fine scissors.
4. Approximating these muscles with three 6-0-ethilon sutures.
5. Using 'Z' plasty on the inner aspect of the mucosa to reduce constriction of the free border.

Some of the methods for assessment of the repaired cleft are Direct surface assessment [4, 5], two-dimensional photographs, [6, 7] three-dimensional imaging [8–10] or a combination of these methods. Objective analysis is commonly performed using an anthropometric method. Esthetic evaluation of the repaired cleft lip has been done by considering landmarks related to the nose, the body of the lip and the vermillion.

Anthropometry and clinical examination were considered best to evaluate the morphology of the repaired cleft lip and nose. With advances in science and technology, the use of the computer and digital imaging now plays a major role in any scientific evaluation, so also in the evaluation of morphology of a repaired cleft lip. Coghlan et al. [11–13] described various studies on computer-based analysis to assess the facial symmetry of treated cleft lip and palate patients. The validity of these tools has been confirmed.

In this study, using ImageJ software (<http://rsb.info.nih.gov/nih-image>), (<http://rsb.info.nih.gov/ij/download.html>) as a quantitative assessment tool, we objectively evaluated the esthetic outcome of the unilateral cleft lip, treated by the same surgeon over a seven year period using Dr. Adenwalla's modification of Millard's rotation advancement technique.

Materials and Methods

The photographic slides of eleven cases of treated unilateral cleft lip by Dr. Adenwalla's modification of Millard's Rotation and Advancement were retrieved from the records. The photographs were taken at least one year post operatively. Photographs that satisfied the following criteria for standardization (frontal view with both ears visible to minimize rotation and least nostril show to minimize tilt) were selected, scanned and acquired into the imageJ program with a measuring plugin. Farkas' [14, 15] cleft-related anthropometric lip landmarks were marked on the enlarged images of the lip area (Fig. 1).

Using these points, the anthropometric measurements of the lip were made using linear measuring tool in ImageJ software (Figs. 2, 3). These linear measurements give an indication of the horizontal and vertical lip length, the

Fig. 1 *ac* (alar curvature) The point most lateral in the curved line of each ala towards the cheek junction. *sn* (subnasale) Midline junction of the columella and lip. *cph* (crista philtri) Lateral high points of the Cupid's bow. *ls* (labiale superius) Lowest point of the Cupid's bow. *ch* (cheilion) Point at the lateral labial commissure

vertical vermillion length as well as the width of the Cupid's bow and nostril floor.

Methodology

These parameters are used to work out a Cleft Lip Component Symmetry Index that could be used to compare objectively the various components of the lip and nose [16].

Cleft Lip Component Symmetry Index

$$= \frac{\text{Cleft side component value}}{\text{Normal side component value}} \times 100$$

Fig. 2 *ch–cph* Point at the lateral labial commissure to the lateral high points of the Cupid's bow (Horizontal lip length). *cph–ac* Point at the lateral high points of the Cupid's bow to the point most lateral point in the curved line of each ala toward the cheek junction (Vertical lip length). *cph–sn* Lateral high points of the Cupid's bow to the midline junction of the columella and lip (Vertical lip length). *cph–ls* Lateral high points of the Cupid's bow to the low point of the Cupid's bow (Width of the Cupid's bow). *cph–redline point* (⊥) Lateral high points of the Cupid's bow to the redline point perpendicular to it (Vertical vermillion length). *sn–ac* Midline junction of the columella and lip to the most lateral point in the curved line of each ala toward the cheek junction (Nostril floor width)

Fig. 3 Measuring the parameters using imageJ software

A value of 100 would indicate perfect symmetry of the lip. A value of either less or more would mean a degree of asymmetry [16].

Data analysis was done using t test and correlation. *P* value was set at 95% confidence interval and below 0.05 was considered significant. *P* value above 0.05 showed that there is no statistical difference between the treated and the normal side.

Results

The sample included 8 males and 3 females with age ranging from 1 to 5 years. Nine patients had cleft on the left side and two with cleft on the right. All the parameters were measured using ImageJ software. The values of treated cleft side and the normal side were used to work out a Cleft Lip Component Symmetry Index that could be used to compare objectively of the various components of the lip and nose (Table 1).

Results of the data analysis were tabulated (Tables 2, 3) (Graph 1).

The mean ratio for the horizontal lip length (ch–cph), vertical lip lengths (cph–ac) (cph–sn), the cupid's bow width (cph–ls), vermillion length (cph–redline point (⊥)), and nostril floor width (ac–sn) are 104.64 (SD: 10.74) (*P*-value: 0.183); 94.29 (SD: 11.84) (*P*-value: 0.466); 101.96 (SD: 16.26) (*P*-value: 0.697); 96.25 (SD: 10.96) (*P*-value: 0.283); 95.30 (SD: 13.60) (*P*-value: 0.279); 112.80 (SD: 25.67) (*P*-value: 0.129) respectively

Discussion

Though there are many surgical techniques to repair the cleft lip, the Millard's procedure has found universal

Table 1 Cleft lip component symmetry index for the parameters

Patient	ch–cph	cph–ac	cph–sn	cph–ls	cph–Red line point (⊥)	ac–sn
1	100.5	97.82	98.46	109.09	94.84	158.13
2	113.98	93.04	87.67	94.52	86.88	117.9
3	98.03	95.52	100	94.11	105.88	77.1
4	116.16	102.631	75.9	93.33	120	122.66
5	100.8	100	108.23	117.5	100	108.33
6	89.51	80.58	140.84	83.92	104.47	102.72
7	117	116.25	113.53	107.27	102.7	101.28
8	90	93.57	107.14	82	94.33	71.68
9	104.3	118.03	90	97.6	89.09	113.3
10	113.08	83	103.4	88.37	100	146.26
11	101.4	89.77	106.5	90.9	88.88	121.3

Table 2 *t*-Test for the parameters

Parameter	Mean	SD	<i>t</i> -Value	<i>P</i> value
ch–cph	104.64	10.474	1.433	0.183
cph–ac	94.29	11.84	−0.758	0.466
cph–sn	101.96	16.26	0.44	0.697
cph–ls	96.25	10.96	−1.136	0.283
cph–redline point (⊥)	95.30	13.60	−1.146	0.279
ac–sn	112.8	25.67	1.654	0.129

Table 3 Correlation for the parameters

Parameter	Correlation	<i>P</i> value
ch–cph	0.881	0.000**
cph–ac	0.807	0.003*
cph–sn	0.889	0.000**
cph–ls	0.906	0.000**
cph–redline point (⊥)	0.906	0.000**
ac–sn	0.212	0.532

** Significant at 1% level

* Significant at 5% level

acceptance. Dr. Adenwalla of Trichur has modified or refined the Millard's technique to prevent the vermillion notching that can occur when using the standard technique. Since then Dr. Adenwalla's modified Millard's rotation advancement technique has been found to address all the aspects of lip symmetry.

Over the last two decades, there has been increasing interest in the short and long term assessment of the quality of cleft care. The surgical outcome can be analyzed subjectively or objectively. A variety of methods have been used in assessment of the repaired cleft associated deformity. These include direct surface assessment, two-dimensional

Graph 1 Mean ratio with standard deviation for the parameters

photographs, three-dimensional imaging or a combination of these methods [17].

Although direct facial assessment has allowed angular and linear measurement to be established, the method is very time consuming and does not permit the clinician to revisit the data without the presence of the patient. On the other hand, most investigators have used two-dimensional photography, because it provides a simple and quick method of depicting the face. These photographic techniques, however, present the three-dimensional face in two dimensions. Standardization of the views necessary for comparison of assessment is difficult, especially in younger subjects. As a result, attempts have been made to develop three-dimensional methods that represent the human face more realistically. However, the natural facial texture of the normal face has not been factored in previous imaging systems. This deficiency has been addressed in a recent three-dimensional imaging system with the added advantage that it also captures the facial image in 50 seconds [17].

With advances in science and technology the use of the computer and digital imaging now plays a major role in any scientific evaluation. It has been used with good effect in the evaluation of the morphology of a repaired cleft lip. Coghlan et al. [11–13] have objectively analyzed symmetry of treated facial cleft patients using follow up photographs and computer analysis. Hurwitz et al. [18] compared the accuracy of NIH-Image based anthropometry with direct measurements of 22 linear distances on the lip and nose. Their study showed that there was strict linear correlation between computer aided and direct measurements.

Objective evaluation of the overall results of cleft lip repair can be difficult, because esthetic appearance is largely a subjective phenomenon. However, the symmetry of the architecture of the lip and nose can be evaluated by mathematical means [16]. In our study, frontal view photographs were retrieved from the records and reformatted using the Image analysis software to evaluate symmetry of the surgically treated unilateral cleft lip. The linear

measurements of heights and widths of certain important features of the lip and nose were used to assess symmetry.

When photographs are taken from varying distances with lenses of different focal lengths the magnification of the final image is unknown. These are then considered unsuitable for absolute measurements, unless standardized procedures are followed to ensure a consistent, known magnification. When calculating proportions the magnification factor is not relevant. Even in the absence of standardized views it is possible to measure proportional ratios [6]. In this study, we objectively evaluated symmetry of treated unilateral cleft lip for esthetic outcome by taking proportions between treated side and normal side [19]. The imageJ software (<http://rsb.info.nih.gov/ij/download.html>) helped us to magnify all the images by a factor of ten and then analyzed the proportions between certain predetermined points on the lip and nose.

Lee TJ [5] studied the growth ratios of vertical, horizontal, and nostril sill dimensions by comparing the cleft side and the non cleft side of the same patient after Modified Rotation Advancement flap repair in unilateral cleft lip patients. He found that the repaired unilateral cleft retained its vertical and horizontal dimensions, determined at the time of the initial repair. Only the ratio of the nostril sill width between cleft side and non-cleft sides changed with time. In our study photographs of 11 patients, with a minimum of one-year follow up, were transferred and worked on using ImageJ software and the resultant data obtained was analyzed.

Image analysis showed that horizontal lip length, vertical lip length from the top of the Cupid's bow to subnasale and nostril floor width were slightly increased on the treated side of the lip when compared to the normal side of the lip. Similarly the vertical lip length from the top of the Cupid's bow to the alar base, the Cupid's bow width and vermillion length on the treated side were slightly decreased when compared to the normal side.

Even though there were some differences, it was not statistically significant. The horizontal and vertical lip lengths, the Cupid's bow width, vermillion length and nostril floor width were almost symmetrical. Hence it does not reject the null hypothesis.

Facial plastic surgeons have always relied on photography for preoperative and postoperative analysis. The current standards for analysis and documentation of facial structures have been determined by obtaining measurements and parameters from two-dimensional photographs or radiographs. However, the face and neck are three-dimensional structures and require manipulation in three planes within the constraints of esthetics, stability, and function. Recent advances in technology have made three-dimensional imaging and analysis possible. The ability to capture images in three dimensions has opened up new

avenues for observation and has increased the ability to analyze changes after surgery. Recently various studies have focused on descriptions of imaging techniques, validation of measurements, and the creation of normative data and its applications in surgery. So that, the availability of three-dimensional imaging and software systems presents new opportunities for the facial cosmetic surgeon to plan, execute, and assess the outcomes in patients undergoing surgery of the face and neck [20].

Conclusion

Image analysis showed that horizontal lip length, vertical lip length from the top of the Cupid's bow to subnasale and nostril floor width were slightly increased on the treated side of the lip when compared to the normal side and the vertical lip length from the top of the Cupid's bow to the alar base, Cupid's bow width and vermillion length on the treated side were slightly decreased when compared to the normal side. Even though there were some differences, it was not statistically significant. The horizontal and vertical lip lengths, Cupid's bow width, vermillion length and nostril floor width were almost symmetrical. So this technique can be considered to provide an esthetically good outcome.

In conclusion, computer assisted anthropometric analysis of photographs using imageJ software demonstrates that Dr. Adenwalla's modification of Millard's rotation and advancement technique produces esthetic lip symmetry consistently. This technique deserves to be widely reported.

Recent Innovation

Recently, the electromagnetic digitizer used and provides the three-dimensional coordinates of landmarks. This method therefore, could couple the benefits of conventional anthropometry and computerized systems and thus provide a simple, fast and direct computerized anthropometry [21].

References

- Burt JD, Byrd HS (2000) Cleft lip: unilateral primary deformities. *Plast Reconstr Surg* 105:1043–1054
- Millard RD Jr (1976) Cleft craft: classification of clefts, vol 1. Little Brown, Boston, pp 41–57
- Nakajima H, Imanishi N, Aiso S (2002) Facial artery in the upper lip and nose: anatomy and a clinical application. *Plast Reconstr Surg* 109:855–860
- Lazarus DD, Hudson DA, Van Zyl JE (1998) Repair of unilateral cleft lip: a comparison of five techniques. *Ann Plast Surg* 41: 587–594
- Lee TJ (1999) Upper lip measurements at the time of surgery and follow up after modified rotation advancement flap repair in unilateral cleft lip patients. *Plast Reconstr Surg* 104:911–915
- Kohout MP, Aljaro LM, Farkas LG (1998) Photogrammetric comparison of two methods for synchronous repair of bilateral cleft lip and nasal deformity. *Plast Reconstr Surg* 102:1339–1349
- Cutting CB, Dayan JH (2003) Lip height and lip width after extended Mohler unilateral cleft lip repair. *Plast Reconstr Surg* 111:17–23
- Bush K, Antonyshyn Oleh (1996) Three-dimensional facial anthropometry using a laser surface scanner: validation of the technique. *Plast Reconstr Surg* 98:226–235
- Yamada T, Mori Y, Minami K, Mishima K (1999) Computer aided three-dimensional analysis of nostril forms: application in normal and operated cleft lip patients. *J Craniomaxillofac Surg* 27(6):345–353
- Yamada T, Mori Y, Sugahara T (2002) Three-dimensional facial morphology, following primary cleft lip repair using the triangular flap with or without rotation advancement. *J Craniomaxillofac Surg* 30:337–342
- Coghlan BA, Boorman JG (1996) Objective evaluation of the Tajima secondary cleft lip nose correction. *Br J Plast Surg* 49: 457–461
- Coghlan BA, Latiung JK, Pigott RW (1993) A computer aided method of measuring nasal symmetry in the cleft lip nose. *Br J Plast Surg* 46:18–21
- Coghlan BA, Matthews B, Pigott RW (1987) A computer based method of measuring facial asymmetry. Results from an assessment of the repair of cleft lip deformities. *Br J Plast Surg* 40: 371–376
- Duffy S, Noar JH, Evans RD (2000) Three-dimensional analysis of the child cleft face. *Cleft palate Craniofac J* 37:137–144
- Farkas LG, Hajnisk Posnick JC (1993) Anthropometric and anthroposcopic findings of the nasal and facial region in cleft patients before and after primary lip and palate repair. *Cleft Palate Craniofac J* 30(1):1–12
- Amaratunga S (1988) Comparison of Millard's and LeMesurier's method of repair of the complete unilateral cleft lip using a new symmetry index. *J Oral Maxillofac Surg* 46:353–356
- Al-Omari I, Millett DT, Ayoub A, Bock M (2003) An appraisal of three methods of rating facial deformity in patients with repaired complete unilateral cleft lip and palate. *Cleft Palate Craniofac J* 40:530–537
- Hurwitz DJ, Ashby ER, Llull R (1999) Computer-assisted anthropometry for outcome assessment of cleft lip. *Plast Reconstr Surg* 103:1608–1623
- Kawai T, Natsume N, Shibata H, Yamamoto T (1990) Three-dimensional analysis of facial morphology using moiré stripes: Part I. Method. *Int J Oral Maxillofac Surg* 19:356
- Honrado CP, Larrabee WF Jr (2004) Update in three-dimensional imaging in facial plastic surgery. *Curr Opin Otolaryngol Head Neck surg* 12(4):327–331
- Ferrario VF, Sforza C, Tartaglia GM (2003) A quantitative three-dimensional assessment of abnormal variations in facial soft tissues of adult patients with cleft lip and palate. *Cleft Palate Craniofac J* 40(5):544–549