On the substrate specificity of a damage-specific DNA binding protein from human cells

---

Ross S. Feldberg

Department of Biology, Tufts University, Medford, MA 02155, USA

Received 27 November 1979

ABSTRACT

The nature of the lesion recognized by a damage-specific DNA binding protein from human cells was investigated by examining the substrate specificity of this protein. Protein-recognizable damage was introduced into both T7 DNA and Poly d(A-T) at relatively low UV fluences. In addition, the protein demonstrated binding to both nitrous acid- and bisulfite-treated DNA, but not to DNA crosslinked with trioxsalen plus near-UV nor to non-irradiated uracil-containing DNA. These results suggest that this protein could be recognizing minor helix distortions in DNA rather than any one single lesion.

INTRODUCTION

The role of DNA damage in mutagenesis and carcinogenesis has been indicated in a variety of studies (1,2). Workers utilizing ultraviolet (UV) radiation as a damaging agent have generally focused on the cis, syn cyclobutyl pyrimidine dimer photoproduct and have established the quantitative significance and biological importance of this lesion (3,4). In recent years, however, a number of studies on the biological responses to UV-irradiation have implied that lesions other than the pyrimidine dimer also contributed to the observed cellular responses (5). Using UV-irradiated chick cells in culture, Paterson, et al. (6) found that even when essentially all the pyrimidine dimers were removed by photoreactivation, the cells demonstrated high levels of unscheduled DNA synthesis and repair replication. Similar results were reported by Lehmann and Stevens (7) who employed the inhibition of replicative DNA synthesis in UV-irradiated chick cells as a probe for DNA damage. The results obtained in both sets of experiments suggest the presence of some non-photoreparable lesion in UV-irradiated DNA. A similar interpretation could account for the discrepancy between the kinetics of dimer elimination and of repair synthesis in UV-irradiated human cells reported by Ehmann, et al. (8). Measuring a very different biological response, Lin and Howard-Flanders (9) found that photoreactivation decreased

UV-induced recombination in heteroimmune infections, but not in homoimmune infections of λ lysogens. These results indicate that, under non-replicating conditions, recombination is being induced by non-pyrimidine dimer lesions.

In previous work (10,11), I have reported on the isolation of a damage-specific DNA binding protein from human placenta which binds tightly to UV-irradiated DNA, and to photoreactivated UV-DNA but not to DNA in which thymine dimers have been introduced by near-UV irradiation in the presence of the photosensitizer, acetophenone. These results indicated that the protein was directed toward some photoproduct other than the cyclobutyl pyrimidine dimer. Although a number of endonucleases directed toward non-dimer damage in UV-irradiated DNA have been isolated (12-16), these enzymes are for the most part active only with heavily-irradiated DNA. In contrast, the damage-specific DNA binding protein binds tightly to DNA irradiated with low UV-doses. In this report, I examine the affinity of this protein toward DNA treated with several different damaging agents in order to gain some insight into the nature of the lesion being recognized.

MATERIALS AND METHODS

Binding Protein Preparation and Assay. The damage-specific DNA binding protein was isolated from the insoluble fraction of a sodium cholate extract (17) of human placenta by a modification of the procedure of Feldberg and Grossman and corresponds in purity to Fraction V (10). Filter binding assays were carried out as previously described (10). Because the purified DNA-binding protein contained levels of protein below the detectable limit, protein titration curves are expressed in μ l of protein added rather than μ g.

Nucleic Acids. Phage T7 DNA labeled with [^{3} H]-thymidine was prepared by the method of Richardson (18). [^{3} H]-uracil labeled DNA from the *B. subtilis* phage PBS2 was prepared as described by Friedberg, *et al.* (19). Poly d(A-T) was synthesized according to the procedure of Schachman *et al.* (20) using [^{3} H]-dATP to give a product of specific activity 5×10^7 cpm/ μ mole. Analysis of the neutral sucrose density gradient sedimentation profile of this polymer yielded a number-average molecular weight of 9×10^6 daltons.

Damage Introduction.

1. UV-Irradiation. A 15 watt germicidal lamp was employed. The fluence was measured with an International Light IL-500 radiometer with a SEE400 detector and either a NBS 254 or NBS 313 filter.

2. Nitrous Acid. T7 DNA was diluted to 1 μ g/ml in 0.2 M sodium acetate, pH 4.3, and was then incubated at 30°C for 3 hr in the absence

(control) or presence of 0.05 M NaNO_2 . The pH was then adjusted to 7.6 with 1/5 volume of 1 M Tris-base and the DNA was dialyzed overnight against 0.02 M Tris-HCl, 0.01 M NaCl, 0.50 mM EDTA (pH 8.0).

3. Trioxsalen Treatment. Photochemical crosslinking with near-UV irradiation was induced using a medium-pressure mercury lamp with a glass envelope (emission bands at 313 nm and 365 nm) and trioxsalen (4,5',8-trimethylpsoralen) at a ratio of trioxsalen to T7 DNA of 153:1. Irradiation was carried out at 10°C for 20 min at a 313 nm fluence of 6.0 J/M²/sec. The DNA was then dialyzed in the dark against 0.01 M Tris-HCl, 0.001 M EDTA (pH 8.0).

4. Bisulfite Treatment. T7 DNA was diluted to 1 µg/ml in either 2 M NaHSO_3 , 0.50 M sodium acetate, pH 5.4 (bisulfite-treated) or in 2 M NaCl, 0.50 M sodium acetate, pH 5.4 (control) and then incubated at 86°C for 15 min., quick-cooled, and dialyzed against 0.01 M Tris-HCl, 0.5 mM EDTA (pH 9.0).

Sucrose Density Gradient Centrifugation. Analysis of the bisulfite-, nitrous acid-, trioxsalen- and control-treated DNA samples was carried out by sedimentation through 5 ml 5-20% sucrose gradients in 0.30 M NaOH, 0.70 M NaCl, 0.001 M EDTA. Centrifugation was carried out in an SW 50.1 rotor in a Beckman L5-50 centrifuge at 4°C to a preset $\omega^2 t$ of 24×10^{10} rad²/sec (45,000 rpm; 3 hr). Twenty-four 7 drop fractions were collected from each gradient with 2 ml cold 1 N HCl and filtration through Millipore GF/A glass fiber filters. Filters were washed with 4 ml 1 N HCl, 1 ml 95% ethanol, dried under a heat lamp and counted.

The poly d(A-T) was analyzed by sedimentation through a neutral 5 ml 5-20% sucrose gradient (1 M NaCl, 0.10 M NaCl, 0.10 M Tris-HCl, 0.001 M EDTA, pH 8) to a preset $\omega^2 t$ of 34×10^{10} rad²/sec (22,500 rpm; 17 hrs). Number average molecular weights were determined from alkaline and neutral gradients using T7 DNA or ϕ X-174 RFI DNA as standards by the method of Charlesby (21).

MATERIALS

Bacillus subtilis strain SB19E and PBS2 phage were a gift of Dr. A. Price. Tritium-labelled thymidine (6.7 Ci/mmole) and [³H]-uracil (23Ci/mmole) were obtained from New England Nuclear, Boston, Mass., 8-[³H]-dATP from ICN and DNA Polymerase I from Boehringer-Mannheim.

RESULTS

Dose-Response of Binding. The effect of increasing UV fluence on the retention of nucleic acid in the filter binding assay is shown in Fig. 1. With protein present in excess, it is possible to use these data to determine the average number of protein binding sites per polymer, m , assuming that the damage is introduced randomly and that one damage site per polymer is sufficient for protein binding and subsequent retention on the filter. A value for m can be obtained from the Poisson distribution

$$m = \ln (1/P_0)$$

where P_0 is the fractional amount of polymer not retained on the filter at a given fluence. For T7 DNA, the rate of damage introduction thus calculated is 0.15 damage sites per 10 J/m^2 per 10,000 base pairs. If more than one damage site were necessary to ensure full retention of the polymer on the filter, this value would be an underestimate. With poly d(A-T) as a substrate, the rate of damage introduction is 0.08 sites per 10 J/m^2 per 10,000 base pairs.

Nitrous Acid Treatment. In addition to 254 nm radiation, several other agents were found to introduce protein-recognized lesions into T7 DNA. Nitrous acid is a powerful mutagen, deaminating guanine to xanthine, adenine to hypoxanthine and cytosine to uracil as well as introducing crosslinks of unknown structure (22). Fig. 2 is a protein titration curve for T7 DNA

Fig. 1. Dose-response curves for protein-nucleic acid interaction in the filter binding assay. (A) T7 DNA; 6500 cpm per assay, 1.0×10^{-4} μmole nucleotides. (B) Poly d(A-T); 8000 cpm per assay, 12×10^{-5} μmole nucleotides.

Fig. 2. Protein titration of T7 DNA. $\square$ — $\square$ Irradiated with 18 J/m^2 254 nm light, $\triangle$ — $\triangle$ Treated with 0.05 M NaNO_2 at pH 4.3, 3 hrs. $\circ$ — $\circ$ Incubated at pH 4.3, 3 hrs. Each assay contained 7200 cpm; 11×10^{-5} μmole nucleotides.

either irradiated to 18 J/m^2 or treated at pH 4.3 and 37°C for 3 hr in the presence or absence of 0.05 M NaNO_2 . From the fraction of DNA not bound at protein saturation and the Poisson distribution I have calculated that under these conditions nitrous acid introduces an average of 0.7 protein-recognizable lesions per T7 molecule. Alkaline sucrose density gradient centrifugation (data not shown) revealed that the pH 4.3-treated control DNA contained 1.6 alkali-labile sites per T7 molecule, presumably apurinic sites (23). Since this control DNA was a very poor substrate for the binding protein, these results indicate that the protein does not bind to apurinic sites in DNA.

Trioxsalen-Crosslinked DNA. To examine the possibility that the protein recognized DNA crosslinks, I employed the photochemical crosslinking agent, trioxsalen (4,5',8-trimethylpsoralen). In the presence of near-UV light, this agent forms monoadducts with thymine residues and cross-linked diadducts between thymines on opposite strands of the DNA (24). After incubation of the T7 DNA with trioxsalen, extensive crosslinking was found in the irradiated sample, but not in the dark control nor in the sample irradiated with near-UV

light in the absence of trioxsalen (Fig. 3). When these samples were examined for protein binding (Table 1), it was found that although near-UV light alone introduced protein-recognizable damage, irradiation in the presence of trioxsalen gave no additional binding. I determined whether the lack of protein binding to the cross-linked DNA could have been due to either a gross helix distortion or to interference by the trioxsalen monoadducts by reirradiating this DNA with 254 nm light and reassaying it. As seen in Table 1, this sample was as good a substrate for the binding protein as was irradiated non-crosslinked DNA.

Bisulfite Treated DNA. Bisulfite is a highly specific deaminating agent which converts cytosine to uracil (25). To retain the double helical character of the DNA necessary for protein interaction, the bisulfite treatment was carried out at temperatures slightly below the T_m of the DNA. Under these conditions, the DNA was partially melted, a necessity for bisulfite action, but the strands were not completely separated. DNA treated in this way was found to be a good substrate for the binding protein, and an average of 0.60 protein-recognizable sites were introduced per T7 molecule (Fig. 4). Alkaline sucrose gradient centrifugation of the control incubated and

Fig. 3. Alkaline sucrose gradient sedimentation of T7 DNA treated with: trioxsalen + near-UV (Δ — Δ); near-UV alone ($\blacksquare$ — $\blacksquare$); trioxsalen in dark ($\circ$ — $\circ$) or no treatment ($\bullet$ — $\bullet$). Direction of sedimentation is from left to right.

Table 1. Binding Protein Recognition of Trioxsalen/Near-UV Treated T7 DNA.

	Trioxsalen	Near-UV	%DNA Bound to Filter ^a	
			—	+254 nm ^b (18 J/m ²)
1.	+	-	10%	46%
2.	-	+	20%	58%
3.	+	+	20%	58%
4.	-	-	2%	47%

^a Assays carried out with a saturating level of binding protein.

^b Irradiation with 254 nm light carried out following trioxsalen treatment and dialysis.

bisulfite incubated DNA (data not shown) indicated an average of 2.4 alkali-labile sites introduced per T7 molecule in both cases. The low level of binding to the control incubated DNA again demonstrates that the protein is unable to recognize apurinic lesions in DNA. The predominant product in

Fig. 4. Protein titration of T7 DNA, □—□ Irradiated with 18 J/m² 254 nm light. △—△ Treated with 2 M NaHSO₃, pH 5.4, 86°C, 15 min. ○—○ Treated with 2 M NaCl, pH 5.4, 86°C, 15 min. Each assay contained 5600 cpm; 9×10^{-5} µmoles nucleotides.

bisulfite-treated DNA will be a uracil opposite a guanine in a base mismatch. Since uracil is not a normal constituent of DNA, and since a number of enzymes which can excise it from double or single-stranded DNA have now been described (26), it seemed critical to determine whether the protein might simply be recognizing uracil *per se*.

Uracil-Containing Nucleic Acid. DNA from the uracil-containing *B. subtilis* phage, PBS2, was found to show little binding with the binding protein unless first irradiated with 254 nm light (Fig. 5). The high levels of filter binding in the absence of any added protein were found only with this DNA and were due to single stranded regions as indicated by the observation that a limited nuclease S1 digestion which decreased the total acid-precipitable counts by 20% reduced the background filter binding to 9%.

DISCUSSION

The presence of damage, other than the cyclobutyl pyrimidine dimer, in UV-irradiated DNA has been demonstrated by a number of studies on the biological responses to UV-irradiation. In addition, the discovery of nucleases

Fig. 5. Protein titration curve of [^3H]-U-PBS2 DNA. $\circ$ - $\circ$ Nonirradiated. Δ - Δ Irradiated to 160 J/m^2 254 nm light.

from both bacterial and eukaryotic sources directed toward non-dimer damage indicates that such damage can be removed in vivo. From the data presented here, I am able to draw several conclusions about the nature of the non-dimer lesion recognized by a human damage-specific DNA binding protein.

From our earlier work (10,11) with photoreactivated UV-irradiated DNA and with photosensitized near-UV irradiated DNA, we were able to conclude that the damage-specific binding protein showed no binding to the cyclobutyl pyrimidine dimer. In addition, the heat stability of the photoproduct and the inability of either pancreatic DNase or Micrococcal nuclease to introduce protein-recognizable lesions, indicated that the damage was neither a photohydrate nor single-strand breaks in the DNA backbone. I am now able to eliminate several other forms of DNA damage as possible substrates for the binding protein. Apurinic sites, introduced into the acid-incubated "control" DNAs in both the nitrous acid and the sodium bisulfite experiments, clearly did not act as substrates for the binding protein (Figs. 2 and 4). In addition, neither trioxsalen crosslinks nor trioxsalen monoadducts were recognized by the binding protein.

What then might we conclude about the lesion(s) recognized by the protein? First, it is clear that the damage is introduced at relatively low doses of UV-irradiation. From Fig. 1 a rate of damage-introduction into T7 DNA of 0.15 sites per 10 J/m^2 per 10,000 base pairs can be calculated. Comparison of this value to the rate of pyrimidine dimer introduction of 1.2 dimers per 10 J/m^2 per 10,000 base pairs as determined by Sancar and Rupert (27) suggests that the non-dimer lesion could play a biologically important role at "physiological" doses of UV-light. Traditionally, DNA damage has been classified in terms of the agent responsible for its introduction. However, this classification scheme obscures the fact that many agents introduce multiple forms of damage, as well as the fact that different agents can introduce similar or identical forms of damage. A more satisfying classification scheme has been proposed by Cerutti (28) and is based on the consequences of the lesion on the local DNA structure. This scheme divides DNA damage into three classes:

- Class I. Monofunctional base lesions causing negligible helix distortion.
- Class II. Mono- and difunctional base lesions causing minor helix distortion.
- Class III. Mono- and difunctional base lesions causing major helix distortions.

It is within the context of this scheme that the rather broad specificity of the binding protein can be best understood. In addition to UV-irradiated DNA, the protein is able to recognize damage in X-irradiated (10), nitrous acid treated and bisulfite treated DNA. In addition, we have found that the protein can bind to irradiated poly dT annealed to nonirradiated poly dA, as well as to irradiated poly dA annealed to nonirradiated poly dT (in preparation). Of these various treatments, the most specific is sodium bisulfite, which, under the conditions employed, would have resulted in the deamination of cytosine to yield a uracil mismatched opposite a guanine. The possibility that the protein simply recognizes the unusual base uracil in DNA is unlikely since nonirradiated PBS2 DNA was not a substrate. Taken together, these results suggest that, rather than recognizing any single lesion per se, the binding protein may recognize a helix distortion of the type introduced by a base mismatch.

ACKNOWLEDGEMENT

This work was supported by grant CA-19419 from the National Cancer Institute, DHEW, and by a Cottrell Research Grant from the Research Corporation. The technical assistance of Mrs. J.L. Lucas and the critical comments of Ms. J. Carew are gratefully acknowledged.

REFERENCES

1. Setlow, R.B. (1978) *Nature* 271, 713-717.
2. Trosko, J.E. and Chang, C-C (1978) *Quart. Rev. Biol.* 53, 115-141.
3. Fisher, G.J. and Johns, H.E. (1976) in *Photochemistry and Photobiology of Nucleic Acids*, S.Y. Wang, Ed., Vol. I, pp. 169-224, Academic Press, New York.
4. Jagger, J. (1976) in *Photochemistry and Photobiology of Nucleic Acids*, S.Y. Wang, Ed., Vol. II, pp. 147-186, Academic Press, New York.
5. Rahn, R.O. (1979) in *Photochemical and Photobiological Reviews*, K.C. Smith, Ed., Vol. 4, pp. 267-330, Plenum Press, New York.
6. Paterson, M.C., Lohman, P.H.M., DeWeerd-Kastelein, E.A. and Westerveld, A. (1974) *Biophys. J.* 14, 454-466.
7. Lehmann, A.R. and Stevens, S. (1975) *Biochim. Biophys. Acta* 402, 179-187.
8. Ehmann, U.K., Cook, K.H. and Friedberg, E.C. (1978) *Biophys. J.* 22, 249-264.
9. Lin, P.-F. and Howard-Flanders, P. (1976) *Molec. gen. Genet.* 146, 107-115.
10. Feldberg, R.S. and Grossman, L. (1976) *Biochemistry* 15, 2402-2408.
11. Feldberg, R.S. (1978) in *DNA Repair Mechanisms: ICN-UCLA Symposia on Molecular and Cellular Biology*, P.C. Hanawalt, E.C. Friedberg and C.F. Fox, Eds., Vol. IX pp. 39-42, Academic Press, New York.
12. Ness, I.F. and Nissen-Meyer, J. (1978) *Biochim. Biophys. Acta* 520, 111-121.
13. Bacchetti, S. and Benne, R. (1975) *Biochim. Biophys. Acta* 390, 285-297.
14. Gates, F.N. and Linn, S. (1977) *J. Biol. Chem.* 252, 1647-1653.
15. Van Lancker, J.L. and Tomura, T. (1974) *Biochim. Biophys. Acta* 353, 99-114.

16. Teebor, G.W., Duker, N.J. and Becker, F.F. (1977) *Biochim. Biophys. Acta* 477, 125-131.
17. Furbish, F.S., Blair, H.E., Shiloach, J., Penchev, P.G. and Brady, R.O. (1977) *Proc. Natl. Acad. Sci. U.S.* 74, 3560-3563.
18. Richardson, C.C. (1966) *J. Mol. Biol.* 15, 49-61.
19. Friedberg, E.C., Ganesan, A.K. and Minton, K. (1975) *J. Viro.* 16, 315-321.
20. Schachman, H.K., Adler, J., Radding, C.M., Lehman, I.R. and Kornberg, A. (1960) *J. Biol. Chem.* 235, 3242-3249.
21. Charlesby, A. (1954) *Proc. Roy. Soc. A* 224, 120-128.
22. Schuster, H. (1960) *Z. Naturforsch.* 156, 304-309.
23. Lindahl, T. and Nyberg, B. (1972) *Biochemistry* 11, 3610-3618.
24. Song, P.S. and Tapley, K.J. (1979) *Photochem. Photobiol.* 29, 1177-1179.
25. Hayatsu, H. (1976) in *Progr. Nuc. Acid Res. Mol. Biol.*, W.E. Cohn, Ed., Vol. 16, pp. 75-124, Academic Press, New York.
26. Lindahl, T. (1979) in *Progr. Nuc. Acid Res. Mol. Biol.*, W.E. Cohn, Ed., Vol. 22, pp. 135-192, Academic Press, New York.
27. Sancar, A. and Rupert, C.S. (1978) *Nature* 272, 471-472.
28. Cerutti, P. (1975) in *Molecular Mechanisms for Repair of DNA*, Hanawalt, P.C. and Setlow, R.B., Eds., pp. 3-12. Plenum Press, New York.