POSTER PRESENTATION

Open Access

A model of healthy aging and motor inhibition in the basal ganglia

Kyle Lyman^{1*}, Joaquin Anguera², David Terman¹

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

The basal ganglia have received a great deal of attention from modeling studies because of their putative role in normal motor function including action selection and sequencing, as well as their disturbances in Parkinson's disease. However, their interaction is complex, and is thought to be more sophisticated than can be accounted for by simple box and plot diagrams.

In the present study, we have examined trial by trial variations in response times in the stop signal task for both young (YA) and older (OA) adults. We compare previously reported YA data [1] with data collected from 18 older adults performing the same task. Unlike YA, the OA do not show a specific pattern of response times that is modulated by stop signal success in the previous trial. Additionally, the OA did not show an effect of inter trial interval (ITI) as was observed in YA. Based on our behavioral data [1], we concluded that a task switching interpretation was the most likely explanation of the YA results.

We also compare the electrophysiological data collected during the ITI from OA and YA performing the stop signal task. In YA, the power spectral data revealed a midline frontal theta effect with more desynchronization following a stop signal trial. As in [1], we concluded that this evinced a task set switching interpretation as YA subjects went from a 'ready to inhibit' mode to a 'ready to respond' mode. In the OA data presented here, we failed to observe a similar frontal theta effect, which was expected given the lack of a behavioral result in the OA subjects.

To test hypotheses about the role of the basal ganglia in task switching in young and older adults, we have developed a conductance-based network model of the subthalamic nucleus, internal and external segments of

the globus pallidus. The model incorporates cellular properties described in previous modeling studies [3] based on experimental findings. First we establish network properties which lead to patterns of beta activity that are consistent with observed phenomenon in human subjects [2]. Next, we determine which network parameters affect trial to trial variations in response times within the model. Using this approach, we examine how known changes in healthy older adult basal ganglia may affect task performance. Similarly, we discuss the implications for Parkinson's disease.

Author details

¹Math Department, Ohio State University, Columbus, OH 43201, USA.

²Department of Neurology and Physiology, University of California San Francisco, San Francisco, CA, USA.

Published: 18 July 2011

References

1. Anguera JA, Lyman K, Gazzaley A: **Neural correlates of response inhibition after-effects.** *Human Brain Mapping Conference* Barcelona Spain; 2010.
2. Swann N, Tandon N, Canolty R, Ellmore TM, McEvoy LK, Dreyer S, DiSano M, Aron AR: **Intracranial EEG reveals a time- and frequency-specific role for the right inferior frontal gyrus and primary motor cortex in stopping initiated responses.** *J Neurosci* 2009, **29**(40):12675-12685.
3. Terman D, Rubin JE, Yew AC, Wilson CJ: **Activity patterns in a model for the subthalamopallidal network of the basal ganglia.** *J Neurosci* 2002, **22**(7):2963-2976.

doi:10.1186/1471-2202-12-S1-P331

Cite this article as: Lyman et al.: A model of healthy aging and motor inhibition in the basal ganglia. *BMC Neuroscience* 2011 **12**(Suppl 1):P331.

* Correspondence: KyleLyman@gmail.com

¹Math Department, Ohio State University, Columbus, OH 43201, USA

Full list of author information is available at the end of the article