Nuclease digestion promotes structural rearrangements in H1-depleted chromatin

Wolfgang O. Weischet⁺ and K.E. Van Holde

Physiologisch-Chemisches Institut, Universitäts-Krankenhaus Eppendorf, 2000 Hamburg 20, GFR,
and Department of Biochemistry and Biophysics, Oregon State University, Corvallis, OR 97331, USA

Received 4 July 1980

ABSTRACT

Digestion of H1-depleted chromatin with micrococcal nuclease at an ionic strength of 0.35M gives rise to structural rearrangements indicating nucleosomal sliding. The ionic strength necessary to reveal this effect is significantly lower than that required in the absence of an accompanying digestion. As an explanation, a model is presented in which the progressing terminal degradation of oligomeric nucleosomes is made responsible for promoting structural rearrangements.

INTRODUCTION

In a recent paper Tatchell and Van Holde (1) published a minimal model of the protein-DNA interactions within chromatin core particles. It abstains from any detailed proposals of particular histone-DNA contacts (cf. e.g. Mirzabekov et al. (2)), designating only specific regions of strong and weak interactions between the core histones and the core DNA. Despite the generality of the concept the model has been quite successful in explaining nucleosome reconstitution and thermal denaturation experiments (1). In the present paper we shall apply these ideas to the interpretation of digestion patterns which result from cleavage of H1-depleted chromatin by staphylococcal nuclease. In addition, this work is related to recent reports of the salt-effected rearrangements of the nucleosomal structure following removal of histone H1, in which unusually tight clusters of core particles have been reported (3,4,5). Although using various techniques, the common result of these accounts indicates the ability of histone core complexes to assume new positions along the DNA under certain conditions. It has been shown that such rearrangement must be distinguished from dissociation and re-

association of the histones (6) thus giving rise to the notion of "nucleosomal sliding" or "rolling" (4,5,7). The stabilizing forces within such core clusters show some properties of hydrophobic interactions (5). In a previous paper we have compiled experimental evidence that the so-called compact oligomers (8) constitute at least part of these nucleosomal clusters (4). These particles exhibit a very regular folding of the DNA around a number of stacked histone cores and an exceptional resistance against cleavage by micrococcal nuclease (8). They are, in fact, artifacts formed in H1-depleted chromatin in the presence of ionic strengths around 0.5M - 0.65M. Their DNA content is smaller than that of an equal number of core particles without intervening spacer DNA; the repeat length can be represented by $(n \times 120 + 20 \text{ or } 30) \text{ bp}$ (8), where n is the number of cores. These facts make them an easily distinguishable probe for nucleosomal rearrangements and candidates for the chromatin fragments remaining after prolonged nuclease digestion.

MATERIALS AND METHODS

Intact calf thymus nuclei were completely and exclusively depleted of histone H1 by the acid depletion method (9) as described earlier (10). H1-depleted chromatin was prepared by dropwise adding suspensions of such nuclei into 5 mM EDTA, pH 7.6 at 37°C and subsequent chilling on ice. After about 90 min in EDTA the released chromatin was transferred into 15 mM Tris HCl, pH 7.6, 0.7 mM EDTA by repeated withdrawal and addition of solvent. Centrifugation was omitted to allow maximum swelling of the chromatin gel. After 18 h at 4°C the gel was cut into pieces which were transferred into, and equilibrated with, buffers of various ionic strengths for 4 h at 4°C.

Chromatin digestions were performed at concentrations of about 2.5 mg/ml using micrococcal nuclease (E.C. 3.1.4.7, Worthington) at concentrations ranging from 70 to 300 units/ml. Standard digestion buffer was 15 mM Tris HCl, pH 7.6, 0.5 mM CaCl_2 supplemented with NaCl as indicated. DNA was extracted from samples digested for various times in the manner described by Britten et al. (11) and used in gel electrophoresis performed according to Loening (12). Acid solubility: Defined as acid sol-

uble are the DNA fragments soluble in 0.3M perchloric acid, 0.3M NaCl (13). The concentrations were determined spectrophotometrically.

RESULTS

When H1-depleted chromatin is digested with micrococcal nuclease in the presence of 350 mM NaCl a number of oligomeric fragments remain intact even after prolonged digestion as indicated by the various DNA bands in Figure 1A. The extensive lack of DNA smaller than 120 bp in this picture should not disguise the fact that the digestion at the latest time point analyzed has almost reached its limit (14) as indicated by a relative

Fig.1: A. Time course of a digestion of H1-depleted chromatin in standard medium supplemented with 350 mM NaCl by 200 units/ml micrococcal nuclease at 37°C. The tracks correspond to 10, 20, 30, 45, 60, 90, 120, 180, 240 min of digestion. Arrows point to the bands of the compact dimer, trimer, and tetramer. Size markers are PM2/Hae III fragments (cf. ref. 8). B. Comparative digestion of identical material in absence of salt with 170 u/ml micrococcal nuclease. Samples were taken after 15, 30, 60, 120, 240 min. Sizes are given in base pairs.

amount of acid soluble material of 42% after 4 h of digestion. We have previously reported such an effect of moderate salt concentration on digestions with micrococcal nuclease and suggested that increased ionic strengths reduce the ratio of tail-clipping over endonucleolysis (10) as does a decrease in temperature (15).

The resistant DNA bands after exhaustive digestion for 4 h are centered around 145 bp, ~270 bp, ~390 bp, and ~510 bp. The combination of nuclease resistance and the peculiar band sizes following the rule $(120n + 30)\text{bp}$ is of considerable diagnostic value as has been carried out previously in detail while studying nucleosomal rearrangements (4). In this paper it has been shown that such a series of DNA bands is generated from a similar series of oligomeric nucleosomes. They have been identified mainly as compact oligomers (8) according to their DNA size, sedimentation coefficient and DNA fragment pattern after re-digestion with DNase I. Furthermore, it has been shown that compact oligomers are formed newly in H1-depleted chromatin during the exposure to high salt concentrations. Basing on these results, in particular on the simple probe for compact oligomers, we will focus in the following on the mode of their formation.

A necessary control concerning the de novo appearance of compact oligomers is shown in Figure 1B. There the resistant DNA is analyzed after digestion of H1-depleted chromatin in the absence of salt. As mentioned previously, and in accord with the literature (5), no DNA bands larger than 145 bp remain after rigorous digestion indicating the absence of compact oligomers under these circumstances.

The formation of the compact oligomers as shown in Figure 1A and that reported previously occur under different conditions, however. In the latter case an exposure of the H1-depleted chromatin to ionic strengths equal to or higher than 500 mM for 20 h at 4°C was required to generate traces of compact oligomers. With longer incubations their amount increased. In contrast, the experiment reported in Figure 1A shows unambiguously that in presence of the nuclease and at 37°C an ionic strength of 350 mM suffices to cause a similar effect. So, it is either the elevated temperature or the presence of the nuclease which replaces, in part, the salt.

In order to decide between these two alternatives we performed the following experiment. A batch of H1-depleted chromatin was cut into ten equal parts which then were pairwise equilibrated with 15 mM Tris HCl, pH 7.6, containing 50, 100, 200, 300, and 350 mM NaCl. One sample of each pair was digested with micrococcal nuclease for up to 4 h at 37°C in the presence of salt while the other one was merely incubated at this temperature for the same duration. The non-digested samples contained 0.7 mM EDTA instead of Ca^{++} . After 4 h they were cooled on ice and transferred into the standard digestion medium without added salt. Equilibration was allowed for 20 h at 4°C. This procedure freezes any formation of compact oligomers but does not reverse it, as shown elsewhere (4). Subsequently these samples were digested in the absence of salt at 37°C. Time courses of these digestions are shown in Figure 2 in parallel to time courses of the digestions in the presence of salt. The results are the following: The pre-incubation at various ionic strengths at 37°C has no effect on the band pattern produced by the digestion under standard conditions. The time courses look identical whether the chromatin has been exposed to 350 mM NaCl or 100 mM NaCl (Figure 2A,C,E,G) or 50 mM or 0 NaCl (data not shown). In particular, the dimer band vanishes when trimmed to between 300 bp and 280 bp and does not persist appreciably longer than 30 min under the conditions chosen. There are no traces of higher oligomer bands and the DNA left after 2 h of digestion is all equal to or shorter than core size (145 bp). Thus, incubating H1-depleted chromatin for 4 h at up to 350 mM NaCl does not form any compact oligomers even at 37°C. Increased temperature alone is not capable of partly replacing the salt necessary for the formation of nuclease resistant chromatin fragments larger than core particles.

This, by exclusion, leaves the presence of the micrococcal nuclease as cause for augmenting the salt effect in the formation of the compact oligomers. The question remaining is how strong that augmentation is; in other words, what is the minimum salt concentration required for the structural rearrangement? A partial answer is also found by inspecting Figure 2. From all the tracks resulting from digestions in presence of salt only

Fig.2: Comparison of the DNA band patterns after digestion in the presence of salt (B,D,F,H) and in absence of salt but after a 4 h incubation at the corresponding NaCl concentration (A,C,E,G) (cf. text). A,B: 100 mM NaCl; C,D: 200 mM; E,F: 300 mM; G,H: 350 mM. The digestions were monitored between 15 min and 4 h. Enzyme concentrations: standard medium: 140 u/ml micrococcal nuclease; B: 70 u/ml; D: 140 u/ml; F and H: 290 u/ml.

those corresponding to 300 and 350 mM NaCl show traces of resistant oligomers. Obviously, 300 mM NaCl is close to the lower limit of ionic strength allowing the rearrangements under the conditions chosen. However, the experiment does not exclude that the amount of nuclease present, i.e. the rate of digestion, may have an effect on the assessment of the limiting salt concentration.

A close inspection of the time course in Figure 1A reveals some additional facts worth mentioning with respect to a mechanistic understanding of the formation of the compact oligomers. The repeat length of H1-depleted calf thymus chromatin is 192 ± 4 bp, like that for native chromatin (10). Thus, the approximate total decrease in DNA size during digestion as apparent from Figure 1A is 110 bp for the dimer, 190 bp for the trimer, and 260 bp for the tetramer. Neglecting any fine structure the gross shift in size of the dimer band (where it can be followed best) appears as a gradual one not passing through

stages where there is more than one resolvable band. Obviously, this fact seems better compatible with the idea of a sequential processing of a particular dimeric nucleosome than with the consecutive appearance and disappearance of various dimer species differing in their resistance against micrococcal nuclease.

DISCUSSION

That increased salt concentrations alone can cause rearrangements of the structure of H1-depleted or deficient chromatin seems a settled fact (3-7). While the lower concentration limit for this effect is not unambiguously established at the present time, two approaches for titrating this limit using rather simple assays yield the same value: 500 mM (4,5). They both use pre-incubations of chromatin samples in mono-valent salt solutions, one at close to 0°C (4) the other at 37°C (5), and a subsequent digestion with micrococcal nuclease as a probe for rearrangements having occurred at high ionic strength. The results shown in Figure 2 are fully compatible with the latter report in that they demonstrate that 350 mM at 37°C is not sufficient to cause rearrangements.

The main result of the experiments described here is the conclusion that the action of the micrococcal nuclease on H1-depleted chromatin can promote structural rearrangements ("sliding") to take place which would only have occurred at considerably higher ionic strengths in the absence of nucleolytic degradation. The parameter used here for measuring rearrangements is the appearance of compact oligomers, particles which are conspicuous for the short lengths of DNA they contain and their great stability against further cleavage by micrococcal nuclease (8). Thus, the purpose of this discussion will be to present a model explaining the formation of these resistant particles through the action of micrococcal nuclease in the presence of a moderate salt concentration. The elements which enter into this hypothesis are the following:

(a) The distribution of weak and strong binding of the histones along the core DNA - as found in the reconstitution experiments of Tatchell and Van Holde (1). It extends over a total of about 166 bp, the maximum length of DNA found to reconstitute

in a precisely oriented manner onto the histone octamer. There is an extended region of strong contacts between ± 53 bp. (The origin of the reference system is located on the DNA at the point on the putative dyad axis of the core particle; cf. ref. 16. This corresponds approximately to 73 bp from each DNA end of the classical core particle.) Further contacts exist pair-wise as points of strong binding located at ± 72 bp and some of considerably weaker binding at ± 83 bp. These locations, at the same time, correspond formally to halts in the trimming of H1-deficient chromatin fragments by micrococcal nuclease (10).

(b) The interpretation will remain within the frame of the full core hypothesis (10): The inner histones of a core particle still arranged within high molecular weight chromatin occupy about 166 bp. Thus, a dimer not containing readily accessible spacer DNA comprises about 332 bp. This idea, resulting from extended studies of H1-depleted nuclei and chromatin from calf thymus, is comparable with the statements of Spadafora et al. (5) who locate the "normal" dimer band for rat liver chromatin between 380 and 320 bp. Their criterion for structural rearrangements is the appearance of bands roughly following a 140 bp repeat pattern.

(c) The histone cores have a tendency to interact with one another and assume a tightly packed arrangement. This binding shows properties of hydrophobic interactions (5). However, in native and H1-depleted chromatin, at low salt, neighboring histone cores do not interact in this way; they are in a metastable state. Contrarily, members of the family of neighboring histone cores which exert the hydrophobic interactions are the stacked compact oligomers. They have all the histone-DNA binding contacts an equivalent number of 145 bp core particles have and the DNA at the interfaces is shared between histone octamers (8). In gel electrophoresis under denaturing conditions, the major compact dimer product comigrates with the 26th band of a DNase I digest of compact oligomers. Using an average spacing of the bands of 10.4 b (17) we get 270.4 b for its DNA size. Acting on ^{32}P -end labeled compact dimers, DNase I produces the same band pattern in the appropriate region as with core particles. These considerations lead to a schematic representation of the histone

binding contacts in compact dimers as shown in Figure 3 (lower right). Mirzabekov et al. (2) found that in core particles only the 5' strands of the 20 terminal base pairs are associated with histones. Adopting this finding for the compact dimers, one would expect that in the central 20 bp region the two strands of the DNA each interact with histones from one octamer. This yields complete coverage and gives, thereby, some reason for the notable resistance of this region against micrococcal nuclease.

(d) The energy-barrier for the transition between the normal and the stacked configurations of neighboring cores is salt-dependent. Possible forces preventing the stacking are repulsion of histone octamers through superficial charges and the tightness of the histone-DNA interactions which virtually fixes the

Fig.3: Pathways in the processing of a dimeric nucleosome without spacer DNA (upper left) to a compact dimer (lower right) in the presence of high ionic strength. Shown is the DNA in a linearized form with regions of extended interaction (hatched area) and of local strong (bar and triangle) and weak (bar) interactions between the DNA and the core histones according to Tatchell and Van Holde (1). The superscribed numbers give approximate distances in base pairs.

histones on the DNA under low salt conditions. Both forces are diminished by increasing salt concentrations.

Using these elements, what predictions can be made about events during the digestion of H1-depleted chromatin with micrococcal nuclease? Due to the preference of this enzyme for spacer DNA the digest after some initial phase will consist mainly of monomers and those oligomers which contain no intervening spacer DNA (15). As an example for an oligomer we shall consider a dimer. Such a particle is represented diagrammatically in Figure 3; it has a length of 332 bp and all DNA-histone contacts still intact. As pointed out by Noll and Kornberg (15) these structures are mainly further degraded by the exonucleolytic action of micrococcal nuclease. Upon degradation at the ends by not more than 10 bp each only two weak contacts are lost while the length is reduced to 312 bp. This end trimming corresponds to the one which in the case of the monomer yields the very stable "core particle" of 145 bp. A further size reduction by 20 bp corresponds formally to a transition between a core particle (145 bp) and the next smaller species with 125 bp. The latter tend to precipitate (15). By analogy, the model assumes these dimers also to be removed from the equilibrium in solution at low ionic strength. Therefore, under normally prevailing conditions of digestion one would expect the dimer band to vanish after degradation to sizes around 290 bp, which is as experimentally found (cf. Figure 1B). However, in the presence of salt, where structural rearrangements are made possible, these 290 bp particles can attain another energy minimum by sliding of the histone cores with respect to the DNA. Formally the isomerization corresponds to the gain of one strong protein-DNA interaction in addition to the hydrophobic protein-protein interactions. Thus, the pressure towards isomerization should be considerable. Furthermore, the energy barrier for this process should be relatively low due to the absence of one strong DNA binding interaction in the starting material. Thus the model explains why this rearrangement occurs at lower salt concentrations than are necessary for sliding in the absence of the nuclease. The product, besides having gained thermodynamic stability, has assumed the typical histone-DNA interaction of

the compact particles and will therefore be very resistant against further nuclease action, except for the DNA ends. Trimming will bring the DNA to a final and stable size of 270 bp. To complete the scheme in Figure 3 it should be pointed out that the same end point can be reached by an isomerization of a symmetrically trimmed normal dimer. From what has been said above the probability for existence of such structures should be very small but the trend towards isomerization should be even larger than that experienced by the less "crippled" dimers.

We emphasize that one need not think of the digestion as "driving" the isomerization at all. The conformational transitions could be completely reversible at certain ionic strengths. But in sliding of the cores into closer proximity (as with the 292 bp particle in Figure 3) weakly bound ends become exposed. In the presence of nuclease they will be digested. This locks the oligomer in the compact conformation, for to go back would require breaking a strong contact.

However, two observations speak against this interpretation. First, preincubation of chromatin at high ionic strength (in absence of nuclease) reveals an energy sink for compact configurations (4,5), presumably due to core-core interactions. It is hard to see why such interactions should be completely unfavorable at somewhat lower ionic strengths (350 mM vs 500 mM). Second, as argued above, the nuclease may lower the energy barrier for the reaction towards the compact structure but it does not for the reverse direction. In this interpretation it does "drive" the isomerization.

Overall, the model gives a rationale for the appearance of a final and rather stable DNA band around 270 bp as the ultimate product of dimer degradation in the presence of salt. The same model mechanism can be applied to any oligomer provided it has a finite frequency of occurring without intervening spacer DNA in the H1-depleted chromatin. Therefore, the band intensities in Figure 1A for compact dimer, trimer, and tetramers reflect the decreasing probability of finding two, three, and four full core particles sufficiently close to one another in native chromatin (that is, without intervening spacer DNA) (10).

The experiment summarized in Figure 2 confines the onset of

the nuclease promoted structural rearrangement to about 300 mM NaCl, i.e. to concentrations higher than those existing in nuclei in vivo. In addition, we have found this phenomenon with H1-depleted nuclei under the same conditions only in rare cases when there was concomitant evidence for nuclear lysis (data not shown). It seems, therefore, not feasible to apply such mechanisms for promoted sliding to H1-deficient regions of chromatin in vivo.

ACKNOWLEDGEMENTS

This work was supported by the Deutsche Forschungsgemeinschaft through a grant to Dr. W.H. Strätling. KVH acknowledges a Research Professorship by the American Cancer Society and support from the Public Health Service (GM-22916).

⁺Reprint requests should be addressed to:

Institut für Physiologische Chemie, Physikalische Biochemie und Zellbiologie der Universität München, Goethestr. 33,
8000 München 2, Federal Republic of Germany

REFERENCES

- 1 Tatchell, K. and Van Holde, K.E. (1979) *Biochemistry* 18, 2871-2880
- 2 Mirzabekov, A.D., Schick, V.V., Belyavsky, A.V. and Bavykin, S.G. (1978) *Proc. Natl. Acad. Sci. USA* 75, 4184-4188
- 3 Steinmetz, M., Streeck, R.E. and Zachau, H.G. (1978) *Eur. J. Biochem.* 83, 615-628
- 4 Weischet, W.O. (1979) *Nucleic Acids Res.* 7, 291-304
- 5 Spadafora, C., Oudet, P. and Chambon, P. (1979) *Eur. J. Biochem.* 100, 225-235
- 6 Germond, J.E., Bellard, M., Oudet, P. and Chambon, P. (1976) *Nucleic Acids Res.* 3, 3173-3192
- 7 Beard, P. (1978) *Cell* 15, 955-967
- 8 Tatchell, K. and Van Holde, K.E. (1978) *Proc. Natl. Acad. Sci. USA* 75, 3583-3587
- 9 Lawson, G.M. and Cole, R.D. (1979) *Biochemistry* 18, 2160-2166
- 10 Weischet, W.O., Allen, J.R., Riedel, G. and Van Holde, K.E. (1979) *Nucleic Acids Res.* 6, 1843-1862
- 11 Britten, R.J., Graham, D.E. and Neufeld, B.R. (1974) *Methods Enzymol.* 29, 363-418
- 12 Loening, U.E. (1967) *Biochem. J.* 102, 251-257
- 13 Lohr, D., Kovacic, R.T. and Van Holde, K.E. (1977) *Biochemistry* 16, 463-471
- 14 Clark, R.J. and Felsenfeld, G. (1971) *Nature (London)* New Biol. 229, 101-106

- 15 Noll, M. and Kornberg, R.D. (1977) J. Mol. Biol. 109, 393-404
- 16 Finch, J.T. and Klug, A. (1978) Cold Spring Harbor Symp. Quant. Biol. 42, 1-9
- 17 Lutter, L.C. (1979) Nucleic Acids Res. 6, 41-56