Contributors: Both authors participated in the design and analysis and in writing the paper. They have both seen and approved the final version. JOB will act as guarantor for the paper.

Funding: No external funding.

Competing interests: None declared.

- 1 Law J, Boyle J, Harris F, Harkness A, Nye C. Screening for speech and language delay: a systematic review of the literature. *Health Technol Assess* 1998;2(9).
- 2 Toppelberg CO, Shapiro T. Language disorders: a 10-year research update review. *J Am Acad Child Adolesc Psychiatry* 2000;39:143-52.
- 3 Beitchman JH. Summary of the practice parameters for the assessment and treatment of children and adolescents with language and learning disorders. *J Am Acad Child Adolesc Psychiatry* 1998;37:1117-9.
- 4 Fenson L, Dale PS, Reznick JS, Bates E, Thal DJ, Pethick S. Variability in early communicative development. *Monogr Soc Res Child Dev* 1994;59: 1-173.
- 5 Crosley CJ. Speech and language disorders. In: Swaiman KF, Ashwai S, eds. *Pediatric neurology, principles and practice*. Minneapolis: Mosby, 1999:568-75.

- 6 Chevre-Muller C, Rigoard M-Th. Child speech and language assessment and measurement. *Acta Otorhinolaryngol Belg* 2000;54:419-25.
- 7 Wetherby AM, Allen L, Cleary J, Kublin K, Goldstein H. Validity and reliability of the communication and symbolic behavior scales developmental profile with very young children. *J Speech Lang Hear Res* 2002;45:1202-18.
- 8 Coplan J, Gleason JR. Quantifying language development from birth to three years using the early language milestone development scale. *Pediatrics* 1990;86:963-71.
- 9 Cronin VS. The syntagmatic-paradigmatic shift and reading development. *J Child Lang* 2002;29:189-204.
- 10 Gillberg C. Clinics in developmental medicine. In: Aicardi J, Bax M, Gillberg C, Ogier H, eds. *Diseases of the nervous system in childhood*. Cambridge: Mac Keith Press, 1998:850-1.
- 11 Berman S. Language disorders. In: Berman S, ed. *Pediatric decision making*. St Louis: Mosby, 1996:40-2.
- 12 Cantwel DP, Baker L. *Psychiatric and developmental disorders in children with communication disorders*. Washington, DC: American Psychiatric Press, 1991.

(Accepted 8 December 2003)

Interactive case report

Treatment of nausea and vomiting during pregnancy: presentation

Nicola Harker, Alan Montgomery, Tom Fahey

Dean Lane Family Practice, Bedminster, Bristol
Nicola Harker
general practitioner

Division of Primary Health Care, University of Bristol
Alan Montgomery
lecturer in primary care research

Tayside Centre for General Practice, University of Dundee, Dundee DD2 4AD

Tom Fahey
professor of primary care medicine

Correspondence to: T Fahey t.p.fahey@dundee.ac.uk

Ms Reynolds, a 25 year old woman, primiparous with one miscarriage, presented to her general practitioner when eight weeks pregnant complaining of nausea and vomiting. These symptoms were associated with a feeling of light headedness. Ms Reynolds had no relevant medical history and had not suffered from nausea and vomiting in her two previous pregnancies. Initially she was treated with prochlorperazine, which she took for five days, but this did not relieve her symptoms.

She presented again two weeks later, still vomiting up to four times a day, with associated nausea and light headedness. Her blood pressure was 120/75 mm Hg sitting and standing, and urine analysis showed no abnormality.

Ms Reynolds was happy to be pregnant and was living with her boyfriend. Her daughter was 5 years old and fit and well. They lived in a council owned flat with

Questions

1 What would you say to the patient about the safety of treatments (both conventional and complementary) in pregnancy?

2 What study design might provide evidence for their relative efficacy in an individual patient?

3 How would you rate the relevance of such evidence to another patient?

4 How many of your patients are using complementary therapies, and what percentage of pregnant women do you think use complementary therapies such as herbal treatments or supplements during pregnancy?

Please respond through bmj.com

The safety of all non-prescription medicines is an issue for pregnant women

no specific social problems, but she commented that her symptoms were making it difficult for her to cope with looking after her daughter. Having tried prochlorperazine without success, she was reluctant to try further conventional drugs and asked about alternatives.

Competing interests: None declared.

We welcome contributions of interactive case reports. Cases should raise interesting clinical, investigative, diagnostic, and management issues but not be so rare that they appeal to only a minority of readers. Full details of criteria are available at: bmj.com/cgi/content/full/3267/7389/564/DC1

This is the first of a three part case report where we invite readers to take part in considering the diagnosis and management of a case using the rapid response feature on bmj.com Next week we will report the case progression and in five weeks' time we will report the outcome and summarise the responses