Published in final edited form as:

J Bone Miner Res. 2012 January ; 27(1): 26–37. doi:10.1002/jbmr.518.

Delayed Short Course Treatment with Teriparatide (PTH1–34) Improves Femoral Allograft Healing by Enhancing Intramembranous Bone Formation at the Graft-Host Junction

Masahiko Takahata¹, Edward M. Schwarz^{1,2,3}, Tony Chen^{1,2}, Regis J O’Keefe^{1,3}, and Hani A. Awad^{1,2,3,4}

¹University of Rochester, Department of Biomedical Engineering, Rochester, NY

²University of Rochester, The Center for Musculoskeletal Research, Rochester, NY

³University of Rochester, Department of Orthopaedics, Rochester, NY

Abstract

Clinical management of critical bone defects remains a major challenge. Despite pre-clinical work demonstrating teriparatide (PTH_{1–34}) effectiveness in small animals, inconclusive data from clinical trials have raised questions of dose and regimen. To address this, we completed a comprehensive study in the murine femoral allograft model, to assess the effects of dose (0.4, 4, and 40 µg/kg/day) and various treatment regimens on radiographic, histologic and biomechanical healing at 2, 4, and 9 weeks. Only the high dose (40 µg/kg) of PTH_{1–34} demonstrated significant effects when given daily over 9 weeks. Remarkably, equivalent biomechanical results were obtained with delayed, short treatment from 2 to 6 weeks that did not induce a significant increase in endochondral bone formation and callus volume. In contrast, PTH_{1–34} treatment from 1 to 5 weeks post-op demonstrated similar osteogenic effects as immediate-daily treatment for 9 weeks, but failed to achieve a significant increase in biomechanics at 9 weeks. Micro-CT and histologic analyses demonstrated that the 2-week delay in treatment allowed for timely completion of the endochondral phase, such that the prominent effects of PTH_{1–34} were enhanced intramembranous bone formation and remodeling at the graft-host junction. These findings support the potential use of PTH_{1–34} as an adjuvant therapy for massive allograft healing, and suggest that there may be an ideal treatment window in which a short course is administered following the endochondral phase to promote osteoblastic bone formation and remodeling to achieve superior union with modest callus formation.

Keywords

Bone; Allograft; Parathyroid Hormone (PTH); Biomechanics; Micro-Computed Tomography (micro-CT)

Introduction

Although most orthopaedic fractures heal, the clinical management of critical (>3cm) segmental defects continues to face major challenges for both amputation and limb salvage approaches, which have high rates of self-reported disability (40% to 50%) and continue to worsen over time.^(1,2) Despite lacking osteogenic and remodeling capacity, devitalized cortical allografts remain a major option for reconstructive surgery because no equivalent

⁴Corresponding Author: Hani A. Awad, PhD, Department of Biomedical Engineering, The Center for Musculoskeletal Research, University of Rochester – Box 665, 601 Elmwood Avenue, Rochester, NY 14642, (585) 273-5268, hani_awad@urmc.rochester.edu.

alternatives exist. However, the limited new bone formation and lack of remodeling associated with massive allograft healing are directly associated with the 23–43% clinical failure rate caused by non-union (27–34%), late graft fracture (24–27%), and infection (9–16%).(3) Thus, the quest for a practical adjuvant therapy for massive allograft surgery remains a high priority.

Intermittent administration of PTH_{1–34}, an active recombinant human peptide sequence of parathyroid hormone or (PTH_{1–34}), has been proven to increase skeletal bone mass in osteoporotic patients.(4) Based on these anabolic properties, several groups have demonstrated that PTH_{1–34} is effective in small animal models of fracture healing.(5–10) Moreover, clinical case reports of off label use of PTH_{1–34} to heal delayed and non-unions have suggested that the drug could be effective as an adjuvant for challenging bone healing situations. (11,12) Unfortunately, a recent phase 2 clinical trial of distal radial fractures in postmenopausal women failed to meet its primary prospective endpoint with 40 µg/day teriparatide, although the study did show that 20 µg/day teriparatide accelerated the time to radiographic healing from 9.1 to 7.4 weeks versus saline controls ($p = 0.006$). (13) The efficacy of the therapy in this study however was confounded by a fracture model that would normally heal in the saline-treated group irrespective of treatment, which raises several questions that warrant further investigation of PTH_{1–34} dose and regimen in challenging preclinical models of bone healing.

One preclinical model of challenging bone healing that has emerged as a useful tool to understand the biological issues associated with delayed healing and non-unions is the murine femoral allograft model.(14) Similar to the clinical situation, femoral allografts in mice heal only via creeping fracture callus from the host that is mediated by endochondral ossification, and there is very little remodeling of the necrotic graft due to the absence of osteoclasts on the cortical surface of the allograft. Furthermore, murine allografted femurs achieve <50% of the torsional biomechanical properties of normal, ungrafted controls,(15) which is the primary outcome measure of reconstructive surgery for segmental defects in preclinical models. Live bone graft studies have demonstrated that the lack of intramembranous bone formation is due to the absence of periosteal cells, which are responsible for the angiogenic, osteogenic and remodeling responses during allograft healing.(16,17) Although these signals can be supplied exogenously via recombinant virus mediated gene transfer or cell therapy,(16,18,19) PTH_{1–34} adjuvant therapy offers the potential for a more practical solution for this problem.

In order to better understand the potential of PTH_{1–34} as an adjuvant therapy for structural allograft healing several questions about optimal dosing regimen must be addressed. With regards to dose, there is a significant discrepancy (approximately 100 fold) between the doses used in the rodent models (typically 40 µg/kg/day),(5–10) versus the approved clinical dose of 20 µg per day total (which in a 50 to 80kg person translates to 0.25 to 0.4 µg/kg/day). This discrepancy suggests that either the higher metabolic activity in small mammals requires higher dosing (on a per weight basis), and/or that efficacious dosing for bone healing is greater than that of anabolic therapy for generalized osteopenia. Moreover, questions about the ideal duration and timing of PTH_{1–34} therapy regimen for bone healing following reconstructive surgery have not been addressed. While osteoporosis therapy is indicated up to 18-months, it is unlikely that PTH_{1–34} treatment beyond a few months could significantly influence the dynamics of fracture healing. Given the potential safety concerns associated with long-term use of the drug, in addition to its high cost and the inconvenience of daily subcutaneous injections, identifying the minimum treatment duration required for efficacy is critical for this indication of PTH_{1–34} therapy. Finally, there is a question as to the ideal timing of PTH_{1–34} therapy following surgery. Since PTH has multiple mechanisms of action during bone repair via direct and indirect effects on multiple cell types during the

initial inflammatory, intermediate osteogenic and late remodeling stages of bone healing, it is important to elucidate the effective window of therapy that achieves the maximum benefit on allograft healing. Therefore, a systematic study to investigate the appropriate clinically-translatable dosing regimen of PTH₁₋₃₄ needs to be conducted. To this end, we examined the effects of various doses, timing, and duration of PTH₁₋₃₄ therapy on radiographic, histologic and biomechanical properties of murine femoral allograft healing.

Methods & Materials

Preparation of Decellularized Allografts

Devitalized femoral allografts were prepared from femurs of 9 week old female ICR mice as previously described.⁽¹⁴⁾ The donor femurs were cut to 4 mm mid-diaphyseal allografts, bone marrow was flushed using saline from a syringe, then the allografts were bathed in 70% ethanol for 3 hours, washed in phosphate buffered saline (PBS) before storing at -70°C for at least 1 week before they were used.

Femoral Allograft Reconstruction Surgery

All animal studies were performed in accordance with protocols approved by the University of Rochester's Committee on Animal Resources. Femoral allograft surgeries were performed as previously described.⁽¹⁴⁾ Briefly, devitalized ICR allografts were implanted into a 4 mm defect created in the mid-diaphysis of the left femur of recipient 9 week old female C57Bl/6 mouse using a 10mm diameter, 0.15mm thick diamond-sintered rotary saw (Brasseler USA, Inc, Savannah, GA), and secured in place with an intramedullary stainless steel pin (Figure 1A). Daily subcutaneous injections of teriparatide (human PTH₁₋₃₄) (Forteo™, Eli Lilly and Co., Indianapolis, IN) or saline (control) was performed according to the dosing regimens described in Figure 1. Group names were defined by dose, timing of initial treatment and duration as follows. Three daily doses of PTH₁₋₃₄ were investigated (L = low dose of 0.4 μg/kg/day; M = middle dose of 4 μg/kg/day; and H = high dose of 40 μg/kg/day). The treatment regimen was defined by the timing of the initial dose (0 or one day after the surgery; and 1, 2 or 4 weeks postop), and the timing of treatment withdrawal (2, 3, 4, 5, 6, 8 or 9 weeks post-op).

Experiment 1: Dose Response Study

Daily doses of PTH₁₋₃₄ (0.4, 4, and 40 μg/kg/day) or saline were administered to the mice (n=15 per group) via subcutaneous injections from one day after the surgery for 4 weeks (Figure 1B). The mice were then maintained without therapy for another 5 weeks before sacrificed at 9 weeks post-surgery. Healing of grafted femurs was monitored by weekly X-rays (Supplemental Figure 1), and the grafted femurs were harvested and subjected to micro-computed tomography (micro-CT) analysis. Then, 10 femurs were used for torsion testing, and the remaining 5 femurs were used for histology.

Experiment 2: Duration and Timing Study

Administration of high dose (40 μg/kg/day) PTH₁₋₃₄ was commenced either 1 day after the surgery (immediately), or at 1, 2 or 4 weeks post-surgery. The dosing periods were either 2 or 4 or 9 weeks (Figure 1C). The saline treated group as a control and the H-0-4 group in experiment 1 were included in this analysis. A total of 15 mice in each group were sacrificed at 9 weeks post-surgery and subjected to micro-CT (n=15), biomechanical testing (n=10), and histology (n=5) as depicted in Figure 1C.

Experiment 3: PTH effects on gene expression during allograft healing

To elucidate the early mechanistic differences between immediate and delayed PTH₁₋₃₄ therapy on allograft healing we evaluated the healing response at 2 or 4 weeks post-surgery. Three PTH₁₋₃₄ dosing regimens starting either 1 day after the surgery (Immediately), 1 week after surgery (1w delayed), or 2 weeks after surgery (2w delayed) were compared to saline treatment control group as depicted in Figure 1D. A total of 10 mice in each group were assessed by micro-CT and histology (n=5 per group per time point). In addition, 3 mice per group were sacrificed at various times (0, 3, 7, 10, 14, 21, 28 days) post-surgery for mRNA expression analysis.

Micro Computed Tomography Imaging of Bone Formation and Union

The femurs were harvested from euthanized mice, disarticulated at the hip and knee joints, and the intramedullary pins were carefully removed. The specimens for biomechanical testing were stored at -20°C, and the specimens for histology were fixed in 10% neutral buffered formalin and stored in 70% ethanol, before micro CT imaging. The specimens were scanned at 12.5 microns isotropic resolution using the VivaCT 40 (Scanco Medical AG, Bassersdorf, Switzerland) as previously described.(11) From these 3D images, the callus bone volume was measured by manual segmentation, followed by standardized thresholding at a grayscale corresponding to 750 mgHA/cm³ based on a phantom of known HA concentrations. The Union Ratio, a measure of allograft osseointegration, which is based on the minimum graft surface area fraction upon which mineralized callus had formed, was also calculated as recently described.(11)

Biomechanical Testing and Failure Mode Analysis

Immediately following micro-CT imaging, the torsional biomechanical properties of the grafted femurs were determined as previously described,(20) using an EnduraTec TestBench™ system (200 N.mm torque cell; Bose Corporation, Minnetonka, MN) at a rate of 1°/sec. Yield torque (T_{Yield}), ultimate torque (T_{Ult}), torsional rigidity (TR) and work to failure were determined for each specimen. After torsion testing, the specimens were x-rayed to analyze the mode of failure as previously described.(20)

Histology and Histomorphometric Analyses

After micro-CT, femurs were decalcified in EDTA. At least, two nonconsecutive 3-μm paraffin embedded mid-sagittal sections were stained with either alcian blue/hematoxylin/orange G (AB/OG) or tartrate resistant acid phosphatase (TRAP). Cartilage area and osteoclast number (Oc.N) were measured in week 2 and week 4 samples in experiment 3 as described previously.(14) Briefly, cartilage area was measured by alcian blue stained area. Oc.N was determined from sections stained for TRAP by counting the number of TRAP positive cells. Oc.N inside the callus area (Oc.N_{callus}) and on the surface of allograft (Oc.N_{graft}) was evaluated separately. The mean values from the two nonconsecutive mid-sagittal sections represented the value for one mouse.

Gene expression analysis

The mRNA expression levels of various genes within the tissue at graft-host junction were measured using real time, reverse transcription-polymerase chain reaction (RT-PCR). The 3mm long callus tissue that formed at graft-host junctions was harvested at postoperative days 3, 5, 7, 10, 14, 21, and 28 as shown in Figure 1D. Tissues were flushed with DEPC treated phosphate-buffered saline to wash out bone marrow cells. The samples were frozen in liquid nitrogen and homogenized using a dismembrator. Total RNA was extracted from the samples using the TRIzol Reagent (Invitrogen Corp., Carlsbad, CA). For cDNA synthesis, 1 μg RNA was reverse-transcribed using iScript cDNA synthesis kit (Bio-Rad

Laboratories, Hercules, CA). Real-time PCR was performed using a 20- μ L reaction volume. Signals were detected using the PerfeCTa SYBR Green FastMix (Quanta BioSciences, Inc., Gaithersburg, MD) with gene-specific primers. Gene-specific primers of type II collagen (*colIIa1*), type X collagen (*colXa1*), vascular endothelial growth factor-a (*vegfa*), matrix metalloproteinase 9 (*mmp9*), type I collagen (*colIa1*), osteocalcin (*oc*), cathepsin K (*ctsk*), tartarate resistant acid phosphatase (*trap*) and β -actin as a housekeeping gene are shown in supplemental Table 1. The relative mRNA expression of each targeted gene was normalized by the cycle threshold values of β -actin.

Statistical Analysis

One-way analysis of variance (ANOVA) and Tukey's or Dunnett's post-hoc multiple comparison tests were performed to assess differences among PTH₁₋₃₄ treatment groups and compared to the control groups, respectively. P-values ≤ 0.05 were used to detect significant differences.

Results

Dose Effects of PTH₁₋₃₄ on murine femoral allograft healing

In order to establish the effective dose of the therapy for murine allograft reconstruction we injected animals with 0 (saline control), 0.4, 4, or 40 μ g/kg/day of PTH₁₋₃₄ for 4 weeks following surgery. As restoration of biomechanical integrity of the injured long bone is the primary goal of structural allografting, we assessed this outcome 9 weeks after surgery to allow for adequate healing for torsion testing. (20) High-dose PTH₁₋₃₄ treatment (H-0-4) increased the grafted femur yield torque (T_{Yield}), ultimate torque (T_{Ult}), and torsional rigidity (TR) to 2.6, 2.2, and 1.9 times those in the control group (Table 1). In contrast, middle- (M-0-4) and low-dose (L-0-4) PTH₁₋₃₄ showed equivalent values to those in the control group. Assessment of the mode of failure confirmed these results, as control and low dose groups resulted in preunion at a rate of 40% and 50%, respectively, but most importantly never achieved mature union by the end of the study. In contrast, none of the allografts in the high dose group resulted in preunion, whereas 10% of those achieved mature union (Supplemental Table 2)

Effects of treatment duration of PTH₁₋₃₄ on biomechanical parameters of femoral allograft healing

To assess the differential effects of duration of PTH₁₋₃₄ therapy on allograft healing in our model, we evaluated high dose treatment regimens commencing immediately out to 2, 4, or 9 weeks post-surgery, and evaluated the biomechanical properties at 9 weeks. The H-0-9 group served as the positive control for this study, as all tested parameters of healing were significantly ($p < 0.05$) greater than saline treated controls. Contrary to our expectation, we found that immediate treatment for only 2 weeks (H-0-2) was sufficient to significantly increase the grafted femur T_{Ult} to 2.4 times that in the control group ($p < 0.05$), and was not significantly different than the H-0-9 group (Table 1). We also observed a trend in which H-0-2 group demonstrated a 3- fold increase in T_{Yield} and 1.6-fold increase in TR of grafted femur over saline control, although these differences were not statistically significant. The 4-week (H-0-4) and 9-week (H-0-9) treatment groups were equivalent in biomechanical properties to 2 weeks-long (H-0-2) treatment, indicating that longer treatments do not further improve allograft healing in terms of bone strength. The conclusion was also supported by the failure mode analysis, which demonstrated that the additional 5 weeks of PTH₁₋₃₄ treatment in the H-0-9 group resulted in only a 20% increase in the incidence of mature union compared the H-0-4 group (Supplemental Table 2).

Effects of delayed PTH₁₋₃₄ treatment on biomechanical parameters of femoral allograft healing

To assess the efficacy of delayed PTH₁₋₃₄ treatment on allograft healing in our model, we evaluated the effects of commencing the treatment after 1, 2, or 4 weeks post-reconstruction for 2 or 4 weeks thereafter. We found the treatment 4 weeks after surgery (H-4-8) to be completely ineffective, as no differences were observed compared to saline-treated controls. Interestingly, 1 or 2 week-delayed treatments resulted in equivalent biomechanical properties to the positive control H-0-9, regardless of the duration of treatment. However, the most remarkable results from this study came from the H-2-6 treatment group, which demonstrated significant increases in all parameters tested compared to saline controls, and achieved equivalent biomechanical outcomes to the H-0-9 treatment group.

While 9 weeks post-surgery is ideal for assessing drug effects on torsional biomechanics in this model (11), this time point proved to be too challenging for quantitative radiology and histomorphometry, due to the extensive remodeling that prohibits accurate segmentation of the graft from the host bone, and remodeling of callus and new periosteal bone. Additionally, unsegmented micro-CT analyses of the grafted femurs demonstrated that no differences in bone volume or bone mineral density at 9 weeks could be observed among the groups (data not shown). However, the 3D micro-CT rendering of the grafted femurs demonstrated gaps and irregular cortical surfaces at the graft-host junction of the control, L-0-4, and M-0-4 groups (Figure 2) that were not observed in the H-0-4 group, demonstrating that only the high dose has effects on bone healing in this model.

Effects of PTH₁₋₃₄ treatments on radiographic and histologic parameters of femoral allograft healing

In order to better understand the mechanism responsible for these drug effects on biomechanical properties, we analyzed micro-CT images and histology at 2 and 4 weeks post-surgery (Figure 1D). These analyses were limited to only the different high dose treatment regimens, since the low and middle dose treatments were not biomechanically efficacious. Interestingly, while immediate PTH₁₋₃₄ treatment significantly increased callus volume at 2 weeks (Figure 3C and H) compared to controls (Figure 3A and H), this increase was not observed in the 1 week delayed treatment group (Figure 3E and H), and was not significant at 4 weeks due to remodeling. No treatment related effects on the callus BMD were detected at either 2 or 4 weeks post surgery (Figure 3I). However, both the immediate and 1 week delayed treatment groups displayed significant increases in their Union Ratios by 4 weeks (Figure 3J), which demonstrates that this parameter of osseointegration is not related to callus volume, as we have previously reported.(11,15) The results of this experiment also demonstrated that while a 2-week delay in the PTH₁₋₃₄ treatment did not increase the Union Ratio per se compared to saline controls (Figure 3J), micro CT data at 9 weeks showed remarkable end to end integration in the treated specimens, regardless of the delay, which underscores that the long term healing is independent of callus size.

Histological analyses of the healing allografts harvested 2 and 4 weeks post-surgery also provided evidence for PTH₁₋₃₄-induced osseointegration independent of callus size (Figure 4). At 2 weeks, there was a small, albeit statistically insignificant, increase in the cartilage area with the immediate PTH₁₋₃₄ treatment compared to the saline controls and 1 week delayed treatment (Figure 4A–C; Table 2). Moreover, there was no significant evidence of persistent cartilage at 4 weeks (Figure 4D–G; Table 2), regardless of the treatment. Furthermore, there were mild increases in the osteoclast number density at the surface of the treated allografts at 2 weeks, which increased by 4 weeks in all PTH₁₋₃₄ treatment groups (Figure 4, Table 2). Furthermore, the osteoclast number density in the callus was significantly higher than that on the allograft surface at 2 weeks, regardless of the treatment.

At 4 weeks, the osteoclast number density remained increased in the treated groups, but decreased significantly in the saline treated controls (Table 2).

Effects of PTH₁₋₃₄ treatment regimen on gene expression during femoral allograft healing

To further elucidate the mechanism responsible for delayed PTH₁₋₃₄ effects on allograft healing, we performed a time course experiment to assess changes in mRNA expression of markers of chondrocytes, angiogenesis, osteoblasts and osteoclasts over 28 days of healing (Figure 5). The results of these experiments corroborated several of the histologic findings. First, only the immediate PTH₁₋₃₄ treatment had significant effects on chondrogenic (*colla1*) and hypertrophic (*colXa1*) gene expression that disappeared by 10 to 14 days, which is consistent with the small increases in cartilage area at 2 weeks and the absence of persistent cartilage at 4 weeks. Immediate PTH₁₋₃₄ treatment also increased the expression of *mmp9* and *vegf-a*, which are associated with cartilage removal and angiogenesis during fracture healing, at 14 days post surgery (17,21). Most importantly, we found that all regimens of high dose PTH₁₋₃₄ significantly ($p<0.05$) increased markers of bone formation (*colla1*, and *oc*) on days 21 and 28 post-surgery, which is consistent with the sustained bone formation at 4 weeks (Supplemental Figures 2 and 3).

Discussion

Despite its clinical effectiveness in the treatment of osteoporosis (22,23), reports of clinical off-label use to resolve fibrotic nonunion fractures (24), and a growing body of preclinical data from various animal studies that raised expectations of its efficacy in accelerating bone repair (5–7,9,10,25–28), parathyroid hormone (PTH) therapy has yet to be widely clinically adopted for the management of nonoperative fractures or the more challenging cases of fibrotic nonunions. There are several reasons that contribute to the slow translation of PTH to the clinical or surgical management these indications. First, there have been safety concerns regarding the range of effective PTH₁₋₃₄ doses in the animal studies of fracture repair (10–200 µg/kg/day), which correspond to tens or hundreds of multiples of the FDA approved doses of 20 or 40 µg/day (0.25 to 0.5 µg/kg/day for an 80 kg person) for treatment of osteoporosis. The high doses, in combination with long treatments, have been linked to high incidence of osteosarcomas in rats (23,29), which despite evidence of safety from surveillance studies of patients previously treated with PTH₁₋₃₄, led the FDA to approve the drug with strong recommendations to limit its prescription to women and men with substantial osteoporosis and susceptibility to fracture, and placed a black box exclusion warning for patients at high risk for osteosarcoma (23). This regulatory barrier is compounded by the absence of long-term patent protection, and hence the lack of commercial incentive to perform expensive safety and efficacy clinical trials for challenging fractures (23). Moreover, there is no consensus on the outcome variables that could be used to definitively establish efficacy in human patients (30). A case-in-point is a recent study that indicated that the clinical value of PTH (teriparatide) treatment in distal radial fractures (assessed by the time to healing based on radiographic evidence of cortical union, patient-related wrist evaluation score, and early callus formation) was limited to marginal acceleration of healing that did not provide strong cost-risk-benefit justification (13,30). The inconclusive findings of this human trial in contrast to the efficacy evidence in animal studies underscore its limitations: i) the lack of validated thresholds of bone healing based on non- or minimally-invasive assessment, and ii) the fact that these fractures still healed with placebo in just weeks, which makes establishing the case for treatment efficacy difficult. Finally, there remain many unanswered questions about the appropriate timing of commencement and withdrawal of the treatment, which have not been thoroughly investigated in preclinical studies. Hence, the motivation for the current study was to address the latter questions of dose, timing, and duration of treatment, in hopes of identifying in a

preclinical model a short treatment window that could help abate the more serious concerns about osteosarcoma risk in non-life threatening indications such as fracture repair, which could then incentivize definitive large animal studies and clinical trials of safety and efficacy.

Since allografts are typically complicated by nonunion and therefore are more challenging than osteotomy or closed fracture models, we elected to investigate PTH₁₋₃₄ treatment regimens in a previously established mouse femoral reconstruction model (11,14,20). We have recently shown in this model that daily PTH₁₋₃₄ (40 µg/kg/day) commencing immediately after devitalized allograft reconstruction for 6 weeks induced trabeculated bone callus formation and remarkable graft-host integration, which were correlated with more than doubling of the biomechanical parameters of torsional rigidity and yield torque compared to saline treated controls (15). In the current study, we first asked whether lower doses of PTH₁₋₃₄ could be as effective as higher dose in enhancing allograft osseointegration and biomechanics. Consistent with several previous studies in rodent models of fracture repair (5–7,9,10,25–28), our findings demonstrated that only the high daily dose (40 µg/kg/day) showed remarkable effects on torsional properties of grafted femurs when administered for 9 weeks, while the lower doses of 0.4 and 4 µg/kg/day did not provide any benefits compared to controls. Rodent models typically report the use of doses as high as 10 to 200 µg/kg/day. While the difference between the effective clinical dose for osteoporosis and the much higher doses for rodent models of fracture repair are thought to arise from differences in metabolism and clearance of PTH, and typically require treatment periods amounting to >50% of the animal life span (31), conclusive statements about the effectiveness of lower dose might still require investigation in larger animal species that might better approximate the metabolism of the drug in humans, and a clinically relevant treatment window.

Our studies showed that immediate PTH treatment substantially improved early cartilaginous template formation in the fracture callus and subsequent mineralization, leading to improved healing at the host-allograft junction. This finding is supported by several previous reports that describe the main effects of intermittent PTH treatment on fracture repair to be through enhancement of the early proliferative response of chondroprogenitor and osteoprogenitor cells (9,10,31).

We also observed that short-term treatments (2 weeks), commencing immediately or with a 1- or 2-week delay following reconstruction, were just as effective in enhancing the biomechanical properties at 9 weeks as the continuous daily treatment for 9 weeks. This data suggests that short-term treatment during the first 4 weeks after surgery is sufficient to improve graft-host union from fibrous union to osseous union and that establishing osseous union is a primary mechanism of PTH₁₋₃₄ induced increase in mechanical strength of grafted femur. These observations have great clinical relevance. The effectiveness of the short treatments could abate concerns related to osteosarcoma risk with prolonged treatment. Furthermore, given that PTH therapy is delivered via inconvenient subcutaneous injections and is very expensive, short treatment could enhance patient compliance and balance cost-benefit concerns. Moreover, since a likely clinical scenario that could benefit from PTH therapy might involve patients in whom bone repair may be delayed (31), the effectiveness of the delayed treatment in our murine model should encourage further investigation in larger species.

The observation that 1- or 2-week delayed short-term treatment was as effective as immediate treatment suggests that PTH₁₋₃₄ might have different actions depending on the overlap with the different stages of healing and the cells involved. Our results corroborate previous findings that suggest that immediate treatment with PTH₁₋₃₄ might have multiple

effects that include suppressing inflammation upon surgery (32), early stimulation of proliferation of mesenchymal and periosteal stem cells (10), and enhanced early differentiation of chondroprogenitors leading to robust “endochondral ossification” bone repair (9,31), which is consistent with our observations of increased callus size in association with PTH₁₋₃₄ treatments. On the other hand, 1- or 2-week delayed PTH therapy in which the treatment missed part or all of the endochondral ossification phase still induced persistent osteoblastic and osteoclastic activity up to 4 weeks. This was manifested by remarkable increases in *colla1* and *oc* gene expression at day 21 which began to decline by day 28. Interestingly, the 4-week delay in the treatment could not improve the biomechanical properties of the allografted femurs compared to controls. These findings suggest that bone repair could still be accomplished, even if treatment is delayed, via intramembranous ossification, and further suggest that the first 3 weeks could be the critical window for PTH₁₋₃₄ therapy. Given that bone formation peaks at around 2 weeks after surgery and is down-regulated by 4 weeks after surgery in control mice that underwent femoral allograft surgery, PTH₁₋₃₄ therapy would be effective as long as active bone formation sustains after surgery but becomes ineffective after repair reaction is attenuated. Some clinical case reports of off-label use of PTH₁₋₃₄ demonstrated that patients with prolonged fracture non-union would benefit from anabolic therapy months to years after fracture (11,33,34), to our knowledge, however, there are no definitive data as to by when after surgery PTH₁₋₃₄ would be effective for bone repair. Therefore, it should be tested in larger animal models or clinically how long after surgery PTH₁₋₃₄ could be expected to show positive effect on bone repair. Although we failed to observe significant differences in osteoclast marker gene expression (*ctsk* and *trap*) during this critical phase of allograft healing, this was not surprising, as the significant PTH₁₋₃₄ effects on osteoclasts were on their location rather than total numbers. As the osteoclasts on the graft surface are primarily involved in generating a new marrow space between the necrotic and new bone, while the osteoclasts in the callus are remodeling the osteoid into lamellar bone, this finding suggests that PTH₁₋₃₄ effects are pertinent for callus remodeling, which was less of a factor in delayed treatment. Thus, it is unlikely that the mechanism responsible for delayed-PTH₁₋₃₄ treatment effects on structural allografting involves increased bone resorption. Future studies should address the effects of PTH₁₋₃₄ on the different cellular compartments involved in the repair and remodeling of the allograft.

Our findings provide strong evidence in support of the efficacy of delayed, short PTH₁₋₃₄ treatment in a mouse model of challenging bone repair. A recent study investigated the effects of different PTH administration regimes applied at different stages of fracture healing in an osteoporotic (ovariectomized or Ovx) rat model of tibial osteotomy healing (35), and similarly reported that the treatment improved fracture repair compared to untreated controls when PTH was administered either immediately or 7 days post osteotomy, independent of the administration frequency, but not after a 14 day delay. Collectively, our findings and others' should motivate further investigation of the efficacy of PTH treatment in challenging bone repair scenarios in larger species and clinical studies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank Dr. Christopher Beck for his help with the statistical analysis, Ryan Tierny and the histology core for their excellent technical assistance, Michael Thullen for assistance with micro CT, and Jacy Krystal Bulaon for help with the biomechanical testing. This work was funded in part by grants from the Aircast Foundation and grants from the National Institutes of Health (AR056696, AR054041, DE019902).

References

1. Busse JW, Jacobs CL, Swiontkowski MF, Bosse MJ, Bhandari M. Complex limb salvage or early amputation for severe lower-limb injury: a meta-analysis of observational studies. *J Orthop Trauma*. 2007; 21(1):70–6. [PubMed: 17211275]
2. Bosse MJ, MacKenzie EJ, Kellam JF, Burgess AR, Webb LX, Swiontkowski MF, Sanders RW, Jones AL, McAndrew MP, Patterson BM, McCarthy ML, Trivison TG, Castillo RC. An analysis of outcomes of reconstruction or amputation after leg-threatening injuries. *N Engl J Med*. 2002; 347(24):1924–31. [PubMed: 12477942]
3. Wheeler DL, Enneking WF. Allograft bone decreases in strength in vivo over time. *Clin Orthop Relat Res*. 2005; (435):36–42. [PubMed: 15930919]
4. Neer RM, Arnaud CD, Zanchetta JR, Prince R, Gaich GA, Reginster JY, Hodsman AB, Eriksen EF, Ish-Shalom S, Genant HK, Wang O, Mitlak BH. Effect of parathyroid hormone (1–34) on fractures and bone mineral density in postmenopausal women with osteoporosis. *N Engl J Med*. 2001; 344(19):1434–41. [PubMed: 11346808]
5. Komatsubara S, Mori S, Mashiba T, Nonaka K, Seki A, Akiyama T, Miyamoto K, Cao Y, Manabe T, Norimatsu H. Human parathyroid hormone (1–34) accelerates the fracture healing process of woven to lamellar bone replacement and new cortical shell formation in rat femora. *Bone*. 2005; 36(4):678–87. [PubMed: 15781006]
6. Andreassen TT, Ejersted C, Oxlund H. Intermittent parathyroid hormone (1–34) treatment increases callus formation and mechanical strength of healing rat fractures. *J Bone Miner Res*. 1999; 14(6):960–8. [PubMed: 10352105]
7. Andreassen TT, Willick GE, Morley P, Whitfield JF. Treatment with parathyroid hormone hPTH(1–34), hPTH(1–31), and monocyclic hPTH(1–31) enhances fracture strength and callus amount after withdrawal fracture strength and callus mechanical quality continue to increase. *Calcif Tissue Int*. 2004; 74(4):351–6. [PubMed: 15255072]
8. Andreassen TT, Fledelius C, Ejersted C, Oxlund H. Increases in callus formation and mechanical strength of healing fractures in old rats treated with parathyroid hormone. *Acta Orthop Scand*. 2001; 72(3):304–7. [PubMed: 11480610]
9. Nakazawa T, Nakajima A, Shiomi K, Moriya H, Einhorn TA, Yamazaki M. Effects of low-dose, intermittent treatment with recombinant human parathyroid hormone (1–34) on chondrogenesis in a model of experimental fracture healing. *Bone*. 2005; 37(5):711–9. [PubMed: 16143574]
10. Nakajima A, Shimoji N, Shiomi K, Shimizu S, Moriya H, Einhorn TA, Yamazaki M. Mechanisms for the enhancement of fracture healing in rats treated with intermittent low-dose human parathyroid hormone (1–34). *J Bone Miner Res*. 2002; 17(11):2038–47. [PubMed: 12412812]
11. Reynolds DG, Shaikh S, Papuga MO, Lerner AL, O’Keefe RJ, Schwarz EM, Awad HA. μ CT-based measurement of cortical bone graft-to-host union. *J Bone Miner Res*. 2009; 24 (5):899–907. [PubMed: 19063685]
12. Bukata SV, Puzas JE. Orthopedic uses of teriparatide. *Curr Osteoporos Rep*. 2010; 8(1):28–33. [PubMed: 20425088]
13. Aspenberg P, Genant HK, Johansson T, Nino AJ, See K, Krohn K, Garcia-Hernandez PA, Recknor CP, Einhorn TA, Dalsky GP, Mitlak BH, Fierlinger A, Lakshmanan MC. Teriparatide for acceleration of fracture repair in humans: a prospective, randomized, double-blind study of 102 postmenopausal women with distal radial fractures. *J Bone Miner Res*. 2010; 25(2):404–14. [PubMed: 19594305]
14. Tiyyapatanaputi P, Rubery PT, Carmouche J, Schwarz EM, O’Keefe RJ, Zhang X. A novel murine segmental femoral graft model. *J Orthop Res*. 2004; 22(6):1254–60. [PubMed: 15475206]
15. Reynolds DG, Takahata M, Lerner AL, O’Keefe RJ, Schwarz EM, Awad HA. Teriparatide therapy enhances devitalized femoral allograft osseointegration and biomechanics in a murine model. *Bone*. 2011; 48(3):562–70. [PubMed: 20950720]
16. Zhang X, Xie C, Lin AS, Ito H, Awad H, Lieberman JR, Rubery PT, Schwarz EM, O’Keefe RJ, Goldberg RE. Periosteal progenitor cell fate in segmental cortical bone graft transplantations: implications for functional tissue engineering. *J Bone Miner Res*. 2005; 20 (12):2124–37. [PubMed: 16294266]

17. Colnot C. Skeletal cell fate decisions within periosteum and bone marrow during bone regeneration. *J Bone Miner Res.* 2009; 24(2):274–82. [PubMed: 18847330]
18. Ito H, Koefoed M, Tiyyapatanaputi P, Gromov K, Goater JJ, Carmouche J, Zhang X, Rubery PT, Rabinowitz J, Samulski RJ, Nakamura T, Soballe K, O'Keefe RJ, Boyce BF, Schwarz EM. Remodeling of cortical bone allografts mediated by adherent rAAV-RANKL and VEGF gene therapy. *Nat Med.* 2005; 11(3):291–297. [PubMed: 15711561]
19. Koefoed M, Ito H, Gromov K, Reynolds DG, Awad HA, Rubery PT, Ulrich-Vinther M, Soballe K, Guldberg RE, Lin AS, O'Keefe RJ, Zhang X, Schwarz EM. Biological effects of rAAV-caAlk2 coating on structural allograft healing. *Molecular therapy.* 2005; 12 (2):212–8. [PubMed: 16043092]
20. Reynolds DG, Hock C, Shaikh S, Jacobson J, Zhang X, Rubery PT, Beck CA, O'Keefe RJ, Lerner AL, Schwarz EM, Awad HA. Micro-computed tomography prediction of biomechanical strength in murine structural bone grafts. *J Biomech.* 2007; 40(14):3178–86. [PubMed: 17524409]
21. Pacicca DM, Patel N, Lee C, Salisbury K, Lehmann W, Carvalho R, Gerstenfeld LC, Einhorn TA. Expression of angiogenic factors during distraction osteogenesis. *Bone.* 2003; 33 (6):889–98. [PubMed: 14678848]
22. Jilka RL, Weinstein RS, Bellido T, Roberson P, Parfitt AM, Manolagas SC. Increased bone formation by prevention of osteoblast apoptosis with parathyroid hormone. *J Clin Invest.* 1999; 104(4):439–46. [PubMed: 10449436]
23. Tashjian AH Jr, Gagel RF. Teriparatide [human PTH(1–34)]: 2.5 years of experience on the use and safety of the drug for the treatment of osteoporosis. *J Bone Miner Res.* 2006; 21(3):354–65. [PubMed: 16491282]
24. Puzas JE, Houck J, Bukata SV. Accelerated fracture healing. *J Am Acad Orthop Surg.* 2006; 14(10 Suppl):S145–51. [PubMed: 17003188]
25. Alkhiary YM, Gerstenfeld LC, Krall E, Westmore M, Sato M, Mitlak BH, Einhorn TA. Enhancement of experimental fracture-healing by systemic administration of recombinant human parathyroid hormone (PTH 1–34). *J Bone Joint Surg Am.* 2005; 87(4):731–41. [PubMed: 15805200]
26. Kaback LA, Soung DY, Naik A, Geneau G, Schwarz EM, Rosier RN, O'Keefe RJ, Drissi H. Teriparatide (1–34 human PTH) regulation of osterix during fracture repair. *J Cell Biochem.* 2008
27. Manabe T, Mori S, Mashiba T, Kaji Y, Iwata K, Komatsubara S, Seki A, Sun YX, Yamamoto T. Human parathyroid hormone (1–34) accelerates natural fracture healing process in the femoral osteotomy model of cynomolgus monkeys. *Bone.* 2007; 40 (6):1475–82. [PubMed: 17369013]
28. Pettway GJ, Schneider A, Koh AJ, Widjaja E, Morris MD, Meganck JA, Goldstein SA, McCauley LK. Anabolic actions of PTH (1–34): use of a novel tissue engineering model to investigate temporal effects on bone. *Bone.* 2005; 36(6):959–70. [PubMed: 15878317]
29. Vahle JL, Long GG, Sandusky G, Westmore M, Ma YL, Sato M. Bone neoplasms in F344 rats given teriparatide [rhPTH(1–34)] are dependent on duration of treatment and dose. *Toxicol Pathol.* 2004; 32(4):426–38. [PubMed: 15204966]
30. Aspenberg P, Johansson T. Teriparatide improves early callus formation in distal radial fractures. *Acta Orthop.* 2010; 81(2):234–6. [PubMed: 20367417]
31. Barnes GL, Kakar S, Vora S, Morgan EF, Gerstenfeld LC, Einhorn TA. Stimulation of fracture-healing with systemic intermittent parathyroid hormone treatment. *J Bone Joint Surg Am.* 2008; 90(Suppl 1):120–7. [PubMed: 18292366]
32. Caruso A, Amico-Roxas M, Cutuli V, Prato A, Fiore CE, Clementi G. Parathyroid hormone fragment 1–34 inhibits drug-induced inflammation in various experimental models. *Eur J Pharmacol.* 1991; 198(1):85–8. [PubMed: 1717293]
33. Oteo-Alvaro A, Moreno E. Atrophic humeral shaft nonunion treated with teriparatide (rh PTH 1–34): a case report. *Journal of shoulder and elbow surgery/American Shoulder and Elbow Surgeons ... [et al].* 2010; 19(7):e22–8.
34. Rubery PT, Bukata SV. Teriparatide may accelerate healing in delayed unions of type III odontoid fractures: a report of 3 cases. *Journal of spinal disorders & techniques.* 2010; 23 (2):151–5. [PubMed: 20051918]

35. Komrakova M, Stuermer EK, Werner C, Wicke M, Kolios L, Sehmisch S, Tezval M, Daub F, Martens T, Witzelhausen P, Dullin C, Stuermer KM. Effect of human parathyroid hormone hPTH (1–34) applied at different regimes on fracture healing and muscle in ovariectomized and healthy rats. *Bone*. 2010; 47(3):480–92. [PubMed: 20580683]

\$watermark-text

\$watermark-text

\$watermark-text

Figure 1. Experimental model and design

(A) Schematic diagram of murine femur allograft surgery. Allografts were harvested from ICR mice and devitalized. Processed allograft were then implanted to 4mm-diaphyseal defects of left femurs of C57bl/6 mice and secured with intramedullary pins. Panels B–D show the experimental groups and treatment schedules for the dose response study, duration and timing study, and molecular time course study respectively. Group name indicates: dose of daily PTH₁₋₃₄ injection (L: low, M: middle, and H: high)-timing of commencing treatment (week)-timing withdrawal of treatment (week). Mice were sacrificed at week 9 in experiments 1 and 2 and at various time points up to 28 days in experiment 3.

Figure 2. Effects of dose and regimen of PTH₁₋₃₄ treatment on graft-host osseointegration at 9 weeks postsurgery

Micro-CT rendering of grafted femurs (full views in top panel and coronal sections in bottom panel) at 9 weeks postsurgery in experiments 1 and 2 demonstrated that allograft healing and incorporation is enhanced by various regimens of high dose (40 $\mu\text{g/kg/day}$) PTH₁₋₃₄ therapy (D–J), while low and middle doses (0.4 and 4 $\mu\text{g/kg/day}$) still showed discontinuities at the graft-host junctions (B and C) similar to untreated controls (A). Delayed PTH₁₋₃₄ treatment up to 1 or 2 weeks post surgery still improved allograft integration with the host, regardless of the duration of treatment (G–J), and were grossly comparable to immediate treatment. Only the 4 weeks-delayed PTH₁₋₃₄ treatment (K) did not show significant effects on graft-host union compared to untreated controls.

Figure 3. Effects of treatment regimen of high dose PTH₁₋₃₄ on graft-host osseointegration at 2 and 4 weeks postsurgery

Representative micro CT rendering of femoral allografts and surrounding callus at 2 weeks (A, C, E) and 4 weeks (B, D, F, and G) depicting 3-D frontal view, coronal cross sectional view, segmented callus, and the segmented allograft, respectively, of grafts treated with saline (controls) or 40 μ g/kg/day PTH₁₋₃₄ commencing immediately or after a 1- or 2-week delay. The callus volume (H) and bone mineral density (I) are quantified and plotted for the various treatment regimens. Allograft images are artificially colored in blue and red to calculate the union ratio (J), where blue represents bare allograft surface not in contact with surrounding mineralized callus, while red areas indicate regions of graft union with mineralized callus from the host. Bar plots (H–J) represent means and SEM. Asterisks indicate significant differences versus controls ($p < 0.05$).

Figure 4. Effects of treatment regimen of high dose PTH₁₋₃₄ on cartilage and bone formation, and osteoclast activity at 2 and 4 weeks postsurgery

Micrographs showing representative sagittal sections of grafted femurs treated with saline (A and D) and 40 $\mu\text{g/kg}$ PTH₁₋₃₄ commencing immediately postsurgery (B and E), after a 1-week delay (C and F), or a 2-week delay (G), respectively. Insets on the left hand side show higher magnification (100X) of the graft-host junction (Boxed area) stained with alcian blue/hematoxylin/orangeG, , while insets on the right hand side show the same view stained with TRAP at 100X and 200X, respectively. Note that cartilage area, which reflects chondrogenesis during endochondral bone formation stage, was increased by immediate PTH₁₋₃₄ treatment (B) compared to untreated control (A) and the 1-week delayed treatment (C). Note also that compared to controls (D) graft-host integration was improved at 4 weeks by PTH₁₋₃₄ treatment regardless of the timing of the treatment (E–G). While a number of osteoclasts were observed in the callus area at 2-weeks in all the groups including control, the number of osteoclasts in callus area decreased from 2 to 4 weeks with the progression of callus remodeling. At 4-weeks, only a few osteoclasts could be seen in the callus area in the control specimens, whereas a much greater number of osteoclasts could still be observed in callus area in the PTH₁₋₃₄ treatment groups. Osteoclasts were rarely seen on the surface of devitalized allograft (not shown), but some could be observed at the region of allograft where callus enveloped.

Figure 5. Analysis of mRNA expression levels at the allograft-host bone junctions over time up to 28 days postsurgery

Gene expression was quantified using real-time RT-PCR for selected markers of chondrogenesis (*col1a1*), hypertrophy and remodeling (*col10a1*, *vegf-a*, and *mmp9*), osteogenesis (*col1a1* and *oc*), and osteoclastogenesis (*ctsk* and *trap*). Bar plots represent means and SEM ($n=3$ per time point). Asterisks indicate significant differences versus controls ($p < 0.05$).

Table 1

Biomechanical (torsional) properties of the grafted femurs at 9 weeks, presented as mean $\pm$ standard deviation. Asterisks indicate significant differences from the Saline control when $p < 0.05$ (p -values shown in parenthesis)

	Yield Torque (T_{Yield})	Ultimate Torque (T_{Ult})	Torsional Rigidity	Work to T_{Ult}
	N.mm	N.mm	N.mm/rad.mm ⁻¹	N.mm.Rad/mm
Saline Control	2.9 $\pm$ 2.7	5.3 $\pm$ 5.0	411.4 $\pm$ 395.8	0.08 $\pm$ 0.08
<u>Dose and Duration Effects for Immediate PTH₁₋₃₄ Treatment</u>				
L-0-4	3.0 $\pm$ 4.1 ($p=1.00$)	4.1 $\pm$ 5.8 ($p=0.99$)	312.9 $\pm$ 334.1 ($p=0.99$)	0.06 $\pm$ 0.09 ($p=0.99$)
M-0-4	2.7 $\pm$ 2.0 ($p=1.00$)	6.1 $\pm$ 4.7 ($p=1.00$)	383.5 $\pm$ 283.7 ($p=1.00$)	0.10 $\pm$ 0.07 ($p=1.00$)
H-0-2	8.7 $\pm$ 2.7 ($p=0.07$)	12.6 $\pm$ 4.1* ($p<0.05$)	672.1 $\pm$ 228.0 ($p=0.51$)	0.22 $\pm$ 0.09 ($p=0.10$)
H-0-4	7.5 $\pm$ 3.7 ($p=0.23$)	11.7 $\pm$ 4.4 ($p=0.11$)	777.1 $\pm$ 262.1 ($p=0.15$)	0.19 $\pm$ 0.18 ($p=0.28$)
H-0-9	10.5 $\pm$ 7.4* ($p<0.01$)	13.3 $\pm$ 7.1* ($p<0.05$)	707.2 $\pm$ 350.7 ($p=0.36$)	0.24 $\pm$ 0.13* ($p<0.05$)
<u>High Dose, 1- week Delayed PTH₁₋₃₄ Treatment</u>				
H-1-3	6.7 $\pm$ 4.8 ($p=0.46$)	8.3 $\pm$ 4.9 ($p=0.87$)	583.2 $\pm$ 227.9 ($p=0.91$)	0.10 $\pm$ 0.05 ($p=1.00$)
H-1-5	8.1 $\pm$ 5.0 ($p=0.13$)	10.3 $\pm$ 5.6 ($p=0.31$)	695.5 $\pm$ 401.5 ($p=0.41$)	0.19 $\pm$ 0.08 ($p=0.25$)
<u>High Dose, 2- week Delayed PTH₁₋₃₄ Treatment</u>				
H-2-4	8.1 $\pm$ 7.1 ($p=0.06$)	12.5 $\pm$ 8.5* ($p=0.05$)	644.6 $\pm$ 466.7 ($p=0.64$)	0.22 $\pm$ 0.13 ($p=0.07$)
H-2-6	9.9 $\pm$ 6.6* ($p<0.05$)	14.0 $\pm$ 7.7* ($p<0.01$)	840.7 $\pm$ 423.7 ($p=0.07$)	0.26 $\pm$ 0.15* ($p<0.01$)
<u>High Dose, 4- week Delayed PTH₁₋₃₄ Treatment</u>				
H-4-8	2.8 $\pm$ 3.1 ($p=1.00$)	4.7 $\pm$ 4.3 ($p=1.00$)	228.3 $\pm$ 292.7 ($p=0.89$)	0.15 $\pm$ 0.19 ($p=0.79$)

Table 2

Histomorphometry analysis of allograft Healing. Values are mean±SD (n=5 for each group at week 2 and n=4 for each group at week 4). Asterisks indicate significant differences from saline controls ($p<0.05$).

Parameter	Post-surgery week 2				Post-surgery week 4			
	Saline Control	PTH ₁₋₃₄ - Immediate	PTH ₁₋₃₄ - 1w delayed	PTH ₁₋₃₄ - Immediate	Saline Control	PTH ₁₋₃₄ - Immediate	PTH ₁₋₃₄ - 1w delayed	PTH ₁₋₃₄ - 2w delayed
Cartilage area (mm ²)	0.40±0.23	1.09±0.85	0.63±0.39	0.09±0.18	0.02±0.03	0.09±0.18	0.05±0.04	0.04±0.03
Osteoclast number on graft surface (/section)	2.3±1.9	8.6±3.7*	5.3±4.3	16.4±7.6	5.0±4.5	16.4±7.6	20.6±8.7*	17.8±8.3
Osteoclast number in callus area (/mm ²)	55.0±12.9	59.4±14.7	64.7±17.0	33.2±14.6	3.7±3.9	33.2±14.6	33.6±22.4	38.78±17.2*