Published in final edited form as:

Pharmacol Biochem Behav. 2012 February ; 100(4): 705–711. doi:10.1016/j.pbb.2011.09.015.

Expression profiling in neuropsychiatric disorders: emphasis on glutamate receptors in bipolar disorder

Stephen D. Ginsberg^{1,3,4,*}, Scott E. Hemby⁵, and John F. Smiley²

¹Center for Dementia Research, Nathan Kline Institute, Orangeburg, NY

²Program in Cognitive Neuroscience and Schizophrenia, Nathan Kline Institute, Orangeburg, NY

³Department of Psychiatry, New York University Langone Medical Center, New York, NY

⁴Department of Physiology & Neuroscience, New York University Langone Medical Center, New York, NY

⁵Departments of Physiology and Pharmacology, Wake Forest University School of Medicine, Winston-Salem, NC

Abstract

Functional genomics and proteomics approaches are being employed to evaluate gene and encoded protein expression changes with the tacit goal to find novel targets for drug discovery. Genome-wide association studies (GWAS) have attempted to identify valid candidate genes through single nucleotide polymorphism (SNP) analysis. Furthermore, microarray analysis of gene expression in brain regions and discrete cell populations has enabled the simultaneous quantitative assessment of relevant genes. The ability to associate gene expression changes with neuropsychiatric disorders, including bipolar disorder (BP), and their response to therapeutic drugs provides a novel means for pharmacotherapeutic interventions. This review summarizes gene and pathway targets that have been identified in GWAS studies and expression profiling of human postmortem brain in BP, with an emphasis on glutamate receptors (GluRs). Although functional genomic assessment of BP is in its infancy, results to date point towards a dysregulation of GluRs that bear some similarity to schizophrenia (SZ), although the pattern is complex, and likely to be more complementary than overlapping. The importance of single population expression profiling of specific neurons and intrinsic circuits is emphasized, as this approach provides informative gene expression profile data that may be underappreciated in regional studies with admixed neuronal and non-neuronal cell types.

Keywords

expression profiling; glutamate receptors; genome-wide association studies; major depressive disorder; molecular fingerprint; postmortem human brain; schizophrenia; single cell analysis

© 2011 Elsevier Inc. All rights reserved.

Correspondence: Stephen D. Ginsberg, Ph.D., Center for Dementia Research, Nathan Kline Institute, New York University Langone Medical Center, 140 Old Orangeburg Road, Orangeburg, NY 10962, 845-398-2170 (phone), 845-398-5422 (FAX), ginsberg@nki.rfmh.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Neuropsychiatric disorder overview

Psychotic disorders are characterized clinically by hallucinations and/or delusions and are categorized as primary psychiatric/psychotic disorders due to precipitants including substance abuse, psychoactive medications, and head trauma or secondary to conditions that affect the central nervous system (CNS) including brain tumors, cerebral infarcts, epilepsy, HIV-AIDS, multiple sclerosis, and neurodegenerative disorders, among others. Disorders in which psychosis is a primary symptom include schizophrenia (SZ), schizoaffective disorder, and schizophreniform disorder as well as delusional disorder and brief and shared psychotic disorders. In addition, psychoses can present in major depressive disorder (MDD) and in bipolar disorder (BP). Unfortunately, psychotic disorders significantly impair the ability of the affected individual to communicate effectively, have clear mentation, and behave in a socially appropriate and consistent manner, which negatively impacts the ability to conduct a normal lifestyle. SZ-related disorders affect approximately 2.4 million people, or 1.1% of the USA population >18 years old each year (Gejman et al., 2010; Regier et al., 1993) while the spectrum of BP affect approximately 5.7 million people, or 2.6% of the USA population >18 years old each year (Kessler et al., 2005a, 2005b; Vieta and Morralla, 2010). These disorders place a profound emotional, financial, and social burden on the individual, their families, and society at large.

To date, the majority of medications receiving FDA approval for neuropsychiatric disorders are modifications of drugs that were developed several decades ago based upon clinical observations, as well as trial and error. Unfortunately, *bona fide* targets for neuropsychiatric and neurodegenerative disorders are relatively scarce, and there is a paucity of proof-of-concept studies for novel drugs in psychiatric and neurological practice. Clearly, there is an urgent need to identify relevant neurobiological mechanisms that can be translated into treatments for amelioration of psychotic disorders. SZ, MDD, and BP are posited to be multigenic, multifactorial diseases that likely converge in final common pathogenic mechanisms that manifest as psychosis.

Current understanding of the mechanisms of psychotic disorders emanate in part from the identification of genetic loci in defined homogeneous populations (ideally replicated in an unrelated cohort) which harbor disease-causing and disease-modifying genes. Discovery of affected genotypes and kindreds has proven beneficial for elucidation of the molecular pathogenesis for these complex neuropsychiatric disorders, and will likely be invaluable tools for genetic counseling as well as diagnosis and selection of medications for affected individuals as the field of translational molecular psychiatry matures into clinical practice. The utilization of human postmortem brain tissues from affected individuals in combination with their respective clinical diagnoses and pathological records relative to appropriate age-matched controls provide a powerful means for investigating disease-related processes related to psychosis. However, mechanisms underlying psychosis remain poorly understood in SZ, BP, and related disorders, effectively impeding the development of relevant and valid animal and cellular models. Traditionally, animal and cellular models have proven useful in relating the role of specific genes, gene targets, and affected pathways with various behavioral indices associated with psychotic disorders. Thus, there are currently few alternatives for studying the etiology and mechanisms for these diseases besides the use of well-characterized human postmortem brains, which places a huge demand on the maintenance and growth of high quality tissue repositories and brain banks that specialize in accrual of brains from neuropsychiatric patients as well as unaffected age-matched normal subjects.

Bipolar disorder

Individuals diagnosed with BP present initially with either a manic or depressive episode. A diagnosis of BP type I occurs if a distinct mood change occurs (elevated, expansive, or irritable) for a period of at least a week, or shorter depending if hospitalization is necessitated (DSM-IV criterion A) (Baldessarini et al., 2010; DSM-IV, 2000). A manic episode involves at least three or more symptoms including altered speech patterns, decreased sleep, distractibility, excessive talking, grandiosity, increased goal directed activity, inflated self esteem, overindulgence in pleasurable activities, and racing thoughts (DSM-IV criterion B) (DSM-IV, 2000). A diagnosis of BP type II, a depressive state, typically occurs with a relatively short duration and severity of manic symptoms, and the occurrence of at least one depressive episode. Therefore, a diagnosis of BP may require several years of monitoring, because a manic episode may not occur for quite some time after initial presentation. A gradation and combination of these diverse symptoms has made diagnosis even more difficult, and categories such as bipolar spectrum disorder (BPS) and subthreshold BP are recognized by the World Health Organization. Current estimated worldwide prevalence of BP include an aggregate lifetime prevalence of 2.4% for BPS, 1.4% for subthreshold BP, 0.6% for BP type I, and 0.4% for BP type II (Merikangas et al., 2011). Estimated one year prevalence of 1.5% for BPS, 0.8% for subthreshold BP, 0.4% for BP type I, and 0.3% for BP type II (Merikangas et al., 2011). Unfortunately, suicide rates of BP patients are among the highest of any neuropsychiatric disorder, including an approximate 10-fold increase in the risk of suicide above the general population (Cassidy, 2011). Persons with BP also tend to abuse drugs and alcohol, increasing suicide and overall mortality levels (Baldessarini et al., 2010; McDonald et al., 2011; Oquendo et al., 2010). Neuroimaging studies implicate volumetric changes in several brain relevant limbic brain regions including the amygdala, anterior cingulate, and hippocampus in BP, suggesting some form of connectivity-based degeneration in BP, but these findings are inconsistent and difficult to differentiate from similar changes seen in other neuropsychiatric disorders (Cherlyn et al., 2010; Friedman et al., 2006; Usher et al., 2010). In summary, treating BP is a public health issue that requires improved diagnosis and treatment, particularly for those individuals that may attempt suicide, and a pharmacogenomic approach awaits identification of suitable targets for development.

Genome-wide association studies (GWAS) in bipolar disorder: identification of several glutamatergic neurotransmission susceptibility candidates

Twin studies have established that BD is a highly heritable condition (Barnett and Smoller, 2009), and research aimed at finding susceptibility genes have been undertaken worldwide. GWAS investigations are well-recognized high-throughput functional genomics assessments that evaluate genes and single nucleotide polymorphisms (SNPs) from affected and unaffected individuals. GWAS studies are useful to identify candidate genes that can be evaluated further in microarray and qPCR based studies. There have been several GWAS studies performed on cohorts of subjects with BP, including large multi-institution evaluations with thousands of subjects with BP compared to age-matched non-psychiatric controls (Belmonte Mahon et al., 2011; Scott et al., 2009; Sklar et al., 2008). Several BP susceptibility loci have been found, including myosin 5B (MYO5B), voltage-dependent calcium channel, L-type, $\alpha 1C$ subunit (CACNA1C) and ankyrin 3 (ANK3), the latter in which the specific identified SNP (rs10994336) has been replicated in several independent studies (Ferreira et al., 2008; Ruberto et al., 2011; Scott et al., 2009; Sklar et al., 2008; Takata et al., 2011). Unfortunately, the majority of risk factors for BP remain unknown, likely due to the heterogeneity of this multigenic neuropsychiatric disorder. This was

evidenced in a recent GWAS analysis of three large cohorts (two from the USA and one from Germany) where the collaborative research group failed to identify any genome-wide significance for any SNPs, even with a breakout of two BP phenotypes, age at onset and psychotic symptoms, which can be interrogated with relatively high power to detect genetic variation (Belmonte Mahon et al., 2011).

Despite these tangible setbacks, there is a serviceable website with a current, annotated database of genetic studies of BP (<http://bioprogramming.bsd.uchicago.edu/BDStudies/>) (Piletz et al., 2011) that includes several candidate genes related to glutamatergic neurotransmission that are worthy of mention. Both positive and negative findings are cited. Specific glutamatergic neurotransmission candidate susceptibility genes include citron (CIT) positive (Lyons-Warren et al., 2005) and negative (Yosifova et al., 2009), D-amino acid oxidase (DAO) positive (Fallin et al., 2005; Prata et al., 2008) and negative (Shi et al., 2008), and the nitric oxide synthase (NOS) genes NOS1 (neuronal) positive (Fallin et al., 2005; Yosifova et al., 2009) and negative (Buttenschon et al., 2004; Gratacos et al., 2009; Okumura et al., 2010), and NOS3 (endothelial) positive (Reif et al., 2006) and negative (Gratacos et al., 2009; Sklar et al., 2002). GluRs identified as BP candidate genes include the ionotropic α -amino-3-hydroxy-5-methylisoxazolepropionic acid (AMPA) receptor subunits GRIA1 positive (Kerner et al., 2009; Shi et al., 2008) and negative (Gratacos et al., 2009), GRIA2 positive (Perlis et al., 2009) and negative (Shi et al., 2008; Sklar et al., 2002), kainate (KA) receptor subunit GRIK4 positive (Pickard et al., 2008; Pickard et al., 2006) and negative (Gratacos et al., 2009), N-methyl-D-aspartate (NMDA) receptor subunits GRIN1 positive (Mundo et al., 2003; Shi et al., 2008; Yosifova et al., 2009) and negative (Georgi et al., 2006), GRIN2A positive (Itokawa et al., 2003) and negative (Gratacos et al., 2009; Shi et al., 2008), GRIN2B positive (Avramopoulos et al., 2007; Fallin et al., 2005; Lorenzi et al., 2010; Martucci et al., 2006; Zhao et al., 2011) and negative (Gratacos et al., 2009; Shi et al., 2008), GRIN2C positive (Shi et al., 2008), and GRIN2D positive (Shi et al., 2008), and metabotropic glutamate receptors (mGluRs) GRM1 positive (Baum et al., 2008; Frank et al., 2011) and negative (Fan et al., 2010; Shi et al., 2008), GRM3 positive (Fallin et al., 2005; Sklar et al., 2008) and negative (Gratacos et al., 2009; Marti et al., 2002; Shi et al., 2008; Yosifova et al., 2009), GRM4 positive (Fallin et al., 2005) and negative (Shi et al., 2008; Sklar et al., 2002), GRM7 positive (Gratacos et al., 2009) and negative (Baum et al., 2008; Shi et al., 2008; Sklar et al., 2002; Yosifova et al., 2009), among other glutamatergic neurotransmission candidates (Cherlyn et al., 2010; Piletz et al., 2011). A high percentage of these glutamatergic neurotransmission/GluR candidate genes also overlap in susceptibility with SZ (Cherlyn et al., 2010; Fallin et al., 2005; Frank et al., 2011; Marti et al., 2002; Martucci et al., 2006; Pickard et al., 2006), highlighting the complex, multigenic, multifactorial nature of these neuropsychiatric disorders and the difficulty teasing apart genes that might discriminate these two conditions. As evidenced by a nearly equal amount of positive and negative association findings of glutamatergic and GluR genes, the apparent lack of agreement across cohorts and studies emphasizes the high likelihood that susceptibility alleles are moderate in effect size and require very large, homogeneous populations for presence detection.

Expression profiling in neuropsychiatric disorders

Due to a paucity of transformative therapy for neuropsychiatric disorders, the majority of current drugs being approved in the present environment are modifications of therapeutics that were identified by investigators in previous generations. Actual targets for neuropsychiatric disorders are difficult to validate, and proof-of-concept studies for novel drugs are rare in molecular psychiatry. The implementation of gene expression profiling methods, including microarray technology and hopefully next generation sequencing technologies in the near future, enables researchers to generate a relative quantitative index

of multiple genes simultaneously from postmortem human brain tissue samples as well as optimally prepared brain tissues or cells from relevant animal and/or cellular models in either a high-throughput or moderate-throughput format (Ginsberg and Mirnics, 2006; Hakak et al., 2001; Mimmack et al., 2002; Mirnics et al., 2000). Molecular fingerprinting is performed by extracting RNAs from identified regions or cells, followed by RNA amplification, and hybridizing labeled RNA to desired array platforms. Our research group favors the study of individual populations of neurons for microarray analysis, as this approach enables an assessment of gene expression alterations that occur in one relevant cell type {e.g., entorhinal cortex stellate cells (Fig. 1), neocortical pyramidal and GABAergic interneurons, cholinergic basal forebrain neurons, and CA1 pyramidal neurons, among others} without the confound of admixed adjacent neuronal and non-neuronal (e.g., astrocytes, microglia, oligodendrocytes, vascular, and epithelial) cell types (Ginsberg et al., 2006, 2010, 2011a, 2011b; Hemby et al., 2002, 2003). Assessing gene expression patterns in targeted neuronal populations is important since reliance on regional assessment of expression may emphasize genes contained in the majority of the neurons and/or those in highest abundance in the region, or those in glial and vascular epithelial cell populations. Regional assessment does not adequately reflect the alterations in gene expression occurring in target neurons due to masking and/or dilution effects caused by changes in the total neuronal population of that region. Such differences are highlighted in Figure 1 where gene expression is compared between microdissected neurons and the region from which the neurons were extracted in normal subjects, providing a clear example of differential expression patterns between the two approaches. These results suggest that, when targeting neurons in heterogeneous neuronal populations for analysis, it is informative to refine the level of molecular analysis to a single neuronal population as well as to analyze the surrounding tissue sample. Quantification of hybridization signal intensity is performed to assess the relative expression level of each feature on an array. Gene expression is determined using bioinformatics software and statistical approaches, enabling coordinate expression level analyses which can be correlated with other variables including age, gender, and antipsychotic medication levels, among others (Altar et al., 2009; Deep-Soboslay et al., 2011; Ginsberg et al., 2010; Ginsberg and Mirnics, 2006). As the majority of neuropsychiatric disorders, including BP, are treated with a combination of antipsychotic and mood stabilizing medications, it is often difficult to determine whether gene expression level changes in postmortem brain tissues are caused organically by the disorder or whether they are a side effect of chronic drug treatments. Unfortunately, antipsychotic medications such as haloperidol and clozapine and mood stabilizers including lithium, valproate, and carbamazepine are notorious for interacting with a wide number of targets and downstream signaling pathways and may have neurodegenerative effects independent of the neuropsychiatric disorder they are being used to treat (Altar et al., 2003, 2009; Bachmann et al., 2005; Gould et al., 2004; Lewis, 2011). The implementation of animal models (principally nonhuman primates and rodents) for drug treatment studies has been helpful to ascertain specific effects of antipsychotic/mood stabilizer treatment on gene and encoded protein expression within a few systems (notably catecholaminergic, cholinergic, GABAergic, and glutamatergic) (Cheng et al., 2008; Fasulo and Hemby, 2003; Fatemi et al., 2006; Lewis et al., 2008; Melchitzky and Lewis, 2008; O'Connor et al., 2007b). For example, select AMPA GluRs are differentially regulated in monkeys chronically treated with antipsychotic drugs, as clozapine treatment significantly decreases GRIA1 and increases GRIA3 mRNA levels, whereas both clozapine and haloperidol increase expression of GRIA2 (O'Connor et al., 2007a, 2007b). Importantly, these AMPA receptor alterations were found pyramidal neurons microdissected from layers II/III and V of dorsolateral prefrontal cortex, but not when regional assessment of dorsolateral prefrontal cortex was performed (O'Connor et al., 2007a, 2007b), indicating these changes are likely to be pyramidal neuron specific. In contrast, delivery of the mood stabilizers lithium, valproate, and carbamazepine has been shown to downregulate mRNA and protein levels of the KA

receptor GRIK2 in astrocytes but not in neurons (Li et al., 2009), indicating the importance of assaying both neuronal and non-neuronal populations in neuropsychiatric disorders and when considering the possibly confounding effects of antipsychotic/mood stabilizing therapeutics. Notably, moderate- and high-throughput, fairly unbiased mRNA methodologies including microarrays provide a cost-effective, simultaneous assessment of gene expression levels. These approaches are somewhat technically challenging, and require high quality RNA to measure the relatively small (20-40%) changes in brain gene expression that typify neuropsychiatric disease (Altar et al., 2009; Atz et al., 2007; Eberwine et al., 2001; Ginsberg, 2008; Ginsberg and Mirnics, 2006; Jurata et al., 2004; Mirnics et al., 2006). They also require validation by independent methods, including real-time quantitative PCR (qPCR) (Bustin, 2010; Nolan et al., 2006) and/or *in situ* hybridization (Bernard et al., 2011; Hashimoto et al., 2008; Mirnics et al., 2000, 2006) as well as analytical measurements of their respective encoded protein products.

Gene expression analysis in bipolar disorder

Published microarray analyses of neuropsychiatric disorders have focused mainly on employing postmortem brain samples from SZ subjects, and studies employing brains accrued from MDD subjects are emerging. To date, only a few postmortem brain studies have evaluated expression level changes in BP. Although a comprehensive assessment of all the genes altered in BP is beyond the scope of this review, classes of transcripts that display differential regulation (principally downregulation) in BP versus age-matched controls include 14-3-3 family genes (e.g., YWHAQ and YWHAZ; which also overlap with changes observed in SZ) (Altar et al., 2009; Elashoff et al., 2007; Klempan et al., 2007; Konradi et al., 2004; Sequeira et al., 2007; Sibille et al., 2004), neuropeptides {e.g., cholecystokinin (CCK), tachykinin, precursor 1 (TAC1), proenkephalin (PENK)}, protein kinases {e.g., calcium/calmodulin-dependent protein kinase kinase 2, beta (CAMKK2), and cyclin-dependent kinase 5 regulatory subunit 1 (CDK5R1)}, ubiquitin pathway genes, and G-protein coupled receptor signalling, among others (Altar et al., 2009; Le-Niculescu et al., 2007; Ogden et al., 2004; Ryan et al., 2006). In addition, several studies that employ subregional microarray/qPCR analyses are highlighted because they illustrate the power of discriminating gene expression levels from presumably affected cell types and/or regions. Specifically, a study of cell cycle regulation markers in SZ and BP within subregions of the hippocampal formation indicates complex, region-specific alterations in genes associated with DNA repair mechanisms. For example, mRNA for the methyl-CpG binding domain protein (MBD4) is upregulated in the stratum oriens region of the CA1 and CA2/CA3 hippocampal sectors (Benes et al., 2009). Moreover, related genes involved in the transcriptional complex, G1 and G2 checkpoints, and neurotrophin/neurogenesis classes are dysregulated in a disease-specific, regionally-specific manner, including downregulation of neuregulin 1 (NRG1), neurotrophin-3 (NT-3), and several fibroblast growth factor subtypes (FGF3 and FGF9) within CA1 of BP subjects, whereas upregulation of neuropilin 1 (NRP1), the neurotrophin-3 receptor TrkC, and the roundabout axon guidance receptor subtype 1 (ROBO1) is found in the same subjects within the CA2/CA3 sector (Benes et al., 2009). These hippocampal subregional results are consistent with *post hoc* analyses of regional hippocampal investigations that show upregulation of 19/44 genes in the apoptotic pathway in BP (Benes et al., 2006). Within the brainstem, a regional study of the locus coeruleus combining laser capture microdissection (LCM) with microarray analysis found several genes related to glutamatergic neurotransmission in astrocytes, including glial high affinity glutamate transporter member 2 (SLC1A2), glial high affinity glutamate transporter member 3 (SLC1A3), and glutamine synthetase (GLUL) differentially regulated in subjects with MDD compared to BP (Bernard et al., 2011). Specifically, dysregulation of these genes were found via microarray analysis in MDD but not BP subjects, which was validated via qPCR and *in situ* hybridization (Bernard et al., 2011). These observations point to the importance

of studying neuronal as well as non-neuronal populations in multiple interconnected brain regions (i.e., not simply restricting analyses to forebrain structures such as the hippocampal formation and dorsolateral prefrontal cortex) and enabling comparisons between and across neuropsychiatric disorders including BP, MDD, and SZ.

Glutamatergic neurotransmission gene expression analysis in bipolar disorder

Glutamatergic signaling, in particular through ionotropic AMPA and NMDA receptors, plays a crucial role in excitatory neurotransmission, synaptic function, neuroplasticity, and development. Dysfunction of the glutamatergic system, notably within excitatory circuits driven by AMPA and NMDA receptors, as well as within non-neuronal cells such as astrocytes and oligodendrocytes, has been implicated in the pathogenesis of several neuropsychiatric disorders, including BP, MDD, and SZ (Javitt, 2007; Marsden, 2011; Sequeira and Turecki, 2006). Although relatively little information has been published on expression level differences of transcripts related to glutamatergic neurotransmission and/or GluRs to date in BP relative to the ongoing work being performed in SZ and MDD, the data that is available is worth summarizing. Specifically, high-density oligonucleotide microarray analyses within postmortem BP brains have identified differential regulation of several genes, including GluRs. Specifically, downregulation of the KA receptor GRIK1 and mGluR receptor GRM1 are observed in a study where a total of 53 genes are differentially regulated (Iwamoto et al., 2004; Kato et al., 2007). Importantly, GRIK1 downregulation has been replicated in an independent study where the research group hypothesized that a significant portion of the expression level changes occur on glial cells (Choudary et al., 2005), consistent with a larger body of regional microarray studies in SZ where glial alterations have been reported by several research groups (Hakak et al., 2001; Haroutunian et al., 2007; Mirmics et al., 2000; Tkachev et al., 2003). Cell type-specific decrements in GluRs within BP and SZ have been investigated, including significant loss of KA receptors GRIK1 and GRIK1 and NMDA receptor GRIN2A in GAD67-positive GABAergic interneurons within the cingulate cortex of subjects with BP and SZ (Woo et al., 2004, 2007). These studies point to the importance of evaluating specific cell types within forebrain regions to determine specific changes in relevant classes of transcripts. Importantly, these observations in BP are consistent with regional and single population expression profiling studies in SZ that indicate frank glutamatergic dysfunction as a mechanism underlying pathophysiology (Hemby et al., 2002; Mirmics et al., 2000). Specifically, downregulation of select AMPA and NMDA receptor subunits via microarray analysis are consistent with *in situ* hybridization and PCR-based observations of decreased GluR and glutamate transporter expression within the hippocampus as well as frontal and temporal neocortex (Beneyto and Meador-Woodruff, 2008; Eastwood and Harrison, 2005, 1999; Humphries et al., 1996; McCullumsmith et al., 2007; Mueller and Meador-Woodruff, 2004; Rao et al., 2010; Uezato et al., 2009). Downregulation of the GluR-interacting partners SAP102 and α -internexin have also been reported in BP (McCullumsmith et al., 2007; Pennington et al., 2008). The general finding of decreased GluR expression is consistent with increased glutamate levels in BP cerebral cortex (Hashimoto et al., 2007; Kim and Webster, 2010). Overall, downregulation of ionotropic (as well as metabotropic) GluRs on both pyramidal neurons and calbindin-immunoreactive, parvalbumin-immunoreactive, and calretinin-immunoreactive GABAergic interneurons may have profound downstream effects upon excitatory neurotransmission and subsequent behavioral and cognitive sequelae.

Future studies are warranted in postmortem BP brain tissues to characterize further potential expression level dysregulation of markers of glutamatergic neurotransmission and GluRs.

Importantly, next generation technologies are now being employed in conjunction with the use of postmortem brains obtained from neuropsychiatric patients (including those with BP) to comprehend the functional genomics/genetic variant imbalances that underlie these disorders from a population-based perspective (Lin et al., 2011; Tang et al., 2011). Recent studies include high-throughput microRNA (miRNA) profiling of human miRNAs, which revealed dysregulation of 7 miRNAs in SZ subjects and an additional 15 miRNAs in BP subjects (Kim et al., 2010). Interestingly, all 22 of these miRNAs were determined via *in silico* mining techniques to regulate brain specific genes that are classified within neurodevelopment and behavior hubs (Kim et al., 2010), potentially impacting the development of SZ and BP. Copy number variation (CNV) analysis in rare variants has also generated significant interest for studying the complex genetic susceptibility of multigenic diseases, including neuropsychiatric disorders (Chen et al., 2010; Grozeva et al., 2010; Saus et al., 2010). Preliminary candidate genes with likely CNVs related to SZ, MDD, and BP include mGluR GRM7 (Saus et al., 2010), consistent with a GWAS study in BP (Gratacos et al., 2009). Next generation deep sequencing and epigenetic approaches will likely help the neuropsychiatric disorder field by uncovering rare genetic variants which may be exploited to understand mechanisms underlying the disease process. Although these new functional genomics technologies generate significant excitement, each has caveats and limitations that have to be acknowledged (Chen et al., 2010; Tang et al., 2011), particularly to their applicability to populations of subjects, and integration of these results with existing microarray, GWAS, and biochemical studies are critical for mechanistic studies of these complex disorders. Moreover, current microarray and qPCR-based studies illustrate the importance of studying subregions and individual cell types for future gene discovery studies in neuropsychiatric disorders even though these approaches are more technically challenging than regional and global tissue based endeavors (Ginsberg et al., 2011a).

Conclusions

The search for genes underlying the pathophysiology BP has been complicated by a limited understanding of pathogenesis, a paucity of animal models, and the genetic and phenotypic complexity of this debilitating neuropsychiatric disorder. This review has illustrated small, but significant breakthroughs in GWAS and expression profiling pertinent to glutamatergic neurotransmission and GluR expression. Glutamatergic dysfunction has been implicated in SZ, MDD, and BP, and differentiating these changes poses a challenge for the field. We posit that expression profiling via subregional and/or homogeneous population approaches will be more informative than global or regional studies with admixed neuronal and non-neuronal cell types that may have complex alterations occurring throughout disease onset and progression. Strategic emphasis is required to continue efforts for the systematic accrual of clinically characterized BP brain tissue and age-matched unaffected control subjects to further neuropathological analyses as well as proposed state-of-the-art connectome and circuit driven approaches that enable highly targeted functional genomic and proteomic assessments. In conclusion, GWAS and population-directed microarray and qPCR interrogative strategies along with exciting next generation sequencing technologies may help develop novel agents that target systems and/or circuits adversely affected specifically during BP pathogenesis, potentially reducing the problems of drug interactions and unwanted side effects that currently plague the treatment approach.

Acknowledgments

We thank Melissa J. Alldred, Ph.D., Cynthia Bleiwas, Helen M. Chao, Ph.D., Shaoli Che, M.D., Ph.D., Irina Elarova, Shaona Fang, Khadija Figarsky, and Arthur Saltzman for their assistance. Support for this project comes from the NIMH (MH085208 and MH086385; SDG and JFS). We also express our appreciation to the families of the patients studied here who made this research possible.

References

- Altar, CA.; Martin, A.; Thurkauf, A. Antipsychotic Agents. In: Abraham, DJ., editor. *Burger's Medicinal Chemistry and Drug Discovery*. Hoboken: Wiley & Sons; 2003. p. 599-672.
- Altar CA, Vawter MP, Ginsberg SD. Target identification for CNS diseases by transcriptional profiling. *Neuropsychopharmacology*. 2009; 34:18–54. [PubMed: 18923405]
- Atz M, Walsh D, Cartagena P, Li J, Evans S, Choudary P, Overman K, Stein R, Tomita H, Potkin S, Myers R, Watson SJ, Jones EG, Akil H, Bunney WE Jr, Vawter MP. Methodological considerations for gene expression profiling of human brain. *J Neurosci Methods*. 2007; 163:295–309. [PubMed: 17512057]
- Avramopoulos D, Lasseter VK, Fallin MD, Wolyniec PS, McGrath JA, Nestadt G, Valle D, Pulver AE. Stage II follow-up on a linkage scan for bipolar disorder in the Ashkenazim provides suggestive evidence for chromosome 12p and the GRIN2B gene. *Genet Med*. 2007; 9:745–751. [PubMed: 18007143]
- Bachmann RF, Schloesser RJ, Gould TD, Manji HK. Mood stabilizers target cellular plasticity and resilience cascades: implications for the development of novel therapeutics. *Mol Neurobiol*. 2005; 32:173–202. [PubMed: 16215281]
- Baldessarini RJ, Vieta E, Calabrese JR, Tohen M, Bowden CL. Bipolar depression: overview and commentary. *Harv Rev Psychiatry*. 2010; 18:143–157. [PubMed: 20415631]
- Barnett JH, Smoller JW. The genetics of bipolar disorder. *Neuroscience*. 2009; 164:331–343. [PubMed: 19358880]
- Baum AE, Akula N, Cabanero M, Cardona I, Corona W, Klemens B, Schulze TG, Cichon S, Rietschel M, Nothen MM, Georgi A, Schumacher J, Schwarz M, Abou Jamra R, Hofels S, Propping P, Satagopan J, Detera-Wadleigh SD, Hardy J, McMahon FJ. A genome-wide association study implicates diacylglycerol kinase eta (DGKH) and several other genes in the etiology of bipolar disorder. *Mol Psychiatry*. 2008; 13:197–207. [PubMed: 17486107]
- Belmonte Mahon P, Pirooznia M, Goes FS, Seifuddin F, Steele J, Lee PH, Huang J, Hamshire ML, Depaulo JR Jr, Kelsoe JR, Rietschel M, Nothen M, Cichon S, Gurling H, Purcell S, Smoller JW, Craddock N, Schulze TG, McMahon FJ, Potash JB, Zandi PP. Genome-wide association analysis of age at onset and psychotic symptoms in bipolar disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2011; 156:370–378. [PubMed: 21305692]
- Benes FM, Lim B, Subburaju S. Site-specific regulation of cell cycle and DNA repair in post-mitotic GABA cells in schizophrenic versus bipolars. *Proc Natl Acad Sci USA*. 2009; 106:11731–11736. [PubMed: 19564623]
- Benes FM, Matzilevich D, Burke RE, Walsh J. The expression of proapoptosis genes is increased in bipolar disorder, but not in schizophrenia. *Mol Psychiatry*. 2006; 11:241–251. [PubMed: 16288314]
- Beneyto M, Meador-Woodruff JH. Lamina-specific abnormalities of NMDA receptor-associated postsynaptic protein transcripts in the prefrontal cortex in schizophrenia and bipolar disorder. *Neuropsychopharmacology*. 2008; 33:2175–2186. [PubMed: 18033238]
- Bernard R, Kerman IA, Thompson RC, Jones EG, Bunney WE, Barchas JD, Schatzberg AF, Myers RM, Akil H, Watson SJ. Altered expression of glutamate signaling, growth factor, and glia genes in the locus coeruleus of patients with major depression. *Mol Psychiatry*. 2011; 16:634–646. [PubMed: 20386568]
- Bustin SA. Why the need for qPCR publication guidelines?--The case for MIQE. *Methods*. 2010; 50:217–226. [PubMed: 20025972]
- Buttenshon HN, Mors O, Ewald H, McQuillin A, Kalsi G, Lawrence J, Gurling H, Kruse TA. No association between a neuronal nitric oxide synthase (NOS1) gene polymorphism on chromosome 12q24 and bipolar disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2004; 124B:73–75. [PubMed: 14681919]
- Cassidy F. Risk factors of attempted suicide in bipolar disorder. *Suicide Life Threat Behav*. 2011; 41:6–11. [PubMed: 21309819]

- Chen X, Li X, Wang P, Liu Y, Zhang Z, Zhao G, Xu H, Zhu J, Qin X, Chen S, Hu L, Kong X. Novel association strategy with copy number variation for identifying new risk Loci of human diseases. *PLoS ONE*. 2010; 5:e12185. [PubMed: 20808825]
- Cheng MC, Liao DL, Hsiung CA, Chen CY, Liao YC, Chen CH. Chronic treatment with aripiprazole induces differential gene expression in the rat frontal cortex. *Int J Neuropsychopharmacol*. 2008; 11:207–216. [PubMed: 17868501]
- Cherlyn SY, Woon PS, Liu JJ, Ong WY, Tsai GC, Sim K. Genetic association studies of glutamate, GABA and related genes in schizophrenia and bipolar disorder: a decade of advance. *Neurosci Biobehav Rev*. 2010; 34:958–977. [PubMed: 20060416]
- Choudary PV, Molnar M, Evans SJ, Tomita H, Li JZ, Vawter MP, Myers RM, Bunney WE Jr, Akil H, Watson SJ, Jones EG. Altered cortical glutamatergic and GABAergic signal transmission with glial involvement in depression. *Proc Natl Acad Sci USA*. 2005; 102:15653–15658. [PubMed: 16230605]
- Deep-Soboslay A, Benes FM, Haroutunian V, Ellis JK, Kleinman JE, Hyde TM. Psychiatric brain banking: three perspectives on current trends and future directions. *Biol Psychiatry*. 2011; 69:104–112. [PubMed: 20673875]
- DSM-IV Diagnostic and Statistical Manual of Mental Disorders. 4th. Arlington: American Psychiatric Association; 2000.
- Eastwood SL, Harrison PJ. Decreased expression of vesicular glutamate transporter 1 and complexin II mRNAs in schizophrenia: further evidence for a synaptic pathology affecting glutamate neurons. *Schizophr Res*. 2005; 73:159–172. [PubMed: 15653259]
- Eastwood SL, Harrison PJ. Detection and quantification of hippocampal synaptophysin messenger RNA in schizophrenia using autoclaved, formalin-fixed, paraffin wax-embedded sections. *Neuroscience*. 1999; 93:99–106. Order. [PubMed: 10430474]
- Eberwine J, Kacharina JE, Andrews C, Miyashiro K, McIntosh T, Becker K, Barrett T, Hinkle D, Dent G, Marciano P. mRNA expression analysis of tissue sections and single cells. *J Neurosci*. 2001; 21:8310–8314. [PubMed: 11606616]
- Elashoff M, Higgs BW, Yolken RH, Knable MB, Weis S, Webster MJ, Barci BM, Torrey EF. Meta-analysis of 12 genomic studies in bipolar disorder. *J Mol Neurosci*. 2007; 31:221–243. [PubMed: 17726228]
- Fallin MD, Lasseter VK, Avramopoulos D, Nicodemus KK, Wolyniec PS, McGrath JA, Steel G, Nestadt G, Liang KY, Hagan RL, Valle D, Pulver AE. Bipolar I disorder and schizophrenia: a 440-single-nucleotide polymorphism screen of 64 candidate genes among Ashkenazi Jewish case-parent trios. *Am J Hum Genet*. 2005; 77:918–936. [PubMed: 16380905]
- Fan J, Ionita-Laza I, McQueen MB, Devlin B, Purcell S, Faraone SV, Allen MH, Bowden CL, Calabrese JR, Fossey MD, Friedman ES, Gyulai L, Hauser P, Ketter TB, Marangell LB, Miklowitz DJ, Nierenberg AA, Patel JK, Sachs GS, Thase ME, Molay FB, Escamilla MA, Nimgaonkar VL, Sklar P, Laird NM, Smoller JW. Linkage disequilibrium mapping of the chromosome 6q21-22.31 bipolar I disorder susceptibility locus. *Am J Med Genet B Neuropsychiatr Genet*. 2010; 153B:29–37. [PubMed: 19308960]
- Fasulo WH, Hemby SE. Time-dependent changes in gene expression profiles of midbrain dopamine neurons following haloperidol administration. *J Neurochem*. 2003; 87:205–219. [PubMed: 12969267]
- Fatemi SH, Reutiman TJ, Folsom TD, Bell C, Nos L, Fried P, Pearce DA, Singh S, Siderovski DP, Willard FS, Fukuda M. Chronic olanzapine treatment causes differential expression of genes in frontal cortex of rats as revealed by DNA microarray technique. *Neuropsychopharmacology*. 2006; 31:1888–1899. [PubMed: 16407901]
- Ferreira MA, O'Donovan MC, Meng YA, Jones IR, Ruderfer DM, Jones L, Fan J, Kirov G, Perlis RH, Green EK, Smoller JW, Grozeva D, Stone J, Nikolov I, Chambert K, Hamshere ML, Nimgaonkar VL, Moskvina V, Thase ME, Caesar S, Sachs GS, Franklin J, Gordon-Smith K, Ardlie KG, Gabriel SB, Fraser C, Blumenstiel B, Defelice M, Breen G, Gill M, Morris DW, Elkin A, Muir WJ, McGhee KA, Williamson R, MacIntyre DJ, MacLean AW, St CD, Robinson M, Van Beck M, Pereira AC, Kandaswamy R, McQuillin A, Collier DA, Bass NJ, Young AH, Lawrence J, Ferrier IN, Anjorin A, Farmer A, Curtis D, Scolnick EM, McGuffin P, Daly MJ, Corvin AP, Holmans PA, Blackwood DH, Gurling HM, Owen MJ, Purcell SM, Sklar P, Craddock N. Collaborative genome-

- wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nat Genet.* 2008; 40:1056–1058. [PubMed: 18711365]
- Frank RA, McRae AF, Pocklington AJ, van de Lagemaat LN, Navarro P, Croning MD, Komiyama NH, Bradley SJ, Challiss RA, Armstrong JD, Finn RD, Malloy MP, MacLean AW, Harris SE, Starr JM, Bhaskar SS, Howard EK, Hunt SE, Coffey AJ, Ranganath V, Deloukas P, Rogers J, Muir WJ, Deary IJ, Blackwood DH, Visscher PM, Grant SG. Clustered coding variants in the glutamate receptor complexes of individuals with schizophrenia and bipolar disorder. *PLoS ONE.* 2011; 6:e19011. [PubMed: 21559497]
- Friedman JN, Hurley RA, Taber KH. Bipolar disorder: imaging state versus trait. *J Neuropsychiatry Clin Neurosci.* 2006; 18:296–301. [PubMed: 16963579]
- Gejman PV, Sanders AR, Duan J. The role of genetics in the etiology of schizophrenia. *Psychiatr Clin North Am.* 2010; 33:35–66. [PubMed: 20159339]
- Georgi A, Jamra RA, Schumacher J, Becker T, Schmael C, Deschner M, Hofels S, Wulff M, Schwarz M, Klopp N, Illig T, Propping P, Cichon S, Nothen MM, Rietschel M, Schulze TG. No association between genetic variants at the GRIN1 gene and bipolar disorder in a German sample. *Psychiatr Genet.* 2006; 16:183–184. [PubMed: 16969270]
- Ginsberg SD. Transcriptional profiling of small samples in the central nervous system. *Methods Mol Biol.* 2008; 439:147–158. [PubMed: 18370101]
- Ginsberg SD, Alldred MJ, Che S. Gene expression levels assessed by CA1 pyramidal neuron and regional hippocampal dissections in Alzheimer's disease. *Neurobiol Dis.* 2011a in press.
- Ginsberg SD, Alldred MJ, Counts SE, Cataldo AM, Neve RL, Jiang Y, Wu J, Chao MV, Mufson EJ, Nixon RA, Che S. Microarray analysis of hippocampal CA1 neurons implicates early endosomal dysfunction during Alzheimer's disease progression. *Biol Psychiatry.* 2010; 68:885–893. [PubMed: 20655510]
- Ginsberg SD, Che S, Wu J, Counts SE, Mufson EJ. Down regulation of trk but not p75NTR gene expression in single cholinergic basal forebrain neurons mark the progression of Alzheimer's disease. *J Neurochem.* 2006; 97:475–487. [PubMed: 16539663]
- Ginsberg SD, Mirnics K. Functional genomic methodologies. *Prog Brain Res.* 2006; 158:15–40. [PubMed: 17027690]
- Ginsberg SD, Mufson EJ, Alldred MJ, Counts SE, Wu J, Nixon RA, Che S. Upregulation of select rab GTPases in cholinergic basal forebrain neurons in mild cognitive impairment and Alzheimer's disease. *J Chem Neuroanat.* 2011b; 42:102–110. [PubMed: 21669283]
- Gould TD, Quiroz JA, Singh J, Zarate CA, Manji HK. Emerging experimental therapeutics for bipolar disorder: insights from the molecular and cellular actions of current mood stabilizers. *Mol Psychiatry.* 2004; 9:734–755. [PubMed: 15136794]
- Gratacos M, Costas J, de Cid R, Bayes M, Gonzalez JR, Baca-Garcia E, de Diego Y, Fernandez-Aranda F, Fernandez-Piqueras J, Guitart M, Martin-Santos R, Martorell L, Menchon JM, Roca M, Saiz-Ruiz J, Sanjuan J, Torrens M, Urretavizcaya M, Valero J, Vilella E, Estivill X, Carracedo A. Identification of new putative susceptibility genes for several psychiatric disorders by association analysis of regulatory and non-synonymous SNPs of 306 genes involved in neurotransmission and neurodevelopment. *Am J Med Genet B Neuropsychiatr Genet.* 2009; 150B:808–816. [PubMed: 19086053]
- Grozeva D, Kirov G, Ivanov D, Jones IR, Jones L, Green EK, St Clair DM, Young AH, Ferrier N, Farmer AE, McGuffin P, Holmans PA, Owen MJ, O'Donovan MC, Craddock N. Rare copy number variants: a point of rarity in genetic risk for bipolar disorder and schizophrenia. *Arch Gen Psychiatry.* 2010; 67:318–327. [PubMed: 20368508]
- Hakak Y, Walker JR, Li C, Wong WH, Davis KL, Buxbaum JD, Haroutunian V, Fienberg AA. Genome-wide expression analysis reveals dysregulation of myelination-related genes in chronic schizophrenia. *Proc Natl Acad Sci U S A.* 2001; 98:4746–4751. [PubMed: 11296301]
- Haroutunian V, Katsel P, Dracheva S, Stewart DG, Davis KL. Variations in oligodendrocyte-related gene expression across multiple cortical regions: implications for the pathophysiology of schizophrenia. *Int J Neuropsychopharmacol.* 2007; 10:565–573. [PubMed: 17291370]
- Hashimoto K, Sawa A, Iyo M. Increased levels of glutamate in brains from patients with mood disorders. *Biol Psychiatry.* 2007; 62:1310–1316. [PubMed: 17574216]

- Hashimoto T, Arion D, Unger T, Maldonado-Aviles JG, Morris HM, Volk DW, Mirnics K, Lewis DA. Alterations in GABA-related transcriptome in the dorsolateral prefrontal cortex of subjects with schizophrenia. *Mol Psychiatry*. 2008; 13:147–161. [PubMed: 17471287]
- Hemby SE, Ginsberg SD, Brunk B, Arnold SE, Trojanowski JQ, Eberwine JH. Gene expression profile for schizophrenia: discrete neuron transcription patterns in the entorhinal cortex. *Arch Gen Psychiatr*. 2002; 59:631–640. [PubMed: 12090816]
- Hemby SE, Trojanowski JQ, Ginsberg SD. Neuron-specific age-related decreases in dopamine receptor subtype mRNAs. *J Comp Neurol*. 2003; 456:176–183. [PubMed: 12509874]
- Humphries C, Mortimer A, Hirsch S, de Belleroche J. NMDA receptor mRNA correlation with antemortem cognitive impairment in schizophrenia. *NeuroReport*. 1996; 7:2051–2055. [PubMed: 8905723]
- Itokawa M, Yamada K, Iwayama-Shigeno Y, Ishitsuka Y, Detera-Wadleigh S, Yoshikawa T. Genetic analysis of a functional GRIN2A promoter (GT)_n repeat in bipolar disorder pedigrees in humans. *Neurosci Lett*. 2003; 345:53–56. [PubMed: 12809987]
- Iwamoto K, Kakiuchi C, Bundo M, Ikeda K, Kato T. Molecular characterization of bipolar disorder by comparing gene expression profiles of postmortem brains of major mental disorders. *Mol Psychiatry*. 2004; 9:406–416. [PubMed: 14743183]
- Javitt DC. Glutamate and schizophrenia: phencyclidine, N-Methyl-d-Aspartate receptors, and dopamine-glutamate Interactions. *Int Rev Neurobiol*. 2007; 78:69–108. [PubMed: 17349858]
- Jurata LW, Bukhman YV, Charles V, Capriglione F, Bullard J, Lemire AL, Mohammed A, Pham Q, Laeng P, Brockman JA, Altar CA. Comparison of microarray-based mRNA profiling technologies for identification of psychiatric disease and drug signatures. *J Neurosci Methods*. 2004; 138:173–188. [PubMed: 15325126]
- Kato T, Kakiuchi C, Iwamoto K. Comprehensive gene expression analysis in bipolar disorder. *Can J Psychiatry*. 2007; 52:763–771. [PubMed: 18186176]
- Kerner B, Jasinska AJ, DeYoung J, Almonte M, Choi OW, Freimer NB. Polymorphisms in the GRIA1 gene region in psychotic bipolar disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2009; 150B:24–32. [PubMed: 18484081]
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005a; 62:593–602. [PubMed: 15939837]
- Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005b; 62:617–627. [PubMed: 15939839]
- Kim AH, Reimers M, Maher B, Williamson V, McMichael O, McClay JL, van den Oord EJ, Riley BP, Kendler KS, Vladimirov VI. MicroRNA expression profiling in the prefrontal cortex of individuals affected with schizophrenia and bipolar disorders. *Schizophr Res*. 2010; 124:183–191. [PubMed: 20675101]
- Kim S, Webster MJ. The stanley neuropathology consortium integrative database: a novel, web-based tool for exploring neuropathological markers in psychiatric disorders and the biological processes associated with abnormalities of those markers. *Neuropsychopharmacology*. 2010; 35:473–482. [PubMed: 19829293]
- Klempman TA, Sequeira A, Canetti L, Lalovic A, Ernst C, Ffrench-Mullen J, Turecki G. Altered expression of genes involved in ATP biosynthesis and GABAergic neurotransmission in the ventral prefrontal cortex of suicides with and without major depression. *Mol Psychiatry*. 2007
- Konradi C, Eaton M, MacDonald ML, Walsh J, Benes FM, Heckers S. Molecular evidence for mitochondrial dysfunction in bipolar disorder. *Arch Gen Psychiatry*. 2004; 61:300–308. [PubMed: 14993118]
- Le-Niculescu H, McFarland MJ, Mamidipalli S, Ogden CA, Kuczenski R, Kurian SM, Salomon DR, Tsuang MT, Nurnberger JI Jr, Niculescu AB. Convergent functional genomics of bipolar disorder: from animal model pharmacogenomics to human genetics and biomarkers. *Neurosci Biobehav Rev*. 2007; 31:897–903. [PubMed: 17614132]
- Lewis DA. Antipsychotic medications and brain volume: do we have cause for concern? *Arch Gen Psychiatry*. 2011; 68:126–127. [PubMed: 21300942]

- Lewis DA, Hashimoto T, Morris HM. Cell and receptor type-specific alterations in markers of GABA neurotransmission in the prefrontal cortex of subjects with schizophrenia. *Neurotox Res.* 2008; 14:237–248. [PubMed: 19073429]
- Li B, Zhang S, Li M, Zhang H, Hertz L, Peng L. Down-regulation of GluK2 kainate receptor expression by chronic treatment with mood-stabilizing anti-convulsants or lithium in cultured astrocytes and brain, but not in neurons. *Neuropharmacology.* 2009; 57:375–385. [PubMed: 19596362]
- Lin M, Pedrosa E, Shah A, Hrabovsky A, Maqbool S, Zheng D, Lachman HM. RNA-Seq of human neurons derived from iPS cells reveals candidate long non-coding RNAs involved in neurogenesis and neuropsychiatric disorders. *PLoS ONE.* 2011; 6:e23356. [PubMed: 21915259]
- Lorenzi C, Delmonte D, Pirovano A, Marino E, Bongiorno F, Catalano M, Colombo C, Bramanti P, Smeraldi E. Searching susceptibility loci for bipolar disorder: a sib pair study on chromosome 12. *Neuropsychobiology.* 2010; 61:10–18. [PubMed: 19923861]
- Lyons-Warren A, Chang JJ, Balkissoon R, Kamiya A, Garant M, Nurnberger J, Scheftner W, Reich T, McMahon F, Kelsoe J, Gershon E, Coryell W, Byerley W, Berrettini W, Depaulo R, McInnis M, Sawa A. Evidence of association between bipolar disorder and Citron on chromosome 12q24. *Mol Psychiatry.* 2005; 10:807–809. [PubMed: 15983625]
- Marsden WN. Stressor-induced NMDAR dysfunction as a unifying hypothesis for the aetiology, pathogenesis and comorbidity of clinical depression. *Med Hypotheses.* 2011; 77:508–528. [PubMed: 21741771]
- Marti SB, Cichon S, Propping P, Nothen M. Metabotropic glutamate receptor 3 (GRM3) gene variation is not associated with schizophrenia or bipolar affective disorder in the German population. *Am J Med Genet.* 2002; 114:46–50. [PubMed: 11840505]
- Martucci L, Wong AH, De Luca V, Likhodi O, Wong GW, King N, Kennedy JL. N-methyl-D-aspartate receptor NR2B subunit gene GRIN2B in schizophrenia and bipolar disorder: Polymorphisms and mRNA levels. *Schizophr Res.* 2006; 84:214–221. [PubMed: 16549338]
- McCullumsmith RE, Kristiansen LV, Beneyto M, Scarr E, Dean B, Meador-Woodruff JH. Decreased NR1, NR2A, and SAP102 transcript expression in the hippocampus in bipolar disorder. *Brain Res.* 2007; 1127:108–118. [PubMed: 17113057]
- McDonald LJ, Griffin ML, Kolodziej ME, Fitzmaurice GM, Weiss RD. The impact of drug use in social networks of patients with substance use and bipolar disorders. *Am J Addict.* 2011; 20:100–105. [PubMed: 21314751]
- Melchitzky DS, Lewis DA. Dendritic-targeting GABA neurons in monkey prefrontal cortex: comparison of somatostatin- and calretinin-immunoreactive axon terminals. *Synapse.* 2008; 62:456–465. [PubMed: 18361442]
- Merikangas KR, Jin R, He JP, Kessler RC, Lee S, Sampson NA, Viana MC, Andrade LH, Hu C, Karam EG, Ladea M, Medina-Mora ME, Ono Y, Posada-Villa J, Sagar R, Wells JE, Zarkov Z. Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Arch Gen Psychiatry.* 2011; 68:241–251. [PubMed: 21383262]
- Mimmack ML, Ryan M, Baba H, Navarro-Ruiz J, Iritani S, Faull RL, McKenna PJ, Jones PB, Arai H, Starkey M, Emson PC, Bahn S. Gene expression analysis in schizophrenia: reproducible up-regulation of several members of the apolipoprotein L family located in a high-susceptibility locus for schizophrenia on chromosome 22. *Proc Natl Acad Sci U S A.* 2002; 99:4680–4685. [PubMed: 11930015]
- Mirnic K, Levitt P, Lewis DA. Critical appraisal of DNA microarrays in psychiatric genomics. *Biol Psychiatry.* 2006; 60:163–176. [PubMed: 16616896]
- Mirnic K, Middleton FA, Marquez A, Lewis DA, Levitt P. Molecular characterization of schizophrenia viewed by microarray analysis of gene expression in prefrontal cortex. *Neuron.* 2000; 28:53–67. [PubMed: 11086983]
- Mueller HT, Meador-Woodruff JH. NR3A NMDA receptor subunit mRNA expression in schizophrenia, depression and bipolar disorder. *Schizophr Res.* 2004; 71:361–370. [PubMed: 15474907]
- Mundo E, Tharmalingham S, Neves-Pereira M, Dalton EJ, Macciardi F, Parikh SV, Bolonna A, Kerwin RW, Arranz MJ, Makoff AJ, Kennedy JL. Evidence that the N-methyl-D-aspartate subunit

- 1 receptor gene (GRIN1) confers susceptibility to bipolar disorder. *Mol Psychiatry*. 2003; 8:241–245. [PubMed: 12610658]
- Nolan T, Hands RE, Bustin SA. Quantification of mRNA using real-time RT-PCR. *Nat Protoc*. 2006; 1:1559–1582. [PubMed: 17406449]
- O'Connor JA, Hemby SE. Elevated GRIA1 mRNA expression in Layer II/III and V pyramidal cells of the DLPFC in schizophrenia. *Schizophr Res*. 2007a; 97:277–288. [PubMed: 17942280]
- O'Connor JA, Muly EC, Arnold SE, Hemby SE. AMPA receptor subunit and splice variant expression in the DLPFC of schizophrenic subjects and rhesus monkeys chronically administered antipsychotic drugs. *Schizophr Res*. 2007b; 90:28–40. [PubMed: 17141476]
- Ogden CA, Rich ME, Schork NJ, Paulus MP, Geyer MA, Lohr JB, Kuczenski R, Niculescu AB. Candidate genes, pathways and mechanisms for bipolar (manic-depressive) and related disorders: an expanded convergent functional genomics approach. *Mol Psychiatry*. 2004; 9:1007–1029. [PubMed: 15314610]
- Okumura T, Kishi T, Okochi T, Ikeda M, Kitajima T, Yamanouchi Y, Kinoshita Y, Kawashima K, Tsunoka T, Inada T, Ozaki N, Iwata N. Genetic association analysis of functional polymorphisms in neuronal nitric oxide synthase 1 gene (NOS1) and mood disorders and fluvoxamine response in major depressive disorder in the Japanese population. *Neuropsychobiology*. 2010; 61:57–63. [PubMed: 20016223]
- Oquendo MA, Currier D, Liu SM, Hasin DS, Grant BF, Blanco C. Increased risk for suicidal behavior in comorbid bipolar disorder and alcohol use disorders: results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *J Clin Psychiatry*. 2010; 71:902–909. [PubMed: 20667292]
- Pennington K, Beasley CL, Dicker P, Fagan A, English J, Pariante CM, Wait R, Dunn MJ, Cotter DR. Prominent synaptic and metabolic abnormalities revealed by proteomic analysis of the dorsolateral prefrontal cortex in schizophrenia and bipolar disorder. *Mol Psychiatry*. 2008; 13:1102–1117. [PubMed: 17938637]
- Perlis RH, Smoller JW, Ferreira MA, McQuillin A, Bass N, Lawrence J, Sachs GS, Nimgaonkar V, Scolnick EM, Gurling H, Sklar P, Purcell S. A genomewide association study of response to lithium for prevention of recurrence in bipolar disorder. *Am J Psychiatry*. 2009; 166:718–725. [PubMed: 19448189]
- Pickard BS, Knight HM, Hamilton RS, Soares DC, Walker R, Boyd JK, Machell J, Maclean A, McGhee KA, Condie A, Porteous DJ, St Clair D, Davis I, Blackwood DH, Muir WJ. A common variant in the 3'UTR of the GRIK4 glutamate receptor gene affects transcript abundance and protects against bipolar disorder. *Proc Natl Acad Sci USA*. 2008; 105:14940–14945. [PubMed: 18824690]
- Pickard BS, Malloy MP, Christoforou A, Thomson PA, Evans KL, Morris SW, Hampson M, Porteous DJ, Blackwood DH, Muir WJ. Cytogenetic and genetic evidence supports a role for the kainate-type glutamate receptor gene, GRIK4, in schizophrenia and bipolar disorder. *Mol Psychiatry*. 2006; 11:847–857. [PubMed: 16819533]
- Piletz JE, Zhang XT, Ranade R, Liu C. Database of genetic studies of bipolar disorder. *Psychiatr Genet*. 2011; 21:57–68. [PubMed: 21085053]
- Prata D, Breen G, Osborne S, Munro J, St Clair D, Collier D. Association of DAO and G72(DAOA)/G30 genes with bipolar affective disorder. *Am J Med Genet B Neuropsychiatr Genet*. 2008; 147B:914–917. [PubMed: 18165970]
- Rao JS, Harry GJ, Rapoport SI, Kim HW. Increased excitotoxicity and neuroinflammatory markers in postmortem frontal cortex from bipolar disorder patients. *Mol Psychiatry*. 2010; 15:384–392. [PubMed: 19488045]
- Regier DA, Narrow WE, Rae DS, Manderscheid RW, Locke BZ, Goodwin FK. The de facto US mental and addictive disorders service system. Epidemiologic catchment area prospective 1-year prevalence rates of disorders and services. *Arch Gen Psychiatry*. 1993; 50:85–94. [PubMed: 8427558]
- Reif A, Strobel A, Jacob CP, Herterich S, Freitag CM, Topner T, Mossner R, Fritzen S, Schmitt A, Lesch KP. A NOS-III haplotype that includes functional polymorphisms is associated with bipolar disorder. *Int J Neuropsychopharmacol*. 2006; 9:13–20. [PubMed: 15967063]

- Ruberto G, Vassos E, Lewis CM, Tatarelli R, Girardi P, Collier D, Frangou S. The cognitive impact of the ANK3 risk variant for bipolar disorder: initial evidence of selectivity to signal detection during sustained attention. *PLoS ONE*. 2011; 6:e16671. [PubMed: 21304963]
- Ryan MM, Lockstone HE, Huffaker SJ, Wayland MT, Webster MJ, Bahn S. Gene expression analysis of bipolar disorder reveals downregulation of the ubiquitin cycle and alterations in synaptic genes. *Mol Psychiatry*. 2006; 11:965–978. [PubMed: 16894394]
- Saus E, Brunet A, Armengol L, Alonso P, Crespo JM, Fernandez-Aranda F, Guitart M, Martin-Santos R, Menchon JM, Navines R, Soria V, Torrens M, Urretavizcaya M, Valles V, Gratacos M, Estivill X. Comprehensive copy number variant (CNV) analysis of neuronal pathways genes in psychiatric disorders identifies rare variants within patients. *J Psychiatr Res*. 2010; 44:971–978. [PubMed: 20398908]
- Scott LJ, Muglia P, Kong XQ, Guan W, Flickinger M, Upmanyu R, Tozzi F, Li JZ, Burmeister M, Absher D, Thompson RC, Francks C, Meng F, Antoniadis A, Southwick AM, Schatzberg AF, Bunney WE, Barchas JD, Jones EG, Day R, Matthews K, McGuffin P, Strauss JS, Kennedy JL, Middleton L, Roses AD, Watson SJ, Vincent JB, Myers RM, Farmer AE, Akil H, Burns DK, Boehnke M. Genome-wide association and meta-analysis of bipolar disorder in individuals of European ancestry. *Proc Natl Acad Sci USA*. 2009; 106:7501–7506. [PubMed: 19416921]
- Sequeira A, Klempan T, Canetti L, French-Mullen J, Benkelfat C, Rouleau GA, Turecki G. Patterns of gene expression in the limbic system of suicides with and without major depression. *Mol Psychiatry*. 2007; 12:640–655. [PubMed: 17353912]
- Sequeira A, Turecki G. Genome wide gene expression studies in mood disorders. *Omics*. 2006; 10:444–454. [PubMed: 17233556]
- Shi J, Badner JA, Hattori E, Potash JB, Willour VL, McMahon FJ, Gershon ES, Liu C. Neurotransmission and bipolar disorder: a systematic family-based association study. *Am J Med Genet B Neuropsychiatr Genet*. 2008; 147B:1270–1277. [PubMed: 18444252]
- Sibille E, Arango V, Galfalvy HC, Pavlidis P, Erraji-Benchekroun L, Ellis SP, Mann JJ. Gene expression profiling of depression and suicide in human prefrontal cortex. *Neuropsychopharmacology*. 2004; 29:351–361. [PubMed: 14603265]
- Sklar P, Gabriel SB, McInnis MG, Bennett P, Lim YM, Tsan G, Schaffner S, Kirov G, Jones I, Owen M, Craddock N, DePaulo JR, Lander ES. Family-based association study of 76 candidate genes in bipolar disorder: BDNF is a potential risk locus. Brain-derived neurotrophic factor. *Mol Psychiatry*. 2002; 7:579–593. [PubMed: 12140781]
- Sklar P, Smoller JW, Fan J, Ferreira MA, Perlis RH, Chambert K, Nimgaonkar VL, McQueen MB, Faraone SV, Kirby A, de Bakker PI, Ogdie MN, Thase ME, Sachs GS, Todd-Brown K, Gabriel SB, Sougnez C, Gates C, Blumenstiel B, Defelice M, Ardlie KG, Franklin J, Muir WJ, McGhee KA, MacIntyre DJ, McLean A, VanBeck M, McQuillin A, Bass NJ, Robinson M, Lawrence J, Anjorin A, Curtis D, Scolnick EM, Daly MJ, Blackwood DH, Gurling HM, Purcell SM. Whole-genome association study of bipolar disorder. *Mol Psychiatry*. 2008; 13:558–569. [PubMed: 18317468]
- Takata A, Kim SH, Ozaki N, Iwata N, Kunugi H, Inada T, Ujike H, Nakamura K, Mori N, Ahn YM, Joo EJ, Song JY, Kanba S, Yoshikawa T, Kim YS, Kato T. Association of ANK3 with bipolar disorder confirmed in East Asia. *Am J Med Genet B Neuropsychiatr Genet*. 2011; 156:312–315. [PubMed: 21438140]
- Tang B, Thornton-Wells T, Askland KD. Comparative linkage meta-analysis reveals regionally-distinct, disparate genetic architectures: application to bipolar disorder and schizophrenia. *PLoS ONE*. 2011; 6:e19073. [PubMed: 21559500]
- Tkachev D, Mimmack ML, Ryan MM, Wayland M, Freeman T, Jones PB, Starkey M, Webster MJ, Yolken RH, Bahn S. Oligodendrocyte dysfunction in schizophrenia and bipolar disorder. *Lancet*. 2003; 362:798–805. [PubMed: 13678875]
- Uezato A, Meador-Woodruff JH, McCullumsmith RE. Vesicular glutamate transporter mRNA expression in the medial temporal lobe in major depressive disorder, bipolar disorder, and schizophrenia. *Bipolar Disord*. 2009; 11:711–725. [PubMed: 19839996]
- Usher J, Leucht S, Falkai P, Scherk H. Correlation between amygdala volume and age in bipolar disorder - a systematic review and meta-analysis of structural MRI studies. *Psychiatry Res*. 2010; 182:1–8. [PubMed: 20226638]

- Vieta E, Morralla C. Prevalence of mixed mania using 3 definitions. *J Affect Disord.* 2010
- Woo TU, Shrestha K, Amstrong C, Minns MM, Walsh JP, Benes FM. Differential alterations of kainate receptor subunits in inhibitory interneurons in the anterior cingulate cortex in schizophrenia and bipolar disorder. *Schizophr Res.* 2007; 96:46–61. [PubMed: 17698324]
- Woo TU, Walsh JP, Benes FM. Density of glutamic acid decarboxylase 67 messenger RNA-containing neurons that express the N-methyl-D-aspartate receptor subunit NR2A in the anterior cingulate cortex in schizophrenia and bipolar disorder. *Arch Gen Psychiatry.* 2004; 61:649–657. [PubMed: 15237077]
- Yosifova A, Mushiroda T, Stoianov D, Vazharova R, Dimova I, Karachanak S, Zaharieva I, Milanova V, Madjirova N, Gerdjikov I, Tolev T, Velkova S, Kirov G, Owen MJ, O'Donovan MC, Toncheva D, Nakamura Y. Case-control association study of 65 candidate genes revealed a possible association of a SNP of HTR5A to be a factor susceptible to bipolar disease in Bulgarian population. *J Affect Disord.* 2009; 117:87–97. [PubMed: 19328558]
- Zhao Q, Che R, Zhang Z, Wang P, Li J, Li Y, Huang K, Tang W, Feng G, Lindpaintner K, He L, Shi Y. Positive association between GRIN2B gene and bipolar disorder in the Chinese Han Population. *Psychiatry Res.* 2011; 185:290–292. [PubMed: 20537720]

Research Highlights

GWAS studies identified glutamate receptor candidate genes for bipolar disorder.

Alterations in glutamate receptor genes were reviewed in bipolar disorder.

Expression profiling of specific neurons and intrinsic circuits is emphasized.

Figure 1.

Representative expression profiling of entorhinal stellate neurons (top panel) and regional entorhinal cortex (lower panel) obtained from normal control human postmortem tissue using the Incyte UniGEM V platform. Several genes expressed in stellate cells were also expressed in entorhinal cortex at differing abundance levels (e.g., compare B2, C5, and B10 between the two panels). mRNAs detectable in stellate cells and not in the regional entorhinal cortex dissection was exemplified in G7 and E4. In contrast, several genes present in the entorhinal cortex dissection were below the limit of detection in stellate cells, likely due to the heterogeneity of admixed cell types within the entorhinal cortex (e.g., compared B5, D4, and H10 between panels). Color coded expression levels depict red as high expression and blue as low expression.