Clinical, Dermatoscopic, and Microscopic Findings of Infestation With *Sarcoptes scabiei* var *hominis*

JOHN R. GRIFFIN, MD, AND CATHERINE C. NEWMAN, MD

Department of Dermatology, Mayo Clinic, Rochester, MN

Infestation of the epidermis with the mite *Sarcoptes scabiei* var *hominis* causes human scabies. This condition affects humans worldwide without respect to age, sex, race, or socioeconomic status. Epidemics are known to occur in overcrowded situations and in long-term care facilities. Transmission of the mite is by direct contact or via fomites. The primary symptom of infestation is intense pruritus with nocturnal exacerbation. This is due to type IV immune response to antigens deposited within the epidermis; therefore, the initial manifestations may not be noted until several weeks later. Immunocompetent patients typically have a low mite burden, and definitive diagnosis by

demonstration of the mite may be challenging. We present clinical, dermatoscopic, and microscopic findings of infestation to serve as a reminder of the various clues that may be sought to confirm this important diagnosis. Of particular interest is the usefulness of the noninvasive technique, dermatoscopy. With a dermatoscope (essentially a magnifying glass with indwelling light source), one can visualize the mite in vivo and note the characteristic “hang-glider” appearance of the mite and its “jet-plane” trail of a burrow to establish the diagnosis.

© 2011 Mayo Foundation for Medical Education and Research