Published in final edited form as:

Neurosci Lett. 2012 January 24; 507(2): 137–142. doi:10.1016/j.neulet.2011.12.005.

Studies of Protein Aggregation in A53T α -Synuclein Transgenic, Tg2576 Transgenic, and P246L Presenilin-1 Knock-in Cross Bred Mice

Kristel L. Emmer, Jason P. Covy, and Benoit I. Giasson

Department of Pharmacology, University of Pennsylvania, Philadelphia, PA

Abstract

Synucleinopathies are a group of neurodegenerative disorders, including Parkinson disease, associated with neuronal amyloid inclusions comprised of the presynaptic protein α -synuclein (α -syn); however the biological events that initiate and lead to the formation of these inclusions are still poorly understood. There is mounting evidence that intracellular α -syn aggregation may proceed via a seeding mechanism and could spread between neurons through a prion-like mechanism that may involve other amyloidogenic proteins. Several lines of evidence suggest that A β peptides and/or extracellular A β deposits may directly or indirectly promote intracellular α -syn aggregation. To assess the effects of A β peptides and extracellular A β deposits on α -syn aggregate formation, transgenic mice (line M83) expressing A53T human α -syn that are sensitive to developing α -syn pathological inclusions were cross bred to Tg2576 transgenic mice that generated elevated levels of A β peptides and develop abundant A β plaques. In addition these mice were bred to mice with the P264L presenilin-1 knock-in mutation that further promotes A β plaque formation. These mice demonstrated the expected formation of A β plaques; however despite the accumulation of hyperphosphorylated α -syn dystrophic neurites within or surrounding A β plaques, no additional α -syn pathologies were observed. These studies show that A β amyloid deposits can cause the local aggregation of α -syn, but these did not lead to more extensive α -syn pathology.

Keywords

Aggregation; amyloid; pathology; Parkinson disease; α -synuclein; transgenic

Introduction

Synucleinopathies are a group of neurodegenerative diseases associated with neuronal, and in some cases oligodendritic, amyloid inclusions comprised of the presynaptic protein α -synuclein (α -syn) [11,20,47,51]. Parkinson disease (PD), the most common known synucleinopathy, is characterized by the loss of dopaminergic neurons in the substantia nigra pars compacta and the formation of α -syn inclusions, known as Lewy bodies (LBs) and Lewy neurites (LNs), in some of the remaining dopaminergic neurons [12,20,51]. PD is a

© 2011 Elsevier Ireland Ltd. All rights reserved.

Address correspondence to: Dr. Benoit I. Giasson, Department of Pharmacology, University of Pennsylvania School of Medicine, 3620 Hamilton Walk, 125 John Morgan Building, Philadelphia, PA 19104-6084; Tel: 215-573-6012; Fax: 215-573-2236; giassonb@mail.med.upenn.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

progressive movement disorder [15,45], however it is associated with a range of nonmotor symptoms [4,40] and many other affected neuronal populations outside of the substantia nigra contribute to the progression of disease [2,7,11,12].

A β senile deposits or plaques, which consist primarily of aggregated A β peptides that vary between 39-43 amino acids in length, are one of the hallmark lesions of Alzheimer disease [19,43,48,53]. A β peptides are secreted from cells following cleavage of the trans-membrane A β precursor protein (β APP) at the A β N-terminal (β -secretase cleavage) and C-terminal (γ -secretase cleavage) [43,53]. A β deposits can be observed in several other neurodegenerative diseases, including dementia with Lewy bodies (DLB) and LB variant of Alzheimer disease (LBVAD), where concomitant α -syn intraneuronal inclusions are present [11,20,38,51].

The most direct and compelling evidence for a fundamental role of α -syn in the pathogenesis of synucleinopathies is the causal relationship between genetic mutations and disease [6,8,34,51]. The missense mutation (c.G209A) in the α -syn gene (*SNCA*) resulting in the amino acid substitution A53T was first identified in a large Italian family (Contursi) and three small Greek families with autosomal dominant PD [41], and this mutation enhances the propensity of α -syn to form amyloid [5,18]. However, the biological events that initiate and lead to the formation of α -syn inclusions are still poorly understood. Several lines of evidence suggest that extracellular A β deposits may directly or indirectly promote intracellular α -syn aggregation. Besides the frequent co-occurrence of α -syn inclusions and A β deposits in the brains of patients with PD, DLB or LBVAD [11,20,38,51], α -syn inclusions are commonly observed in patients with familial Alzheimer disease where genetic defects in the *APP*, *presenilin-1* (PS1) or *presenilin-2* (PS2) genes affect biological pathways that promote the formation of A β aggregates [23,27,35,46,55]. PS1 and PS2 are enzymatic components of the transmembrane γ -secretase complex [29,39] that cleaves β APP. Over 100 mutations in the *PS1* and *PS2* genes have been identified in familial Alzheimer disease and these mutations result in increased production of the longer A β 1-42(43) species [13,43,53]. A β 1-42(43) peptides have been shown to have a greater propensity to form amyloidogenic fibrils compared to the shorter A β 1-40 peptide [25]. In addition, A β 1-42 is deposited early and selectively in senile plaques [24], but the nature and mechanism of A β toxicity are still debated [1,3,30,42,53].

To investigate the possibility that A β peptides or amyloid plaques may promote/initiate the aggregation of α -syn, α -syn transgenic mice (line M83) expressing A53T human α -syn that are sensitive to developing α -syn pathological inclusions [16] were cross bred to the previously characterized transgenic mice that overexpress human β APP (695 amino acid splice form) with the “Swedish” double mutation K670M/N671L (line Tg 2576) that develop abundant age-dependent A β plaques [22,28]. In addition these mice were bred to mice with the P264L PSI knock-in mutation that increase A β 1-42 production and further promote A β plaque formation [10,44].

Material and Methods

Antibodies

pSer129 is a mouse monoclonal antibody specific to α -syn phosphorylated at S129 [52]. Syn505 and Syn506 are conformational anti- α -syn mouse monoclonal antibodies that preferentially detect α -syn in pathological inclusions [50]. Syn 211 is a mouse monoclonal antibody specific for human α -syn [17]. Rabbit anti-A β antibody was purchased from Cell Signaling Technologies (Danver, MA). The anti- A β mouse monoclonal antibody 6E10 was purchased from Covance (Princeton, NJ). Karen, a goat polyclonal anti-N-terminal APP antibody and NAB228, a monoclonal antibody raised against A β 1-11 synthetic peptide [32],

were generous gifts from Dr. Virginia Lee. Affinity purified mouse anti-actin (clone C4) monoclonal antibody was purchased from Millipore (Billerica, MA).

Transgenic Mice

The previously described M83 A53T human α -syn [16] and Tg2576 β APP [22] transgenic mouse lines were used in these studies. In addition, P264L PS1 knock-in mice were previously described [10,44]. For genotyping, genomic DNA samples were isolated from mouse tails with proteinase K digestion followed by purification with the Wizard® SV Genomic DNA Purification System (Promega, Madison, WI). The α -syn and β APP transgenes were screened by Southern blot analysis with 32 P-labeled oligonucleotide-primed specific DNA probes respectively, as previously described [16,22]. The presence of the P264L PSI mutation that results in a larger DNA product was screened by PCR using the forward primer GCTGGAGCAATGCTGTGTTA and the reverse primer GAGATGGCTTACGGGTTGAG [10].

Western blot analysis

Brain tissues were harvested and lysed in 3% SDS/50 mM Tris, pH 6.8 by sonication and heating to 100°C for 10 min. Total protein extracts were quantified using the BCA assay using bovine serum albumin as the standard. Equal amounts of protein extracts (5 μ g) were separated by electrophoresis onto 15% polyacrylamide gels for α -syn analysis or 8% polyacrylamide gels for β APP analysis. Immunoblotting was performed as previously described [52].

Immunohistochemical Analysis

Mice were sacrificed with CO₂ euthanization as approved by the University of Pennsylvania Institutional Animal Care and Use Committee and perfused with PBS/heparin, followed by perfusion with either 70% ethanol/150mM NaCl or PBS buffered formalin. The brain and spinal cord were then removed, fixed and processed for paraffin infiltration as previously described [9]. Embedded tissue paraffin blocks were cut into 7 μ m sections. Immunostaining of the sections was performed as previously described [9]. Immunocomplexes were visualized with the chromogen 3,3'-diaminobenzidine and the sections were counterstained with hematoxylin.

Double-Labeling Immunofluorescence Analysis of Mouse Brain Tissue

Paraffin-embedded tissue sections were hydrated and stained with primary antibodies as previously described [52]. Sections were incubated with goat anti-mouse secondary conjugated to Alexa 594 and goat anti-rabbit secondary conjugated to Alexa 488 (Invitrogen, Eugene, OR) and counterstained with 4',6-diamidino-2-phenylindole (DAPI). The sections were coverslipped with Fluoromount-G (SouthernBiotech, Birmingham, AL) and visualized using an Olympus BX51 microscope mounted with a DP71 Olympus digital camera to capture images.

Results and Discussion

The A53T human α -syn transgenic mice (line M83) are prone to develop an age-dependent lethal motor phenotype associated with the formation of neuronal α -syn inclusions in the brain and spinal cord, however heterozygous mice from this transgenic line do not develop these phenotypic and cellular pathological changes until at least 22 months [16]. Consequently, this line provides a unique opportunity to study the effects of A β expression on the initiation and promotion of α -syn aggregation. To this end, A53T human α -syn transgenic mice (line M83) were crossbred with Tg2576 β APP transgenic mice to generate mice that are heterozygous for each gene. In addition, these mice were further bred with

knock-in mice carrying the P264L PSI mutation (Fig. 1A). The expression of human α -syn and human β APP for the respective genotypes was further confirmed by Western blot analysis (Fig. 1 B and C). The Tg 2576 transgenic mice start to develop extracellular amyloid deposits around 10-12 month and the presence of these inclusion increase rapidly thereafter [10,22,28]. The analysis of M83/Tg2576 bigenic mice from 7-22 months (N = 22) demonstrated the predicted profile of A β deposits with some initial detection of plaques being at 11-12 months and increasing abundance with age (Fig. 2). Further histological analysis revealed the presence of abundant α -syn dystrophic neurites within and surrounding many A β plaques (Fig. 2C-F) and many of these neurites accumulated α -syn phosphorylated at S129 (Fig. 2F), a modification that is persistent in pathological inclusions [14,52]. However, no additional α -syn pathologies was observed beyond these changes.

α -Syn/Tg2576 bigenic mice were further crossbred mice with the mice carrying the P264L PSI knock-in mutation. The P264L PSI knock-in mutation increases the amount A β 1-42 production, promoting A β plaque formation [10,44]. These mice were studied between 5-14 months of age (N =18). Similar to previously reported Tg2576/ P264L PSI mice [10,44], this triple crossed line developed amyloid plaques earlier than the α -syn/Tg2576 bigenic mice, starting around 6 months of age with an increasing abundance of plaques at older time points (Fig. 3). Many of these plaques displayed abundant α -syn dystrophic neurites that were hyperphosphorylated at S129 (Fig. 3 C and D), but further α -syn pathology was not observed.

Several lines of evidence suggest that extracellular A β deposits may directly or indirectly promote intracellular α -syn aggregation. Besides, the frequent co-occurrence of A β deposits and α -syn inclusions in the brains of patients with various synucleinopathies such as PD, DLB or LBVAD [11,20,38,51], α -syn inclusions are frequently observed in brains from patients with familial Alzheimer disease where genetic defects in the *APP*, *presenilin-1* (PS1) or *presenilin-2* (PS2) genes affect biological pathways that promote A β aggregation [23,27,35,46,55]. In addition, in vitro studies showed that A β peptides have the ability to promote polymerization of α -syn [26,36] and cell culture studies indicated that the addition of A β 1-42 can also promote the formation of α -syn inclusions [36]. Moreover, in studies of bigenic mice over-expressing β APP (line J9) and wild-type α -syn (line D), it was reported that the presence of A β (or perhaps other APP products) significantly enhances the formation of filamentous α -syn aggregates [36]. J9 transgenic β APP mice express human APP that is modified with both the “Swedish” (K670N/M271I) mutation and the “Indiana” (V717F) mutation resulting in an increase in total A β production as well as the amount of the longer forms of the A β 1-42/43 peptides with amyloid deposit starting at 5-7 months [37]. Similarly, cross breeding Tg2756 mice on a P264L PSI background has been shown to increase A β 1-42 production [10,44], which is responsible for the earlier formation of amyloid plaques.

Previous analysis of Tg2576 β APP transgenic mice showed the accumulation of α -syn dystrophic neurites within amyloid plaques [49,54]. In addition, similar α -syn dystrophic neurites were reported with Tg2576 β APP transgenic mice crossed to transgenic L286V mutant PSI mice [31]. The presence of the L286V PSI mutation accelerated the formation of amyloid plaques and the presence of α -syn dystrophic neurites within the plaques.

Given mounting evidence that α -syn aggregation may proceed intracellular via a seeding mechanism, and even between neurons through a prion-like spreading disease mechanism [21,33], it was of interest to assess whether α -syn dystrophic neurites resulting from amyloid plaques, and perhaps other changes induced by A β expression, could lead to further α -syn pathology in M83 A53T human α -syn transgenic mice that are prone to develop α -syn inclusions. Tg2576/M83 bigenic mice on a wild-type or P264L PSI background developed

abundant age-dependent α -syn dystrophic pathologies within and surrounding A β amyloid plaques, but it is surprising that additional α -syn pathologies did not present beyond these changes. It is possible that the aggregated forms of α -syn within these neurites are not compatible with the seeding of α -syn amyloid formation. However, it is also possible that additional cellular insults are required for the spreading of α -syn to occur in vivo. Nevertheless, the relationship between the formation of α -syn pathological inclusions and alteration in A β aggregation and/or processing remain enigmatic and further studies will be required to understand their possible association.

Acknowledgments

We would like to thank Michael Heenan for technical assistance.

Role of the funding source

This study was supported by the National Institute of Neurological Disorders and Stroke (NS053488). The funding sponsor has not been involved in the design, execution of the study, in the manuscript drafting or in the decision to submit the paper for publication.

Abbreviations used

βAPP	A β precursor protein
DLB	dementia with Lewy bodies
LBVAD	LB variant of Alzheimer disease
LB	Lewy body
LN	Lewy neurite
α-syn	α -synuclein

References

1. Atwood CS, Obrenovich ME, Liu T, Chan H, Perry G, Smith MA, Martins RN. Amyloid-beta: a chameleon walking in two worlds: a review of the trophic and toxic properties of amyloid-beta. *Brain Res. Brain Res. Rev.* 2003; 43:1–16. [PubMed: 14499458]
2. Braak H, Bohl JR, Muller CM, Rub U, de Vos RA, Del TK. Stanley Fahn Lecture 2005: The staging procedure for the inclusion body pathology associated with sporadic Parkinson's disease reconsidered. *Mov Disord.* 2006; 21:2042–2051. [PubMed: 17078043]
3. Canevari L, Abramov AY, Duchen MR. Toxicity of amyloid beta peptide: tales of calcium, mitochondria, and oxidative stress. *Neurochem. Res.* 2004; 29:637–650. [PubMed: 15038611]
4. Chaudhuri KR, Healy DG, Schapira AH. Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol.* 2006; 5:235–245. [PubMed: 16488379]
5. Conway KA, Harper JD, Lansbury PT. Accelerated in vitro fibril formation by a mutant alpha-synuclein linked to early-onset Parkinson disease. *Nat. Med.* 1998; 4:1318–1320. [PubMed: 9809558]
6. Cookson MR. The biochemistry of Parkinson's disease. *Annu. Rev. Biochem.* 2005; 74:29–52. [PubMed: 15952880]
7. Cronford ME, Chang L, Miller BL. The neuropathology of Parkinsonism: an overview. *Brain Cogn.* 1995; 28:321–341. [PubMed: 8546858]
8. Dawson TM, Ko HS, Dawson VL. Genetic animal models of Parkinson's disease. *Neuron.* 2010; 66:646–661. [PubMed: 20547124]
9. Duda JE, Giasson BI, Gur TL, Montine TJ, Robertson D, Biaggioni I, Hurtig HI, Stern MB, Gollomp SM, Grossman M, Lee VMY, Trojanowski JQ. Immunohistochemical and biochemical

- studies demonstrate a distinct profile of alpha-synuclein permutations in multiple system atrophy. *J. Neuropathol. Exp. Neurol.* 2000; 59:830–841. [PubMed: 11005264]
10. Flood DG, Reaume AG, Dorfman KS, Lin YG, Lang DM, Trusko SP, Savage MJ, Annaert WG, De Strooper B, Siman R, Scott RW. FAD mutant PS-1 gene-targeted mice: increased A beta 42 and A beta deposition without APP overproduction. *Neurobiol. Aging.* 2002; 23:335–348. [PubMed: 11959395]
 11. Forman MS, Lee VM, Trojanowski JQ. Nosology of Parkinson's disease: looking for the way out of a quackmire. *Neuron.* 2005; 47:479–482. [PubMed: 16102530]
 12. Forno LS. Neuropathology of Parkinson's disease. *J. Neuropathol. Exp. Neurol.* 1996; 55:259–272. [PubMed: 8786384]
 13. Fraser PE, Yang DS, Yu G, Levesque L, Nishimura M, Arawaka S, Serpell LC, Rogaeva E, George-Hyslop P. Presenilin structure, function and role in Alzheimer disease. *Biochim. Biophys. Acta.* 2000; 1502:1–15. [PubMed: 10899427]
 14. Fujiwara H, Hasegawa M, Dohmae N, Kawashima A, Masliah E, Goldberg MS, Shen J, Takio K, Iwatsubo T. α -synuclein is phosphorylated in synucleinopathy lesions. *Nat. Cell Biol.* 2002; 4:160–164. [PubMed: 11813001]
 15. Gelb DJ, Oliver E, Gilman S. Diagnostic criteria for Parkinson disease. *Arch. Neurol.* 1999; 56:33–39. [PubMed: 9923759]
 16. Giasson BI, Duda JE, Quinn SM, Zhang B, Trojanowski JQ, Lee VMY. Neuronal α -synucleinopathy with severe movement disorder in mice expressing A53T human α -synuclein. *Neuron.* 2002; 34:521–533. [PubMed: 12062037]
 17. Giasson BI, Jakes R, Goedert M, Duda JE, Leight S, Trojanowski JQ, Lee VMY. A panel of epitope-specific antibodies detects protein domains distributed throughout human alpha-synuclein in Lewy bodies of Parkinson's disease. *J. Neurosci. Res.* 2000; 59:528–533. [PubMed: 10679792]
 18. Giasson BI, Uryu K, Trojanowski JQ, Lee VMY. Mutant and wild type human alpha-synucleins assemble into elongated filaments with distinct morphologies in vitro. *J. Biol. Chem.* 1999; 274:7619–7622. [PubMed: 10075647]
 19. Glenner GG, Wong CW. Alzheimer's disease: initial report of the purification and characterization of a novel cerebrovascular amyloid protein. *Biochem. Biophys. Res. Commun.* 1984; 120:885–890. [PubMed: 6375662]
 20. Goedert M. Alpha-synuclein and neurodegenerative diseases. *Nat. Rev. Neurosci.* 2001; 2:492–501. [PubMed: 11433374]
 21. Goedert M, Clavaguera F, Tolnay M. The propagation of prion-like protein inclusions in neurodegenerative diseases. *Trends Neurosci.* 2010; 33:317–325. [PubMed: 20493564]
 22. Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, Yang F, Cole G. Correlative memory deficits, A beta elevation, and amyloid plaques in transgenic mice. *Science.* 1996; 274:99–102. [PubMed: 8810256]
 23. Ishikawa A, Piao YS, Miyashita A, Kuwano R, Onodera O, Ohtake H, Suzuki M, Nishizawa M, Takahashi H. A mutant PSEN1 causes dementia with Lewy bodies and variant Alzheimer's disease. *Ann. Neurol.* 2005; 57:429–434. [PubMed: 15732120]
 24. Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. Visualization of A beta 42(43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42(43). *Neuron.* 1994; 13:45–53. [PubMed: 8043280]
 25. Jarrett JT, Berger EP, Lansbury PT Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry.* 1993; 32:4693–4697. [PubMed: 8490014]
 26. Jensen PH, Hojrup P, Hager H, Nielsen MS, Jacobsen L, Olesen OF, Gliemann J, Jakes R. Binding of A β to α - and β -synucleins: identification of segments in α -synuclein/NAC precursor that binds A β and NAC. *Biochem. J.* 1997; 323(Pt 2):539–546. [PubMed: 9163350]
 27. Jimenez-Escrig A, Rabano A, Guerrero C, Simon J, Barquero MS, Guell I, Ginestal RC, Montero T, Orensanz L. New V272A presenilin 1 mutation with very early onset subcortical dementia and parkinsonism. *Eur. J. Neurol.* 2004; 11:663–669. [PubMed: 15469450]

28. Kawarabayashi T, Younkin LH, Saido TC, Shoji M, Ashe KH, Younkin SG. Age-dependent changes in brain, CSF, and plasma amyloid (beta) protein in the Tg2576 transgenic mouse model of Alzheimer's disease. *J. Neurosci.* 2001; 21:372–381. [PubMed: 11160418]
29. Kimberly WT, Wolfe MS. Identity and function of gamma-secretase. *J. Neurosci. Res.* 2003; 74:353–360. [PubMed: 14598311]
30. Koo EH, Kopan R. Potential role of presenilin-regulated signaling pathways in sporadic neurodegeneration. *Nat. Med.* 2004; 10(Suppl):S26–S33. [PubMed: 15272268]
31. Kurata T, Kawarabayashi T, Murakami T, Miyazaki K, Morimoto N, Ohta Y, Takehisa Y, Nagai M, Ikeda M, Matsubara E, Westaway D, Hyslop PS, Harigaya Y, Kamiya T, Shoji M, Abe K. Enhanced accumulation of phosphorylated alpha-synuclein in double transgenic mice expressing mutant beta-amyloid precursor protein and presenilin-1. *J. Neurosci. Res.* 2007; 85:2246–2252. [PubMed: 17526016]
32. Lee EB, Skovronsky DM, Abtahian F, Doms RW, Lee VM. Secretion and intracellular generation of truncated Abeta in beta-site amyloid-beta precursor protein-cleaving enzyme expressing human neurons. *J. Biol. Chem.* 2003; 278:4458–4466. [PubMed: 12480937]
33. Lee SJ, Desplats P, Sigurdson C, Tsigelny I, Masliah E. Cell-to-cell transmission of non-prion protein aggregates. *Nat. Rev. Neurol.* 2010; 6:702–706. [PubMed: 21045796]
34. Lesage S, Brice A. Parkinson's disease: from monogenic forms to genetic susceptibility factors. *Hum. Mol. Genet.* 2009; 18:R48–R59. [PubMed: 19297401]
35. Lippa CF, Fujiwara H, Mann DM, Giasson B, Baba M, Schmidt ML, Nee LE, O'Connell B, Pollen DA, George-Hyslop P, Ghetti B, Nochlin D, Bird TD, Cairns NJ, Lee VMY, Iwatsubo T, Trojanowski JQ. Lewy bodies contain altered alpha-synuclein in brains of many familial Alzheimer's disease patients with mutations in presenilin and amyloid precursor protein genes. *Am. J. Pathol.* 1998; 153:1365–1370. [PubMed: 9811326]
36. Masliah E, Rockenstein E, Veinbergs I, Sagara Y, Mallory M, Hashimoto M, Mucke L. beta - Amyloid peptides enhance alpha -synuclein accumulation and neuronal deficits in a transgenic mouse model linking Alzheimer's disease and Parkinson's disease. *Proc. Natl. Acad. Sci. U. S. A.* 2001; 98:12245–12250. [PubMed: 11572944]
37. Mucke L, Masliah E, Yu GQ, Mallory M, Rockenstein EM, Tatsuno G, Hu K, Kholodenko D, Johnson-Wood K, McConlogue L. High-level neuronal expression of abeta 1-42 in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J. Neurosci.* 2000; 20:4050–4058. [PubMed: 10818140]
38. Norris EH, Giasson BI, Lee VM. Alpha-synuclein: normal function and role in neurodegenerative diseases. *Curr. Top. Dev. Biol.* 2004; 60:17–54. [PubMed: 15094295]
39. Periz G, Fortini ME. Functional reconstitution of gamma-secretase through coordinated expression of presenilin, nicastrin, Aph-1, and Pen-2. *J. Neurosci. Res.* 2004; 77:309–322. [PubMed: 15248287]
40. Poewe W. Non-motor symptoms in Parkinson's disease. *Eur. J. Neurol.* 2008; 15(Suppl 1):14–20. [PubMed: 18353132]
41. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H, Rubenstein J, Boyer R, Stenroos ES, Chandrasekharappa S, Athanassiadou A, Papapetropoulos T, Johnson WG, Lazzarini AM, Duvoisin RC, Di Iorio G, Golbe LI, Nussbaum RL. Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science.* 1997; 276:2045–2047. [PubMed: 9197268]
42. Selkoe DJ. The origins of Alzheimer's disease. *JAMA.* 2000; 283:1615–1617. [PubMed: 10735401]
43. Selkoe DJ. Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev.* 2001; 81:741–766. [PubMed: 11274343]
44. Siman R, Reaume AG, Savage MJ, Trusko S, Lin YG, Scott RW, Flood DG. Presenilin-1 P264L knock-in mutation: differential effects on abeta production, amyloid deposition, and neuronal vulnerability. *J. Neurosci.* 2000; 20:8717–8726. [PubMed: 11102478]
45. Simuni, T.; Hurtig, HI. Parkinson's disease: the clinical picture.. In: Clark, CM.; Trojanowski, JQ., editors. *Neurodegenerative dementias*. McGraw-Hill; New York: 2000. p. 193-203.

46. Snider BJ, Norton J, Coats MA, Chakraverty S, Hou CE, Jervis R, Lendon CL, Goate AM, McKeel DW Jr, Morris JC. Novel presenilin 1 mutation (S170F) causing Alzheimer disease with Lewy bodies in the third decade of life. *Arch. Neurol.* 2005; 62:1821–1830. [PubMed: 16344340]
47. Spillantini MG, Schmidt ML, Lee VMY, Trojanowski JQ, Jakes R, Goedert M. Alpha-synuclein in Lewy bodies. *Nature.* 1997; 388:839–840. [PubMed: 9278044]
48. Taylor JP, Hardy J, Fischbeck KH. Toxic proteins in neurodegenerative disease. *Science.* 2002; 296:1991–1995. [PubMed: 12065827]
49. Terai K, Iwai A, Kawabata S, Tasaki Y, Watanabe T, Miyata K, Yamaguchi T. beta-amyloid deposits in transgenic mice expressing human beta-amyloid precursor protein have the same characteristics as those in Alzheimer's disease. *Neuroscience.* 2001; 104:299–310. [PubMed: 11377835]
50. Waxman EA, Duda JE, Giasson BI. Characterization of antibodies that selectively detect alpha-synuclein in pathological inclusions. *Acta Neuropathol.* 2008; 116:37–46. [PubMed: 18414880]
51. Waxman EA, Giasson BI. Molecular mechanisms of alpha-synuclein neurodegeneration. *Biochim. Biophys. Acta.* 2008; 1792:616–624. [PubMed: 18955133]
52. Waxman EA, Giasson BI. Specificity and regulation of casein kinase-mediated phosphorylation of alpha-synuclein. *J. Neuropathol. Exp. Neurol.* 2008; 67:402–416. [PubMed: 18451726]
53. Wilson CA, Doms RW, Lee VMY. Intracellular APP processing and A β production in Alzheimer's disease. *J. Neuropath. Exp. Neurol.* 1999; 58:787–794. [PubMed: 10446803]
54. Yang F, Ueda K, Chen P, Ashe KH, Cole GM. Plaque-associated alpha-synuclein (NACP) pathology in aged transgenic mice expressing amyloid precursor protein. *Brain Res.* 2000; 853:381–383. [PubMed: 10640638]
55. Yokota O, Terada S, Ishizu H, Ujike H, Ishihara T, Nakashima H, Yasuda M, Kitamura Y, Ueda K, Checler F, Kuroda S. NACP/alpha-synuclein, NAC, and beta-amyloid pathology of familial Alzheimer's disease with the E184D presenilin-1 mutation: a clinicopathological study of two autopsy cases. *Acta Neuropathol.* 2002; 104:637–648. [PubMed: 12410385]

Highlights

Affects of A β amyloid on the formation of α -synuclein aggregation was studied in mice.

Double M83/Tg2576 and triple M83/Tg2576/P246L presenilin-1 mice were generated.

Mice were evaluated for A β and α -synuclein pathology.

Hyperphosphorylated α -synuclein dystrophic neurites accumulated within A β plaques.

No further α -synuclein pathologies were observed.

Figure 1. PCR and Western blot analysis confirming the genotypes of the mice

(A) Ethidium bromide stained TBE/1.5% agarose gel loaded with PCR products amplified as described in “Material and Methods” from the mice genotyped indicated above each lane.

(B) Western blot analysis with polyclonal antibody Karen specific for β APP. (C) Western blot analysis with antibody Syn 211 specific for human α -syn. (D) Western blots were also probed with an actin antibody to assess equal protein loading. The mobility of protein markers in kDa are indicated on the left.

Figure 2. Representative Images of A β Plaques and α -Syn Dystrophic Neurites in Tg2576/M83 Bigenic Mice

Immunocytochemistry and immunofluorescence analyses were performed as described in “Material and Methods”. Abundant A β plaques stained with antibodies 6E10 (A) or NAB228 (B) in the cortex and hippocampus of a 21-month old Tg2576/M83 bigenic mouse. Accumulation of α -syn dystrophic neurites surrounding A β plaques detected with antibodies Syn 505 (C,D) and pSer129 (E). Double immunofluorescence of A β plaques with a rabbit anti-A β antibody (green) and antibody Syn 506 (red) showing accumulation of α -syn dystrophic neurites surrounding amyloid plaques (F). For immunocytochemistry sections were counterstained with hematoxylin and for immunofluorescence sections were counterstained with DAPI. Arrows indicate α -syn dystrophic neurites. Bar = 200 μ m in A, B, 100 μ m in C, F and 50 μ m in D, E.

Figure 3. Representative Images of Aβ Plaques and α-Syn Dystrophic Neurites in M83/Tg2576/P264L PSI Transgenic Mice

Immunocytochemistry analysis was performed as described in “Material and Methods”. Abundant Aβ plaques stained with antibodies 6E10 (A) or NAB228 (B) in the cortex and hippocampus of a 12-month old Tg2576/M83/PSI P264L transgenic mouse. Accumulation of α-syn dystrophic neurites surrounding Aβ plaques stained with antibodies Syn 505 (C) and pSer129 (D). Arrows indicate α-syn dystrophic neurites. Sections were counterstained with hematoxylin. Bar = 200 μm in A, B and 100 μm in C, D.