Published in final edited form as:

Acta Neurochir Suppl. 2011 ; 111: 31–35. doi:10.1007/978-3-7091-0693-8_6.

A mouse model of intracranial aneurysm—technical considerations

Yoshiteru Tada, MD, PhD^{1,2,3}, Yasuhisa Kanematsu, MD, PhD^{3,4}, Miyuki Kanematsu, MD⁵, Yoshitsugu Nuki, MD, PhD^{1,2}, Elena I Liang, BS^{1,2}, Kosuke Wada, MD^{1,2}, Hiroshi Makino, MD^{1,2}, and Tomoki Hashimoto, MD^{1,2}

¹Department of Anesthesia and Perioperative Care, University of California, San Francisco

²Center for Cerebrovascular Research, University of California, San Francisco, U.S.A

⁴Department of Neurosurgery, Institute of Health Biosciences, The University of Tokushima, Tokushima, Japan

⁵Department of Thoracic, Endocrine Surgery and Oncology, Institute of Health Biosciences, The University of Tokushima, Tokushima, Japan

Summary

Intracranial aneurysms can be induced by a single stereotaxic injection of elastase into the cerebrospinal fluid at the right basal cistern in hypertensive mice. This mouse model produces large aneurysm formations with an incidence of 60–80% within 3–4 weeks. Intracranial aneurysms in this model recapitulate key pathological features of human intracranial aneurysms. Several technical factors are critical for the successful induction of intracranial aneurysm in this model. Precise stereotaxic placement of the needle tip into the cerebrospinal fluid space is especially important. Aneurysm formations in this model can serve as a simple and easily interpretable outcome for future studies that utilize various inhibitors, knockout mice, or transgenic mice to test roles of specific molecules and pathways in the pathophysiology of intracranial aneurysms.

Keywords

Intracranial aneurysm; animal model; macrophage; matrix metalloproteinase; hypertension; inflammation; mouse

Introduction

Despite recent advances in diagnosis and treatment, the mechanisms for formation, growth, and subsequent rupture of intracranial aneurysms are not well understood (9). In order to understand the pathophysiology of intracranial aneurysms and identify therapeutic targets for the prevention of aneurysmal rupture, an animal model of intracranial aneurysm would be useful. Recently, we have developed a new mouse model of intracranial aneurysm that recapitulates key pathological features of human intracranial aneurysms (7). This model

Corresponding author: Tomoki Hashimoto, MD, Department of Anesthesia and Perioperative Care, University of California, San Francisco, 1001 Potrero Avenue, No. 3C-38, San Francisco, CA 94110, USA. Phone: (415) 206-8958. Fax: (415) 206-8170; hashimot@anesthesia.ucsf.edu.

³These authors (YT and YK) contributed equally to this study.

Conflict of interest statement

Authors do not have conflict of interest.

produces large intracranial aneurysms within a short incubation time of two to three weeks. In this article, we will discuss technical considerations that are critical for the successful application of this model for future studies.

Overview of elastase-induced intracranial aneurysm in hypertensive mice

In this model, two well-known clinical factors associated with human intracranial aneurysms—hypertension and the degeneration of elastic lamina—are combined to induce intracranial aneurysm formation in mice (7). Hypertension is associated with the formation and rupture of intracranial aneurysm. Degeneration of elastic lamina is a key pathological change of intracranial aneurysm. Fragmentation of elastic lamina alone is considered as pre-aneurysmal change that can eventually mature into an aneurysm. In this model, hypertension is induced by continuous infusion of angiotensin II; degradation of elastic lamina is induced by a single injection of elastase into the cerebrospinal fluid using a stereotaxic method. This model produces large intracranial aneurysms at an incidence of 60–80% within three to four weeks. Aneurysms are formed in the Circle of Willis and its major branches. Figure 1 shows a control brain (A) and representative aneurysms (B, C) from this model.

There are dose-dependent effects of angiotensin II and elastase on the incidence of aneurysm in this model. Neither angiotensin II nor elastase alone could induce aneurysms, indicating that synergistic effects of elastase and hypertension are needed for the aneurysm formation in our model. When hypertension-induced vascular remodeling is coupled with elastase-induced inflammation, abnormal vascular remodeling may occur. Intracranial aneurysm formation may represent maladaptive vascular remodeling that is mediated by hemodynamic changes and inflammation (4).

Histologically, intracranial aneurysms formed in this model closely resemble human intracranial aneurysms. Part of the aneurysmal wall in this model is often thickened. Although endothelial and smooth muscle cell layers are sometimes fragmented, they are generally present throughout the aneurysmal wall, showing that the intracranial aneurysms formed in this model are true aneurysms, but not pseudo-aneurysms.

Aneurysms formed in this model are generally large and easily distinguishable from the normal arteries without using histological assessment. Aneurysm formations in this model can serve as a simple and easily interpretable outcome for future studies that utilize various inhibitors, knockout mice, or transgenic mice to test roles of specific molecules and pathways in the pathophysiology of intracranial aneurysms. For example, using this model, we have shown critical roles of matrix metalloproteinases (MMPs) in the formation of intracranial aneurysm (7).

Angiotensin II-induced hypertension

Angiotensin II infusion through implanted osmotic pump is a well-established, well-characterized model of hypertension in mice (2, 5, 11). A horizontal skin incision approximately 1 cm long should be made on one side of the back at 1 cm cephalad from the base of tail. The small pocket between the skin and muscle layers that will contain the osmotic pump can be made using a pair of forceps. The osmotic pump can be implanted into this pocket, and the skin should be closed with sutures. This procedure takes only 5–10 minutes with a minimum amount of training. Proper suturing of the incision is very important since the wound dehiscence can result in unexpected explantation of the osmotic pump. Frequent monitoring of the wound site is needed, and the additional suture can be added as needed.

Using angiotensin II infusion, a rise in systolic blood pressure can be detected as early as one week, and hypertension can be maintained at least up to six weeks. Norepinephrine infusion or deoxycorticosterone acetate (DOCA)-salt treatment can be also used to induce hypertension in mice (5, 11). DOCA-salt hypertension may be useful when non-hemodynamic effects of angiotensin II need to be avoided. Angiotensin II can exert various effects on the vasculature in addition to its hypertensive effect, including induction of reactive oxygen species and promotion of inflammation (5). It should be noted that in our experience, the rise in blood pressure in these two methods was milder than that of angiotensin II-induced hypertension; hypertension became less significant after three weeks using norepinephrine infusion or deoxycorticosterone acetate (DOCA)-salt treatment. Angiotensin II-induced hypertension may be more consistent and sustained.

Blood pressure measurement

To monitor successful induction and maintenance of systemic hypertension, blood pressure measurement is critical. Since repeated measurements of blood pressure are needed, the tail cuff method is used in our laboratory (ML125M, ADInstruments). In our experience, this method provides accurate estimates for systolic blood pressure. Our preliminary data showed a highly significant correlation between the systolic blood pressure values using the tail cuff method and the mean blood pressure values obtained from the direct measurement using a femoral artery catheter.

Blood pressure measurement of conscious mice requires a series of training of mice. Some strains of mice, such as C57BL/6J mice, appear to have high locomotor activity and resistance to restraint, even after training for blood pressure measurements (2). Alternatively, blood pressure measurement can be performed on anesthetized mice. Systolic blood pressure can be reliably measured when mice are under a steady state of isoflurane anesthesia (5, 7). This would require a continuous monitoring of isoflurane concentration in the anesthesia chamber or mask. We have conducted an experiment comparing systolic blood pressure values between conscious state and anesthetized state.

In three groups of mice, blood pressure was measured in both conscious and anesthetized state. Mice received either phosphate buffered saline, angiotensin II at 500 ng/kg/min, or angiotensin II at 1000 ng/kg/min through osmotic pumps ($n = 3$ in each group). After two consecutive days of acclimation training, mice underwent “conscious” blood pressure measurement. Immediately after the blood pressure measurements were taken in the conscious state, mice were anesthetized with isoflurane, and blood pressure was measured at the steady state of isoflurane anesthesia (after 15 minutes of equilibration with isoflurane concentration at 1.5%). As shown in Figure 2, there was a close correlation between systolic blood pressure measured under conscious state and that under anesthetized state ($R^2 = 0.96$, $P < .05$). As we have expected, systolic blood pressure measured under anesthetized state was slightly lower than that under conscious state ($y = 1.05x - 17$), reflecting blood pressure lowering effect of isoflurane. We have previously shown that isoflurane anesthesia does not mask effects of anti-hypertensive agents (5). These data showed that systolic blood pressure measurement under a steady isoflurane anesthesia can be reliably used to monitor changes in blood pressure in mice.

Stereotaxic injection of elastase

Detailed method for the single stereotaxic injection of elastase into the cerebrospinal fluid at the right basal cistern has been described previously (7). Briefly, the tip of a 26-gauge needle (10 μ L syringe Model 701 with 26g 2 inch Hamilton Replacement needle Point style 3, Fisher Scientific) was stereotaxically placed in the right basal cistern using the coordinates obtained from Mouse Brain Atlas (2.5 mm posterior to the bregma, 1.0 mm

lateral to the midline, and 5.0 mm ventral to the skull surface). Elastase (E7885, porcine pancreatic elastase, lyophilized powder, 20mg, Sigma Aldrich) was dissolved in PBS. 2.5 μ L of the elastase solution (0 – 35 milli-units) was injected at 0.2 μ L/min. This part of the procedure will take approximately 30–40 minutes. Significant variations may exist in the potency of elastase activity in different lots or different preparations of commercially available elastase (1, 3). It may be prudent to use elastase from the same lot in each set of experiments.

In our experience, we found that the stereotaxic coordinates need to be adjusted depending on the operator and the age of mice. Therefore, a series of test injections using dye (bromophenol blue) was extremely important to ensure the correct placement of the needle tip at the right basal cistern. The coordinates we described in the original paper (7) should be regarded as a guideline. Each researcher should perform a series of test injections to adjust for the appropriated coordinates before starting the actual experiments. Figure 3A shows a result of the typical successful test injections. The mouse was sacrificed immediately after injection; bromophenol blue dye was observed along the right half of Circle of Willis. It appears that at least 10 test injections are needed before each operator establishes the appropriate coordinates that yield stable aneurysm induction. We routinely perform a test injection once every 10 mice to ensure the correct needle placement.

The depth of the needle is another critical factor for the successful induction of aneurysm in this model. If the needle depth is too shallow, the injection of elastase into the brain parenchyma results in intracerebral hemorrhage similar to the collagenase-induced intracerebral hemorrhage (Figure 3B) (6, 8, 10). If the needle tip is too close to the surface of the brain, the injection of elastase into the cerebrospinal fluid can result in diffuse hemorrhage from the surface of the brain, even when the needle tip is inside the subarachnoid space (Figure 3C). This might be due to compartmentalization of the subarachnoid space.

We recommend that the operator should find the depth at which the needle tip touches the surface of the dura matter or skull base. Then, the operator can use the depth that is approximately 0.3–0.4 mm shallower than the dura or skull base for subsequent test injections using dye. The spread of dye along the right half of Circle of Willis should be confirmed (Figure 3A). If the dura matter is intensely stained with the dye, subdural injection should be suspected. After establishing the depth for successful injections, the operator should perform test injections of elastase and sacrifice the animals one day after injection. Lack of diffuse bleeding from the surface of the brain or intracerebral hemorrhage would indicate the successful placement of the needle (Figure 3D).

The surgical procedures for the aneurysm induction seem to cause significant peri-operative stress to mice. Water intake of some of the postoperative mice seemed to have decreased during the first two post-operative days. Therefore, we inject 500 μ L of saline intraperitoneally once a day for the first two post-operative days in all mice.

Summary

Elastase-induced intracranial aneurysm in hypertensive mice represents an easily reproducible animal model that yields a high incidence of large intracranial aneurysm within a practical incubation time. There are several technical issues that are critical for the successful use of this model. This model can be used to study molecular pathways that are potentially involved in the pathophysiology of intracranial aneurysms.

Acknowledgments

Sources of Funding

This study was funded by NIH R01NS055876 (TH) and NIH P01NS044155 (TH)

References

1. Carsten CG 3rd, Calton WC, Johanning JM, Armstrong PJ, Franklin DP, Carey DJ, Elmore JR. Elastase is not sufficient to induce experimental abdominal aortic aneurysms. *J Vasc Surg.* 2001; 33:1255–1262. [PubMed: 11389426]
2. Daugherty A, Manning MW, Cassis LA. Angiotensin II promotes atherosclerotic lesions and aneurysms in apolipoprotein E-deficient mice. *J Clin Invest.* 2000; 105:1605–1612. [PubMed: 10841519]
3. Ding YH, Danielson MA, Kadirvel R, Dai D, Lewis DA, Cloft HJ, Kallmes DF. Modified technique to create morphologically reproducible elastase-induced aneurysms in rabbits. *Neuroradiology.* 2006; 48:528–532. [PubMed: 16708202]
4. Hashimoto T, Meng H, Young WL. Intracranial aneurysms: links between inflammation, hemodynamics and vascular remodeling. *Neurol Res.* 2006; 28:372–380. [PubMed: 16759441]
5. Kanematsu Y, Kanematsu M, Kurihara C, Tsou TL, Nuki Y, Liang EI, Makino H, Hashimoto T. Pharmacologically Induced Thoracic and Abdominal Aortic Aneurysms in Mice. *Hypertension.* 2010; 55:1267–1274. [PubMed: 20212272]
6. Kitaoka T, Hua Y, Xi G, Hoff JT, Keep RF. Delayed argatroban treatment reduces edema in a rat model of intracerebral hemorrhage. *Stroke.* 2002; 33:3012–3018. [PubMed: 12468805]
7. Nuki Y, Tsou TL, Kurihara C, Kanematsu M, Kanematsu Y, Hashimoto T. Elastase-induced intracranial aneurysms in hypertensive mice. *Hypertension.* 2009; 54:1337–1344. [PubMed: 19884566]
8. Rosenberg GA, Mun-Bryce S, Wesley M, Kornfeld M. Collagenase-induced intracerebral hemorrhage in rats. *Stroke.* 1990; 21:801–807. [PubMed: 2160142]
9. Shi C, Awad IA, Jafari N, Lin S, Du P, Hage ZA, Shenkar R, Getch CC, Bredel M, Batjer HH, Bendok BR. Genomics of human intracranial aneurysm wall. *Stroke.* 2009; 40:1252–1261. [PubMed: 19228845]
10. Tang J, Liu J, Zhou C, Alexander JS, Nanda A, Granger DN, Zhang JH. Mmp-9 deficiency enhances collagenase-induced intracerebral hemorrhage and brain injury in mutant mice. *J Cereb Blood Flow Metab.* 2004; 24:1133–1145. [PubMed: 15529013]
11. Weiss D, Kools JJ, Taylor WR. Angiotensin II-induced hypertension accelerates the development of atherosclerosis in apoE-deficient mice. *Circulation.* 2001; 103:448–454. [PubMed: 11157699]

Figure 1. Representative aneurysms

A. Control brain. B. Aneurysms were found at the bifurcation at the right middle cerebral artery and the internal carotid artery and at the proximal part of right middle cerebral artery. C. Aneurysm at the right anterior cerebral artery. RACA: right anterior cerebral artery, LACA: left anterior cerebral artery, RMCA: right middle cerebral artery.

Figure 2. Correlation of systolic blood pressure between conscious and anesthetized state

There was a close correlation between systolic blood pressure measured in a conscious state and systolic blood pressure measured in an anesthetized state ($y = 1.05x - 17$, $R^2 = 0.96$, $P < .05$).

Figure 3. Inspection of brain tissues 1 day after injection of dye or elastase

A. Test injection of dye. Immediately after a stereotaxic injection of bromophenol blue (2.5 μ L) into the cerebrospinal fluid at the right basal cistern, the mouse brain was extracted. The dye was observed along the right half of Circle of Willis. **B. Intracerebral hemorrhage due to elastase injection into the brain parenchyma.** If the needle depth is too shallow, an injection of elastase results in intracerebral hemorrhage. **C. Diffuse bleeding from the brain surface.** If the needle tip is too close to the surface of the brain, the injection of elastase into the cerebrospinal fluid can result in diffuse hemorrhage from the surface of the brain, even when the needle tip is inside the subarachnoid space. **D. Successful injection of elastase.** When the brain was inspected 1 day after elastase injection, there was no diffuse bleeding from the surface of the brain or intracerebral hemorrhage, indicating a successful placement of the needle