Published in final edited form as:

Life Sci. 2011 November 7; 89(19-20): 748–754. doi:10.1016/j.lfs.2011.08.016.

BB rat *Gimap* gene expression in sorted lymphoid T and B cells

Daniel H. Moralejo^{a,b}, Jessica M. Fuller^{c,d}, Elizabeth A. Rutledge^e, Brian Van Yserloo^d, Ruth A. Ettinger^d, Richard Jensen^d, William Osborne^f, Anne Kwitek^g, and Åke Lernmark^{b,c}

^aDept. of Comparative Medicine, University of Washington 1959 N.E. Pacific St., Box 357710, Seattle, Washington, 98195, USA

^bDept. of Veterinary Sciences, University of Texas M.D. Anderson Cancer Center, 650 Cool Water Drive, Bastrop, Texas, 78602, USA

^cDept. of Clinical Sciences, Lund University, Clinical Research Center, Entrance 72, Building 91:10, Malmö, 20502, Sweden

^dDept. of Medicine, University of Washington, 1959 N.E. Pacific St., Box 357710, Seattle, Washington, 98195, USA

^eSalish Kootenai College, Department of Life Sciences, 58138 Hwy 93, Pablo, MT 59855, USA

^fDept. of Pediatrics, University of Washington, 815 Mercer St., Box 358055, Seattle, Washington, 98109, USA

^gDept. of Internal Medicine, University of Iowa, 375 Newton Rd, 3111B MERF, Iowa City, IA, 52242, USA

Abstract

Aims—The *Gimap* gene family has been shown to be integral to T cell survival and development. A frameshift mutation in *Gimap5*, one of seven members of the *Gimap* family, results in lymphopenia and is a prerequisite for spontaneous type 1 diabetes (T1D) in the BioBreeding (BB) rat. While not contributing to lymphopenia, the *Gimap* family members proximal to *Gimap5*, encompassed within the *Iddm39* quantitative trait locus (QTL), have been implicated in T1D. We hypothesized that expression of the *Gimap* family members within the *Iddm39* QTL, during thymocyte development as well as in peripheral T and B cells contribute to T1D.

Main methods—Cell sorted subpopulations were analyzed by quantitative real time (qRT) PCR.

Key findings—*Gimap4* expression was reduced in DR.^{lyp/lyp} rat double negative, double positive and CD8 single positive (SP) thymocytes while expression of *Gimap8*, *Gimap6*, and *Gimap7* was reduced only in CD8 SP thymocytes. Interestingly, expression of the entire *Gimap* gene family was reduced in DR.^{lyp/lyp} rat peripheral T cells compared to non-lymphopenic, non-diabetic DR.^{+/+} rats. With the exception of *Gimap6*, the *Gimap* family genes were not expressed in B cells from spleen and mesenteric lymph node (MLN). Expression of *Gimap9* was only detected in hematopoietic cells of *non* B cell lineage such as macrophage, dendritic or NK cells.

© 2011 Elsevier Inc. All rights reserved.

Corresponding author: William Osborne, University of Washington, Department of Pediatrics, 815 Mercer Street, Box 358055, Seattle, WA 98109, USA. Phone: 206-685-3315, Fax: (206) 543-3567, wosborne@u.washington.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Significance—These results suggest that lack of the *Gimap5* protein in the DR.^{lyp/lyp} congenic rat was associated with impaired expression of the entire family of *Gimap* genes and may regulate T cell homeostasis in the peripheral lymphoid organs.

Keywords

Gimap; lymphopenia; autoimmunity; type 1 diabetes

Introduction

The GTPase immunity-associated protein (*Gimap*) gene family in the rat is comprised of eight genes located within 150 Kb on rat chromosome (RNO) 4 (Krucken, et al. 2004; Rutledge, et al. 2009). The seven functional *Gimap* genes in rat; *Gimap8*, *Gimap9*, *Gimap4*, *Gimap6*, *Gimap7*, *Gimap1* and *Gimap5* are conserved across multiple vertebrates including human and mouse as well as in *Arabidopsis thaliana* (MacMurray, et al. 2002; Reuber and Ausubel 1996; Schnell, et al. 2006). These small GTPases regulate pro- and anti- apoptotic T cell pathways (Carter, et al. 2007; Dalberg, et al. 2007; Hernández-Hoyos, et al. 1999; Krücken, et al. 2005; Nitta, et al. 2006; Nitta and Takahama 2007; Pandarpurkar, et al. 2003; Schnell, et al. 2006) as well as thymocyte maturation and differentiation (Filen, et al. 2009; Nitta, et al. 2006) and are characterized by novel guanosine diphosphate/guanosine triphosphate (AIG1) and coiled-coil domains, common binding motifs and variable C termini (Krucken, et al. 2004; MacMurray, et al. 2002).

Peripheral T cell lymphopenia in the BioBreeding (BB) diabetes prone (DP) rat is linked to a single nucleotide deletion in *Gimap5* (*Iddm2*; *lyp*) resulting in loss of *Gimap5* protein expression that is a prerequisite for onset of spontaneous type 1 diabetes (T1D) in the DP rat (Fuller, et al. 2006; Hornum, et al. 2002; MacMurray, et al. 2002; Michalkiewicz, et al. 2004). While the *Gimap5* mutation was not associated with T1D in humans (Payne, et al. 2004), the uncommon SNP in the polyadenylation signal of human *GIMAP5* was found to be associated with risk for high IA-2 autoantibody levels in T1D patients (Shin, et al. 2007) implicating a possible role in islet autoimmunity. The common SNP at this location was associated with systemic lupus erythematosus underscoring the possible contribution of *GIMAP5* to autoimmunity (Hellquist, et al. 2007).

Gimap8, *Gimap9*, *Gimap4*, and *Gimap7* are transcribed in the same direction while *Gimap6* runs opposite to the rest of the family. Each gene is separated by approximately 12 Kb of non-coding sequence. *Gimap8*, unique with three AIG1 binding domains (Dion, et al. 2005; Krucken, et al. 2004; Nitta, et al. 2006; Rutledge, et al. 2009), localizes to the endoplasmic reticulum, Golgi and mitochondria and may play an anti-apoptotic role in immune cells (Krücken, et al. 2005). *Gimap6* and *Gimap7* are transmembrane proteins containing weak endoplasmic reticulum (ER) localization signals (Krucken, et al. 2004). *Gimap4*, a cytosolic lymphoid specific signaling molecule, is associated with β -selection during maturation of CD4⁺CD8⁺ double negative (DN) thymocytes into CD4⁺CD8⁺ double positive (DP) thymocytes (Fehlingvon Boehmer 1997; Schnell, et al. 2006) positive selection of T cells and accelerated T cell apoptosis (Nitta, et al. 2006; Poirier, et al. 1999; Schnell, et al. 2006).

Our previous study of the *Gimap* family showed that all seven members are differentially expressed in DR.^{+/+} and DR.^{lyp/lyp} spleen, thymus and mesenteric lymph node (MLN) (Rutledge, et al. 2009). However, as whole organs of mixed cell populations were studied, it remains uncertain to what extent the observed differences were due to the cellular composition of these organs which are markedly affected by the reduction of T cells and a concordant increase in B cells and monocytes. To investigate these *Gimap* gene expression variations in sorted subpopulations of cells, we performed quantitative real time (qRT) PCR

analysis on purified T and B cells from DR.^{+/+} and DR.^{lyp/lyp} rat thymus, spleen and MLN. We chose to study lymphocytes from MLN in DR.^{lyp/lyp} rats because their defective intestinal permeability may permit passage of environmental antigens initiating an autoimmune response, triggering T1D and changes in expression of *Gimap* genes (Graham, et al. 2004; Malaisse, et al. 2004; Vaarala 2008). Clarifying *Gimap* gene expression in specific populations of cells will aid in our understanding of the role of the *Gimap* family not only in thymocyte development, T cell survival in the periphery, or both, but also in T1D.

Materials and Methods

DR.lyp Rats

The DR.*lyp* (BBDR.BBDP-(*D4Rhw17-SS99306861*)(*D4Rhw11-D4Rhw10*)/*Rhw*) congenic rat line was derived from animals with two independent recombination events developed from our previously described introgression of the lymphopenia locus by cyclic cross-intercross breeding of BBDP with BBDR rats (Bieg, et al. 1998). The first recombination event was flanked by simple sequence length polymorphism (SSLP) marker D4Rhw11 (75.81 Mb) and the second flanked by SSLP marker D4Rhw10 (77.81 Mb) (Fuller, et al. 2006). Thus, the DP DNA in the DR.*lyp* rat line encompasses the *lyp* critical interval from D4Rhw6 (76.83 Mb) to *IIsnp3* (77.16 Mb) (MacMurray, et al. 2002). In addition, the DR.*lyp* congenic rat line used in the present study also contains BBDP DNA at D4Rat102 (66.22 Mb) and D4Rat26 (69.18 Mb). The DR.*lyp* congenic rat line is kept in heterozygous sister-brother breeding and produces Mendelian proportions of the DR.^{lyp/lyp} (25%), DR.^{lyp/+} (50%) and DR.^{+/+} (25%) genotypes. One hundred percent of DR.^{lyp/lyp} rats are lymphopenic and diabetic.

Housing

Rats were housed in a specific pathogen-free facility at the University of Washington, Seattle, Washington, on a 12-h light/dark cycle with 24-h access to food (Harlan Teklad, Madison, WI) and water. The University of Washington Rodent Health Monitoring Program was used to track infectious agents via a quarterly sentinel monitoring system. The University of Washington institutional animal use and care committee approved all of the protocols used in the present study.

Genotyping Chromosome 4

Between 25 and 30 days of age, 2 mm tail snips were obtained and DNA isolated as previously described (Fuller, et al. 2009). The isolated DNA samples were diluted to 25 ng/ μ L in ddH₂O and 2 μ L of this genomic DNA used in the following 10 μ L reaction: 2 μ L 5X reaction buffer (Promega, Madison, WI), 0.2 μ L 10 mmol/L dNTP (Promega), 1.0 μ L 1 μ mol/L IRDye 700 labeled primer (LiCor Biosciences, Lincoln, NE), 0.5 μ L 20 μ mol/L unlabeled reverse primer (Invitrogen, Carlsbad, CA), 0.20 μ L GoTaq DNA Polymerase (Promega) and 4.1 μ L ddH₂O. The PCR protocol was 95 °C for 5 min, 95 °C for 20 sec, 62 °C for 20 sec, 72 °C for 30 sec, steps 2–4 repeated 30 times, then 72 °C for 3 min. PCR products were diluted to 25% with STOP solution (LiCor Biosciences) and analyzed using a NEN Global IR2 DNA Analyzer System (Model 4200S-2) with a 6.5% gel matrix (LiCor Biosciences).

Cell Preparation and Purification

Rat thymocytes, spleen, and MLN were minced in PBS containing 10% FCS. The single cell suspensions from spleen were washed in lysing buffer (BD Pharm Lyse™) (BD Biosciences, San Diego, CA) to eliminate red blood cells. B cell suspensions were purified

by positive selection of CD45⁺ and CD45RA⁺ cells with the respective biotinylated monoclonal antibody (mAb) from BD Bioscience Pharmingen (San Diego, CA) using streptavidin-conjugated MACS microbeads according to the manufacturer's instructions Miltenyi MicroBead (Invitrogen, Carlsbad, CA).

FACS Sorting Of Lymphocyte Subsets

Single cell suspensions of rat thymocytes, spleen and MLN cells were washed three times in PBS containing 10% FCS and then incubated for 30 min at 4°C with primary mAbs in the dark. Cells were washed twice and after the final wash filtered through a 40-µm cell-strainer immediately before cell sorting. The following mouse anti rat mAbs obtained from BD Bioscience Pharmingen, were used R-phycoerythrin (R-PE)-labeled anti-CD3 (G4.18), Cy-chrome labeled anti-CD4 (OX35), FITC labeled anti-CD8 (OX8), and FITC labeled anti-alpha-beta T cell receptor (R73). Cell populations were sorted on a FACS Aria cell sorter (BD Biosciences, Mountain View, CA). Purities were checked by flow cytometry to be between 90 and 99.9%.

RNA Isolation

FACS sorted thymus, spleen and MLN cells from 45–50 day old rats were homogenized in RNA lysis solution (Stratagene, La Jolla, CA or Qiagen, Valencia, CA) using a Kinematica Polytron PT 10/35 (Brinkmann, Westbury, NY) then aliquoted to minimize degradation from freezing and thawing. Total RNA was isolated using either RNeasy (Qiagen) or Absolutely RNA Miniprep Kit (Stratagene) followed by treatment with DNase according to the manufacturer's instructions.

Quantitative RT-PCR

Quantitative RT-PCR was performed on an Mx4000® Multiplex QPCR System (Stratagene) with samples run in duplicate using approximately 10 ng of total RNA. Twenty-five µL reactions were run using a Brilliant® Single-Step qRT-PCR Kit (2.5 µL 10x core RT-PCR buffer, 5.0 mmol/L MgCl₂, 400 nmol/L each primer, 200 nmol/L each probe, 0.8 mmol/L dNTP mix, 75 nmol/L passive reference dye, 2.0 units Stratascript RT, 1.25 units SureStart *Taq* DNA-polymerase). The PCR cycling conditions were as follows; 45°C for 30 min, 95°C for 10 min, and 40 cycles of 95°C for 30 sec, 60°C for 1 min. Resulting Ct values were converted to copies, normalized to GAPDH and expressed as the mean ± standard error of the mean (SEM). Probes were positioned in the 3' regions of the transcripts where there is more variation between the different *Gimap* genes and subjected to BLAST alignment to ensure specificity. The primers and probes used for each gene are detailed elsewhere (Rutledge, et al. 2009). Primers were obtained from Integrated DNA Technologies (Coralville, IA) or Qiagen Operon (Valencia, CA). Fluorescently labeled probes were obtained from Integrated DNA Technologies. Representative qRT-PCR products for each gene were run on an agarose gel to check for primer pair binding specificity. These PCR amplicons (used to convert to copy number) were quantitated in triplicate wells with the PicoGreen® RNA Quantitation Kit (Molecular Probes, Eugene, OR) using standards supplied by the manufacturer. Results from each assay were validated and normalized against GAPDH. The standard curves showed the following reaction efficiencies; *Gimap8*: 96% ± 3 SD, *Gimap9*: 94% ± 1 SD, *Gimap4*: 92% ± 2 SD, *Gimap6*: 90% ± 3 SD, *Gimap7*: 91% ± 4 SD, *Gimap1*: 94% ± 2 SD, *Gimap5*: 92% ± 2 SD, *Lr8*: 94% ± 3 SD, and GAPDH: 92% ± 1 SD.

Data analysis

Analyses of *Gimap* gene expression were carried out using linear mixed effect models (Pinheiro and Bates 2000) with S-PLUS (Insightful Corp., Seattle, WA). Comparisons of

expression between cell types consisted of twelve rats with two repeated measures for each rat. Comparison between DR.^{lyp/lyp} and DR.^{+/+} genotypes consisted of six rats from three litters consisting of one DR.^{lyp/lyp} and one DR.^{+/+} rat in each litter and two repeated measures for each rat. All of these analyses contained random effects for each of the three litters and two measures for each rat nested within those litters. Likelihood ratio tests were used to test the need for random intercepts. The significance for the fixed effects were assessed by *F* test for the coefficient or the group of coefficients (Lehmann 1986). A Bonferroni correction was applied to *p*-values to adjust for multiple comparisons and adjusted values were considered significant at *p* < 0.05.

Results

Thymic T cells

Expression of the five *Gimap* genes within the *Iddm39* QTL; *Gimap8*, *Gimap9*, *Gimap4*, *Gimap6* and *Gimap7*, was compared to *Gimap1*, *Gimap5* and *Lr8*, a gene unassociated with the *Gimap* family but located within the 33 Kb lymphopenia critical interval (Fig. 1). The number of thymocytes, sorted based on double staining with CD4 (T helper) and CD8 (T cytotoxic) monoclonal antibodies, did not differ between DR.^{+/+} and DR.^{lyp/lyp} rats (Fig. 2). The sorted cells showed a high degree (>90%) of purity (Fig. 2, Panels A&B). Expression of *Gimap4* in CD4–CD8– DN, CD4 single positive (SP), and CD8 SP thymocytes differed between DR.^{+/+} and DR.^{lyp/lyp} rats (*p*<0.05 for all) while expression of *Gimap8* (*p*<0.05), *Gimap6* (*p*<0.05) and *Gimap7* (*p*<0.001) only differed in CD8 SP cells (Fig. 3). It should be noted that all statistical tests have been corrected for both litter and multiple comparisons. Within the *lyp* critical interval, *Gimap5* expression was reduced in DR.^{lyp/lyp} rat CD4 and CD8 SP thymocytes when compared to DR.^{+/+} (*p*<0.05 for both). *Gimap9*, *Gimap1* and *Lr8* were not appreciably expressed at any stage of T cell development in the thymus.

Spleen and Mesenteric Lymph Node Peripheral T cells

Peripheral T cells from spleen and MLN were triple stained with CD3 (T cell receptor), CD4 and CD8 monoclonal antibodies. The number of sorted cells gated for CD3 and then sorted based on staining with CD4 and CD8 did not differ between DR.^{+/+} and DR.^{lyp/lyp} rats and showed a high degree of purity (Fig. 2, panel C and D). Correcting for litter and multiple testing, we found that the expression of all *Gimap* genes in CD4+ splenic T cells did not differ between DR.^{+/+} and DR.^{lyp/lyp} rats (Fig. 4). In contrast, the expression of all of the *Gimap* genes within the *Iddm39* QTL was reduced in DR.^{lyp/lyp} CD8+ splenic T cells.

The expression of *Gimap4* and *Gimap9* genes in the MLN CD4+ T cells showed no difference between DR.^{lyp/lyp} and DR.^{+/+} rats, however, there was a reduction in *Gimap8*, *Gimap6* and *Gimap7*. The MLN CD8+ T cells showed a reduction in expression of *Gimap8*, *Gimap4* and *Gimap7* while there was no difference in *Gimap9* and *Gimap6* (Fig. 4). As in the thymus, *Gimap1* and *Lr8* in the *lyp* critical interval did not express appreciably in either DR.^{+/+} or DR.^{lyp/lyp} rat peripheral T cells (Fig. 4).

Spleen and MLN derived B cells

Hematopoietic B and non B cells (including T cells, macrophages, dendritic or NK cells) from spleen and MLN were double stained with CD45 (hematopoietic) and CD45RA (B cell) monoclonal antibodies and sorted based on staining with CD45RA (+/–). With the exception of *Gimap6*, the *Gimap* family in did not express appreciably in DR.^{+/+} or DR.^{lyp/lyp} rat B cells (CD45+CD45RA+) from spleen or MLN (Fig. 5). Expression of the *Gimap* genes within the *Iddm39* QTL; *Gimap8*, *Gimap9*, *Gimap4*, *Gimap6* and *Gimap7* was reduced in non B cells (CD45+CD45RA–) from DR.^{lyp/lyp} rat spleen and MLN. The comparisons which remained statistically significant after correction for litter and multiple

testing are indicated (Fig. 5). Interestingly, *Gimap9* was expressed in both DR.^{+/+} and DR.^{lyp/lyp} rat *non* B cells suggesting that *Gimap9* is expressed in hematopoietic cells of *non* T or B cell lineage such as NK, macrophage or dendritic cells. Within the *lyp* critical interval (Fig. 1), *Gimap5* showed a similar pattern with little or no expression in B cells and a reduced expression in DR.^{lyp/lyp} rat *non* B cells ($p < 0.05$). As in the thymus, spleen and MLN, *Gimap1* did not express appreciably in either DR.^{+/+} or DR.^{lyp/lyp} rat B or *non* B cells (Fig. 5).

Discussion

The *Gimap5* mutation within the *lyp* critical interval has been documented to result in peripheral T cell lymphopenia in the DR.^{lyp/lyp} rat (Dion, et al. 2005; Hornum, et al. 2002; MacMurray, et al. 2002). In our previous qRT-PCR analysis we showed reduced expression of all seven *Gimap* genes; *Gimap8*, *Gimap9*, *Gimap4*, *Gimap6*, *Gimap7*, *Gimap1* and *Gimap5* in DR.^{lyp/lyp} rat spleen and MLN when compared to DR.^{+/+} while only four; *Gimap9*, *Gimap4*, *Gimap1*, and *Gimap5* were reduced in thymus (Rutledge, et al. 2009). The present study resolves these previously observed expression differences to specific cell types both during thymocyte development and to mature T cells in the periphery. As lymphopenia impacts the cellular composition of the spleen and MLN, but not necessarily the thymus, the present qRT-PCR analysis of sorted cells further substantiates the *Gimap5* frameshift mutation as the primary defect in the periphery. Three of the *Gimap* genes within the *Iddm39* QTL; *Gimap8*, *Gimap6* and *Gimap7* as well as the *Gimap5* gene itself were expressed in DN and DP thymocytes with no significant difference in expression between DR.^{+/+} and DR.^{lyp/lyp} rats. Interestingly, while these genes were turned on during maturation of DN to DP thymocytes, no impact of the *Gimap5* mutation on gene expression was observed, as was seen in the periphery. Upon transition from DP to SP thymocytes, expression of four of the *Iddm39* genes; *Gimap8*, *Gimap4*, *Gimap6* and *Gimap7* as well as *Gimap5* increased in CD4 SP thymocytes and showed the highest overall expression as compared to the other three stages of thymocyte development however only *Gimap4* and *Gimap5* showed expression variation between DR.^{+/+} and DR.^{lyp/lyp} rats. These same five *Gimap* genes were poorly expressed in CD8 SP thymocytes and all showed a significant difference in expression between DR.^{+/+} and DR.^{lyp/lyp} rats. We believe this apparent reduction in expression of *Gimap8*, *Gimap4*, *Gimap6*, *Gimap7* and *Gimap5* in DR.^{lyp/lyp} CD8⁺ cells before leaving the thymus is a significant observation that may aid in understanding the possible role of the *Gimap* genes in T cells as well as their possible role in the pathogenesis of T1D.

The possible role of the *Gimap* genes in either protection from T cell apoptosis, such as *Gimap5* (Nitta, et al. 2006; NittaTakahama 2007; Pandarpurkar, et al. 2003) and *Gimap8* (Krücken, et al. 2005), or to induce apoptosis such as *Gimap4* (Nitta, et al. 2006) would perhaps explain the severe lack of CD8⁺ T cells in peripheral blood and tissues. *Gimap4* and *Gimap5* interact with the Bcl-2 family of signal transduction molecules to regulate the mitochondrial mediated T cell apoptosis pathway (Nitta, et al. 2006). The lower observed expression of *Gimap4* in the DR.^{lyp/lyp} CD4⁺ SP thymocytes would potentially explain that more CD4⁺ and less CD8⁺ T cells are detected in peripheral blood and tissues. It is however not understood why CD8⁺ cells are expressing less of all the *Gimap* genes, whether from DR.^{+/+} or DR.^{lyp/lyp} rats, or how this low expression impacts the ability of CD8⁺ T cells to survive in the periphery.

The major finding from the analysis of CD3⁺ spleen and MLN T cells sorted for either CD4 or CD8 positivity was that all the *Gimap* genes, except *Gimap9* and *Gimap1*, whether in the *Iddm39* QTL or the *lyp* critical interval showed reduced expression in the cells from the DR.^{lyp/lyp} rats. While the *Gimap* genes showed the highest expression levels overall in cells

sorted from the spleen and MLN, the difference in *Gimap* gene expression between DR.^{+/+} and DR.^{lyp/lyp} appeared more pronounced than in CD4 or CD8 SP thymocytes. The study of lymphocytes from MLN is justified because it is well known that the defective intestinal permeability associated with DR.^{lyp/lyp} rats may permit passage of environmental antigens initiating an autoimmune response and triggering T1D (Graham, et al. 2004; Malaisse, et al. 2004; Vaarala 2008). It should be noted that the same number of cells were sorted from both the spleen and the MLN whether the cells were from a DR.^{+/+} (typical sorting time for CD8+ T cells was 90 minutes) or DR.^{lyp/lyp} (typical sorting time for CD8+ T cells was 4 hours) rat. The higher level of expression of all the *Gimap* genes, except *Gimap9* and *Gimap1*, in peripheral compared to thymic CD8+ cells suggests that only the highest *Gimap* expressing cells survive the transition from the thymus to the periphery.

Our observation in sorted peripheral CD4+ and CD8+T cells was that the loss of the *Gimap5* protein is associated with a reduction in expression of all the *Gimap* family members (except *Gimap1* and *Gimap9*) suggesting that despite an increase in expression from the thymus to the periphery, CD8+ T cells from the DR.^{lyp/lyp} rat are prone to die. It is tempting to speculate that the coordinate expression of *Gimap8*, *Gimap4*, *Gimap6*, *Gimap7* and *Gimap5* may be due to the absence of the *Gimap5* protein. However, analyses of the *Gimap5* amino acid sequence show no indication that this protein would represent a transcription factor. The alternative explanation would be that *Gimap8*, *Gimap4*, *Gimap6*, *Gimap7* and *Gimap5* share a common transcriptional element. Bioinformatic analyses of our own and available database sequences have failed to detect a common promoter region or other sequences that would indicate the presence of a common regulator. Further studies are therefore required to uncover the mechanisms by which *Gimap8*, *Gimap4*, *Gimap6*, *Gimap7* and *Gimap5* are regulated in a coordinate fashion in CD4+ and CD8+ cells from DR.^{lyp/lyp} rats.

The nature of the *Gimap5* frame shift mutation leading to the absence of the *Gimap5* protein (MacMurray, et al. 2002; Michalkiewicz, et al. 2004) suggests that the loss of this anti-apoptotic protein contributes to lymphopenia. The importance of the *Gimap5* mutation for the DR.^{lyp/lyp} rat lymphopenia was amply illustrated by the marked reduction in expression of the *Gimap5* in the *non* B cells sorted from the spleen or the MLN. As the *Gimap5* mutation is comparable to a null mutation, i.e. no *Gimap5* protein is made (MacMurray, et al. 2002; Michalkiewicz, et al. 2004) it is of interest to note that the *Gimap5* knock out mouse that we recently reported (Schulteis, et al. 2008) resulted in impaired thymocyte maturation as well as survival of peripheral CD4 and CD8 positive T cells. *Gimap5* deficiency also blocked natural killer and NKT cell differentiation which could be restored on transfer of *Gimap5* deficient bone marrow into a wild-type environment. Although the phenotype of the *Gimap5* knockout mouse is comparable to that of the DR.^{lyp/lyp} rat, it remains to be clarified if the other *Gimap* genes are affected as demonstrated in the present study. As many of these proteins have been implicated in cell survival (Nitta, et al. 2006; Pandarpurkar, et al. 2003; Sandal, et al. 2003), it cannot be excluded that the concomitant reduction in several of the *Gimap* genes outside the *lyp* critical interval may contribute to the severe lymphopenia observed in the periphery (Dion, et al. 2005; Hornum, et al. 2002; MacMurray, et al. 2002).

In conclusion, a frameshift mutation in *Gimap5* results in lymphopenia and precedes onset of spontaneous T1D in the diabetes prone BB rat. Reduced expression of all seven *Gimap* genes in peripheral lymphoid organs suggests that the lack of *Gimap5* protein in the DR.^{lyp/lyp} congenic rat line impairs expression of the entire *Gimap* gene family and regulates T cell homeostasis in the peripheral lymphoid organs. The major findings in the present study both confirm our initial observations as well as resolve *Gimap* gene expression in specific thymic lineages as well as peripheral T cell and B cell subsets. Further elucidating the role of the *Gimap* family, specifically *Gimap5*, in T cell survival, development and/or

activation will aid in our understanding of the pathways in onset of spontaneous T1D in the BB rat and ultimately the pathways leading to development of human disease.

Acknowledgments

This work was supported by the National Institutes of Health (AI42380- D.H.M., J.M.F., R.A.E., A.K., Å.L.) and the Molecular and Genetics Core of the University of Washington Diabetes and Endocrinology Research Center (DK17047- E.A.R., B.V., W.R.O.), a Junior Faculty Award (D.H.M.), a Mentor Based Postdoctoral Fellowship Award (Å.L.) from the American Diabetes Association, the Robert H Williams Endowment at the University of Washington and the National Heart, Lung, and Blood Institute Training Grant T32HL007902 (R.A.J.).

References

- Bieg S, Koike G, Jiang J, Klaff L, Pettersson A, MacMurray AJ, Jacob HJ, Lander ES, Lernmark Å. Genetic isolation of the *Lyp* region on chromosome 4 in the diabetes prone BioBreeding (BB) rat. *Mammalian Genome*. 1998; 9(4):324–6. [PubMed: 9530633]
- Carter C, Dion C, Schnell S, Coadwell WJ, Graham M, Hepburn L, Morgan G, Hutchings A, Pascall JC, Jacobs H, Miller JR, Butcher GW. A natural hypomorphic variant of the apoptosis regulator Gimap4/IAN1. *J Immunol*. 2007; 179(3):1784–95. [PubMed: 17641045]
- Dalberg U, Markholst H, Hornum L. Both Gimap5 and the diabetogenic BBDP allele of Gimap5 induce apoptosis in T cells. *Int Immunol*. 2007; 19(4):447–53. [PubMed: 17369194]
- Dion C, Carter C, Hepburn L, Coadwell WJ, Morgan G, Graham M, Pugh N, Anderson G, Butcher GW, Miller JR. Expression of the Ian family of putative GTPases during T cell development and description of an Ian with three sets of GTP/GDP-binding motifs. *Int Immunol*. 2005; 17(9):1257–68. [PubMed: 16103028]
- Fehling HJ, von Boehmer H. Early alpha beta T cell development in the thymus of normal and genetically altered mice. *Curr Opin Immunol*. 1997; 9(2):263–75. [PubMed: 9099797]
- Filen J-J, Filen S, Moulder R, Tuomela S, Ahlfors H, West A, Kouvonen P, Kantola S, Bjorkman M, Katajamaa M, Rasool O, Nyman TA, Lahesmaa R. Quantitative proteomics reveals GIMAP family proteins 1 and 4 to be differentially regulated during human T helper cell differentiation. *Molecular & Cellular Proteomics*. 2009; 8(1):32–44. [PubMed: 18701445]
- Fuller J, Kwitek AE, Hawkins T, Moralejo DH, Lu W, Tupling TD, MacMurray AJ, Borchardt G, Hasinoff M, Lernmark A. Introgression of F344 rat genomic DNA on BB rat chromosome 4 generates diabetes-resistant lymphopenic BB rats. *Diabetes*. 2006; 55(12):3351–7. [PubMed: 17130479]
- Fuller JM, Bogdani M, Tupling TD, Jensen RA, Pefley R, Manavi S, Cort L, Blankenhorn EP, Mordes JP, Lernmark A, Kwitek AE. Genetic dissection reveals diabetes loci proximal to the gimap5 lymphopenia gene. *Physiol Genomics*. 2009; 38(1):89–97. [PubMed: 19351909]
- Graham S, Courtois P, Malaisse WJ, Rozing J, Scott FW, Mowat AMI. Enteropathy precedes type 1 diabetes in the BB rat. *Gut*. 2004; 53(10):1437–44. [PubMed: 15361491]
- Hellquist A, Zucchelli M, Kivinen K, Saarialho-Kere U, Koskenmies S, Widen E, Julkunen H, Wong A, Karjalainen-Lindsberg M-L, Skoog T, Vendelin J, Cunnihame-Graham DS, Vyse TJ, Kere J, Lindgren CM. The human GIMAP5 gene has a common polyadenylation polymorphism increasing risk to systemic lupus erythematosus (SLE). *J Med Genet*. 2007; 44(5):314–21. [PubMed: 17220214]
- Hernández-Hoyos G, Joseph S, Miller NG, Butcher GW. The lymphopenia mutation of the BB rat causes inappropriate apoptosis of mature thymocytes. *European Journal of Immunology*. 1999; 29(6):1832–41. [PubMed: 10382745]
- Hornum L, Romer J, Markholst H. The diabetes-prone BB rat carries a frameshift mutation in Ian4, a positional candidate of Iddm1. *Diabetes*. 2002; 51(6):1972–9. [PubMed: 12031988]
- Krücken J, Epe M, Benten WPM, Falkenroth N, Wunderlich F. Malaria-suppressible expression of the anti-apoptotic triple GTPase mGIMAP8. *Journal of Cellular Biochemistry*. 2005; 96(2):339–48. [PubMed: 16088918]

- Krucken J, Schroetel RMU, Muller IU, Saldani N, Marinovski P, Benten WPM, Stamm O, Wunderlich F. Comparative analysis of the human gimap gene cluster encoding a novel GTPase family. *Gene*. 2004; 341:291–304. [PubMed: 15474311]
- Lehmann, EL. Testing Statistical Hypotheses. John Wiley and Sons; New York: 1986.
- MacMurray AJ, Moralejo DH, Kwitek AE, Rutledge EA, Van Yserloo B, Gohlke P, Speros SJ, Snyder B, Schaefer J, Bieg S, Jiang J, Ettinger R, Fuller J, Daniels T, Pettersson A, Orlebeke K, Birren B, Jacob H, Lander E, Lernmark A. Lymphopenia in the BB rat model of Type 1 Diabetes is due to a mutation in a novel immune-associated nucleotide (Ian)-related gene. *Genome Res*. 2002; 12(7): 1029–39. [PubMed: 12097339]
- Malaisse WJ, Courtois P, Scott FW. Insulin-dependent diabetes and gut dysfunction: The BB rat model. *Horm Metab Res*. 2004; 36(9):585–94. [PubMed: 15486808]
- Michalkiewicz M, Michalkiewicz T, Ettinger RA, Rutledge EA, Fuller JM, Moralejo DH, VanYserloo B, MacMurray AJ, Kwitek AE, Jacob HJ, Lander ES, Lernmark A. Transgenic rescue demonstrates involvement of the Ian5 gene in T-cell development in the rat. *Physiol Genomics*. 2004; 19(2):228–32. [PubMed: 15328390]
- Nitta T, Nasreen M, Seike T, Goji A, Ohigashi I, Miyazaki T, Ohta T, Kanno M, Takahama Y. IAN family critically regulates survival and development of T lymphocytes. *PLoS Biology*. 2006; 4(4)
- Nitta T, Takahama Y. The lymphocyte guard-IANs: regulation of lymphocyte survival by IAN/ GIMAP family proteins. *Trends in Immunology*. 2007; 28(2):58–65. [PubMed: 17196432]
- Pandarpurkar M, Wilson-Fritch L, Corvera S, Markholst H, Hornum L, Greiner DL, Mordes JP, Rossini AA, Bortell R. Ian4 is required for mitochondrial integrity and T cell survival. *Proc Natl Acad Sci U S A*. 2003; 100(18):10382–7. [PubMed: 12930893]
- Payne F, Smyth DJ, Pask R, Barratt BJ, Cooper JD, Twells RC, Walker NM, Lam AC, Smink LJ, Nutland S, Rance HE, Todd JA. Haplotype tag single nucleotide polymorphism analysis of the human orthologues of the rat type 1 diabetes genes Ian4 (Lyp/Iddm1) and Cblb. *Diabetes*. 2004; 53(2):505–9. [PubMed: 14747305]
- Pinheiro, JC.; Bates, DM. Mixed-Effects Models in S and S-PLUS. Springer-Verlag; New York: 2000.
- Poirier GMC, Anderson G, Huvar A, Wagaman PC, Shuttleworth J, Jenkinson E, Jackson MR, Peterson PA, Erlander MG. Immune-Associated Nucleotide-1 (IAN-1) is a thymic selection marker and defines a novel gene family conserved in plants. *J Immunol*. 1999; 163(9):4960–9. [PubMed: 10528200]
- Reuber TL, Ausubel FM. Isolation of arabidopsis genes that differentiate between resistance responses mediated by the RPS2 and RPM1 disease resistance genes. *Plant Cell*. 1996; 8(2):241–9. [PubMed: 8742710]
- Rutledge EA, Fuller JM, Van Yserloo B, Moralejo DH, Ettinger RA, Gaur P, Hoehna JL, Peterson MR, Jensen R, Kwitek AE, Lernmark A. Sequence variation and expression of the Gimap gene family in the BB Rat. *Exp Diabetes Res*. 2009; 835650:1–10.
- Sandal T, Aumo L, Hedin L, Gjertsen BT, Doskeland SO. Irod/Ian5: an inhibitor of gamma-radiation- and okadaic acid-induced apoptosis. *Mol Biol Cell*. 2003; 14(8):3292–304. [PubMed: 12925764]
- Schnell S, Demolliere C, van den Berk P, Jacobs H. Gimap4 accelerates T-cell death. *Blood*. 2006; 108(2):591–9. [PubMed: 16569770]
- Schulteis RD, Chu H, Dai X, Chen Y, Edwards B, Haribhai D, Williams CB, Malarkannan S, Hessner MJ, Glisic-Milosavljevic S, Jana S, Kerschen EJ, Ghosh S, Wang D, Kwitek AE, Lernmark A, Gorski J, Weiler H. Impaired survival of peripheral T cells, disrupted NK/NKT cell development, and liver failure in mice lacking Gimap5. *Blood*. 2008; 112(13):4905–14. [PubMed: 18796632]
- Shin JH, Janer M, McNeney B, Blay S, Deutsch K, Sanjeevi CB, Kockum I, Lernmark A, Graham J. Groups obotSCDatDliSS. IA-2 autoantibodies in incident type I diabetes patients are associated with a polyadenylation signal polymorphism in GIMAP5. *Genes Immun*. 2007; 8(6):503–12. [PubMed: 17641683]
- Vaarala O. Leaking gut in type 1 diabetes. *Current Opinion in Gastroenterology*. 2008; 24(6):701–6. [PubMed: 19122519]

Fig. 1. The Gimap Gene Family on RNO4

An expanded map of the 2 Mb of DP DNA in the congenic DR.*lyp* rat line is shown along with the *Gimap* family chromosomal locations on RNO4. The 33 Kb *lyp* critical interval is indicated between the SSLP markers D4Rhw6 and IIsnp3. Arrows indicate the orientation of the full length mRNA with exons shown in black and introns in white. *Ian* aliases are in parentheses underneath the corresponding *Gimap* name.

Fig. 2. T cell Profile and Purity

Scatter plots are shown from one DR.+/+ and one DR.lyp/lyp rat FACS sorted T cells from thymus, spleen and MLN following staining with CD4 and CD8 monoclonal antibodies. Cells from spleen and MLN were first gated based on staining with a CD3 monoclonal antibody. The percent gated cells are shown within each quadrant. Panels A–D show the purity of sorted cells from one DR.lyp/lyp rat.

Fig. 3. *Gimap* Gene Expression in *DR*^{+/+} and *DR*^{lvp/lvp} Sorted Thymic Cells

The mean $\pm$ standard error of the mean (SEM) is shown for *Gimap* family expression in CD4 and CD8 FACS sorted thymocytes following normalization to GAPDH. DN is double negative (CD4⁻CD8⁻) and DP is double positive (CD4⁺CD8⁺). Black columns represent *DR*^{+/+} rats (+, n=3) and hatched columns represent *DR*^{lvp/lvp} (l, n=3). Significant differences are * for p<0.05 and ** for p<0.001. Genes appear in the order at which they appear on RNO4.

Fig. 4. Gimap Gene Expression in Peripheral T cells

The mean $\pm$ standard error of the mean (SEM) is shown for *Gimap* family expression in CD4⁺ and CD8⁺ FACS sorted T cells from spleen (Spl) and MLN following normalization to GAPDH. Cells from spleen and MLN were first gated based on staining with a CD3 monoclonal antibody. Black columns represent DR^{+/+} rats (+, n=3) and hatched columns represent DR^{lyp/lyp} (l, n=3). Significant differences are * for $p < 0.05$ and ** for $p < 0.001$. Genes appear in the order at which they appear on RNO4.

Fig. 5. Gimap Gene Expression in Peripheral B cells

The mean $\pm$ standard error of the mean (SEM) is shown for *Gimap* family expression in FACS sorted hematopoietic B cells (CD45+CD45RA+) and *non* B cells (CD45+CD45RA-) from spleen and MLN after normalization to GAPDH. Black columns represent DR.+/+ rats (+, n=3) and hatched columns represent DR. lyp/lyp (l, n=3). Significant differences are * for p<0.05. Genes appear in the order at which they appear on RNO4.