Published in final edited form as:

Biochem Pharmacol. 2011 October 15; 82(8): 873–882. doi:10.1016/j.bcp.2011.06.001.

Role of $\alpha 6$ nicotinic receptors in CNS dopaminergic function; relevance to addiction and neurological disorders

Maryka Quik¹, Xiomara A. Perez¹, and Sharon R. Grady²

¹Center for Health Sciences, SRI International, 333 Ravenswood Ave, California 94025, USA

²Institute for Behavioral Genetics, University of Colorado, 447UCB, Boulder, CO 80409, USA

Abstract

Although a relative newcomer to the nicotinic acetylcholine receptor (nAChR) family, substantial evidence suggests that $\alpha 6$ containing nAChRs play a key role in CNS function. This subtype is unique in its relatively restricted localization to the visual system and catecholaminergic pathways. These latter include the mesolimbic and nigrostriatal dopaminergic systems, which may account for the involvement of $\alpha 6$ containing nAChRs in the rewarding properties of nicotine and in movement. Here, we review the literature on the role of $\alpha 6$ containing nAChRs with a focus on the striatum and nucleus accumbens. This includes molecular, electrophysiological and behavioral studies in control and lesioned animal models, as well as in different genetic models. Converging evidence suggest that the major $\alpha 6$ containing nAChRs subtypes in the nigrostriatal and mesolimbic dopamine system are the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ nAChR populations. They appear to have a dominant role in regulating dopamine release, with consequent effects on nAChR-modulated dopaminergic functions such as reinforcement and motor behavior. Altogether these data suggest that drugs directed to $\alpha 6$ containing nAChRs may be of benefit for the treatment of addiction and for neurological disorders with locomotor deficits such as Parkinson's disease.

Keywords

Addiction; Cyclic voltammetry; Nucleus accumbens; Parkinson's disease; Striatum

1. Introduction

Knowledge concerning the presence and function of nicotinic acetylcholine receptors (nAChRs) throughout biological systems has increased significantly over the last century. Extensive evidence now indicates that multiple nAChR subtypes exist in the mammalian central and peripheral nervous systems, as well as in other tissues. The muscle nAChR, which is composed of 4 distinct subunits ($\alpha 1\beta 1\gamma\delta$), was the first to be studied in detail in the early 1970's because of its relative abundance in *Torpedo* electric organ and the isolation of the selective ligand α -bungarotoxin [1, 2]. This was followed, several decades later, by the discovery of the $\alpha 3\beta 4^*$, $\alpha 4\beta 2^*$ and $\alpha 7$ nAChR populations, which are mainly present in

© 2011 Elsevier Inc. All rights reserved.

Corresponding author; Maryka Quik, Center for Health Sciences, SRI International, 333 Ravenswood Ave, California 94025, USA. Tel. 1-650-859-6112; Fax 1-650-859-5099. maryka.quik@sri.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

neuronal tissues [3–5]. The asterisk indicates the possible presence of other subunits in the receptor complex.

nAChRs expressing the $\alpha 6$ subunit were one of the last to be identified, with cloning of the $\alpha 6$ nAChR subunit only reported in the 1990's [6]. Progress in understanding its functional significance was slow because of difficulties in expressing receptors containing the $\alpha 6$ subunit ($\alpha 6^*$ nAChRs) in cell models and the lack of suitable probes for its characterization. However, subsequent advances in these areas, especially the discovery of the selective antagonist α -conotoxinMII (α -CtxMII) (Cartier et al, 1996), have greatly enhanced our basic knowledge concerning $\alpha 6^*$ nAChRs.

One point of note is that $\alpha 6^*$ nAChRs are not present in the peripheral nervous system and exhibit a relatively restricted distribution in the mammalian CNS [7–12]. This receptor shares the same basic motif as other nAChR subtypes, that is, it is a ligand gated ion channel consisting of five subunits around a central pore that lie within the membrane bilayer. Great strides have been made over the last few years in our understanding of the subunit composition of $\alpha 6^*$ nAChRs, their role in CNS dopaminergic function and the ensuing behavioral consequences. These combined findings, which are the focus of the current review, suggest that $\alpha 6^*$ nAChRs may represent key therapeutic targets for the treatment of addiction and for neurodegenerative movement disorders involving deficits in the nigrostriatal dopaminergic system.

1.1. Localization of $\alpha 6^*$ nAChRs in the brain

In situ hybridization studies done to identify the presence of $\alpha 6$ nAChR subunit mRNA were the first to suggest that receptors containing the $\alpha 6$ subunit exhibited a very restricted localization in both rodent and primate brain. The strongest $\alpha 6$ mRNA signal is in the retina and catecholaminergic nuclei, including the locus coeruleus, ventral tegmental area, and substantia nigra, with a weaker signal in the superior colliculus, medial habenula, interpeduncular nucleus, visual cortex and a few other small nuclei [12–16]. This limited distribution of $\alpha 6^*$ nAChR transcripts contrasts with that of the $\alpha 4^*$ and $\alpha 7$ nAChR mRNAs, which are widespread throughout the brain [13, 14, 17].

The mRNA distribution studies were followed by experiments to localize $\alpha 6^*$ nAChR protein (Table 1). Such work became feasible because of the isolation of α -CtxMII, a novel snail toxin that was first shown to bind to $\alpha 3\beta 2^*$ nAChRs [18] and was subsequently shown to have even higher affinity for $\alpha 6\beta 2^*$ nAChRs [10]. Another breakthrough was the development of $\alpha 6$ nAChR-subunit directed antibodies [19, 20]. The combined use of these two different probes showed that $\alpha 6^*$ nAChR protein is present in the mesolimbic and nigrostriatal dopaminergic pathways, consistent with the existence of $\alpha 6$ mRNA in catecholaminergic nuclei. $\alpha 6^*$ nAChRs are also very prominent in structures linked to the visual system including the superior colliculus, optic tract, olivary pretectal area, and mediolateral and dorsolateral geniculate nuclei, as well as the retina [20–25]. In addition, small populations of $\alpha 6^*$ nAChRs have been identified in the rodent habenulo-interpeduncular pathway [26].

This unique anatomical distribution suggests that $\alpha 6^*$ nAChRs may play a role in behaviors linked to reward, motor activity and vision.

1.2. Composition of $\alpha 6^*$ nAChRs

1.2.1. CNS dopaminergic pathways—Our knowledge concerning the subunit composition of $\alpha 6^*$ nAChRs in dopaminergic pathways has grown considerably as a result of a concerted effort using multiple experimental strategies. Immunoprecipitation studies

with antibodies targeting $\beta 2$ nAChRs, coupled with the use of $\beta 2$ knockout mice, demonstrated the presence of the $\beta 2$ subunit in all $\alpha 6^*$ nAChR complexes in rodent and monkey striatum and/or nucleus accumbens (Fig. 1 and Table 1) [19, 20, 27, 28]. The $\beta 3$ subunit also appears to be expressed in the majority of $\alpha 6^*$ nAChR complexes (Fig. 1 and Table 1). Indirect evidence for this possibility initially stemmed from the observation that $\alpha 6$ and $\beta 3$ mRNAs were uniquely co-localized in the same brain regions [14, 29]. Subsequent dual immunoprecipitation studies using $\alpha 6$ and $\beta 3$ nAChR subunit-directed antibodies confirmed this idea by showing that the $\alpha 6$ and $\beta 3$ subunit proteins coexist within the same receptor complex. Lesion studies further showed that $\alpha 6$ and $\beta 3$ nAChR subunit protein declined in parallel with nigrostriatal damage [19, 20, 27, 28, 30]. It should be noted that there is not a one-to-one correspondence in the decline of receptors expressing these subunits in nAChR knockout mice [29]. This may be due to the development of compensatory mechanisms in knockout mice not normally present in wildtype animals.

In addition to the presence of the $\beta 2$ and $\beta 3$ nAChR subunits, a significant proportion of $\alpha 6^*$ nAChR complexes also include the $\alpha 4$ subunit (Fig. 1 and Table 1) [20, 27, 28, 31]. The relative levels of the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ subtypes in the nigrostriatal and mesolimbic dopamine system are currently under investigation. One study reports the existence of roughly equal amounts of the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ subtypes in the striatum [32], while another suggests a predominance of the $\alpha 6\alpha 4\beta 2\beta 3$ subtype [28]. In ventral striatum, which includes the nucleus accumbens, ventral pallidum and olfactory tubercle, the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ nAChR subtypes are present in similar proportions [28]. Although the mesolimbic dopaminergic system also projects to the frontal cortex, the primary nAChRs in this latter region appear to be the $\alpha 4\beta 2^*$ and $\alpha 7$, with a role for the $\alpha 6\beta 2^*$ subtype currently uncertain [33, 34].

Overall, these data indicate that the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ subtypes are the major subtypes in the nucleus accumbens and striatum, although there is always the possibility that minor novel subtypes remain to be identified. Since $\alpha 6$ containing receptors in the mesolimbic and nigrostriatal systems presently appear to contain a $\beta 3$ subunit, they will be designated as $\alpha 6\beta 2\beta 3^*$ nAChRs in subsequent sections of the review.

1.2.1. Visual system—Although $\alpha 6^*$ nAChRs have been studied most extensively in CNS dopaminergic pathways, it should be noted that they are in fact most densely expressed in areas linked to the visual system, as well as the superior colliculus, optic tract, geniculate nuclei and the retina, including microglial cells [21, 24, 25, 35]. Sequential immunoprecipitation studies with select nAChR subunit antibodies indicate that considerable subtype heterogeneity exists in regions of the mammalian visual system (Fig. 1 and Table 1), with expression of $\alpha 6\beta 2^*$, $\alpha 6\alpha 3\beta 2^*$, $\alpha 6\alpha 2\beta 2^*$, $\alpha 6\beta 2\beta 3^*$, $\alpha 6\alpha 4\beta 2^*$, $\alpha 6\alpha 3\alpha 4\beta 2^*$ identified to date [11, 23, 24]. This diversity in $\alpha 6^*$ nAChR composition in the visual system may be important for the fine-tuning of this essential function under varying physiological and pathological conditions.

2. $\alpha 6\beta 2\beta 3^*$ nAChRs and the control of dopaminergic function

Although the most intense expression of $\alpha 6^*$ nAChRs is in the visual system, very little is known about their function in vision. By contrast, significant work has been done to investigate their involvement in dopaminergic activity in the terminal regions of both the mesolimbic and nigrostriatal pathways, that is, the nucleus accumbens and striatum. Two approaches that have greatly contributed to our understanding of their role in these latter regions are measurement of nAChR-stimulated ^3H -dopamine release from synaptosomes or slices, and electrically-evoked dopamine release from slices using fast scan cyclic voltammetry.

2.1. nAChR-evoked ^3H -dopamine release

The concept that nAChRs could modulate dopamine release initially stemmed from the work of Giorgiueff and coworkers, who investigated nAChR-mediated release of newly synthesized ^3H -dopamine from ^3H -tyrosine [36]. This was followed several years later by experiments showing that nicotine stimulated release of ^3H -dopamine from nucleus accumbens or striatal synaptosomes [37–39]. For about a decade, dopamine release in both the nucleus accumbens and striatum was thought to be regulated only by $\alpha 4\beta 2^*$ nAChRs. However, the discovery of α -CtxMII, which interacts with $\alpha 6\beta 2\beta 3^*$ nAChRs, subsequently showed that activation of this subtype also stimulated ^3H -dopamine release [34, 40, 41]. In fact, the $\alpha 4\beta 2^*$ and $\alpha 6\beta 2\beta 3^*$ nAChR subtypes promote dopamine release evoked by 1 μM nicotine in roughly equal proportions in rodent nucleus accumbens and striatum (Fig. 2) [34, 40, 41]. Furthermore, the $\alpha 6\beta 2\beta 3^*$ nAChR subtype plays a dominant role in nonhuman primate nucleus accumbens and striatum (Fig. 2), with 80% of release mediated by $\alpha 6\beta 2\beta 3^*$ subtypes and only 20% by the $\alpha 4\beta 2^*$ nAChR population [42, 43]. It should be noted that the functional data regarding our knowledge of $\alpha 6\beta 2\beta 3^*$ nAChRs described above, as well as in the next section, was obtained with a single, antagonist α -CtxMII, because this is currently the only selective agent available for their study. Work with other selective antagonists, antibodies or alternate molecular probes, is necessary for confirmation of the role of $\alpha 6\beta 2\beta 3^*$ nAChRs in dopaminergic function.

2.2. Cyclic voltammetry

Another approach that has yielded significant insight concerning the role of $\alpha 6\beta 2\beta 3^*$ nAChRs in dopaminergic function is cyclic voltammetry, in which carbon-fiber microelectrodes are used as chemical sensors to detect electrically evoked dopamine release in tissue slices [44–50]. This approach is distinct from nAChR-stimulated ^3H -dopamine release as it provides a measure of action-potential evoked endogenous dopamine release. Moreover, it has the advantage that it provides information about release with single and multiple electrical pulses, which may more closely resemble tonic and burst stimulation with subsequent reward and movement.

One strategy currently being used to evaluate the role of different nAChR subtypes in modulating dopamine release involves the use of nAChR antagonists. In the mesolimbic dopaminergic system, the $\alpha 6\beta 2\beta 3^*$ nAChR selective antagonist α -CtxMII reduced single-pulse stimulated dopamine release by the same amount as general nAChR antagonists in nucleus accumbens slices from rodents [44] and monkeys [51]. These data indicate that the prime regulator of dopamine release in the mesolimbic dopamine pathway is the $\alpha 6\beta 2\beta 3^*$ nAChR.

The role of the $\alpha 6\beta 2\beta 3^*$ subtype was somewhat less pronounced in the nigrostriatal dopaminergic system. However, it still exerted major effects, regulating about 50% of single-pulse evoked dopamine release in the dorsal striatum of rodents (Fig. 3) [44, 46, 48] and ventral putamen of nonhuman primates (Fig. 3) [47, 50]. Results in nonhuman primate dorsal putamen were similar to those in the mesolimbic dopamine systems with all single-pulse nAChR-mediated evoked dopamine release modulated by $\alpha 6\beta 2\beta 3^*$ nAChRs.

As might be expected, nAChR modulation of burst firing is much more complex, with an increase or no apparent change in dopamine release with either $\alpha 6\beta 2^*$ or $\alpha 4\beta 2^*$ nAChR antagonism [44–47, 50, 52–55]. This may be attributed to the fact that the effects of acetylcholine with burst stimulation involve a complex interplay between nAChR desensitization and activation, as well as muscarinic AChR activation, all of which would be dependent on the concentration of acetylcholine at the synaptic cleft.

Altogether these studies suggest that $\alpha 6\beta 2\beta 3^*$ nAChRs are major modulators of dopaminergic function in both striatum and nucleus accumbens of rodents and monkeys. The relative roles of the $\alpha 6\beta 2\beta 3$ and $\alpha 4\alpha 6\beta 2\beta 3$ subtypes are currently unclear; however, novel approaches involving concatameric $\alpha 6^*$ nAChRs [56] and/or the development of selective drugs should help in this regard. Overall, $\alpha 6\beta 2\beta 3^*$ nAChRs appear to play a prominent role in evoked dopamine release, and consequently drug reinforcement and motor function.

3. Long term nicotine exposure modulates $\alpha 6\beta 2\beta 3^*$ nAChR expression and function

Although acetylcholine or nAChR agonists acutely increase dopamine release, it is well established that continued nicotine exposure rapidly (seconds to minutes) results in a decline in release due to receptor desensitization [57, 58]. Sustained nicotine exposure (days to months) leads to additional long term molecular and cellular changes which may represent mechanisms that underlie nicotine addiction and tolerance [59, 60]. It is therefore important to understand the effects of chronic nicotine treatment on nAChR subtype expression and function.

Nicotine administration is well known to increase $\alpha 4\beta 2^*$ nAChR expression throughout the brain, including the striatum, in experimental animal models and in human smokers [61–69]. Moreover, functional studies in rodents and primate show that the elevated $\alpha 4\beta 2^*$ nAChR levels in striatum may be associated with increases or no change in receptor-mediated function (Table 2) [70, 71]. By contrast, the same long term nicotine treatment paradigms decrease or do not change $\alpha 6\beta 2\beta 3^*$ nAChR expression in rodent and monkey striatum (Table 2) [42, 64, 72–74]. Nicotine-induced changes in $\alpha 6\beta 2\beta 3^*$ nAChR levels generally correlate with declines in ^3H -dopamine release from striatal and nucleus accumbens synaptosomes (Table 2). Subsequent work to distinguish between changes in the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ subtypes showed that chronic nicotine dosing to mice led to selective decline in $\alpha 6\alpha 4\beta 2\beta 3$ nAChRs with some increase in $\alpha 6\beta 2\beta 3$ nAChRs [48]. These results are consistent with data using transfected cell culture systems, which show that $\alpha 6\beta 2$ or $\alpha 6\beta 2\beta 3$ nAChRs can be upregulated with long term nicotine treatment [75–78].

Cyclic voltammetry has also been used to evaluate changes in dopaminergic activity with long term nicotine dosing [46, 48]. Single-pulse evoked dopamine release in the presence of α -CtxMII was still reduced with chronic nicotine treatment. However, nicotine exposure modified the pattern of burst firing observed in the presence of α -CtxMII. In untreated rats, burst firing is enhanced or not changed in the presence of α -CtxMII. By contrast, in nicotine-treated rats, there was a decrease in burst-stimulated dopamine release in the presence of α -CtxMII. The finding that nicotine exposure preferentially downregulates the $\alpha 6\alpha 4\beta 2\beta 3$ subtype in striatum [48], coupled with the release results, suggest that the $\alpha 6\alpha 4\beta 2\beta 3$ nAChR subtype is a primary regulator of evoked dopamine release under physiological (untreated control) conditions. The differential burst-stimulated release pattern with chronic nicotine treatment may be due to a more prominent role of the $\alpha 6\beta 2\beta 3$ subtype after nicotine treatment [48].

Chronic nicotine exposure also altered striatal dopaminergic function in nonhuman primate striatum. However, the changes were different from those obtained in rat striatum and also exhibited regional variations [47]. Nicotine treatment via the drinking water (3 to 4 months) altered both nonburst and burst-stimulated dopamine release in ventral but not dorsal putamen [47]. In addition, chronic nicotine treatment resulted in large declines in dopaminergic function in monkey nucleus accumbens with a complete loss of $\alpha 6\beta 2\beta 3^*$ nAChR- modulated evoked dopamine release and a depression of non-nAChR-modulated

evoked dopamine release [51]. Comparable studies have not yet been done to evaluate the effects of chronic nicotine treatment on dopamine release in rat nucleus accumbens using cyclic voltammetry.

Overall, these findings indicate that chronic nicotine exposure may result in dramatic alterations in $\alpha 6\beta 2\beta 3^*$ nAChR-modulated dopamine release in the striatum and nucleus accumbens. These mechanisms are of relevance to the chronic nicotine exposure that occurs with smoking in humans, and may underlie the reinforcing and addictive properties of smoking.

4. Role of mesolimbic $\alpha 6\beta 2\beta 3^*$ nAChRs in reward and addiction

4.1. Nicotine addiction

An extensive clinical and preclinical literature over the last few decades supports the idea that nicotine in tobacco contributes to its addictive properties [79–85]. Molecular, cellular and behavioral studies using various animal models have provided significant insight into the nAChR subtypes relevant to addiction [81, 86–88]. Initial work with nAChR blockers and selective lesions showed that the reinforcing effects of nicotine were mediated via nAChRs in the mesolimbic dopaminergic pathway [83, 84]. The subsequent use of nAChR knockout and transgenic mice, coupled with receptor re-expression, proved critical in defining the nAChR subtypes involved. Lentiviral vector re-expression of the $\beta 2$ subunit into the ventral tegmental area of $\beta 2$ knockout mice demonstrated a key role for $\beta 2^*$ nAChRs in nicotine reinforcement [89, 90]. Work with genetically engineered mice containing a point mutation in $\alpha 4$ nAChR subunits (Leu9' to Ala9' in the M2 domain) that yielded receptors hypersensitive to nicotine showed that the $\alpha 4\beta 2^*$ nAChR subtype is sufficient for nicotine-induced reward, tolerance, and sensitization [91]. Evidence for a role for $\alpha 6^*$ nAChRs stemmed from studies showing that $\alpha 6$ nAChR knockout mice did not self-administer nicotine, but that re-expression of the $\alpha 6$ subunit restored this behavior [92]. Moreover, intraventricular or local infusion of $\alpha 6\beta 2\beta 3^*$ nAChR selective toxins into the rat nucleus accumbens shell blocked behaviors linked to nicotine reward and withdrawal [93, 94]. Studies with mice expressing hypersensitive $\alpha 6\beta 2\beta 3^*$ nAChRs (with $\alpha 6$ subunit Leu 9' to Ser 9' mutation) but expressing or lacking the $\alpha 4$ subunit further suggest that an $\alpha 6\alpha 4\beta 2^*$ nAChR subtype may play an important role in nicotine reward and reinforcement [95, 96].

Thus, converging data from numerous animal models indicate that $\alpha 6\beta 2\beta 3^*$ nAChR, possibly in concert with other nAChR subtypes [97], are important for nicotine's addictive properties. These findings suggest that selective $\alpha 6\beta 2\beta 3^*$ nAChR agonists and antagonists would represent novel agents for the therapeutic management of nicotine addiction.

4.2. Alcohol addiction

Another widely used neuroactive substance with addictive properties is ethanol. Although its mechanism of action is most likely multifactorial, ethanol appears to exert some of its reinforcing effects by activating dopaminergic neurons in the ventral tegmental area through an interaction at nAChRs [98]. This idea is based in part on studies with varenicline, an agonist/partial agonist that acts at multiple subtypes including $\alpha 3\beta 4^*$, $\alpha 4\beta 2^*$, $\alpha 6\beta 2\beta 3^*$ and $\alpha 7$ nAChRs [98–101]. Varenicline decreases alcohol consumption in rats [102–104], as well as in heavy-drinking smokers [105]. With respect to the nAChR subtype involved, studies with genetically modified nAChR mice indicate that $\alpha 4\beta 2^*$ nAChRs play a role in alcohol addiction [106]. In addition, $\alpha 3\beta 4^*$ partial agonists reduce ethanol consumption and seeking in rats [107]. The $\alpha 6\beta 2\beta 3^*$ subtype has also been implicated based on experiments showing that intracranial injection of α -CtxMII into the ventral tegmental area decreased ethanol-induced accumbal dopamine release [108].

Thus multiple nAChR subtypes in the mesolimbic dopaminergic system may be involved in the activating and reinforcing effects of ethanol. Drugs targeting $\alpha 6\beta 2\beta 3^*$, as well as $\alpha 4\beta 2^*$ and $\alpha 3\beta 4^*$ nAChR subtypes may thus have potential in modulating addictive behaviors associated with ethanol consumption.

5. Role of nigrostriatal $\alpha 6\beta 2\beta 3^*$ nAChRs in motor control

5.1. Studies in nonlesioned animal models

The presence of $\alpha 6\beta 2\beta 3^*$ nAChRs in the nigrostriatal pathway, coupled with the well known association between this system and movement, prompted studies to investigate a role for striatal $\alpha 6\beta 2\beta 3^*$ nAChRs in locomotor activity. The first report to explore this possibility was that of Le Novere and coworkers [109] who showed that $\alpha 6$ antisense oligonucleotide infusion into the CNS decreased nicotine-mediated locomotor activity. In addition, transgenic mice expressing hypersensitive $\alpha 6\beta 2\beta 3^*$ receptors exhibit altered locomotor responsiveness with increased walking, turning, and rearing [95, 110]. These genetically modified mice express an $\alpha 6$ subunit in which Leu280 in the M2 domain ($\alpha 6$ L9'S mice) was mutated to Ser rendering an $\alpha 6^*$ receptor with enhanced sensitivity to nicotine. By contrast, mice that lack the $\alpha 4$ nAChR subunit behaved normally despite the presence of a hyperactive $\alpha 6$ nAChR subunit, suggesting a critical involvement of the $\alpha 6\alpha 4\beta 2\beta 3$ nAChR subtype in motor activity [95].

These combined data indicate that striatal $\alpha 6\alpha 4\beta 2\beta 3$ nAChRs play an important role in movement. They most likely influence motor function in concert with $\alpha 6\beta 2\beta 3$, $\alpha 4\beta 2^*$ and $\alpha 7$ nAChRs in the striatum, as well as other brain areas including the mesolimbic dopaminergic system [28, 111, 112].

5.2. Effect of nigrostriatal damage; relevance to Parkinson's disease

The next question that arises is the effect of neuronal damage on $\alpha 6\beta 2\beta 3^*$ nAChR expression, cellular function and behavior. This is of particular relevance to Parkinson's disease, a neurological movement disorder characterized by severe losses in nigrostriatal dopaminergic neurons. To approach this, work has been done in parkinsonian animal models in which the nigrostriatal pathway is selectively damaged with dopaminergic neurotoxins such as 6-hydroxydopamine or MPTP. Such lesions result in a decrease in $\alpha 6\beta 2\beta 3^*$ nAChR expression and function that closely parallels the decline in dopaminergic terminal integrity [19, 20, 22, 113, 114]. These findings indicate that $\alpha 6\beta 2\beta 3^*$ nAChRs are primarily localized to dopaminergic terminals in the striatum. Nigrostriatal damage also reduced $\alpha 4\beta 2^*$ nAChRs levels and function. However, the decreases in this subtype were much smaller than those in $\alpha 6\beta 2\beta 3^*$ nAChRs suggesting that $\alpha 4\beta 2^*$ nAChRs are also present on other neuronal elements in the striatum [19, 20, 114, 115].

Results in parkinsonian animal models parallel those in Parkinson's disease brains. Large declines are observed in $\alpha 6\beta 2\beta 3^*$ nAChRs in striatum of Parkinson's disease patients, which correlate with the magnitude of the dopamine transporter loss [116–118]. By contrast, striatal $\alpha 4\beta 2^*$ nAChRs are reduced to a lesser extent, in agreement with studies in rodent and nonhuman primate striatum. To further delineate effects on select nAChR populations, competition studies were done with α -CtxMII [E11A] that differentiates between $\alpha 6\beta 2\beta 3^*$ subtypes (Fig. 4). A preferential loss of the striatal $\alpha 6\alpha 4\beta 2\beta 3$ nAChR subtype was observed with mild to moderate nigrostriatal damage, while the $\alpha 6\beta 2\beta 3$ subtype was only decreased with more severe nigrostriatal damage. This was observed across species, that is, in the striatum of parkinsonian rodents and in monkeys, as well as with Parkinson's disease (Fig. 4) [32]. These data suggest that drugs targeting both the $\alpha 6\alpha 4\beta 2\beta 3$ and $\alpha 6\beta 2\beta 3$ nAChR subtypes may be useful with moderate nigrostriatal damage, but that $\alpha 6\beta 2\beta 3$ nAChRs drugs may be more suitable with severe degeneration.

5.3. Role for selective nAChR drugs in Parkinson's disease

Studies have been done in parkinsonian animal models, which suggest that nicotine or nAChR agonist treatment may provide therapeutic benefit in Parkinson's disease. Extensive evidence has shown that nicotine protects against nigrostriatal damage through an interaction at nAChRs in several parkinsonian animal models including mice, rats and monkeys [119–121]. Further studies investigating the nAChR subtypes that contribute to this effect suggest that protection against nigrostriatal damage may be linked to the striatal $\alpha 6\alpha 4\beta 2\beta 3$ nAChR population [122].

There have also been reports that nicotine may reduce a serious side effect of L-dopa treatment for Parkinson's disease, that is, L-dopa-induced abnormal involuntary movements (AIMs) or dyskinesias [123–127]. There are currently only limited treatment options for these abnormal movements of the head, trunk and/or limbs, which develop in most patients on L-dopa therapy [128]. Recent work shows that nicotine treatment ameliorates L-dopa induced dyskinesias in parkinsonian mice, rats and monkeys by acting at nAChRs [129–132]. Because nicotine is nonselective and may result in side effects in addition to the desired response, studies were done to determine whether more selective nAChR agonists improve these abnormal movements in a rat model of L-dopa-induced dyskinesias. The drugs tested include varenicline, an agonist that interacts with multiple nAChRs such as the $\alpha 4\beta 2^*$, $\alpha 6\beta 2\beta 3^*$ and $\alpha 7$ subtypes [99, 133], and iodo-A-85380, an agonist that interacts with $\alpha 4\beta 2^*$ and $\alpha 6\beta 2^*$ nAChRs [134]. Both these compounds reduced L-dopa-induced dyskinesic-like movements by 50% in parkinsonian rats [135]. Since these drugs reduced AIMs more effectively in rats with partial compared to near-complete nigrostriatal damage, these data suggest that presynaptic $\alpha 4\beta 2^*$ and $\alpha 6\beta 2\beta 3^*$ nAChRs on dopaminergic terminals may play a role in reducing L-dopa-induced AIMs but also with a contribution from other nAChRs located throughout the brain.

The mechanisms whereby nAChR agonists improve L-dopa-induced AIMs remain to be elucidated. Current experimental evidence suggests that they most likely involve long term molecular changes in the brain since several days of agonist administration is required to reduce AIMs [135]. The site of action of these long term changes is currently not known. One might anticipate that agonists exert their beneficial effect by initially modulating nigrostriatal function. This may be followed by secondary actions in other CNS pathways, for instance, in the serotonergic system [136–138]. Evidence for this idea is derived from studies showing that L-dopa-induced AIMs are substantially reduced in rodents with serotonergic lesions. The idea that multiple brain regions may subsequently mediate the effects of nAChR activation is also suggested from an extensive literature showing that deep brain stimulation of the subthalamic nucleus is very effective in reducing dyskinesias in Parkinson's disease patients [139].

6. Role for $\alpha 6^*$ nAChRs in the visual system

The intense expression of diverse $\alpha 6^*$ nAChRs in the mammalian visual system and retina (Table 1) suggests that the $\alpha 6^*$ nAChR subtype most likely has a significant role in vision [11, 35]. Experiments done to evaluate the effect of visual deprivation on nAChRs showed no changes in $\alpha 6^*$ nAChR expression in rats reared in the dark [35], although eye enucleation did reduce $\alpha 6^*$ nAChRs in superior colliculus and lateral geniculate nucleus [23]. However, the function of $\alpha 6^*$ nAChRs in vision is currently unknown. Acetylcholine acting at nAChRs has been shown to modulate neuronal excitability in the visual cortex [140], while studies with nAChR blockers suggest there may be a permissive role for cholinergic input in visual function [141].

The intense localization of $\alpha 6^*$ nAChRs in the visual system, coupled with the importance of vision in daily life, suggests that this could represent a promising new research area for future studies.

7. Summary

Accumulating literature points to a significant and possibly predominant role for $\alpha 6\beta 2\beta 3^*$ nAChRs in modulating dopaminergic function in the mesolimbic and nigrostriatal dopaminergic pathways. Since these systems are critically involved in behaviors linked to addiction/reward and the control of movement, $\alpha 6\beta 2\beta 3^*$ nAChRs may represent novel therapeutic targets for addiction and for the treatment of movement disorders such as Parkinson's disease.

Abbreviations

α -CtxMII	α -conotoxinMII
nAChR	nicotinic acetylcholine receptor

Acknowledgments

This work was supported by NIH grants NS42091, NS59910, NS65851, DA003194 and DA012242, grants from the Michael J Fox Foundation and from the California Tobacco Related Disease Research Program 17RT-0119.

References

1. Albuquerque EX, Pereira EF, Alkondon M, Rogers SW. Mammalian nicotinic acetylcholine receptors: from structure to function. *Physiological reviews*. 2009; 89:73–120. [PubMed: 19126755]
2. Changeux JP, Kasai M, Lee CY. Use of a snake venom toxin to characterize the cholinergic receptor protein. *Proceedings of the National Academy of Sciences of the United States of America*. 1970; 67:1241–1247. [PubMed: 5274453]
3. Boulter J, Connolly J, Deneris E, Goldman D, Heinemann S, Patrick J. Functional expression of two neuronal nicotinic acetylcholine receptors from cDNA clones identifies a gene family. *Proc Natl Acad Sci U S A*. 1987; 84:7763–7767. [PubMed: 2444984]
4. Boulter J, Evans K, Goldman D, Martin G, Treco D, Heinemann S, et al. Isolation of a cDNA clone coding for a possible neural nicotinic acetylcholine receptor alpha-subunit. *Nature*. 1986; 319:368–374. [PubMed: 3753746]
5. Boulter J, Luyten W, Evans K, Mason P, Ballivet M, Goldman D, et al. Isolation of a clone coding for the alpha-subunit of a mouse acetylcholine receptor. *J Neurosci*. 1985; 5:2545–2552. [PubMed: 2993547]
6. Lamar E, Miller K, Patrick JW. *Society for Neuroscience Abstracts*. 1990
7. Yang KC, Jin GZ, Wu J. Mysterious alpha6-containing nAChRs: function, pharmacology, and pathophysiology. *Acta Pharmacol Sin*. 2009; 30:740–751. [PubMed: 19498417]
8. Tapper AR. alpha6-containing nicotinic receptors unmasked in vivo: selective dopaminergic neuronal expression and activation in motor and reward pathways. *Cellscience*. 2009; 5:1742–1752.
9. Calabresi P, Di Filippo M. ACh/dopamine crosstalk in motor control and reward: a crucial role for alpha 6-containing nicotinic receptors? *Neuron*. 2008; 60:4–7. [PubMed: 18940582]
10. Quik M, McIntosh JM. Striatal alpha6* nicotinic acetylcholine receptors: potential targets for Parkinson's disease therapy. *J Pharmacol Exp Ther*. 2006; 316:481–489. [PubMed: 16210393]
11. Millar NS, Gotti C. Diversity of vertebrate nicotinic acetylcholine receptors. *Neuropharmacology*. 2009; 56:237–246. [PubMed: 18723036]
12. Champiaux N, Han ZY, Bessis A, Rossi FM, Zoli M, Marubio L, et al. Distribution and pharmacology of alpha 6-containing nicotinic acetylcholine receptors analyzed with mutant mice. *J Neurosci*. 2002; 22:1208–1217. [PubMed: 11850448]

13. Le Novère N, Zoli M, Changeux JP. Neuronal nicotinic receptor alpha 6 subunit mRNA is selectively concentrated in catecholaminergic nuclei of the rat brain. *Eur J Neurosci.* 1996; 8:2428–2439. [PubMed: 8950106]
14. Quik M, Polonskaya Y, Gillespie A, Jakowec M, Lloyd GK, Langston JW. Localization of nicotinic receptor subunit mRNAs in monkey brain by in situ hybridization. *J Comp Neurol.* 2000; 425:58–69. [PubMed: 10940942]
15. Marks M. unpublished.
16. Han ZY, Le Novère N, Zoli M, Hill JA Jr, Champtiaux N, Changeux JP. Localization of nAChR subunit mRNAs in the brain of *Macaca mulatta*. *Eur J Neurosci.* 2000; 12:3664–3674. [PubMed: 11029636]
17. Wada E, Wada K, Boulter J, Deneris E, Heinemann S, Patrick J, et al. Distribution of alpha 2, alpha 3, alpha 4, and beta 2 neuronal nicotinic receptor subunit mRNAs in the central nervous system: a hybridization histochemical study in the rat. *J Comp Neurol.* 1989; 284:314–335. [PubMed: 2754038]
18. Cartier GE, Yoshikami D, Gray WR, Luo S, Olivera BM, McIntosh JM. A new alpha-conotoxin which targets alpha3beta2 nicotinic acetylcholine receptors. *J Biol Chem.* 1996; 271:7522–7528. [PubMed: 8631783]
19. Champtiaux N, Gotti C, Cordero-Erausquin M, David DJ, Przybylski C, Lena C, et al. Subunit composition of functional nicotinic receptors in dopaminergic neurons investigated with knock-out mice. *J Neurosci.* 2003; 23:7820–7829. [PubMed: 12944511]
20. Zoli M, Moretti M, Zanardi A, McIntosh JM, Clementi F, Gotti C. Identification of the nicotinic receptor subtypes expressed on dopaminergic terminals in the rat striatum. *J Neurosci.* 2002; 22:8785–8789. [PubMed: 12388584]
21. Whiteaker P, McIntosh JM, Luo S, Collins AC, Marks MJ. 125I-alpha-conotoxin MII identifies a novel nicotinic acetylcholine receptor population in mouse brain. *Mol Pharmacol.* 2000; 57:913–925. [PubMed: 10779374]
22. Quik M, Polonskaya Y, Kulak JM, McIntosh JM. Vulnerability of 125I-alpha-conotoxin MII binding sites to nigrostriatal damage in monkey. *J Neurosci.* 2001; 21:5494–5500. [PubMed: 11466420]
23. Gotti C, Moretti M, Zanardi A, Gaimarri A, Champtiaux N, Changeux JP, et al. Heterogeneity and selective targeting of neuronal nicotinic acetylcholine receptor (nAChR) subtypes expressed on retinal afferents of the superior colliculus and lateral geniculate nucleus: identification of a new native nAChR subtype alpha3beta2(alpha5 or beta3) enriched in retinocollicular afferents. *Mol Pharmacol.* 2005; 68:1162–1171. [PubMed: 16049166]
24. Cox BC, Marritt AM, Perry DC, Kellar KJ. Transport of multiple nicotinic acetylcholine receptors in the rat optic nerve: High densities of receptors containing alpha6 and beta3 subunits. *J Neurochem.* 2008; 105:1924–1938. [PubMed: 18266937]
25. Liu J, McGlinn AM, Fernandes A, Milam AH, Strang CE, Andison ME, et al. Nicotinic acetylcholine receptor subunits in rhesus monkey retina. *Invest Ophthalmol Vis Sci.* 2009; 50:1408–1415. [PubMed: 18952912]
26. Grady SR, Moretti M, Zoli M, Marks MJ, Zanardi A, Pucci L, et al. Rodent Habenulo-Interpeduncular Pathway Expresses a Large Variety of Uncommon nAChR Subtypes, But Only the {alpha}3{beta}4 and {alpha}3{beta}3{beta}4 Subtypes Mediate Acetylcholine Release. *J Neurosci.* 2009; 29:2272–2282. [PubMed: 19228980]
27. Quik M, Vailati S, Bordia T, Kulak JM, Fan H, McIntosh JM, et al. Subunit composition of nicotinic receptors in monkey striatum: effect of treatments with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine or L-DOPA. *Mol Pharmacol.* 2005; 67:32–41. [PubMed: 15470079]
28. Gotti C, Guiducci S, Tedesco V, Corbioli S, Zanetti L, Moretti M, et al. Nicotinic acetylcholine receptors in the mesolimbic pathway: primary role of ventral tegmental area alpha6beta2* receptors in mediating systemic nicotine effects on dopamine release, locomotion, and reinforcement. *J Neurosci.* 2010; 30:5311–5325. [PubMed: 20392953]
29. Cui C, Booker TK, Allen RS, Grady SR, Whiteaker P, Marks MJ, et al. The beta3 nicotinic receptor subunit: a component of alpha-conotoxin MII-binding nicotinic acetylcholine receptors

- that modulate dopamine release and related behaviors. *J Neurosci.* 2003; 23:11045–11053. [PubMed: 14657161]
30. Gotti C, Moretti M, Clementi F, Riganti L, McIntosh JM, Collins AC, et al. Expression of nigrostriatal $\alpha 6$ -containing nicotinic acetylcholine receptors is selectively reduced, but not eliminated, by $\beta 3$ subunit gene deletion. *Mol Pharmacol.* 2005; 67:2007–2015. [PubMed: 15749993]
 31. Gotti C, Clementi F, Fornari A, Gaimarri A, Guiducci S, Manfredi I, et al. Structural and functional diversity of native brain neuronal nicotinic receptors. *Biochem Pharmacol.* 2009; 78:703–711. [PubMed: 19481063]
 32. Bordia T, Grady SR, McIntosh JM, Quik M. Nigrostriatal damage preferentially decreases a subpopulation of $\alpha 6\beta 2^*$ nAChRs in mouse, monkey and Parkinson's disease striatum. *Mol Pharmacol.* 2007; 72:52–61. [PubMed: 17409284]
 33. Livingstone PD, Wonnacott S. Nicotinic acetylcholine receptors and the ascending dopamine pathways. *Biochem Pharmacol.* 2009; 78:744–755. [PubMed: 19523928]
 34. Grady SR, Murphy KL, Cao J, Marks MJ, McIntosh JM, Collins AC. Characterization of nicotinic agonist-induced [(3)H]dopamine release from synaptosomes prepared from four mouse brain regions. *J Pharmacol Exp Ther.* 2002; 301:651–660. [PubMed: 11961070]
 35. Moretti M, Vailati S, Zoli M, Lippi G, Riganti L, Longhi R, et al. Nicotinic acetylcholine receptor subtypes expression during rat retina development and their regulation by visual experience. *Mol Pharmacol.* 2004; 66:85–96. [PubMed: 15213299]
 36. Giorguieff MF, Le Floc'h ML, Westfall TC, Glowinski J, Besson MJ. Nicotinic effect of acetylcholine on the release of newly synthesized (3H)dopamine in rat striatal slices and cat caudate nucleus. *Brain research.* 1976; 106:117–131. [PubMed: 1268701]
 37. Rapier C, Lunt GG, Wonnacott S. Stereoselective nicotine-induced release of dopamine from striatal synaptosomes: concentration dependence and repetitive stimulation. *J Neurochem.* 1988; 50:1123–1130. [PubMed: 3346670]
 38. Grady S, Marks MJ, Wonnacott S, Collins AC. Characterization of nicotinic receptor-mediated [3H]dopamine release from synaptosomes prepared from mouse striatum. *J Neurochem.* 1992; 59:848–856. [PubMed: 1494911]
 39. Rowell PP, Carr LA, Garner AC. Stimulation of [3H]dopamine release by nicotine in rat nucleus accumbens. *J Neurochem.* 1987; 49:1449–1454. [PubMed: 2889800]
 40. Kaiser SA, Soliakov L, Harvey SC, Luetje CW, Wonnacott S. Differential inhibition by α -conotoxin-MII of the nicotinic stimulation of [3H]dopamine release from rat striatal synaptosomes and slices. *J Neurochem.* 1998; 70:1069–1076. [PubMed: 9489727]
 41. Kulak JM, Nguyen TA, Olivera BM, McIntosh JM. α -conotoxin MII blocks nicotine-stimulated dopamine release in rat striatal synaptosomes. *J Neurosci.* 1997; 7:5263–5270. [PubMed: 9204910]
 42. McCallum S, Parameswaran N, Bordia T, Fan H, McIntosh M, Quik M. Differential regulation of mesolimbic $\alpha 3/\alpha 6\beta 2^*$ and $\alpha 4\beta 2^*$ nAChR sites and function after long-term oral nicotine to monkeys. *J Pharmacol Exp Ther.* 2006; 318:381–388. [PubMed: 16622038]
 43. McCallum SE, Parameswaran N, Bordia T, McIntosh JM, Grady SR, Quik M. Decrease in $\alpha 3/\alpha 6^*$ nicotinic receptors but not nicotine-evoked dopamine release in monkey brain after nigrostriatal damage. *Mol Pharmacol.* 2005; 68:737–746. [PubMed: 15933214]
 44. Exley R, Clements MA, Hartung H, McIntosh JM, Cragg SJ. $\alpha 6$ -Containing Nicotinic Acetylcholine Receptors Dominate the Nicotine Control of Dopamine Neurotransmission in Nucleus Accumbens. *Neuropsychopharmacology.* 2008; 33:2158–2166. [PubMed: 18033235]
 45. Exley R, Cragg SJ. Presynaptic nicotinic receptors: a dynamic and diverse cholinergic filter of striatal dopamine neurotransmission. *Br J Pharmacol.* 2008; 153 Suppl 1:S283–S297. [PubMed: 18037926]
 46. Meyer EL, Yoshikami D, McIntosh JM. The neuronal nicotinic acetylcholine receptors $\alpha 4^*$ and $\alpha 6^*$ differentially modulate dopamine release in mouse striatal slices. *J Neurochem.* 2008; 105:1761–1769. [PubMed: 18248619]

47. Perez X, O'Leary K, Parameswaran N, McIntosh JM, Quik M. Prominent role of $\alpha_3/\alpha_6\beta_2^*$ nAChRs in regulating evoked dopamine release in primate putamen; effect of long-term nicotine treatment. *Mol Pharmacol*. 2009; 75:938–946. [PubMed: 19144785]
48. Perez XA, Bordia T, McIntosh JM, Grady SR, Quik M. Long-term nicotine treatment differentially regulates striatal $\alpha_6\alpha_4\beta_2^*$ and $\alpha_6(\text{non}\alpha_4)\beta_2^*$ nAChR expression and function. *Mol Pharmacol*. 2008; 74:844–853. [PubMed: 18583454]
49. Perez XA, Bordia T, McIntosh JM, Quik M. $\alpha_6\beta_2^*$ and $\alpha_4\beta_2^*$ nicotinic receptors both regulate dopamine signaling with increased nigrostriatal damage: relevance to Parkinson's disease. *Mol Pharmacol*. 2010; 78:71–80.
50. Perez XA, Ly J, McIntosh JM, Quik M. Chronic nicotine exposure depresses dopaminergic function in the nucleus accumbens; primary role for $\alpha_6\beta_2^*$ nicotinic receptors. 2011
51. Perez XA, Ly J, McIntosh JM, Quik M. Chronic nicotine exposure depresses dopaminergic function in the nucleus accumbens; primary role for $\alpha_6\beta_2^*$ nicotinic receptors. unpublished.
52. Rice ME, Cragg SJ. Nicotine amplifies reward-related dopamine signals in striatum. *Nat Neurosci*. 2004; 7:583–584. [PubMed: 15146188]
53. Zhang H, Sulzer D. Frequency-dependent modulation of dopamine release by nicotine. *Nat Neurosci*. 2004; 7:581–582. [PubMed: 15146187]
54. Zhang T, Zhang L, Liang Y, Siapas AG, Zhou FM, Dani JA. Dopamine signaling differences in the nucleus accumbens and dorsal striatum exploited by nicotine. *J Neurosci*. 2009; 29:4035–4043. [PubMed: 19339599]
55. Threlfell S, Clements MA, Khodai T, Pienaar IS, Exley R, Wess J, et al. Striatal Muscarinic Receptors Promote Activity Dependence of Dopamine Transmission via Distinct Receptor Subtypes on Cholinergic Interneurons in Ventral versus Dorsal Striatum. *J Neurosci*. 2010; 30:3398–3408. [PubMed: 20203199]
56. Kuryatov A, Lindstrom JM. Expression of Functional Human $\alpha_6\beta_2\beta_3^*$ in *Xenopus* Oocytes Achieved Through Subunit Chimeras and Concatamers. *Mol Pharmacol*. 2010; 79:126–140. [PubMed: 20923852]
57. Picciotto MR, Addy NA, Mineur YS, Brunzell DH. It is not "either/or": Activation and desensitization of nicotinic acetylcholine receptors both contribute to behaviors related to nicotine addiction and mood. *Prog Neurobiol*. 2008; 84:329–342. [PubMed: 18242816]
58. Buccafusco JJ, Beach JW, Terry AV. Desensitization of nicotinic acetylcholine receptors as a strategy for drug development. *J Pharmacol Exp Ther*. 2009; 328:364–370. [PubMed: 19023041]
59. Govind AP, Vezina P, Green WN. Nicotine-induced upregulation of nicotinic receptors: underlying mechanisms and relevance to nicotine addiction. *Biochem Pharmacol*. 2009; 78:756–765. [PubMed: 19540212]
60. Benowitz NL. Neurobiology of nicotine addiction: implications for smoking cessation treatment. *Am J Med*. 2008; 121:S3–S10. [PubMed: 18342164]
61. Marks MJ, Burch JB, Collins AC. Effects of chronic nicotine infusion on tolerance development and nicotinic receptors. *J Pharmacol Exp Ther*. 1983; 226:817–825. [PubMed: 6887012]
62. Benwell ME, Balfour DJ, Anderson JM. Evidence that tobacco smoking increases the density of $(-)[^3\text{H}]$ nicotine binding sites in human brain. *J Neurochem*. 1988; 50:1243–1247. [PubMed: 3346676]
63. Marks MJ, Pauly JR, Gross SD, Deneris ES, Hermans-Borgmeyer I, Heinemann SF, et al. Nicotine binding and nicotinic receptor subunit RNA after chronic nicotine treatment. *J Neurosci*. 1992; 12:2765–2784. [PubMed: 1613557]
64. McCallum SE, Parameswaran N, Bordia T, Fan H, Tyndale RF, Langston JW, et al. Increases in α_4^* but not α_3^*/α_6^* nicotinic receptor sites and function in the primate striatum following chronic oral nicotine treatment. *J Neurochem*. 2006; 96:1028–1041. [PubMed: 16412091]
65. Pauly JR, Marks MJ, Robinson SF, van de Kamp JL, Collins AC. Chronic nicotine and mecamylamine treatment increase brain nicotinic receptor binding without changing α_4 or β_2 mRNA levels. *J Pharmacol Exp Ther*. 1996; 278:361–369. [PubMed: 8764371]

66. Perry DC, Davila-Garcia MI, Stockmeier CA, Kellar KJ. Increased nicotinic receptors in brains from smokers: membrane binding and autoradiography studies. *J Pharmacol Exp Ther.* 1999; 289:1545–1552. [PubMed: 10336551]
67. Marks M, McClure-Begley T, Whiteaker P, Salminen O, Brown R, Cooper J, et al. Increased Nicotinic Acetylcholine Receptor Protein Underlies Chronic Nicotine-induced Up-Regulation of Nicotinic Agonist Binding Sites in Mouse Brain. *J Pharmacol Exp Ther.* 2011; 337:187–200. [PubMed: 21228066]
68. Schwartz RD, Kellar KJ. In vivo regulation of [3H]acetylcholine recognition sites in brain by nicotinic cholinergic drugs. *J Neurochem.* 1985; 45:427–433. [PubMed: 4009168]
69. Nashmi R, Lester H. Cell autonomy, receptor autonomy, and thermodynamics in nicotine receptor up-regulation. *Biochem Pharmacol.* 2007; 74:1145–1154. [PubMed: 17662697]
70. Nashmi R, Xiao C, Deshpande P, McKinney S, Grady SR, Whiteaker P, et al. Chronic nicotine cell specifically upregulates functional $\alpha 4^*$ nicotinic receptors: basis for both tolerance in midbrain and enhanced long-term potentiation in perforant path. *J Neurosci.* 2007; 27:8202–8218. [PubMed: 17670967]
71. Buisson B, Bertrand D. Nicotine addiction: the possible role of functional upregulation. *Trends Pharmacol Sci.* 2002; 23:130–136. [PubMed: 11879680]
72. Lai A, Parameswaran N, Khwaja M, Whiteaker P, Lindstrom JM, Fan H, et al. Long-term nicotine treatment decreases striatal $\alpha 6^*$ nicotinic acetylcholine receptor sites and function in mice. *Mol Pharmacol.* 2005; 67:1639–1647. [PubMed: 15681595]
73. Mugnaini M, Garzotti M, Sartori I, Pilla M, Repeto P, Heidbreder CA, et al. Selective down-regulation of [(125)I]Y(0)- α -conotoxin MII binding in rat mesostriatal dopamine pathway following continuous infusion of nicotine. *Neuroscience.* 2006; 137:565–572. [PubMed: 16289885]
74. Perry DC, Mao D, Gold AB, McIntosh JM, Pezzullo JC, Kellar KJ. Chronic nicotine differentially regulates $\alpha 6$ - and $\beta 3$ -containing nicotinic cholinergic receptors in rat brain. *J Pharmacol Exp Ther.* 2007; 322:306–315. [PubMed: 17446303]
75. Kuryatov A, Luo J, Cooper J, Lindstrom J. Nicotine acts as a pharmacological chaperone to up-regulate human $\alpha 4\beta 2$ acetylcholine receptors. *Mol Pharmacol.* 2005; 68:1839–1851. [PubMed: 16183856]
76. Sallette J, Pons S, Devillers-Thiery A, Soudant M, Prado de Carvalho L, Changeux JP, et al. Nicotine upregulates its own receptors through enhanced intracellular maturation. *Neuron.* 2005; 46:595–607. [PubMed: 15944128]
77. Tumkosit P, Kuryatov A, Luo J, Lindstrom J. $\beta 3$ subunits promote expression and nicotine-induced up-regulation of human nicotinic $\alpha 6^*$ nicotinic acetylcholine receptors expressed in transfected cell lines. *Mol Pharmacol.* 2006; 70:1358–1368. [PubMed: 16835356]
78. Walsh H, Govind AP, Mastro R, Hoda JC, Bertrand D, Vallejo Y, et al. Upregulation of nicotinic receptors by nicotine varies with receptor subtype. *J Biol Chem.* 2008; 283:6022–6032. [PubMed: 18174175]
79. Polosa R, Benowitz NL. Treatment of nicotine addiction: present therapeutic options and pipeline developments. *Trends Pharmacol Sci.* 2011
80. Rahman S. Brain nicotinic receptors as emerging targets for drug addiction neurobiology to translational research. *Prog Mol Biol Transl Sci.* 2011; 98:349–365. [PubMed: 21199776]
81. Changeux JP. Nicotine addiction and nicotinic receptors: lessons from genetically modified mice. *Nat Rev Neurosci.* 2010; 11:389–401. [PubMed: 20485364]
82. Benowitz NL. Nicotine addiction. *N Engl J Med.* 2010; 362:2295–2303. [PubMed: 20554984]
83. Corrigall WA, Coen KM, Adamson KL. Self-administered nicotine activates the mesolimbic dopamine system through the ventral tegmental area. *Brain research.* 1994; 653:278–284. [PubMed: 7982062]
84. Corrigall WA, Franklin KB, Coen KM, Clarke PB. The mesolimbic dopaminergic system is implicated in the reinforcing effects of nicotine. *Psychopharmacology (Berl).* 1992; 107:285–289. [PubMed: 1615127]

85. Goldberg SR, Henningfield JE. Reinforcing effects of nicotine in humans and experimental animals responding under intermittent schedules of i.v. drug injection. *Pharmacol Biochem Behav.* 1988; 30:227–234. [PubMed: 3051048]
86. Picciotto MR, Corrigall WA. Neuronal systems underlying behaviors related to nicotine addiction: neural circuits and molecular genetics. *J Neurosci.* 2002; 22:3338–3341. [PubMed: 11978809]
87. Fowler CD, Arends MA, Kenny PJ. Subtypes of nicotinic acetylcholine receptors in nicotine reward, dependence, and withdrawal: evidence from genetically modified mice. *Behav Pharmacol.* 2008; 19:461–484. [PubMed: 18690103]
88. Balfour DJ. The neuronal pathways mediating the behavioral and addictive properties of nicotine. *Handb Exp Pharmacol.* 2009:209–233. [PubMed: 19184651]
89. Picciotto MR, Zoli M, Rimondini R, Lena C, Marubio LM, Pich EM, et al. Acetylcholine receptors containing the beta2 subunit are involved in the reinforcing properties of nicotine. *Nature.* 1998; 391:173–177. [PubMed: 9428762]
90. Maskos U, Molles BE, Pons S, Besson M, Guiard BP, Guilloux JP, et al. Nicotine reinforcement and cognition restored by targeted expression of nicotinic receptors. *Nature.* 2005; 436:103–107. [PubMed: 16001069]
91. Tapper AR, McKinney SL, Nashmi R, Schwarz J, Deshpande P, Labarca C, et al. Nicotine activation of alpha4* receptors: sufficient for reward, tolerance, and sensitization. *Science.* 2004; 306:1029–1032. [PubMed: 15528443]
92. Pons S, Fattore L, Cossu G, Tolu S, Porcu E, McIntosh JM, et al. Crucial role of alpha4 and alpha6 nicotinic acetylcholine receptor subunits from ventral tegmental area in systemic nicotine self-administration. *J Neurosci.* 2008; 28:12318–12327. [PubMed: 19020025]
93. Brunzell DH, Boschen KE, Hendrick ES, Beardsley PM, McIntosh JM. alpha-Conotoxin MII-Sensitive Nicotinic Acetylcholine Receptors in the Nucleus Accumbens Shell Regulate Progressive Ratio Responding Maintained by Nicotine. *Neuropsychopharmacology.* 2010; 35:665–673. [PubMed: 19890263]
94. Jackson KJ, McIntosh JM, Brunzell DH, Sanjakdar SS, Damaj MI. The role of {alpha}6-containing nicotinic acetylcholine receptors in nicotine reward and withdrawal. *J Pharmacol Exp Ther.* 2009; 331:547–554. [PubMed: 19644040]
95. Drenan RM, Grady SR, Steele AD, McKinney S, Patzlaff NE, McIntosh JM, et al. Cholinergic Modulation of Locomotion and Striatal Dopamine Release Is Mediated by {alpha}6{alpha}4* Nicotinic Acetylcholine Receptors. *J Neurosci.* 2010; 30:9877–9889. [PubMed: 20660270]
96. Zhao-Shea R, Liu L, Soll LG, Improgo MR, Meyers EE, McIntosh JM, et al. Nicotine-Mediated Activation of Dopaminergic Neurons in Distinct Regions of the Ventral Tegmental Area. *Neuropsychopharmacology.* 2011; 36:1021–1032. [PubMed: 21289604]
97. Besson M, Granon S, Mameli-Engvall M, Cloez-Tayarani I, Maubourguet N, Cormier A, et al. Long-term effects of chronic nicotine exposure on brain nicotinic receptors. *Proceedings of the National Academy of Sciences of the United States of America.* 2007; 104:8155–8160. [PubMed: 17470777]
98. Soderpalm B, Lof E, Ericson M. Mechanistic studies of ethanol's interaction with the mesolimbic dopamine reward system. *Pharmacopsychiatry.* 2009; 42 Suppl 1:S87–s94. [PubMed: 19434560]
99. Mihalak KB, Carroll FI, Luetje CW. Varenicline is a partial agonist at alpha4beta2 and a full agonist at alpha7 neuronal nicotinic receptors. *Mol Pharmacol.* 2006; 70:801–805. [PubMed: 16766716]
100. Quik M. unpublished observation.
101. Grady SR, Drenan RM, Breining SR, Yohannes D, Wageman CR, Fedorov NB, et al. Structural differences determine the relative selectivity of nicotinic compounds for native alpha 4 beta 2*, alpha 6 beta 2*, alpha 3 beta 4* and alpha 7-nicotine acetylcholine receptors. *Neuropharmacology.* 2010; 58:1054–1066. [PubMed: 20114055]
102. Steensland P, Simms JA, Holgate J, Richards JK, Bartlett SE. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, selectively decreases ethanol consumption and seeking. *Proc Natl Acad Sci U S A.* 2007; 104:12518–12523. [PubMed: 17626178]
103. Lof E, Olausson P, Debeczy A, Stomberg R, McIntosh JM, Taylor JR, et al. Nicotinic acetylcholine receptors in the ventral tegmental area mediate the dopamine activating and

- reinforcing properties of ethanol cues. *Psychopharmacology (Berl)*. 2007; 195:333–343. [PubMed: 17703283]
104. Hendrickson LM, Gardner P, Tapper AR. Nicotinic acetylcholine receptors containing the alpha4 subunit are critical for the nicotine-induced reduction of acute voluntary ethanol consumption. *Channels (Austin)*. 2011; 5
 105. McKee SA, Harrison EL, O'Malley SS, Krishnan-Sarin S, Shi J, Tetrault JM, et al. Varenicline reduces alcohol self-administration in heavy-drinking smokers. *Biol Psychiatry*. 2009; 66:185–190. [PubMed: 19249750]
 106. Hendrickson LM, Zhao-Shea R, Pang X, Gardner PD, Tapper AR. Activation of alpha4* nAChRs is necessary and sufficient for varenicline-induced reduction of alcohol consumption. *J Neurosci*. 2010; 30:10169–10176. [PubMed: 20668200]
 107. Chatterjee S, Steensland P, Simms JA, Holgate J, Coe JW, Hurst RS, et al. Partial agonists of the alpha3beta4* neuronal nicotinic acetylcholine receptor reduce ethanol consumption and seeking in rats. *Neuropsychopharmacology*. 2011; 36:603–615. [PubMed: 21048701]
 108. Larsson A, Jerlhag E, Svensson L, Soderpalm B, Engel JA. Is an alpha-conotoxin MII-sensitive mechanism involved in the neurochemical, stimulatory, and rewarding effects of ethanol? *Alcohol*. 2004; 34:239–250. [PubMed: 15902919]
 109. le Novère N, Zoli M, Lena C, Ferrari R, Picciotto MR, Merlo-Pich E, et al. Involvement of alpha6 nicotinic receptor subunit in nicotine-elicited locomotion, demonstrated by in vivo antisense oligonucleotide infusion. *Neuroreport*. 1999; 10:2497–2501. [PubMed: 10574359]
 110. Drenan RM, Grady SR, Whiteaker P, McClure-Begley T, McKinney S, Miwa JM, et al. In vivo activation of midbrain dopamine neurons via sensitized, high-affinity alpha 6 nicotinic acetylcholine receptors. *Neuron*. 2008; 60:123–136. [PubMed: 18940593]
 111. Marubio LM, Gardier AM, Durier S, David D, Klink R, Arroyo-Jimenez MM, et al. Effects of nicotine in the dopaminergic system of mice lacking the alpha4 subunit of neuronal nicotinic acetylcholine receptors. *Eur J Neurosci*. 2003; 17:1329–1337. [PubMed: 12713636]
 112. Jerlhag E, Egencioglu E, Dickson SL, Svensson L, Engel JA. Alpha-conotoxin MII-sensitive nicotinic acetylcholine receptors are involved in mediating the ghrelin-induced locomotor stimulation and dopamine overflow in nucleus accumbens. *Eur Neuropsychopharmacol*. 2008; 18:508–518. [PubMed: 18343642]
 113. Quik M, Polonskaya Y, Gillespie A, Lloyd KG, Langston JW. Differential Alterations in Nicotinic Receptor alpha6 and beta3 Subunit messenger RNAs in Monkey Substantia nigra After Nigrostriatal Degeneration. *Neuroscience*. 2000; 100:63–72. [PubMed: 10996459]
 114. Quik M, Sum JD, Whiteaker P, McCallum SE, Marks MJ, Musachio J, et al. Differential declines in striatal nicotinic receptor subtype function after nigrostriatal damage in mice. *Mol Pharmacol*. 2003; 63:1169–1179. [PubMed: 12695545]
 115. Kulak JM, McIntosh JM, Quik M. Loss of nicotinic receptors in monkey striatum after 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine treatment is due to a decline in alpha-conotoxin MII sites. *Mol Pharmacol*. 2002; 61:230–238. [PubMed: 11752225]
 116. Bohr IJ, Ray MA, McIntosh JM, Chalon S, Guilloteau D, McKeith IG, et al. Cholinergic nicotinic receptor involvement in movement disorders associated with Lewy body diseases. An autoradiography study using [(125)I]alpha-conotoxinMII in the striatum and thalamus. *Exp Neurol*. 2005; 191:292–300. [PubMed: 15649484]
 117. Gotti C, Moretti M, Bohr I, Ziabreva I, Vailati S, Longhi R, et al. Selective nicotinic acetylcholine receptor subunit deficits identified in Alzheimer's disease, Parkinson's disease and dementia with Lewy bodies by immunoprecipitation. *Neurobiol Dis*. 2006; 23:481–489. [PubMed: 16759874]
 118. Quik M, Bordia T, Forno L, McIntosh JM. Loss of alpha-conotoxinMII- and A85380-sensitive nicotinic receptors in Parkinson's disease striatum. *J Neurochem*. 2004; 88:668–679. [PubMed: 14720216]
 119. Quik M, O'Neill M, Perez XA. Nicotine neuroprotection against nigrostriatal damage: importance of the animal model. *Trends Pharmacol Sci*. 2007; 28:229–235. [PubMed: 17412429]

120. O'Neill MJ, Murray TK, Lakics V, Visanji NP, Duty S. The role of neuronal nicotinic acetylcholine receptors in acute and chronic neurodegeneration. *Curr Drug Target CNS Neurol Disord.* 2002; 1:399–411.
121. Picciotto MR, Zoli M. Neuroprotection via nAChRs: the role of nAChRs in neurodegenerative disorders such as Alzheimer's and Parkinson's disease. *Front Biosci.* 2008; 13:492–504. [PubMed: 17981563]
122. Huang LZ, Parameswaran N, Bordia T, Michael McIntosh J, Quik M. Nicotine is neuroprotective when administered before but not after nigrostriatal damage in rats and monkeys. *J Neurochem.* 2009; 109:826–837. [PubMed: 19250334]
123. Schapira AH, Emre M, Jenner P, Poewe W. Levodopa in the treatment of Parkinson's disease. *Eur J Neurol.* 2009; 16:982–989. [PubMed: 19538218]
124. Poewe W. Treatments for Parkinson disease--past achievements and current clinical needs. *Neurology.* 2009; 72:S65–s73. [PubMed: 19221317]
125. Lang AE. When and how should treatment be started in Parkinson disease? *Neurology.* 2009; 72:S39–s43. [PubMed: 19221313]
126. Fahn S. How do you treat motor complications in Parkinson's disease: Medicine, surgery, or both? *Ann Neurol.* 2009; 64:S56–S64. [PubMed: 19127577]
127. Pezzoli G, Zini M. Levodopa in Parkinson's disease: from the past to the future. *Expert Opin Pharmacother.* 2010; 11:627–635. [PubMed: 20163273]
128. Ahlskog JE, Muenter MD. Frequency of levodopa-related dyskinesias and motor fluctuations as estimated from the cumulative literature. *Mov Disord.* 2001; 16:448–458. [PubMed: 11391738]
129. Huang LZ, Bordia T, Quik M. Nicotine treatment reduces L-dopa-induced dyskinetic-like movements in parkinsonian mice. *Society for Neuroscience Abstracts.* 2009
130. Bordia T, Campos C, Huang L, Quik M. Continuous and intermittent nicotine treatment reduces L-3,4-dihydroxyphenylalanine (L-DOPA)-induced dyskinesias in a rat model of Parkinson's disease. *J Pharmacol Exp Ther.* 2008; 327:239–247. [PubMed: 18650244]
131. Quik M, Cox H, Parameswaran N, O'Leary K, Langston JW, Di Monte D. Nicotine reduces levodopa-induced dyskinesias in lesioned monkeys. *Annals of Neurology.* 2007; 62:588–596. [PubMed: 17960553]
132. Bordia T, Campos C, McIntosh JM, Quik M. Nicotinic receptor-mediated reduction in L-dopa-induced dyskinesias may occur via desensitization. *J Pharmacol Exp Ther.* 2010; 333:929–938. [PubMed: 20200117]
133. Rollema H, Coe JW, Chambers LK, et al. Rationale, pharmacology and clinical efficacy of partial agonists of alpha(4)beta(2) nACh receptors for smoking cessation. *Trends Pharmacol Sci.* 2007; 28:316–325. [PubMed: 17573127]
134. Kulak JM, Sum J, Musachio JL, McIntosh JM, Quik M. 5-Iodo-A-85380 binds to alpha-conotoxin MII-sensitive nicotinic acetylcholine receptors (nAChRs) as well as alpha4beta2* subtypes. *J Neurochem.* 2002; 81:403–406. [PubMed: 12064487]
135. Huang LZ, Campos C, Ly J, Carroll FI, Quik M. Nicotinic receptor agonists decrease L-dopa-induced dyskinesias most effectively in moderately lesioned parkinsonian rats. *Neuropharmacology.* 2011; 60:861–868. [PubMed: 21232546]
136. Carta AR, Frau L, Pontis S, Pinna A, Morelli M. Direct and indirect striatal efferent pathways are differentially influenced by low and high dyskinetic drugs: behavioural and biochemical evidence. *Parkinsonism Relat Disord.* 2008; 14 Suppl 2:S165–s168. [PubMed: 18583175]
137. Carta M, Carlsson T, Munoz A, Kirik D, Bjorklund A. Involvement of the serotonin system in l-dopa-induced dyskinesias. *Parkinsonism Relat Disord.* 2008; 14 Suppl 2:S154–S158. [PubMed: 18579429]
138. Cenci MA, Konradi C. Maladaptive striatal plasticity in l-DOPA-induced dyskinesia. *Progress in brain research.* 2010; 183C:209–233. [PubMed: 20696322]
139. Limousin P, Martinez-Torres I. Deep brain stimulation for Parkinson's disease. *Neurotherapeutics.* 2008; 5:309–319. [PubMed: 18394572]
140. Xiang Z, Huguenard JR, Prince DA. Cholinergic switching within neocortical inhibitory networks. *Science.* 1998; 281:985–988. [PubMed: 9703513]

141. Morishita H, Miwa JM, Heintz N, Hensch TK. Lynx1, a cholinergic brake, limits plasticity in adult visual cortex. *Science*. 2010; 330:1238–1240. [PubMed: 21071629]
142. Salminen O, Murphy KL, McIntosh JM, Drago J, Marks MJ, Collins AC, et al. Subunit composition and pharmacology of two classes of striatal presynaptic nicotinic acetylcholine receptors mediating dopamine release in mice. *Mol Pharmacol*. 2004; 65:1526–1535. [PubMed: 15155845]
143. Luetje CW. Getting past the asterisk: the subunit composition of presynaptic nicotinic receptors that modulate striatal dopamine release. *Mol Pharmacol*. 2004; 65:1333–1335. [PubMed: 15155826]

Primary $\alpha 6^*$ nAChR subtypes in the nigrostriatal and mesolimbic dopaminergic systems

Fig. 1.

Putative composition of $\alpha 6^*$ nAChRs in the nigrostriatal and mesolimbic dopaminergic systems. Cumulative results from numerous molecular and functional studies indicate that nAChRs containing the $\alpha 6$ subunit are pentameric channels with the subunits around a central channel or pore that lie within the membrane bilayer. The acetylcholine binding sites (triangles) are located at the interface between an α and β subunit. Although there is the potential for multiple combinations, the major nAChR subtypes in the nigrostriatal and mesolimbic dopaminergic systems appear to be composed of $\alpha 6 \beta 2 \beta 3$ (left) and $\alpha 6 \alpha 4 \beta 2 \beta 3$ (right) subunits (see also Table 1) [7–11, 142, 143]. The possibility exists, however, that other minor $\alpha 6 \beta 2^*$ nAChR subtypes remain to be identified.

Fig. 2.

Prominent contribution of the $\alpha 6\beta 2\beta 3^*$ nAChR subtype in stimulating ^3H -dopamine release from nucleus accumbens and striatal synaptosomes prepared from rodent or nonhuman primate brain. nAChR-evoked ^3H -dopamine was determined as previously described [34, 42, 64]. The results show that the $\alpha 6\beta 2\beta 3^*$ nAChR subtypes make a similar contribution to ^3H -dopamine release (stimulated by 1 μM nicotine) as the $\alpha 4\beta 2^*$ nAChR subtypes ($\alpha 4\beta 2$ and $\alpha 4\alpha 5\beta 2$) in rodent nucleus accumbens and dorsal striatum. Moreover, the $\alpha 6\beta 2\beta 3^*$ nAChR subtypes play a dominant role in modulating ^3H -dopamine in nonhuman primate nucleus accumbens and dorsal putamen, with ~80% of release evoked by the $\alpha 6\beta 2\beta 3^*$ and only ~20% by $\alpha 4\beta 2^*$ ($\alpha 4\beta 2$ and $\alpha 4\alpha 5\beta 2$) nAChR subtypes. Values represent the mean $\pm$

SEM of 5–7 animals. *** $p < 0.001$ indicate significance of difference from control, using Student's t-test.

Rat dorsal striatum

Monkey ventral putamen

Monkey dorsal putamen

Fig. 3. $\alpha 6\beta 2\beta 3^*$ nAChRs are major regulators of nAChR-modulated dopamine release in the nigrostriatal system. Cyclic voltammetry was used to measure electrically stimulated dopamine release in coronal brain slices containing rat dorsal striatum (top panel), monkey ventral putamen (middle panel) or monkey dorsal putamen (bottom panel). Endogenous dopamine release was evoked by a bipolar stimulating electrode and detected with a carbon fiber recording microelectrode as described [47, 49]. Representative traces of single-pulse stimulated total dopamine release in the different tissue slices are shown on the left. The traces in the middle and on the right depict dopamine release in the presence of the $\alpha 6\beta 2\beta 3^*$ nAChR antagonist α -CtxMII (100nM) or the general nAChR blocker mecamylamine (100 μ M), respectively. Quantitative analyses of the data show the relative contribution of the

$\alpha 6\beta 2\beta 3^*$ and $\alpha 4\beta 2^*$ nAChR subtypes ($\alpha 4\beta 2$ and $\alpha 4\alpha 5\beta 2$) in modulating dopamine release. As can be seen, 75–100% of nAChR-modulated evoked dopamine release occurs via the $\alpha 6\beta 2\beta 3^*$ subtype. Values represent the mean $\pm$ SEM of 3–9 animals. *** $p < 0.001$ indicate significance of difference from control, while # $p < 0.05$ indicates significance of difference from α -CtxMII, using a Newman-Keuls multiple comparisons *post hoc* test. Data taken in modified form with permission [47, 49].

Fig. 4.

Preferential loss of the $\alpha 6 \alpha 4 \beta 2 \beta 3$ nAChR subtype in striatum with nigrostriatal damage. The top panel depicts ^{125}I - α -CtxMII competition curves in the presence of varying concentrations of α -CtxMII [E11A] in the striatum of wildtype and $\alpha 4$ nAChR knockout mice. Biphasic curves are observed in wildtype mouse striatum, but a monophasic curve in $\alpha 4$ knockout mice. These findings suggest the presence of two major $\alpha 6 \beta 2 \beta 3^*$ nAChR populations, with the higher affinity site being the $\alpha 6 \alpha 4 \beta 2 \beta 3$ subtype and the lower affinity site the $\alpha 6 \beta 2 \beta 3$ receptor. A point of note is that the K_i of the lower affinity site in wildtype mice was not identical to that in the knockout mice. This may be due to differences in molecular configurations of the $\alpha 6 \beta 2^*$ interfaces, variable post-translational modifications

(phosphorylation or glycosylation) or possibly additional subtype combination with knockout of the $\alpha 4$ subunit. The second panel depicts similar competition experiments using striatum from control and parkinsonian mice. Interestingly, the curves in the lesioned mice are identical to those in the $\alpha 4$ knockout mice, suggesting a loss of the $\alpha 6\alpha 4\beta 2\beta 3$ subtype. The third and fourth panels depict similar competition experiments from striatum of control and parkinsonian monkeys, and from control human and Parkinson's disease cases. The biphasic inhibition curves obtained in control striatum become monophasic with nigrostriatal damage, again suggesting a loss of the $\alpha 6\alpha 4\beta 2\beta 3$ subtype. Values represent the mean $\pm$ SEM of 4–8 mice, 4–6 monkeys, and 5 control and 4 Parkinson's disease cases. Taken in modified form with permission [32].

Table 1Distribution of $\alpha 6^*$ nAChR subtypes in the mammalian CNS

Brain Region	Species	Subtype	Reference
Nigrostriatal dopaminergic system - dorsal striatum	rodent	$\alpha 6\beta 2\beta 3$ $\alpha 6\alpha 4\beta 2\beta 3$ $\alpha 6\beta 2^*$	[1–6]
	monkey	$\alpha 6\beta 2\beta 3$ $\alpha 6\alpha 4\beta 2\beta 3$ $\alpha 6\beta 2^*$	[7]
	human	$\alpha 6\beta 2\beta 3$ $\alpha 6\alpha 4\beta 2\beta 3$ $\alpha 6\beta 2^*$	[1, 8–10]
Mesolimbic dopaminergic system - ventral striatum	rodent	$\alpha 6\beta 2\beta 3$ $\alpha 6\alpha 4\beta 2\beta 3$ $\alpha 6\beta 2^*$	[6]
Retina and optic tract	rodent	$\alpha 6\beta 2^*$ $\alpha 6\alpha 3\beta 2^*$ $\alpha 6\alpha 2\beta 2^*$ $\alpha 6\beta 2\beta 3^*$ $\alpha 6\alpha 4\beta 2^*$ $\alpha 6\alpha 3\alpha 4\beta 2^*$	[11–13]
Habenulo-interpeduncular pathway	rodent	$\alpha 6\beta 2^*$ $\alpha 6\beta 3\beta 4^*$	[14]

Although the $\alpha 6\beta 2\beta 3$ and $\alpha 6\alpha 4\beta 2\beta 3$ nAChRs are the primary subtypes in the mesolimbic and nigrostriatal dopaminergic systems, the $\alpha 6\beta 2^*$ nAChR designation is retained in the event novel subtypes are identified.

Long term oral nicotine treatment decreases or does not change $\alpha 6\beta 2\beta 3^*$ nAChR expression and function in striatum and nucleus accumbens of rats and monkeys. By contrast, this same treatment increases or does not change $\alpha 4\beta 2^*$ nAChR expression and function.

Table 2

nAChR subtype	Species	Region	nAChR binding sites	nAChR-mediated ^3H -DA release	Reference
$\alpha 6\beta 2\beta 3^*$	Rat	striatum	↓	↓	[15–20]
	Monkey	striatum	↓	↓	[19]
		n. accumbens	No change	No change	[18]
$\alpha 4\beta 2^*$	Rat	striatum	↑	No change	[15, 19–27]
	Monkey	striatum	↑	↑	[19]
		n. accumbens	↑	↑	[18]