Published in final edited form as:

J Phys Chem B. 2012 January 26; 116(3): 1134–1142. doi:10.1021/jp208918n.

UV resonance Raman studies of the NaClO₄ dependence of poly-L-lysine conformation and hydrogen exchange kinetics

Lu Ma, Zhenmin Hong, Bhavya Sharma, and Sanford Asher*

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, Tel: (412)-624-8570, Fax: (412)-624-0588

Abstract

We used 204 nm excitation UV Resonance Raman (UVRR) spectroscopy to examine the effects of NaClO₄ on the conformation of poly-L-lysine (PLL). The presence of NaClO₄ induces the formation of α -helix, π -helix/bulge and turn conformations. The dependence of the AmIII₃ frequency on the peptide Ψ Ramachandran angle allows us to experimentally determine the conformational population distributions and the energy landscape of PLL along the Ramachandran Ψ angle. We also used UVRR to measure the NaClO₄ concentration dependence of PLL amide hydrogen exchange kinetics. Exchange rates were determined by fitting the D₂O exchanging PLL UVRR AmII' band time evolution. Hydrogen exchange is slowed at high NaClO₄ concentrations. The PLL AmII' band exchange kinetics at 0.0, 0.2 and 0.35 M NaClO₄ can be fit by single exponentials, but the AmII' band kinetics of PLL at 0.8 M NaClO₄ requires a double exponential fit. The exchange rates for the extended conformations were monitored by measuring the C α -H band kinetics. These kinetics are identical to those of the AmII' band until 0.8 M NaClO₄ whereupon the extended conformation exchange becomes clearly faster than that of the α -helix-like conformations. Our results indicate that ClO₄⁻ binds to the PLL backbone to protect it from OH⁻ exchange catalysis. In addition, ClO₄⁻ binding also slows the conformational exchange between the extended and α -helix-like conformations, probably by increasing the activation barriers for conformational interchanges.

Keywords

Conformational distribution; Energy landscape; Conformational dynamics; Salt binding

INTRODUCTION

An understanding of the mechanism(s) of protein folding remains an important unsolved problem in structural biology.^{1–6} The major underlying assumption is that the protein native structure is the most thermodynamically stable structure.^{7–9} Thus, the protein primary sequence is expected to contain all of the information necessary to both specify the protein native structure and its folding mechanism(s). The well-known Levinthal paradox implies that there exist specific folding intermediates or pathways through which the protein quickly evolves into its final functional structure.¹⁰

The mechanisms of protein folding have been probed experimentally by techniques such as CD, NMR, IR, Raman, fluorescence and a variety of time resolved spectroscopies, mutational studies and hydrogen exchange methods.^{11–16} Most recently, single molecule methods have probed the distributions of the unfolded and folded states by following the

*corresponding author: asher@pitt.edu.

folding reaction of individual molecules.^{17–20} Single molecule methods, such as optical tweezers and AFM, have been used to elucidate protein folding intermediates and the energy landscapes by using mechanical force as a denaturant.^{17,21} Computational modeling has also made significant contributions to the understanding of protein folding.^{22–24}

In the work here, we use UV resonance Raman (UVR) spectroscopy to study the conformational distributions of poly-L-lysine (PLL) in the presence of different NaClO₄ concentrations. ClO₄[–] induces an α -helix conformation at neutral and low pH values, where the lysine side chains are positively charged.^{25,26} Ma et al. recently observed that PLL adopts multiple α -helix-like conformations in the presence of ClO₄[–], including pure α -helix and π -bulge/helix conformations.²⁷

We also determined the dependence of the PLL conformational population distributions and the energy landscapes on NaClO₄ concentration. We measured the hydrogen exchange behavior of the PLL peptide bond NH at pH 2.8 at different NaClO₄ concentrations. The results show that the exchange rates are NaClO₄ concentration dependent and that the hydrogen exchange behavior indicates that the PLL conformations are in rapid equilibrium until the highest NaClO₄ concentrations where two exchange rates are observed.

EXPERIMENTAL

Materials

Poly-L-lysine HCl (MW_{vis} = 20900, DP_{vis} = 127, MW_{MALLS} = 11400, DP_{MALLS} = 69, where DP_{vis} and DP_{MALLS} refer to the degree of polymerization measured by viscosity and multi-angle laser light scattering, respectively) was obtained from Sigma Co. and used without further purification. NaClO₄ was purchased from Sigma Co., and was used without further purification. D₂O (99.9 atom %D) was purchased from Cambridge Isotope Laboratories, Inc. Small aliquots of HCl and DCl solutions were used to adjust the solution pH values. DCl (99 + atom %D) was acquired from Aldrich. The UVR spectra of PLL at different NaClO₄ concentrations were measured at 0.85 mg/ml peptide concentrations at pH 3, and the spectra were normalized to the Raman intensity of the symmetric stretching band of ClO₄[–].

UVR Instrument

We used a Coherent Infinity Nd: YAG laser (Coherent, Inc.) to produce 355 nm light pulses (3rd harmonic) at a 100 Hz repetition rate with a pulse width of 3 ns. This beam was Raman shifted to 204 nm (5th anti-Stokes) using a 1-m tube filled with hydrogen (60 psi), giving 2 mW average power.^{28,29} A Pellin Broca prism was used to select the 204 nm excitation. The sample was circulated in a free surface, temperature controlled stream to avoid heating or accumulation of photochemical degradation products formed by the high peak power laser pulses. The Raman scattered light was imaged into a subtractive double spectrometer.²⁸ The dispersed UV light was detected by the liquid nitrogen cooled, Unichrome coated, back-thinned CCD with a >30% quantum efficiency in the deep UV (Princeton Instruments Spec-10:400B). More details on the UVR apparatus are given by Bykov et al.²⁸

CD Measurements

Circular dichroism (CD) spectra were measured by using a Jasco J-710 spectropolarimeter. The spectra were measured by using a temperature controlled 0.2 mm path length cell with 0.87 mg/ml PLL concentrations.

Hydrogen Exchange Experiment

1 ml pD 2.9 D₂O solutions containing 8.5 mg PLL and 0, 0.2, 0.35 or 0.8 M NaClO₄ concentrations were prepared and mixed rapidly with a 9 ml H₂O solution at pH 2.8 with the same NaClO₄ concentrations. The dead times for all mixing experiment were 25 to 60 s. The time dependent UVRR spectra of the solutions at 0.85 mg/ml PLL concentration containing 0, 0.2, 0.35 or 0.8 M NaClO₄ were collected at 5 s intervals with 5 s data accumulation times. For samples containing NaClO₄, all Raman spectra are normalized to the intensity of the 932 cm⁻¹ ClO₄⁻ Raman band. In the absence of NaClO₄ the AmI (AmI') band intensity was used as the internal intensity standard.

RESULTS AND DISCUSSION

NaClO₄ dependence of 204 nm UVRR spectra of PLL

Previous studies show that NaClO₄ converts unfolded PLL with charged side chains to α -helix-like conformations.²⁷ For example, 0.8 M NaClO₄ at pH 5.5 at 2.5 °C converts normally extended PLL conformations to an ~86% α -helix-like content.³⁰ Fig. 1A shows 204 nm UVRR spectra of PLL at different NaClO₄ concentrations at pH 3 and 20 °C. In the absence of NaClO₄, as observed previously, PLL occurs in an equilibrium between extended polyproline II (PPII) and 2.5₁-helix conformations at low pH.^{27,31} These conformations give rise to AmIII₃ bands at ~1245 cm⁻¹ and ~1268 cm⁻¹, respectively as shown in Fig. 1B. As the NaClO₄ concentration increases the intensities of the AmIII bands between 1190 and 1300 cm⁻¹, the ~1400 cm⁻¹ C α -H b band, and the ~1567 cm⁻¹ AmII band decrease. In addition, the AmI and AmII band frequencies downshift; 0.8 M NaClO₄ causes the AmI and AmII bands to downshift by 12 cm⁻¹, and the AmI bandshape narrows and becomes more symmetric. These spectral changes indicate the formation of α -helical-like conformations as confirmed by the CD measurement in Fig. 2.

We previously believed that NaClO₄ converts charged PLL to a highly α -helical conformation by forming ion-pairs between ClO₄⁻ and the PLL -NH₃⁺ side chains to neutralize the repulsions that destabilize α -helical conformations.^{27,32-34} A simple Debye-Hückel screening mechanism was excluded because high NaCl concentrations negligibly impact the PLL conformation.^{26,30} Ion-pair formation should dramatically decrease side chain electrostatic repulsions. This should decrease the 2.5₁-helix conformation content that is stabilized by side chain electrostatic repulsions.

Indeed our previous study of K10 (Ac-lys10-NH₂) demonstrated that the neutralization of the charged lysine side chains at increased pH significantly decreased the relative fraction of 2.5₁-helix compared to that of the PPII conformation.³⁰ However, surprisingly, the UVRR spectra of PLL showed that NaClO₄ does not decrease the relative fraction of 2.5₁-helix significantly compared to that of the PPII conformation. We frankly do not understand this result.

The C α -H b bands that are resonantly enhanced in the extended PPII, β -sheet, and β -strand-like conformations, are not enhanced in the α -helix-like conformations. As a result, in Fig. 1A, the C α -H b band shows a large intensity decrease relative to that of the AmII and AmIII bands. Using the C α -H b band intensity to indicate the extended PLL concentration content, we subtract the appropriate amount of the extended PLL spectrum from the measured spectra, leaving the UVRR of the NaClO₄-induced α -helix conformation. This subtraction assumes that the extended conformation possesses an invariant UVRR spectrum as the NaClO₄ concentration increases; this assumption is supported by the fact that the UVRR difference spectra between successive additions of NaClO₄ are very similar, and that the system appears 2-state from the Fig. 1 and Fig. 2 isosbestic points in the UVRR and the CD.

Fig. 3 shows the estimated α -helix-like fraction of PLL at different NaClO_4 concentrations obtained from the UVRR. The α -helix-like content fraction of PLL at different NaClO_4 concentrations, f_H , can also be calculated from the Fig. 2 CD spectra by utilizing the equation^{29,35}:

$$f_H = \frac{\theta_{222} - \theta_r}{\theta_H - \theta_r} \quad (1)$$

θ_{222} is the molar ellipticity at 222 nm, which is assumed to be proportional to the helix content, θ_r and θ_H are the molar ellipticities of the extended and α -helix-like conformations. Here, θ_r was experimentally measured as $2854 \text{ deg cm}^2 \text{ dmol}^{-1}$, and $\theta_H = 32200 \text{ deg cm}^2 \text{ dmol}^{-1}$ is the mean residue ellipticity as averaged over several polypeptides.³⁶

The α -helix-like fraction obtained from CD is smaller than that obtained from UVRR (Fig. 3). The 222 nm CD ellipticity results from interpeptide bond exciton coupling of the α -helical conformation electronic dipole transition moments. The ellipticity per residue decreases for short α -helices.^{29,35,37}

In contrast, the Raman intensity is the sum of the intensities from each peptide bond. We calculated the α -helix fraction by assuming that there only exist two conformations, either an extended conformation or an α -helix-like conformation. We used the $\text{C}_\alpha\text{-H}$ b band intensity to calculate the PLL extended conformational fraction. We do not expect phenomena such as hypochromism to alter the extended conformation peptide bond Raman cross sections.

We calculated the UVRR α -helix-like conformation fraction from $f_H = 1 - f_{\text{extended}}$. The α -helix-like conformational fractions calculated from the UVRR include α -helix-like turns that do not give significant contributions to the 222 nm ellipticity. Thus, the Raman indicates a higher α -helix-like fraction than CD (Fig. 3).

Fig. 4 shows the calculated α -helix-like PLL AmIII region UVRR spectra. The AmIII region of the α -helix-like PLL spectra is well modeled by Gaussian bands. The 1110 cm^{-1} band derives from the ClO_4^- asymmetric stretching band, while the AmIII₁ and AmIII₂ bands are located at 1332 cm^{-1} and 1294 cm^{-1} . As shown in Fig. 4 the AmIII₃ band spectral region depends on the NaClO_4 concentration. At 0.1 and 0.2 M NaClO_4 concentrations, the AmIII₃ region resolves into two bands at 1222 cm^{-1} and 1260 cm^{-1} . In contrast, at 0.35, 0.5 and 0.8 M NaClO_4 , the AmIII₃ region resolves into three bands at 1222 cm^{-1} , 1253 cm^{-1} and 1273 cm^{-1} .

NaClO_4 dependence of PLL Ψ angle distribution

Using the method developed by Mikhonin et al.,³⁸ we can calculate the Ramachandran Ψ angle distribution from the AmIII₃ band frequency distribution if we know the peptide bond hydrogen bonding states.^{38–40} Fig. 5 shows the NaClO_4 concentration dependence of the Ψ angle distributions of PLL calculated from the AmIII₃ bands deriving from the calculated α -helix-like spectra of Fig. 4, and from the pure extended conformations of PLL in the absence of NaClO_4 of Fig. 1B.

Assuming α -helix-like intramolecular peptide bond hydrogen bonding, the 1273 cm^{-1} , 1260 cm^{-1} and 1253 cm^{-1} AmIII₃ bands are calculated to derive from conformations with average Ramachandran Ψ angles of -60° , -43° , -36° .³⁸

Thus, the 1273 cm^{-1} and 1253 cm^{-1} AmIII₃ bands derive from π -bulge and pure α -helix conformations, respectively. The 1260 cm^{-1} AmIII₃ band occurs at a frequency between that of the α -helix and π -bulge/helix conformation. It could derive from distorted α -helix or π -helix conformations.

The 1222 cm^{-1} AmIII₃ band could result from Ψ angles, of either 4° or 124° . Because the 1222 cm^{-1} AmIII₃ band does not also show the significant intensity in the $C_\alpha\text{-H b}$ band region that would be expected for this partially extended structure, we exclude the 124° Ψ angle possibility, and assume that the conformation is close to that of an α -helix. This conformation could derive from type I or I' and type II or II' turns from their i+2 residues based on this Ψ value.³⁸ For brevity we label this turn conformation T1. We assume here equal Raman cross sections for α -helix, π -helix and T1 structures.

The extended conformation PLL spectra, measured in the absence of NaClO_4 , shown in Fig. 1B shows three AmIII₃ bands. Two were assigned to the PPII and 2.5₁-helix conformations, as discussed above. The third AmIII₃ band appears as a shoulder at 1214 cm^{-1} . This band which we label T2 could also result from turn conformations, with Ψ angles of either 14° or 114° . However, it is unclear whether this turn conformation shows significant $C_\alpha\text{-H b}$ band intensity because of the overwhelming contributions of the PPII and 2.5₁-helix conformations.

$\Psi = 14^\circ$ could derive from a somewhat distorted type I, II, or II', i+2 residues, or type I' or III', i+1 and i+2 residues. $\Psi = 114^\circ$ could derive from a somewhat distorted type II, i+1 residues or a type VIII, i+2 residue.³⁸ Fig 5 shows the Ψ distribution assuming that the T2 conformation possesses $\Psi = 114^\circ$ and that this conformation shows Raman bands with the same cross sections as the PPII and 2.5₁-helix extended conformations.

At 0.1 M NaClO_4 the Ψ angle distribution show only the distorted α -helix or π -bulge conformations and the T1 and T2 turn structures and the PPII and 2.5₁-helix structures. π -bulge structures are induced by increasing NaClO_4 concentration. Further, as the NaClO_4 concentration increases, the α -helix-like content increases, decreasing the extended conformation content. The relative concentrations of α -helix and π -bulge concentrations increase compared to that of the T1 turn structure.

Conformational free energy landscape of PLL

We can use the calculated conformation population distribution of Fig. 5 to calculate the Gibbs free energy of the equilibrium conformations relative to the PPII conformation along the Ψ Ramachandran angle folding reaction coordinate.^{27,31} Fig. 6 shows the resulting energy landscape at different NaClO_4 concentrations at pH 3 and 20 °C. The energy landscape contains an α -helix-like conformational basin and an extended conformation energy basin.

At 0.1 M NaClO_4 , the α -helical basin shows a double-well with a separate minima from the α -helix and the T1 turn structure. The α -helix and π -bulge conformations lie 1.5 kcal/mol above those of the PPII and 2.5₁-helix conformations. Increasing the NaClO_4 concentration lowers the α -helix-like conformation basin energy. At 0.2 M NaClO_4 , the α -helix-like conformation is 0.58 kcal/mol above the PPII and 2.5₁-helix conformations. At 0.35 M NaClO_4 , the α -helix or π -helix-like energy well splits into an α -helix, π -bulge and T1 triplet of energy minima. The α -helix, π -bulge and T1 triplet energies are similar to those of the PPII and 2.5₁-helix conformations. At 0.8 M NaClO_4 , the α -helix and π -bulge conformational energies are 0.55 and 0.65 kcal/mol below that of extended conformations, consistent with our previous study.²⁷ The relative energy of the T1 conformation energy decreases less than that of the α -helix and π -bulge conformations. The variations in shape

of the PLL Gibbs free energy landscape with changing NaClO₄ concentration indicates that complex conformation changes occur as the NaClO₄ concentration increases.

UVRR spectra of protonated and deuterated PLL

Deuteration of the PLL backbone N–H groups leads to significant PLL UVRR spectral changes³⁰ because the AmII, AmIII and C_α–H b bands involve significant N–H bending; deuteration removes the N–H in plane bending component, leaving an almost pure C–N s AmII' vibration at 1470 cm^{−1} and an AmIII' vibration at 935 cm^{−1} which has a large N–D bending component. Fig. 7 shows the pH 3 UVRR spectrum of PLL in the absence and presence of 0.8 M NaClO₄ in H₂O and D₂O. The AmIII, AmII and C_α–H b bands disappear and are replaced with the very strong AmII' band, and a weak AmIII' band (not shown). The UVRR spectra clearly differentiate between ND vs NH peptide bonds. The AmII' band is strong and well resolved, allowing for accurate quantitative analysis of the relative deuterated fraction.

NaClO₄ dependence of PLL amide hydrogen exchange rate

We examined PLL hydrogen exchange at 0, 0.2, 0.35, 0.8 M NaClO₄ concentrations, by mixing PLL D₂O NaClO₄ solutions rapidly with H₂O NaClO₄ solutions. Fig. 8 shows the evolution of the UVRR on hydrogen exchange of PLL in the absence and presence of 0.8 M NaClO₄. The AmII' band gradually loses intensity upon replacement of N–D by N–H, while the AmII, C_α–H b and AmIII bands gain intensity.

Fig. 9 shows the normalized intensity decay of the AmII' band calculated as:

$$I_{N,AmII'}(t) = \frac{I_{AmII'}(t) - I_{AmII'}(\infty)}{I_{AmII'}(0) - I_{AmII'}(\infty)} \quad (2)$$

where $I_{N,AmII'}(t)$ is the normalized AmII' intensity decay, whose value decreases from 1.0 to 0.0, $I_{AmII'}(t)$ is the experimentally observed AmII' band intensity at exchange time t , $I_{AmII'}(0)$ and $I_{AmII'}(\infty)$ are the AmII' band intensities at $t = 0$ and $t = \infty$. Since the AmII' band shapes at all exchange times for specific NaClO₄ concentration are almost identical, we utilized peak height intensities for analysis.

The 1459 cm^{−1} HOD bending band partially overlaps the AmII' band. Since the H₂O/D₂O rapidly reaches equilibrium, the HOD bending band contribution to the AmII' band intensity is constant at all times during the PLL hydrogen exchange. Thus, the HOD band subtracts off during the data analysis using eqn. 2.

As the NaClO₄ concentration increases, hydrogen exchange slows (Fig. 9). The apparent exchange rate is calculated by fitting the normalized AmII' band intensities. The AmII' intensity decay curves at 0, 0.2, and 0.35 M NaClO₄ concentrations at pH 2.8 are well fit by single exponentials with apparent exchange rates of 0.95, 0.11, 0.063 min^{−1}, respectively. The best single exponential fitting to the AmII' 0.8 M exchange curve gave an apparent exchange rate of 0.014 min^{−1}. However, Fig. 9 shows that the 0.8 M exchange curve does not fit well to a single exponential. Excellent fitting was obtained with a double exponential that give apparent rates of 0.022 min^{−1} and 0.00097 min^{−1} with weighting factors of 0.77 and 0.23 for the two exponents, respectively, as discussed below.

We also monitored the normalized intensity increase upon exchange that occurs for the C_α–H b band ($I_{N,C-H}(t) = (I_{C-H}(t) - I_{C-H}(0)) / (I_{C-H}(\infty) - I_{C-H}(0))$) that is plotted together

with $1-I_{N,AmII'}(t)$ in Fig. 10. At 0.0, 0.2 and 0.35 M NaClO₄ concentrations, the AmII' and C_α-H b band kinetics fully overlap.

We also monitored the normalized intensities of the 1567 cm⁻¹ AmII band and the 1250 cm⁻¹ and 1267 cm⁻¹ components of the AmIII₃ band. At 0.0, 0.2 and 0.35 M NaClO₄ concentrations, the AmII and the AmIII₃ band kinetics overlap those of the AmII' band. However, at 0.8 M NaClO₄, the $I_{N,C-H}(t)$ kinetics are faster than that of $1-I_{N,AmII'}(t)$ as shown in Fig. 10. The AmII band kinetics is very close to that of the AmII' band, while the 1250 cm⁻¹ and 1267 cm⁻¹ AmIII₃ band kinetics fall between the $I_{N,C-H}(t)$ and $1-I_{N,AmII'}(t)$ kinetics (not shown).

In the presence of NaClO₄ PLL has multiple conformations in equilibrium (α , π , turn, PPII, 2.5₁-helix etc.). All conformations with NH peptide bonds contribute to the AmII and AmIII bands, with the α -helical-like conformations contributing with lower Raman cross sections. Uniquely, the C_α-H b band is contributed only by NH peptide bond extended conformations. All ND peptide bonds contribute to the AmII' band, with the α -helical conformation showing a lower Raman cross section.

The exchange rates of peptide bond NH in the extended conformations can be selectively calculated from the C_α-H b band exchange kinetics shown in Fig. 10. At 0.0, 0.2 and 0.35 M, identical exchange rates are observed for the C_α-H b band from the extended structure, as from the AmII' band contributed by the predominant α -helix conformation. In sharp contrast, in 0.8 M NaClO₄, the extended conformation monitored 0.024 min⁻¹ single exponential C_α-H b band calculated exchange rate is almost twice faster than that of the mainly α -helix AmII' band of 0.014 min⁻¹. As discussed above, the hydrogen exchange rates of peptide bond NH in the extended conformations slow as NaClO₄ concentration increases. The Am II and AmIII monitored D/H exchange kinetics show a poorer S/N because the magnitude of the spectral changes are attenuated by the overlapping slow α -helix exchange rates, and by the weaker, broader AmIII bands that contain partially overlapping contributions of α -helix-like and extended conformations.

The H/D exchange rates for PLL in solution should follow the scheme:

where 'Helix' and 'Extended' represent the α -helix-like and extended conformations, $k_{ex,helix}$ and $k_{ex,extended}$ refer to the H/D exchange rate constants of the peptide bond NH in the α -helix-like and extended conformations, respectively, k_α and $k_{-\alpha}$ are rate constants for the conformational transformation between the α -helix-like and extended conformations, and the equilibrium constant $K = k_{-\alpha}/k_\alpha = [extended]/[helix]$. At 0.0 M NaClO₄, only the extended conformation exists. At 0.2, 0.35, and 0.8 M NaClO₄, $K = k_{-\alpha}/k_\alpha$ can be calculated from Table 1, to be 2.0, 0.82, and 0.39, respectively.

The overlapping time dependencies of the normalized intensities of the AmII', C_α-H b, AmII, and the 1250, 1267 cm⁻¹ AmIII₃ band kinetics for 0.2 and 0.35 M NaClO₄ concentrations indicate that the hydrogen exchange rates for the extended and α -helix-like conformations are either identical, or the conformational interconversion rate is significantly faster than our H/D exchange measurement time scale. Since we expect the α -helix-like conformation to have a very slow D/H rate, we must conclude that we are actually

monitoring the D/H exchange of the extended conformation, and that conformational exchange is very fast between the α -helix-like conformation and extended conformations.

However, the 0.8 M NaClO₄ solution PLL sample of Fig. 10 shows faster kinetics for the extended conformation C $_{\alpha}$ -H b band than for the predominantly α -helix AmII' band indicating that the extended structure undergoes D/H exchange faster than the α -helix and that the conformational interconversion rates are slow compared to the D/H exchange rates.

Thus, the conformational interconversion rate of PLL decreases as the NaClO₄ concentration increases. ClO₄⁻ binding probably increases the activation barriers restricting the conformational interconversion rate. Previous studies showed that peptide conformational dynamics could be influenced by the presence of salts.⁴¹ Further, Dzubiella et al. MD simulations showed that the α -helical (un)folding kinetics of a ala-based peptide are slowed by the binding of ions to peptide charged groups.⁴²

The 0.8 M NaClO₄ AmII' band normalized intensity time dependence is not well fit to a single exponential, presumably because the extended conformation and α -helix-like conformations have different exchange rates. The double exponential fit to $1 - I_{N,AmII'}(t)$:

$$1 - I_{N,AmII'}(t) = 1 - ae^{-k_1 t} - (1 - a)e^{-k_2 t} \quad (3)$$

where $k_1 = 0.022 \pm 0.001 \text{ min}^{-1}$, $k_2 = 0.00097 \pm 0.00128 \text{ min}^{-1}$ and $a = 0.77 \pm 0.04$.

The amide NH exchange rate obtained from the excellent single exponential fit to the C $_{\alpha}$ -H b band exchange kinetics is $k = 0.024 \text{ min}^{-1}$, which is very close to the k_1 rate obtained from the AmII' band kinetics. This similarity suggests that the k_1 kinetics derives from the D/H exchange of the extended conformations, leaving the k_2 kinetics to derive from the α -helix-like conformations.

Assuming we have only extended and α -helix-like conformations we expect parameter a to be given as:

$$a = \frac{\sigma_{\text{extended}} f_{\text{extended}}}{\sigma_{\text{extended}} f_{\text{extended}} + \sigma_{\text{helix}} f_{\text{helix}}} \quad (4)$$

where f_{extended} and f_{helix} are the PLL fraction of extended and α -helix-like conformations in the presence of 0.8 M NaClO₄. σ_{helix} and σ_{extended} are the AmII' band cross sections of the α -helix-like and extended conformations.

Assuming that the Raman cross sections of the extended and α -helix-like conformations are identical in the presence and absence of NaClO₄, $\sigma_{\text{extended}}/\sigma_{\text{helix}}$ can be estimated from the measured pH~3 PLL UVRR AmII' band intensities in D₂O in the absence and presence of 0.8 M NaClO₄ (Fig. 7). Using the AmI' band as an internal intensity standard, the measured AmII' band integrated intensity ratio in the absence and presence of 0.8 M NaClO₄, $I_{AmII',0MClO_4}/I_{AmII',0.8MClO_4}$ is measured to be 1.78 ± 0.05 . Since

$$\frac{I_{AmII',0MClO_4}}{I_{AmII',0.8MClO_4}} = \frac{f_{\text{extended},0MClO_4} \sigma_{\text{extended},AmII'} + f_{\text{helix},0MClO_4} \sigma_{\text{helical},AmII'}}{f_{\text{extended},0.8MClO_4} \sigma_{\text{extended},AmII'} + f_{\text{helix},0.8MClO_4} \sigma_{\text{helical},AmII'}} \quad (5)$$

Given our determined PLL extended conformational fractions at 0.0 and 0.8 M NaClO₄ are 100% and 28%, we estimate the AmII' band Raman cross section ratio of the extended conformation to the α -helix-like conformation is 2.55 ± 0.15 . Using this cross section ratio and eqn. 4, we calculate $a = 0.50 \pm 0.02$, which is somewhat less than that of the fitted kinetic result (0.77). This indicates our 2-state H/D exchange model fails to fully account for the PLL exchange kinetics.

NaClO₄ decreases the H/D exchange rates of extended conformation peptide bonds

Fig. 11, which shows the NaClO₄ concentration dependence of the exponential exchange rates, shows that increasing NaClO₄ concentrations rapidly decreases the H/D exchange rate of the peptide bond ND. We can model this dependence at 0, 0.2 and 0.35 M NaClO₄ concentrations where the conformational exchange rates are fast by assuming that ClO₄⁻ is a competitive inhibitor of the PLL H/D exchange. In the Appendix we calculate the impact of ClO₄⁻ binding on the extended PLL D/H exchange rate by modeling our observed exchange rate, k_{obs} of Fig. 11:

$$\frac{k_{obs}}{f_{extended}} = \frac{k_{ex}}{1 + K[ClO_4^-]} \quad (6)$$

Fitting the apparent exchange rate constant, k_{obs} at different NaClO₄ concentrations (Fig. 11) gives an intrinsic D/H exchange rate of $k_{ex} = 0.95 \pm 0.03$, and a ClO₄⁻ association equilibrium constant to extended PLL of $K = 21 \pm 4$. Thus, ClO₄⁻ binds to extended PLL at a site close to the peptide bond N–D to protect it against D/H exchange that is primarily catalyzed by OH⁻. This base catalyzed reaction occurs through abstraction of the amide NH by direct interaction with OH⁻.⁴³ The model above predicts that increasing the pH should speed up the D/H exchange rate because of an increased competing OH⁻ concentration. We find that the D/H exchange of PLL at pH 6.5 in 0.8 M NaClO₄ is faster than the dead time of our exchange measurement (< 1 min); the pH 6.5 in 0.8 M NaClO₄ PLL UVRR is identical to that at pH 2.8, indicating identical PLL conformations.

The ClO₄⁻ induced decrease in the peptide bond NH exchange rates in the extended PLL conformations indicates a specific ClO₄⁻ binding geometry that protects the backbone peptide bond ND from exchange. Fig. 12 shows how ClO₄⁻ ion could interact both with extended conformation peptide bond ND and the neighboring lysine –NH₃⁺ group through hydrogen bonding and electrostatic interactions to transiently form stable ring structures that protects the amide NH from exchange.

CONCLUSIONS

We examined the impact of NaClO₄ on poly-L-lysine's conformational equilibrium and its hydrogen exchange kinetics. The presence of NaClO₄ induces charged PLL to fold into α -helical-like conformations. Extended conformations such as the PPII and 2.5₁-helix conformations dominate the PLL equilibria at low NaClO₄ concentrations. In contrast, at higher NaClO₄ concentrations α -helical-like conformations that consist of turn, α -helix and π -helix/bulge conformations dominate. The conformational population Ψ angle distribution and the Gibbs free energy landscape of PLL were calculated along the Ramachandran Ψ angle folding coordinate.

We find that the D/H exchange kinetics of PLL at 0, 0.2, 0.35 and 0.8 M NaClO₄ concentrations slow as the NaClO₄ concentration increases. The kinetics of the AmII' and C α -H b bands are identical at 0, 0.2 and 0.35 M NaClO₄ concentrations which indicates that

the conformational interconversion rates are significantly faster than our H/D exchange measurement time scale.

At 0.8 M NaClO₄ concentration the AmII' band intensity kinetics requires a double exponential fit. In addition, the C_α-H b band extended structure D/H kinetics are faster than that of the AmII' band indicating that the PLL conformational interconversion rate is slowed by NaClO₄. The exchange rate of the extended structure is much faster than that of the α -helix structure. These results allow us to determine the association constant of perchlorate to the extended PLL conformation, and allow us to propose a binding geometry.

Acknowledgments

This work is supported by NIH Grant No. 1R01EB009089.

References

1. Baldwin RL, Rose GD. Trends Biochem Sci. 1999; 24:26. [PubMed: 10087919]
2. Creighton, TE. Protein Structure. 2. W. H. Freeman; New York: 1997.
3. Dill KA. Biochemistry. 1985; 24:1501. [PubMed: 3986190]
4. Dill KA. Protein Sci. 1999; 8:1166. [PubMed: 10386867]
5. Dobson CM, Sali A, Karplus M. Angew Chem Int Ed. 1998; 37:868.
6. Dill KA, Ozkan SB, Shell MS, Weikl TR. Annu Rev Biophys. 2008; 37:289. [PubMed: 18573083]
7. Anfinsen CB. Science. 1973; 181:223. [PubMed: 4124164]
8. Anfinsen CB, Scheraga HA. Adv Prot Chem. 1975; 29:205.
9. Haber E, Anfinsen CB. J Biol Chem. 1962; 237:1839. [PubMed: 13903380]
10. Levinthal C. J Chim Phys. 1968; 65:44.
11. Royer CA. Chem Rev. 2006; 106:1769. [PubMed: 16683754]
12. Gruebele M. Annu Rev Phys Chem. 1999; 50:485. [PubMed: 15012420]
13. Ervin J, Sabelko J, Gruebele M. J Photochem Photobio B. 2000; 54:1.
14. Amunson KE, Ackels L, Kubelka J. J Am Chem Soc. 2008; 130:8146. [PubMed: 18529000]
15. Barth A, Zscherp C. Q Rev Biophys. 2002; 35:369. [PubMed: 12621861]
16. Buchner GS, Murphy RD, Buchete NV, Kubelka J. Biochim. Biophys. Acta. Proteins and Proteomics. 2011; 1814:1001.
17. Borgia A, Williams PM, Clarke J. Annu Rev Biochem. 2008; 77:101. [PubMed: 18412537]
18. Schuler B. ChemPhysChem. 2005; 6:1206. [PubMed: 15991265]
19. Schuler B, Lipman EA, Eaton WA. Nature. 2002; 419:743. [PubMed: 12384704]
20. Rhoades E, Gussakovsky E, Haran G. Proc Natl Acad Sci U S A. 2003; 100:3197. [PubMed: 12612345]
21. Cecconi C, Shank EA, Bustamante C, Marqusee S. Science. 2005; 309:2057. [PubMed: 16179479]
22. Suarez M, Jaramillo A. J R Soc Interface. 2009; 6:S477. [PubMed: 19324680]
23. Morra G, Meli M, Colombo G. Curr Protein Pept Sci. 2008; 9:181. [PubMed: 18393887]
24. Thirumalai D, O'Brien EP, Morrison G, Hyeon C. Annu Rev Biophys. 39:159. [PubMed: 20192765]
25. Dearborn DG, Wetlaufer DB. Biochem Biophys Res Commun. 1970; 39:314. [PubMed: 5421933]
26. Painter PC, Koenig JL. Biopolymers. 1976; 15:229. [PubMed: 1247653]
27. Ma L, Ahmed Z, Mikhonin AV, Asher SA. J Phys Chem B. 2007; 111:7675. [PubMed: 17567063]
28. Bykov S, Lednev I, Ianoul A, Mikhonin A, Munro C, Asher SA. Appl Spectrosc. 2005; 59:1541. [PubMed: 16390595]
29. Lednev IK, Karnoup AS, Sparrow MC, Asher SA. J Am Chem Soc. 1999; 121:8074.
30. Ma L, Ahmed Z, Asher SA. J Phys Chem B. 2011; 115:4251. [PubMed: 21413713]

31. Mikhonin AV, Myshakina NS, Bykov SV, Asher SA. J Am Chem Soc. 2005; 127:7712. [PubMed: 15913361]
32. Ebert G, Kuroyanagi Y. Polymer. 1982; 23:1147.
33. Ebert G, Kuroyanagi Y. Polymer. 1982; 23:1154.
34. Goto Y, Takahashi N, Fink AL. Biochemistry. 1990; 29:3480. [PubMed: 2162192]
35. Woody, RW. Circular Dichroism and the Conformational Analysis of Biomolecules. Fasman, GD., editor. Plenum Press; New York: 1996. p. 25
36. Reed J, Reed TA. Anal Biochem. 1997; 254:36. [PubMed: 9398343]
37. Ozdemir A, Lednev IK, Asher SA. Biochemistry. 2002; 41:1893. [PubMed: 11827535]
38. Mikhonin AV, Bykov SV, Myshakina NS, Asher SA. J Phys Chem B. 2006; 110:1928. [PubMed: 16471764]
39. Asher SA, Ianoul A, Mix G, Boyden MN, Karnoup A, Diem M, Schweitzer-Stenner R. J Am Chem Soc. 2001; 123:11775. [PubMed: 11716734]
40. Asher SA, Mikhonin AV, Bykov SB. J Am Chem Soc. 2004; 126:8433. [PubMed: 15238000]
41. Dzubiella J. J Phys Chem B. 2010; 114:7098. [PubMed: 20438069]
42. von Hansen Y, Kalcher I, Dzubiella J. J Phys Chem B. 2010; 114:13815. [PubMed: 20942455]
43. Dempsey CE. Prog Nucl Magn Reson Spectrosc. 2001; 39:135.

APPENDIX

NaClO₄ inhibits H/D exchange of extended PLL at low pH

Our results in Fig. 11 that show a slowing H/D exchange rate as the ClO₄⁻ concentration increases indicates that the deuterated amide ND peptide bonds in PLL are protected from exchange by binding of ClO₄⁻ and PLL to form the ND-ClO₄ complexes. The exchange process of the extended conformations is described by

The first and third steps are the association reactions that occur between extended conformations of PLL and ClO₄⁻, where k_a and k_{-a} are the association and dissociation rate constants. k_{ex} is the D to H exchange rate constant. The ClO₄⁻ association and dissociation steps are very fast and the reaction is assumed to be in ClO₄⁻ exchange equilibrium.

The equilibrium association reaction constant between extended PLL and ClO₄⁻ is

$$K = \frac{k_a}{k_{-a}} = \frac{[ND - ClO_4]}{[ND][ClO_4^-]} \quad (A1)$$

So that

$$[ND - ClO_4] = K[ND][ClO_4^-] \quad (A2)$$

$$[ND - ClO_4] + [ND] = K[ND][ClO_4^-] + [ND] \quad (A3)$$

$$[ND] = \frac{[ND] + [ND - ClO_4]}{1 + K[ClO_4^-]} \quad (A4)$$

During the H/D exchange process, at low and moderate ClO_4^- concentrations (0.2 and 0.35 M) we can assume fast conformational exchange such that the extended and α -helix-like conformations are in equilibrium. Thus, the extended conformational fraction is:

$$f_{extended} = \frac{[ND] + [ND - ClO_4]}{[ND] + [ND - ClO_4] + [\alpha - D] + [\alpha - D - ClO_4]} \quad (A5)$$

where $[\alpha - D]$ and $[\alpha - D - ClO_4]$ are the unbound and bound ClO_4^- states of α -helix-like conformations. The measured overall hydrogen exchange rate is

$$-\frac{d([ND] + [ND - ClO_4] + [\alpha - D] + [\alpha - D - ClO_4])}{dt} = k_{ex}[ND] \quad (A6)$$

The hydrogen exchange rate of the extended conformation is obtained from eqn. A6:

$$-\frac{d([ND] + [ND - ClO_4])}{dt} = f_{extended} k_{ex} [ND] \quad (A7)$$

Substituting eqn. A4 into A7 yields

$$-\frac{d([ND] + [ND - ClO_4])}{dt} = f_{extended} k_{ex} \frac{[ND] + [ND - ClO_4]}{1 + K[ClO_4^-]} \quad (A8)$$

Reorganizing A8,

$$-\frac{d([ND] - [ND - ClO_4])}{[ND] - [ND - ClO_4]} = \frac{f_{extended} k_{ex}}{1 + K[ClO_4^-]} dt \quad (A9)$$

Thus, the apparent exchange rate constant of the extended conformation is a function of the ClO_4^- concentration:

$$k_{obs} = \frac{f_{extended} k_{ex}}{1 + K[ClO_4^-]} \quad (A10)$$

Rearrange the eqn. A10, and eqn. 6 is obtained.

Figure 1.

(A) 204 nm UVRR spectra of PLL at pH 3 and 20 °C at 0, 0.1, 0.2, 0.35, 0.5 and 0.8 M NaClO₄ concentrations. (B) Deconvolution of the 20 °C 204 nm UVRR PLL spectrum in the absence of NaClO₄ at pH 3.

Figure 2.
NaClO₄ concentration dependence of 0.87 mg/ml PLL CD spectra at pH 3 and 20 °C.

Figure 3. Comparison of PLL α -helix-like conformation fraction calculated from CD and UVRR at different NaClO_4 concentrations at pH 3 and 20 °C.

Figure 4. NaClO₄ concentration dependence of the AmIII region of the 204 nm UVRR spectra of the calculated α -helix-like PLL conformations at pH 3 and 20 °C. See text for details.

Figure 5.

NaClO₄ concentration dependence of the Ramachandran Ψ angle distribution of PLL at pH 3 and 20 °C, assuming the T2 conformation at $\Psi = 114^\circ$. The possible $\Psi = 14^\circ$ conformation is not shown. The extended structure Ψ angles are assumed to be those of PLL in the absence of NaClO₄ as in Fig. 1B.

Figure 6.

Calculated Gibbs free energy landscape for PLL at different NaClO_4 concentrations at pH 3 and 20 °C. The dotted lines show the turn regions, assuming that the T2 turn occurs at $\Psi = 114^\circ$.

Figure 7. 204 nm UVRR spectra of PLL in the absence and presence of 0.8 M NaClO₄ in H₂O and D₂O at pH 3/pD 3 and at 20 °C.

Figure 8.

(A) pH 2.8 UVRR of PLL at 20 °C hydrogen exchange at 28, 33, 43, and 103 s after H₂O addition in the absence of NaClO₄. (B) pH 2.8 UVRR PLL hydrogen exchange in 0.8 M NaClO₄ obtained at 3, 8, 12, 22, 32, 42, 64, 102 and 150 min after 0.8 M NaClO₄ H₂O solution addition.

Figure 9. Time dependence of normalized AmII' band intensity of PLL due to H-exchange at pH 2.8 at 20 °C at different NaClO₄ concentrations. The thin solid lines show single exponential fitting.

Figure 10.

Comparison of $I_{N,C-H}(t)$ (blue) and $1-I_{N,AmII'}(t)$ (red), exchange kinetics of the C α -H band and AmII' band at pH 2.8 at 20 °C. The black lines are single exponential (0.2, 0.35 M) and double exponential (0.8 M) fitting curves.

Figure 11. NaClO₄ concentration dependence of extended PLL exchange rates. The solid line shows the eqn. 6 fitting result.

Figure 12. Interactions between ClO_4^- and PLL in the PPII conformation to protect the peptide NH from exchange. The ClO_4^- ion binds to the amide NH and neighboring lysine $-\text{NH}_3^+$ group to form a ring structure.