Published in final edited form as:

J Am Chem Soc. 2011 August 24; 133(33): 13169–13175. doi:10.1021/ja2046364.

Charge Delocalization and Enhanced Acidity in Tricationic Superelectrophiles

Rajasekhar Reddy Naredla^a, Chong Zheng^a, Sten O. Nilsson Lill^b, and Douglas A. Klumpp^a

Douglas A. Klumpp: dklumpp@niu.edu

^aDepartment of Chemistry and Biochemistry, Northern Illinois University, DeKalb, Illinois 60115, United States

^bDepartment of Chemistry, University of Gothenburg, SE-412 96 Gothenburg, Sweden

Abstract

This paper presents the results from studies related to the chemistry of tricationic superelectrophiles. A series of triaryl methanols were ionized in Brønsted superacids and the corresponding tricationic intermediates were formed. The trications are found to participate in two types of reactions, both are characteristic of highly charged organic cations. One set of reactions occurs through charge migration. A second set of reactions occurs through deprotonation of an unusually acidic site on the tricationic species. One of the tricationic intermediates has been directly observed by low temperature NMR spectroscopy. These highly charged ions and their reactions have also been studied using density functional theory calculations. As a result of charge migration, electron density at a carbocation site is found to *increase* with progression from monocationic to pentacationic structures.

Introduction

During the mid-1970s, Olah proposed the concept of superelectrophilic activation.¹ Several published studies had previously described the high reactivities of monocationic electrophiles in superacidic media. This suggested that the monocationic electrophiles were interacting with the acidic media to enhance their reactivities. Thus, protonation of a monocationic species such as the nitronium ion (**1**) could produce an ion with an increasing positive charge (**2**, eq 1). In the limiting cases, even dicationic species (**3**) are considered to be involved in the chemistry. While many examples of dicationic superelectrophiles have been described,² few examples of tricationic species have been studied. Most tricationic and tetracation species are formed as resonance stabilized ions.³ Little is known about the reactivities of small organic ions bearing such a large amount of positive charge.⁴ In the following Article, we describe studies in which a reactive carbocation site interacts with two adjacent cationic charges. Reactions of the tricationic species with arenes lead to products consistent with charge migration. Under some conditions, these reactive intermediates undergo cyclizations to produce novel heterocyclic ring systems. The tricationic intermediates are further studied by spectroscopic and theoretical methods.

Correspondence to: Douglas A. Klumpp, dklumpp@niu.edu.

Supporting Information **Available**: Detailed experimental procedures, characterization data, ¹H and ¹³C NMR spectra for new compounds, ¹³C NMR spectra for ions **26** and **27**, computational methods and results, and crystallographic data for compounds **11** and **18**.

(1)

Results and Discussion

Our studies began with an examination of the chemistry of triarylmethanols (**4-8**, **14-15**) by reaction in superacidic $\text{CF}_3\text{SO}_3\text{H}$ (Table 1). The triarylmethanol substrates were prepared readily from reactions of the appropriate ketones or esters with organolithium reagents. Two types of conversions were observed, depending on whether or not benzene was present in the mixture. In the presence of C_6H_6 , alcohol **4** gives product **9** in 89% yield at 60°C (entry 1). If the reaction was done at lower temperatures, then the starting material was recovered. Similar results were obtained with imidazole and benzimidazole-based substrates (**5**, **7**, **8**; entries 2,4,5). When the reaction was done with a substrate (**6**) bearing the 2-naphthyl substituent, nucleophilic attack occurred at the C-1 position of the naphthyl ring to give product **11** in good yield. The structure of compound **11** was established by ^1H and ^{13}C NMR spectra, mass spectrum analysis, and X-ray crystallography.⁵

In the absence of benzene, reactions of the triarylmethanols leads to cyclization products (entries 6-10). For example, compound **4** reacts in $\text{CF}_3\text{SO}_3\text{H}$ at 60°C to provide the pyrido[1,2-*a*]indole derivative (**16**) in 81% yield. Benzimidazole derivatives **5-6** also react to give cyclization products (**17** and **18**). In the reaction of **6**, cyclization occurs regioselectively at the 1-position of the naphthyl ring to give product **18** (confirmed by X-ray crystal structure).⁵ The pyrido[1,2-*a*]indole derivatives **19** and **20** were formed by reactions of compounds **14** and **15**, respectively. Product **20** is the only major product from **15**, indicating a preference for cyclization at the pyridine ring. The vinyl-substituted derivative (**21**) does not undergo cyclization in $\text{CF}_3\text{SO}_3\text{H}$, however a low yield of cyclization product (**22**) may be obtained from reaction in the stronger acid system $\text{CF}_3\text{SO}_3\text{H}\text{-SbF}_5$ (ca. 10:1; eq 2). These types of cyclizations involving *N*-heterocycles are not well known, as the only other example was described by Posselt (involving cyclization between a 4-methoxyphenyl group and a pyridine or pyridinium ring).⁶ Nevertheless, our results indicate that novel heterocyclic ring systems are prepared by the superacid promoted cyclizations of the triarylmethanols.

(2)

When triarylmethanols **23** or **24** are reacted under similar conditions, the carbocations **25** and **26** are produced (Figure 1). However with benzene, no Friedel-Crafts products are formed from the monocation **25** or dication **26**. With simple heating to 60°C (no benzene), cyclization products are also not obtained.⁷ Compound **4** generates the tricationic intermediate **27** in heated superacid. Thus, trication **27** exhibits enhanced reactivity

compared to **25** and **26**. When substrate **4** is reacted with substituted arenes, product formation (**28-30**) occurs with good regioselectivity. Reaction of compound **4** with toluene in $\text{CF}_3\text{SO}_3\text{H}$ gives **28** as the major product, isolated in 72% yield. Gas chromatography analysis of the crude product mixture indicated a *para,ortho/meta* ratio of about 10:1. A competition experiment was also done with substrate **4** in a $\text{CF}_3\text{SO}_3\text{H}$ -promoted reaction with toluene/benzene (1/1 ratio). Analysis of the product mixture showed high selectivity for the toluene with k_t/k_b about 30:1. Similar results are obtained from reactions of **4** with bromobenzene and 4-(3-phenylpropyl)pyridine. Products **29** and **30** are formed in good yields with high regioselectivities.

In addition to arene nucleophiles, we also attempted to capture the electrophilic intermediate(s) with *n*-type nucleophiles, including carbon monoxide, fluoride, and water. Although carbon monoxide often reacts efficiently with carbocationic intermediates,⁸ no adducts of this nucleophile could be isolated in reactions with **4** in superacid. When compound **4** is reacted in $\text{CF}_3\text{SO}_3\text{H}$ and KHF_2 is added, then fluoride adduct **31** may be isolated in 51% yield (eq 3; compound **4** is also recovered). Interestingly, nucleophilic attack occurs at the methine carbon rather than at the ring carbon. A similar reaction of **4** with $\text{CF}_3\text{SO}_3\text{H}$ – followed by water – leads to formation of the starting material **4**, rather than functionalization at the remote position.

(3)

In our mechanistic studies, the intermediates and transition states were studied using density functional theory (DFT). Geometries for these structures were initially optimized in the gas-phase at the M06/6-31G(d) level of theory⁹ using Jaguar.¹⁰ Gibbs' free energy corrections (ΔG) were calculated and structures were verified to be minima or transition states by inspection of the number of imaginary frequencies. In a second step, single-point energies of the optimized structures were calculated using M06/cc-pVTZ(-f),^{11,12} giving a basis set correction term (ΔBS). In a final step, solvent phase geometry optimizations were done. Using a Poisson–Boltzmann solver as implemented in Jaguar,¹³ electrostatic solvation effects from the surroundings were calculated using the SCRF method with $\text{CF}_3\text{SO}_3\text{H}$ as a solvent (dielectric constant=77.4, probe radius=2.5985274).¹⁴ The free energy (ΔG) and basis set correction terms (ΔBS) were added to the solution phase energy. NMR chemical shifts were calculated using B3LYP/IGLO-II as implemented in Gaussian 03,¹⁵⁻¹⁷ on M06/6-31G(d) gas-phase optimized structures with benzene set to the experimental value of 127.7 ppm as the NMR reference. NBO charges were calculated using NBO 5.0 as implemented in Jaguar.¹⁸ To verify that the M06/6-31G(d) compounds were able to properly describe the σ - and π -complexes, tests were also performed using B3LYP-DCP/6-31+G(d,p), a dispersion-corrected DFT method,^{19,20} resulting in very similar optimized structures as with M06/6-31G(d).

In considering the mechanisms of the reactions, we propose a mechanism involving formation of tricationic intermediates with highly delocalized carbocation centers and equilibria with a dicationic species from deprotonation. The alcohol substrates (**4-8**) react with benzene in the superacid and nucleophilic attack occurs at the remote position on the

aryl substituent group (Figure 2). Thus, compound **4** ionizes in superacid and loss of water produces the tricationic superelectrophile **27**. With formation of the carbocation **27**, positive charge is highly delocalized into the adjacent phenyl ring. Nucleophilic attack occurs at the remote position on the phenyl group to give intermediate **32**, when benzene is present in the reaction mixture. Several examples of charge migration and remote functionalization have previously been reported involving dicationic superelectrophiles.²¹

Product **31** is formed by fluoride attack at the methine carbon, suggesting that some positive charge remains at the methine center. This raises an obvious question: why would the *n*-type nucleophiles (fluoride and water) react at the methine carbon while π -type nucleophiles (benzene and substituted arenes) react at the remote *para*-position of the phenyl group? First, there is likely some degree of steric effects influencing the regiochemistry of nucleophilic attack. Nucleophilic attack at the methine carbon (by benzene) would form the highly congested tetraarylmethane. Secondly, formation of the π -complex may enhance charge separation (Figure 3). Formation of the toluene π -complex (**33a**), and subsequent Wheland intermediate or σ -complex (**34a**), effectively moves the positive charge away from the pyridinium rings. Density functional theory (DFT) calculations were done (*vide infra*) and showed the toluene ring bears +0.62 charge in the π -complex (**33a**) and +0.97 charge in the σ -complex (**34a**). In the case of toluene, steric effects in the π -complex may also be important in the high *para* selectivity.²² Although high positional selectivity is often due to a large energy barrier between the π -complex and σ -complex,²³ this does not appear to be the case with **33a** and **34a**. DFT calculations have indicated that the π -complex resides in a very shallow potential energy well, as small perturbations in structure **33a** easily lead to formation of the σ -complex **34a** as found by DFT geometry optimizations. In the competition experiment, high substrate selectivity was observed ($k_t/k_b \sim 30:1$). This value likely reflects the relative stabilities of the σ -complexes with benzene and toluene.²³ When the π - and σ -complexes are examined, the σ -complex from toluene (**34a**) is 7.9 kcal·mol⁻¹ more stable than the π -complex from toluene (**33a**), while the σ -complex from benzene (**34b**) is only 4.6 kcal·mol⁻¹ more stable than its π -complex with benzene (**34a**).

In reactions of compound **6**, products **11** and **18** were formed. While charge migration occurred into the naphthyl ring system, it did not delocalize to the most remote position (across both rings). Ionization of the benzimidazole rings and the hydroxy group leads to trication **35a** (eq 5). Formation of product **11** occurs by nucleophilic attack at the 1-position of the naphthyl group. This implies a greater contribution from resonance structure **35b**, instead of structure **35c** (the structure with greater charge separation).

(5)

In order to gain further insight into these reactions, low temperature ¹³C NMR experiments were done. Compound **4** was ionized in FSO₃H-SbF₅-SO₂ClF at -60°C and the observed spectrum is consistent with the formation of the tricationic species **27** (Table 2). For comparison, the DFT calculated spectrum of trication **27** is in reasonably good agreement with the results from solution (for complete assignments of ¹³C signals, see the Supporting Information). It is notable that the *para* and *ortho* carbons are significantly deshielded. This

is consistent with the charge delocalization driven by charge-charge repulsive effects. The dicationic ion **26** and the trityl cation (**25**)²⁴ were also studied in solution by low temperature ¹³C NMR, with comparison to calculated spectra. Interestingly, the chemical shift at the methine carbon moves progressively down field from the monocationic trityl cation (**25**), to the dication (**26**), and to the trication (**27**). This is an indication that charge delocalization becomes enhanced with increasing charge. The tetra- and pentacationic structures (**36-37**) were also calculated and the results show further deshielding of the *para*-position and shielding of the methine carbon. While the trityl cation has a calculated chemical shift of δ 144 at the *para* position of the phenyl ring, the pentacation **37** has calculated chemical shift of δ 231 at the *para* position of the phenyl ring.

As a measure of charge delocalization in these systems, natural bond order (NBO) charges, bond lengths and dihedral angles were analyzed from the calculated structures. In a series of related structures, ¹³C NMR chemical shifts may be roughly correlated to charge densities.²⁵ Down field chemical shifts (or deshielding) often corresponds to electron deficient carbon atoms.²⁵ Within the isoelectronic series from the trityl cation (**25**) to the pentacation (**37**), experimental and calculated NMR spectral data suggests that the carbocationic center moves progressively from the methine carbon to the *para* position of the phenyl ring. The calculated NBO charges also show this trend (Table 3). Remarkably, there is almost a complete transfer of positive charge from the methine carbon to the *para* carbon. The methine carbon carries a +0.24 NBO charge in the trityl cation (**25**) and a -0.14 NBO charge in the pentacation (**37**), while the *para* carbon carries a -0.15 NBO charge in **25** and a +0.20 NBO charge in **37**. As the positive charge is moved to the *para* position, there is also increasing double bond character between the methine carbon and the phenyl ring. This is apparent from the decreasing bond dihedral angle (δ) and C-C bond length as charge is added to the system. Thus, the trityl cation **25** has a dihedral angle of 33.2° and C-C bond length of 1.441 Å, while the pentacation **37** has a dihedral bond angle of 11.2° and C-C bond length of 1.391 Å. The dihedral angle and NBO charge are strongly correlated to the overall charge on the cationic structure. Increasing charge leads to progressively greater positive charge at the *para* carbon, while simultaneously increasing electron density at the methine carbon (evident from the negative NBO charges). Although it might be expected that a carbocationic center would exhibit *increasing* positive charge as neighboring groups become more electron deficient, the opposite occurs in this case. The increasing charge of the *N*-heterocycles leads to charge migration across the phenyl ring to the remote *para* position (maximizing the charge-charge separation).

As described in Table 1, compound **4** reacts in CF₃SO₃H to give the pyrido[1,2-*a*]indole derivative (**16**) in the absence of an arene nucleophile. Since ionization of **4** should provide trication **27**, theoretical calculations were also done to study the cyclization of the tricationic species **27** (Figure 4). Calculations suggest that direct cyclization of the trication **27** is unlikely, as the transition state **38** leading to **39** is estimated to be 41 kcal/mol above **27**. The cyclized trication **39** itself is 38 kcal/mol higher in energy. A lower energy pathway was found involving initial deprotonation at the pyridinium ring, proceeding through two minima (**40** and **41**) and giving dication **42** in a slightly exothermic step. Cyclization then occurs with a barrier of 9.8 kcal/mol (**43**) and it gives stabilized species **44** and **45**. Upon basic work up, cation **45** provides the observed product **16**. Although triflate normally would not be expected to deprotonate a pyridinium ring, the high charge density of the trication **27** must lead to enhanced acidity of the pyridinium hydrogen. Previous studies have shown superelectrophiles to possess highly acidic protons.²⁶ With trication **27**, the carbocationic site cannot readily lead to a deprotonation equilibrium, so instead the pyridinium ring undergoes deprotonation.

Ionization of compound **4** in $\text{FSO}_3\text{H-SbF}_5\text{-SO}_2\text{ClF}$ gives NMR spectral data that differs somewhat from the calculated spectrum for trication **27**. While experimental NMR spectra often differ slightly from the calculated spectra, the present case may be the result of an equilibrium between trication **27** and the dication **42**. Rapid proton transfer should average the spectral data from **27** and **42** and this would account for the differences between the calculated and experimental spectra for trication **27**. As shown in Table 2, the experimental data for the compound **4** ionization product – presumably trication **27** – showed a methine resonance at δ 176.8 and *para*-carbon at δ 169.4. The calculated spectrum for **27** has a methine resonance at δ 176.0 and *para*-carbon at δ 178.2. When dication **42** was studied computationally, calculated spectrum has a methine resonance at δ 179.1 and *para*-carbon at δ 161.6 (for a full list of peaks, see Supporting Information section). If a numerical average is made from the calculated chemical shift values of trication **27** and dication **42**, then an averaged spectrum should have methine resonance at δ 177.6 and *para*-carbon at δ 169.9. These averaged values are very close to the experimentally determined values from ionization of **4** in magic acid. Moreover, the calculated energies of **27** and **42** are within 0.5 kcal/mol, suggesting a significant concentration of these two species from the low temperature ionization.

This raises an obvious question: could all of the chemistry simply involve equilibria between dications? This question was examined computationally by considering the step in which the alcohol ionizes to the carbocation. Assuming the alcohol group must be protonated for ionization, this means the dicationic intermediate **46** would of necessity be involved in the chemistry. However, calculations indicate that **46** is about 35 kcal/mol less stable than the dication in which both pyridyl rings are protonated (**47**). This strongly suggests the formation of the tricationic oxonium ion **48**, leading to the proposed tricationic superelectrophile **27**.

As described in Table 1, compound **15** reacts in superacid to give the pyrido[1,2-*a*]indole **20**. This product is somewhat surprising, as a dicationic cyclization might be expected to cyclize into the benzothiazole ring. Assuming ionization of **15** leads to the trication **49**, deprotonation could lead to either **50** or **51**, but only dication **50** should lead to product **20** (Figure 5). However, the pK_a data for the parent heterocycles suggest that the benzothiazole ring should be more acidic (pK_a for pyridine 5.2; pK_a for benzothiazole 1.2).²⁷ This apparent inconsistency may be explained by considering the stabilities of the intermediates **52** and **53**. DFT calculations have shown that cyclization through the pyridine ring leads to an intermediate (**52**) that is 4.6 kcal/mol more stable than a similar intermediate with benzothiazole cyclization (**53**). Likewise, product **20** is almost 4 kcal/mol more stable than its isomer **54**.

Conclusions

We have studied the superacid-promoted reactions of a series of triarylmethanols which generate tricationic intermediates. Compared to analogous mono- and dicationic species, the tricationic intermediates exhibit new reactions resulting from the effects of the closely oriented positive charges. These effects include the tendency for cationic charge to

“migrate” across a phenyl ring and the tendency for neighboring protons to exhibit very high acidities. The migration of charge across organic structures is important in material science applications and in the design of organic based electronics. Our results show that densely charged structures may have high charge mobilities. As a consequence, these tricationic superelectrophiles react with arene nucleophiles at a remote site. Without a nucleophilic reactant, cyclizations occur to provide novel *N*-heterocyclic products. The functionalized heterocycles are produced in good to excellent yields. Computational studies indicate that this chemistry arises from the deprotonation of an unusually acidic pyridinium ring.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the support of the National Science Foundation (CHE-0749907 and CRIF MU-0840504) and the NIH-National Institute of General Medical Sciences (GM085736-01A1). S.O.N.L. gratefully acknowledges the financial support from the Åke Wiberg Foundation.

References

1. Olah GA, Germain A, Lin HC, Forsyth D. *J Am Chem Soc.* 1975; 97:2928–2929.
2. Olah, GA.; Klumpp, DA. *Superelectrophiles and Their Chemistry.* Wiley; NY: 2008.
3. (a) Prakash GKS, Rawdah TN, Olah GA. *Angew Chem Int Ed.* 1983; 22:390–401. (b) Ito S, Morita N, Asao T. *Bull Chem Soc Jpn.* 2000; 73:1865–1874. (c) Head NJ, Prakash GKS, Bashir-Hashemi A, Olah GA. *J Am Chem Soc.* 1995; 117:12005–12006. (d) Rathore R, Burns CL, Green IA. *J Org Chem.* 2004; 69:1524–1530. [PubMed: 14987006] (e) Reddy VP, Rasul G, Prakash GKS, Olah GA. *J Org Chem.* 2003; 68:3507–3510. [PubMed: 12713353] (f) See also: Pagni RM. *Tetrahedron.* 1984; 49:4161–4215.
4. For recent examples of reactive tricationic species see: (a) Ohwada T, Yamagata N, Shudo K. *J Am Chem Soc.* 1991; 113:1364–1373. (b) Kurouchi H, Sugimoto H, Otani Y, Ohwada T. *J Am Chem Soc.* 2010; 132:807–815. [PubMed: 20017483] (c) Zhang Y, Sheets MR, Raja EK, Boblak KJ, Klumpp DA. *J Am Chem Soc.* 2011; 133:8467–8469. [PubMed: 21548654]
5. Crystallographic data found in Supporting Information section.
6. (a) Posselt K. *Arzneim-Forsch.* 1978; 28:1056–1065. [PubMed: 582688] (b) See also: Alarcon HAR, Soldatenkov AT, Soldatova SA, Samalyoa AI, Obando HU, Prostakov NS. *Khimiya Geterotsiklicheskikh Soedinenii.* 1993:1233–1238.
7. Decomposition products are obtained.
8. Sommer J, Bukala J. *Acc Chem Res.* 1993; 26:370–376.
9. Zhao Y, Truhlar D. *Theor Chem Acc.* 2008; 120(1-3):215–241.
10. Jaguar, version 7.7. Schrodinger, LLC; New York, NY: 2010.
11. Dunning TH. *J Chem Phys.* 1989; 90:1007–1023.
12. Woon DE, Dunning TH. *J Chem Phys.* 1993; 98:1358–1371.
13. Marten B, Kim K, Cortis C, Friesner RA, Murphy RB, Ringnalda MN, Sitkoff D, Honig B. *J Phys Chem.* 1996; 100:11775–11788.
14. Lira AL, Zolotukhin M, Fomina L, Fomine S. *J Phys Chem A.* 2007; 111:13606–13610. [PubMed: 18052258]
15. Frisch, MJT., et al. *Gaussian 03, Revision D.02.* Gaussian Inc.; Wallingford, CT: 2004.
16. Stephens PJ, Devlin FJ, Chabalowski CF, Frisch MJ. *J Phys Chem.* 1994; 98:11623–11627.
17. Kutzelnigg, W.; Fleischer, U.; Schindler, M. *NMR, Basic Principles and Progress.* Diehl, P.; Fluck, E.; Günther, H.; Kosfield, R.; Seelig, J., editors. Vol. 23. Springer-Verlag; Berlin: 1990. p. 165-262.

18. Glendening, ED.; Badenhoop, JK.; Reed, AE.; Carpenter, JE.; Bohmann, JA.; Morales, CM.; Weinhold, F. Theoretical Chemistry Institute, University of Wisconsin; Madison, WI: 2001. NBO 5.0. <http://www.chem.wisc.edu/~nbo5>
19. Mackie ID, DiLabio GA. *J Phys Chem A*. 2008; 112:10968–10976. [PubMed: 18828578]
20. Nilsson Lill SO. *J Mol Graph Mod*. 2010; 29(2):178–187.
21. (a) Ohwada T, Okabe K, Ohta T, Shudo K. *Tetrahedron*. 1990; 46:7539–7555. (b) Shudo K, Ohta T, Okamoto T. *J Am Chem Soc*. 1981; 103:645–653. (c) Nakamura S, Sugimoto H, Ohwada T. *J Am Chem Soc*. 2007; 129:1724–1732. [PubMed: 17249674] (d) O'Connor MJ, Boblak KN, Topinka MJ, Kindelin PJ, Briski JM, Zheng C, Klumpp DA. *J Am Chem Soc*. 2010; 132:3266–3267. [PubMed: 20178387]
22. Nelson KL, Brown HC. *J Am Chem Soc*. 1951; 73:5605–5607.
23. Olah GA. *Acc Chem Res*. 1971; 4:240–248.
24. Abarca B, Asensios G, Ballesteros R, Luna C. *Tetrahedron Lett*. 1986; 27:5657–5660.
25. (a) Forsyth DA, Sandel BB. *J Org Chem*. 1980; 45:2391–2394. (b) Forsyth DA, Spear RJ, Olah GA. *J Am Chem Soc*. 1976; 98:2512–2518.
26. (a) Nenajdenko VG, Shevchenko NE, Balenkova ES. *Chem Rev*. 2003; 103:229–282. [PubMed: 12517185] (b) Olah GA, Grant JL, Spear RJ, Bollinger JM, Serianz A, Sipos G. *J Am Chem Soc*. 1976; 98:2501. (c) Larsen JW, Bouis PA. *J Am Chem Soc*. 1975; 97:6094–6102. (d) Olah GA, Calin M. *J Am Chem Soc*. 1968; 90:4672–4675. (e) Olah GA, Reddy VP, Rasul G, Prakash GKS. *J Am Chem Soc*. 1999; 121:9994–9998.
27. (a) Perrin, DD. *Dissociation Constants of Organic Bases in Aqueous Solution*. Butterworths; London: 1965. (b) Eicher, T.; Hauptmann, S. *The Chemistry of Heterocycles*. Wiley-VHC; Weinheim: 2003.

Figure 1.
Ionization products (**25-27**) and Friedel-Craft reaction products **28-30**.

Figure 2.

Proposed mechanism for the reactions of trication **27**, leading to product **9**.

Figure 3. Relative energies of Wheland- and pi-complexes with toluene (**33a**, **34a**) and benzene (M06/6-31G* optimized structures, **33b**, **34b**).

Figure 4.
Calculated structures related to the cyclization of trication **27**.

Figure 5.
Cyclization of trication **46** to compound **20** and **51**.

Table 1

Reactions of heterocyclic alcohols in $\text{CF}_3\text{SO}_3\text{H}$ at 60°C (with and without C_6H_6).

Entry	Starting material	Product with C_6H_6	Yield	Entry	Starting material	Product without C_6H_6	Yield
(1)			89%	(6)			81%
(2)			90%	(7)			91%
(3)			91%	(8)			92%
(4)			95%	(9)			88%

Entry	Starting material	Product with C ₆ H ₆	Yield	Entry	Starting material	Product without C ₆ H ₆	Yield
(5)			96%	(10)			82%

Table 2

Observed ^{13}C NMR data from isoelectronic carbocations **25-27** and calculated ^{13}C NMR data for **25-27** and **36-37**.

A, B, C, D	Structure	Charge	^{13}C NMR data from $\text{FSO}_3\text{H-SbF}_5\text{-SO}_2\text{ClF}$	Calculated ^{13}C NMR data: C-methine, C- <i>ipso</i> , C- <i>ortho</i> , C- <i>meta</i> , C- <i>para</i>
CH, CH, CH, CH		1+	211.6 (C+), 144.1, 143.3, 140.6, 131.1	205.1, 138.3, 142.2, 129.5, 144.4
NH ⁺ , CH, CH, CH	26^b	2+	192.7 (C+), 149.9, 148.4, 145.2, 144.1, 137.9, 137.8, 133.0, 132.1, 132.1	184.3, 136.5, 144.3, 134.7, 157.0
NH ⁺ , NH ⁺ , CH, CH	27^a	3+	176.8 (C+), 169.4, 150.8, 149.6, 148.2, 141.2, 140.3, 139.4, 137.2, 135.6	176.0, 138.7, 145.8, 141.9, 178.2
NH ⁺ , NH ⁺ , NH ⁺ , CH	36	4+		150.2, 149.0, 146.0, 148.9, 201.2
NH ⁺ , NH ⁺ , NH ⁺ , NH ⁺	37	5+		125.8, 165.0, 143.9, 155.7, 231.0

^a Spectrum taken at -60°C.

^b Spectrum taken at 0°C.

Table 3

Calculated properties of isoelectronic ions **25-27** and **36-37**.

Structure	Charge	NBO charge		Dihedral angle, δ	C-C Bond length (methine-ipso), Å
		C-methine	C-para		
25	1+	+0.24	-0.15	33.2°	1.441
26	2+	+0.15	-0.10	31.1°	1.429
27	3+	+0.12	0.00	24.0°	1.402
36	4+	-0.01	+0.10	19.5°	1.394
37	5+	-0.14	+0.20	11.2°	1.391