Published in final edited form as:

Behav Neurosci. 2012 February ; 126(1): 186–195. doi:10.1037/a0026460.

AN NMDA ANTAGONIST IN THE MPOA IMPAIRS COPULATION AND STIMULUS SENSITIZATION IN MALE RATS

Anna V. Vigdorichik², Bradley P. Parrish¹, Gwen A. Lagoda², Jenna A. McHenry¹, and Elaine M. Hull^{1,2}

¹Department of Psychology, Florida State University, Tallahassee, FL 32306-4301

²Department of Psychology, State University of New York at Buffalo, Buffalo, New York 14260-4110

Abstract

Systemic injections of an NMDA antagonist have been shown to impair mating in male rats. One site where glutamate and its NMDA receptors may contribute to mating is the medial preoptic area (MPOA), which is vital for male sexual behavior. Glutamate is released in the MPOA during copulation, and especially at the time of ejaculation. We report here that the NMDA antagonist MK-801, microinjected into the MPOA, impaired copulatory behavior in sexually naïve as well as experienced males. In animals tested both as naïve and after sexual experience, drug treatment produced more profound impairment in naïve males. In addition, MK-801, microinjected into the MPOA before each of 7 noncopulatory exposures to receptive female rats, resulted in copulatory impairments on a drug-free test on day 8, relative to aCSF-treated animals; their behavior was similar to that of males that had not been pre-exposed to females. Therefore, NMDA receptors in the MPOA contribute to the control of copulation and stimulus sensitization. Glutamate, acting via NMDA receptors, regulates many neural functions, including neuronal plasticity. This is the first demonstration that a similar mechanism in the MPOA sensitizes male rats to the stimuli from a receptive female, and thereby enhances their behavior.

Keywords

Sexual behavior; Medial Preoptic Area; glutamate; NMDA; MK-801; Dizocilpine

Glutamate, acting on N-methyl-D-aspartate (NMDA) receptors plays a key role in a wide range of physiologic and pathologic processes, including synaptic transmission, neuronal development (Debanne, Daoudal, Sourdet, & Russier, 2003), and multiple aspects of neuronal plasticity and long-term memory (LTM) formation (Barker & Warburton, 2008; Brigman et al., 2010; Delint-Ramirez, Salcedo-Tello, & Bermudez-Rattoni., 2008; Hernandez, Andrzejewski, Sadeghian, Panksepp, & Kelly, 2005; Lee & Everitt, 2008;

Correspondence should be addressed to: Dr. Elaine M. Hull Department of Psychology and Program in Neuroscience 1107 West Call Street Tallahassee, FL 32306-4301 Phone: (850) 645-2389 Fax: (850) 644-7739 hull@psy.fsu.edu.

Publisher's Disclaimer: The following manuscript is the final accepted manuscript. It has not been subjected to the final copyediting, fact-checking, and proofreading required for formal publication. It is not the definitive, publisher-authenticated version. The American Psychological Association and its Council of Editors disclaim any responsibility or liabilities for errors or omissions of this manuscript version, any version derived from this manuscript by NIH, or other third parties. The published version is available at www.apa.org/pubs/journals/bne

Current address for Anna Vigdorichik: 443 Allenhurst Rd, Unit B Buffalo, NY 14226-2871 Current address for Bradley Parrish: Drexel University College of Medicine Professional Studies 245 N. 15th Street, NCB Philadelphia, PA 19102

Current address for Gwen Lagoda: Johns Hopkins School of Medicine Brady Urological Institute 600 North Wolfe Street, Marburg 414 Baltimore, MD 21287

Popescu, Saghyian, & Paré, 2007; Zhang Meng, Li, & Han, 2009). Administration of non-competitive NMDA receptor antagonists such as dizocilpine maleate (MK-801) has been shown to block induction, but not maintenance, of long-term potentiation (LTP) (Gilbert & Mack, 1990).

NMDA receptors also contribute to the facilitative effects of sexual experience on later copulatory proficiency (Fleming & Kucera, 1991; Powell, Dominguez, & Hull, 2003), analogous to their effects on the formation of LTM. However, whereas NMDA antagonists typically affect only the formation, and not the expression, of LTM, systemic administration of an NMDA antagonist did impair the expression of copulation in sexually experienced male rats (Powell et al., 2003). Because glutamate is the major excitatory transmitter throughout the brain, it is not clear where its facilitative effects on male sexual behavior occur.

One possible site where glutamate, acting on NMDA receptors, may normally facilitate male sexual behavior is the medial preoptic area (MPOA). The MPOA is a critical brain area regulating male sexual activity (reviewed in Hull & Rodriguez-Manzo, 2009). The MPOA receives input from (Simerly & Swanson, 1988) and sends projections to (Simerly & Swanson, 1986) other brain areas vital for the initiation and regulation of sexual behavior. Lesions of the MPOA impair mating, and electrical or chemical stimulation of the MPOA enhances copulation and genital reflexes in male rats (reviewed in Hull & Rodriguez-Manzo, 2009). In addition, repeated electrical stimulation of the MPOA resulted in copulation by previously inactive males (Paredes, Haller, Manero, Alvarado, & Ågmo, 1990).

There are several reports of facilitative effects of glutamate in the MPOA on male sexual behavior. In anesthetized male rats, microinjection of glutamate into the MPOA elicited erection (Giuliano et al., 1996) or the urethrogenital reflex (a model of orgasm, Marson & McKenna, 1994). Glutamate is released in the MPOA during copulation and increases dramatically during ejaculation; furthermore, increased glutamate levels, resulting from reverse-dialysis of glutamate uptake inhibitors, facilitated copulation (Dominguez, Gil, & Hull, 2006). Glutamate, reverse-dialyzed into the MPOA, increased dopamine release via a nitric oxide-dependent process (Dominguez, Muschamp, Schmich, & Hull, 2004); MPOA dopamine has also been shown to facilitate male sexual behavior (reviewed in Hull & Rodriguez-Manzo, 2009).

Sexual experience significantly increases the efficiency of copulation. Sexually experienced male rats tend to mount, intromit and ejaculate faster than inexperienced males (Bialy, Rydz, & Kaczmarek, 2000; Dewsbury, 1969; Larsson, 1978). They also copulate more readily following various lesions or castration, or in a novel environment, compared to naïve animals (de Jonge et al., 1989; Lisk & Heimann, 1980; Pfaus & Wilkins, 1995). Even repeated noncopulatory exposure to an estrous female enhances copulation on the first sexual experience, compared to unexposed naïve controls (de Jonge, Oldenburger, Louwerse, & Van de Pol, 1992; Lagoda, Muschamp, & Hull, 2004; Powell, Dominguez, & Hull, 2003). The effects of repeated sexual experience or noncopulatory exposure may result from functional alterations in circuitry mediating male sexual behavior that may in turn enhance responsiveness to sex-related stimuli. However, the neurobiological mechanisms of such stimulus sensitization remain unclear.

The current studies investigated the role of NMDA glutamate receptors in the MPOA in male rat sexual behavior and in stimulus sensitization to sex-related input. In Experiments 1 and 2, we tested the effects of MK-801, microinjected into the MPOA of groups of sexually naïve and sexually experienced males, respectively, before copulation tests. In order to test

directly the effects of sexual experience as well as of the drug, Experiment 3 tested drug effects in the same males, first when they were naïve and then after they had been given sexual experience. Finally, Experiment 4 tested whether NMDA receptors in the MPOA also play a role in stimulus sensitization resulting from repeated noncopulatory exposures to a female. MK-801 was microinjected into the MPOA of sexually naïve males before each of 7 daily noncopulatory exposures to an estrous female rat. Sexual behavior was scored on a drug-free test on the eighth day, thereby avoiding any potential effects of MK-801 on motor behavior.

Methods

Animals

Adult male Long-Evans/Blue Spruce rats weighing 250-300g (Harlan, Indianapolis, IN), were housed individually in large plastic cages in a temperature- and humidity-controlled environment on a 14:10 hour light:dark cycle, with lights off at 11:00 and on at 21:00. Food and water were available *ad libitum*. Animals were weighed daily to ensure health. Ovariectomized female rats of the same strain were used as stimulus animals. All procedures were in accordance with the National Institutes of Health guidelines for the care and use of animals and were approved by the local Institutional Animal Care and Use Committees.

Surgery

Male rats were anesthetized with ketamine hydrochloride (80 mg/kg) and xylazine hydrochloride (10 mg/kg) and then implanted with a 23-g stainless steel guide cannula ending 1 mm above the MPOA. The coordinates with respect to bregma were +2.1 mm AP, +4 mm ML, and 7.3 mm DV (Pellegrino, Pellegrino, & Cushman, 1979). Three jeweler's screws were placed in the skull surrounding the cannula, allowing the cannula to be secured when dental acrylic was added. A stylet was placed into the guide cannula after the dental acrylic dried in order to prevent entry of contaminants. After surgery, rats were immediately administered Buprenex (1.5 mg/0.1 ml), and received ground rat chow mixed with water for 1 day. Each animal was allowed 1 week for recovery before beginning testing. Females of the same strain were anesthetized with ketamine hydrochloride (50 mg/kg) and xylazine hydrochloride (4 mg/kg) and ovariectomized using bilateral flank incisions; Buprenex (1.5 mg/0.1 ml) was administered after the surgery. Females were given at least one week after ovariectomy before being used as stimulus animals. They were injected with 10 µg estradiol benzoate 48 hrs prior to testing and with 500 µg of progesterone 4 hrs before testing. Female receptivity was confirmed with a stud male before testing.

Drug

Dizocilpine maleate (MK-801) was obtained from Sigma-Aldrich (St. Louis, MO). For the drug condition, MK-801 was dissolved in aCSF (Dulbecco's solution) at a concentration of 2.5 µg/µl and administered at 0.5 µl/min for a total of 1.0 µl. The dose was the most effective dose used by Maliszewska-Scislo & Trojnar (2000) and did not appear to impair motor behavior in a preliminary mating test. For the vehicle condition, aCSF was administered at 0.5 µl/min for a total of 1.0 µl.

Procedure

Behavior testing

Copulatory behavior tests were performed in a test room, under dim red light. After several minutes of adaptation, a stimulus female was presented into the male's home cage. The following measures were recorded: mount latency (ML), the time from the introduction of the female to the first mount; intromission latency (IL), the time from the introduction of the

female to the first intromission; ejaculation latency (EL), the time from the first intromission to the first ejaculation; postejaculatory interval (PEI), the time from an ejaculation to the next intromission; total mount frequency (MF); total intromission frequency (IF); total ejaculation frequency (EF); mount frequency before the first ejaculation (MF₁); intromission frequency before the first ejaculation (IF₁); intromission ratio total (IR_T) and intromission ratio before the first ejaculation (IR₁) [$IR = I/(M+I)$]; and inter-intromission interval (III = EL₁/IF₁). If an animal did not ejaculate, MF₁, IF₁, IR₁, and III were not calculated for that animal.

Preliminary testing of doses

In a small number of animals three doses of MK-801 were microinjected into the MPOA, followed by 30 min copulation testing. The treatments were counterbalanced on a weekly basis. The results showed that the highest dose (5.0 µg/µl) produced significant motor and coordination impairment, evidenced by awkward interactions with the female, such as trying to mount from her side. In addition, rats that received this dose became increasingly aggressive. Animals that received the lowest dose (1.0 µg/µl) did not show any abnormal behavior and did not have any motor and coordination impairment. However, this dose did not affect copulation, compared to aCSF treatment. The middle dose (2.5 µg/µl), taken from Maliszewska-Scislo & Trojnar (2000), did not produce any observable motor impairments, but did result in copulatory impairment.

Experiment 1

Twenty sexually naïve male rats were randomly divided into 2 groups, MK-801 (n=10) and control (n=10). Prior to behavioral testing, the stylet was removed, and the injection cannula was inserted into the guide cannula; it extended 1 mm below the end of the guide cannula into the MPOA. 1.0 µl of drug (2.5 µg/µl) or aCSF was infused unilaterally via a syringe pump at the rate of 0.5 µl/min, with the injection cannula remaining in place for at least 60 sec after the completion of each microinjection to facilitate drug diffusion. Each rat moved freely in his home cage during the injection. To ensure complete diffusion of the drug around the injected area, behavioral testing was not started until 10 min after removal of the injection cannula. Three animals were removed from this study due to incorrect cannula placement, resulting in 9 animals remaining in the MK-801 group and 8 animals in the aCSF group.

Experiment 2

Methods of Experiment 2 were the same as in Experiment 1, except that prior to testing, animals were allowed to achieve one ejaculation on each of three twice-weekly experience days. If an animal did not ejaculate on one experience day, he was given an additional experience day, so that all males had achieved 3 ejaculations one week before drug vs. vehicle testing began. Thus, 30 sexually experienced male rats were divided into two groups, MK-801 (n=15) and aCSF (n=15). MK-801 or aCSF was microinjected into the MPOA in counterbalanced order in 2 weekly tests, followed by behavioral testing with receptive female rats.

Experiment 3

Twenty-two sexually naïve animals were tested using methods similar to Experiment 1. Two drug-treated males fought repeatedly with the stimulus female, so their tests were ended early. Seven animals had misplaced cannulae; their data were removed, resulting in the following groups: MK-801 (n=9), aCSF (n=6). The animals were then given up to 4 additional mating experiences over 2 weeks to achieve 3 ejaculations and become sexually experienced. They were then given two additional mating tests. For the first, they received

the opposite treatment from what they had received on their naïve test, and for the second, they received the same treatment as on their naïve test. In other words, if they received drug on the naïve test, they received aCSF on the first post-experience test and drug on the second, and if they received aCSF on the naïve test, they received drug and then aCSF on the two post-experience tests.

Experiment 4

The protocol for this experiment was adapted from Powell et al. (2003) and Lagoda et al., (2004). Sixty sexually naïve male rats were divided into 3 groups, MK-801 (n=20), aCSF (n=20), and control (n=20). Males in the first two groups received daily microinjections of either MK-801 or aCSF for 7 consecutive days; 10 min after the microinjection, each male was exposed to an inaccessible receptive female rat. The female rat was placed in a wire mesh cage (12.5 × 26 × 15 cm) immediately above the male's cage, thus allowing the male to smell, see and hear her, but not copulate with her. The unexposed control group received daily microinjections of aCSF but were never exposed to females. On the 8th day, animals from all 3 groups were given drug-free copulation tests with a receptive female for 30 min after the first intromission, or for 30 min if no intromission occurred. Nine animals were removed due to incorrect cannula placement, thus resulting in the following groups: MK-801 (n=14), aCSF (n=19), unexposed control (n=18).

Because of concerns that daily microinjections of MK-801 might compromise the health of the animals, at the end of this experiment, the body weights of drug-treated animals were statistically compared with those of the animals that received aCSF; no significant difference was found between the body weights of the 2 groups. Since the MPOA also plays an important role in body temperature regulation, we also measured body temperatures of 5 drug-treated and 5 aCSF-treated animals, using a rectal thermometer, following the female exposures. No significant difference was found between temperatures of drug and aCSF animals.

Histology

Upon completion of each experiment, animals were anesthetized with a lethal dose of sodium pentobarbital (65 mg/kg) and decapitated. Brains were removed, frozen to −17°C, and 40 µm coronal sections were cut of the MPOA and any other tissue that contained cannula tracks. Sections were mounted on gelatin-coated slides. Verification of cannula placements was done using a photomagnifier; sections were compared with the images from Brain Maps III (Swanson, 2004). Figure 1 displays cannula placements for all experiments.

Statistics

Experiment 1—Independent-sample t-tests were used to analyze MF, IF, EF and IR₁. In addition, ML, IL, EL, and III were analyzed only from animals that exhibited the relevant behaviors. Chi-square tests were performed on the numbers of animals that mounted, intromitted or ejaculated.

Experiment 2—One-way repeated measures analyses of variance (ANOVA) were used to analyze MF, IF, EF, ML, IL, EL and IR₁. As in Experiment 1, latencies for animals that did not show a particular behavior were not analyzed.

Experiment 3—Multiple statistical analyses were used to examine within- and between-subjects effects. Because very few naïve rats copulated after MK-801 injections, latencies to intromit and ejaculate, inter-intromission interval, and postejaculatory interval could not be analyzed. Chi-square tests were used to compare the numbers of naïve males in the drug and vehicle conditions that mounted, intromitted, or ejaculated.

Two-way ANOVAs compared the effects of drug and experience on frequencies and latencies of mounts, intromissions and ejaculations. For those comparisons, the data from animals tested with aCSF on the naïve test were compared with data from the aCSF treatment after they had gained experience, and data from animals tested initially with drug on the naïve test were compared with data from the drug test after experience. Individual planned contrast comparisons were also used to probe significant drug effects for separate naïve and experienced tests. Additionally, paired t-tests were used to compare drug vs. aCSF in the counterbalanced tests of experienced males. Chi-square analyses compared the numbers of MK-801- and aCSF-treated experienced males that mounted, intromitted, and ejaculated.

Experiment 4—One-way between-subjects ANOVAs and Bonferroni post hoc tests were used to calculate significance of ML, IL, EL, PEI, MF, IF, EF, and IR₁ on all animals on the drug-free test day. Chi-square tests were performed on the numbers of animals that mounted, intromitted, or ejaculated. For all tests $p < .05$ was considered significant.

Results

Experiment 1

Independent t tests revealed significant drug-induced decreases in the numbers of mounts [$t(15) = 3.6, p < .01$] and intromissions [$t(15) = 2.14, p < .05$ (Figure 2A)]. Drug-treated males also had lower intromission ratios than did aCSF-treated animals [$t(15) = 3.3, p < .01$] (Figure 2A). In addition, animals that received MK-801 had significantly longer mount latencies [$t(15) = 2.3, p < .05$] than did aCSF-treated males (Figure 2B). Chi-square analyses revealed that more aCSF-treated animals intromitted [$\chi^2(1) = 3.6, p < .05$] and ejaculated [$\chi^2(1) = 7.2, p < .01$] than animals that received MK-801.

Experiment 2

Repeated-measures ANOVAs revealed significant decreases in numbers of intromissions [$F(1,27) = 4.23, p < .05$] and ejaculations [$F(1,27) = 4.31, p < .05$ (Figure 3A)] in drug-treated males. Intromission ratios were also significantly decreased [$F(1,27) = 7.7, p < .01$] for the drug-treated animals (Figure 3A). In addition, MK-801 increased mount [$F(1,20) = 14.44, p < .01$], intromission [$F(1,18) = 19.96, p < .01$], and ejaculation [$F(1,15) = 8.07, p < .01$] latencies (Figure 3B). Furthermore, fewer drug-treated animals intromitted [$\chi^2(1) = 13.2, p < .01$] and ejaculated [$\chi^2(1) = 9.3, p < .01$], compared to the aCSF-treated animals.

Experiment 3

Drug-treated naïve males were impaired on several measures. Only 3 out of 9 drug-treated naïve males mounted, compared to 6 out of 6 for vehicle-treated animals ($\chi^2 = 4.0, p < .05$); 2 out of 9 intromitted, compared to 4 out of 6 for controls [$\chi^2(1) = 5.44, p < .01$]; and 1 out of 9 ejaculated, compared to 3 of 6 for controls [$\chi^2(1) = 8.60, p < .01$]. There were no significant differences in mount latency or mount, intromission or ejaculation frequencies.

Two-way ANOVAs, comparing data from naïve animals with their matched treatment after experience, revealed several significant main effects, but no significant interactions. There was a significant increase in MF as a result of experience [$F(1,13) = 4.95, p < .05$], but there was no effect of drug (Figure 4A). For IF, MK-801 decreased intromissions [$F(1,13) = 7.46, p < .05$], and experience increased intromissions [$F(1,13) = 16.53, p < .001$] (Figure 4B). Similarly, for EF, MK-801 decreased the number of ejaculations [$F(1,13) = 7.69, p < .02$], whereas experience increased ejaculations [$F(1,13) = 15.04, p < .01$] (Figure 4C). There were trends for decreased intromission latencies in experienced vs. naïve animals [$F(1,6) = 7.69, p = .05$] and for increased ejaculation latencies in drug-treated vs. aCSF-treated

animals [$F(1,3) = 10.45$, $p = .05$]. Planned contrast comparisons of the means for separate naïve and experienced tests revealed that in naïve males MK-801 significantly decreased the numbers of intromissions [$F(1,13) = 9.446$, $p = .009$, Figure 4B] and ejaculations [$F(1,13) = 5.368$, $p = .037$, Figure 4C], compared to aCSF treatments. In experienced males there were no significant effects of MK-801, although there was a trend for MK-801 to decrease the number of ejaculations [$F(1,13) = 3.64$, $p = .079$], compared to aCSF treatment (Figure 4C). Thus, there were greater effects of MK-801 in animals tested while naïve than after they had gained sexual experience.

Paired t tests on post-experience counterbalanced drug and aCSF tests revealed that MK-801 significantly decreased intromission [$t(14) = 3.74$, $p < .01$] and ejaculation [$t(14) = 3.15$, $p < .01$] frequencies [Figure 4D], increased mount latencies [$t(10) = 2.25$, $p < .05$], and produced a trend for increased intromission latencies [$t(10) = 2.03$, $p = .07$] [Figure 4E]. Chi-square tests revealed that sexual experience increased the numbers of drug-treated males that were able to mount ($X^2(1) = 4.1$, $p < .05$), intromit ($X^2(1) = 5.8$, $p < .05$) or ejaculate ($X^2(1) = 8.9$, $p < .01$); experience did not significantly affect numbers of vehicle-treated males that mounted, intromitted, or ejaculated. Thus, there were complex effects of both drug and experience on sexual behavior.

Experiment 4

MK-801 significantly impaired stimulus sensitization. One-way ANOVAs revealed significant main effects of MK-801 on mount [$F(2,51) = 5.83$, $p < .01$], intromission [$F(2,51) = 9.30$, $p < .01$], and ejaculation [$F(2,51) = 5.21$, $p < .01$] frequencies and on intromission ratio 1 [$F(2,46) = 3.23$, $p < .05$]. Bonferroni post hoc tests revealed that aCSF-treated exposed animals had significantly more mounts, intromissions, and ejaculations and higher intromission ratios than the MK-801-treated exposed animals (Figure 5). In addition, control unexposed animals had significantly more mounts than the MK-801-treated exposed animals, but fewer mounts and intromissions than aCSF-treated exposed males.

Chi-square analyses revealed that more aCSF-treated exposed [$X^2(1) = 6.17$, $p < .05$] or control unexposed [$X^2(1) = 9.11$, $p < .01$] animals mounted, compared to MK-801-treated exposed animals. Also significantly more aCSF-treated exposed animals exhibited intromissions than drug-treated exposed [$X^2(1) = 10.45$, $p < .01$] or control unexposed animals [$X^2(1) = 4.94$, $p < .05$]. Finally, more aCSF-treated exposed animals ejaculated, compared to MK-801-treated exposed males [$X^2(1) = 8.97$, $p < .01$].

Discussion

Microinjections of MK-801, a selective and potent noncompetitive antagonist at NMDA receptors (Wong et al., 1986), into the MPOA of male rats decreased the numbers of mounts, intromissions, and/or ejaculations and increased latencies for those measures in sexually naïve and experienced male rats in Experiments 1 and 2. We had expected sexually naïve animals to be more susceptible to the inhibitory effects of MK-801 than sexually experienced animals; however, in Experiments 1 and 2 there was little difference in the effectiveness of the drug in these groups. According to previous reports, sexual behavior in sexually naïve animals is usually disrupted more easily when compared to sexually experienced animals (de Jonge et al., 1989). Also, we have shown in a similar study that sexually naïve animals that received microinjections of the NOS inhibitor L-NAME before a copulation test displayed more deficits in sexual behavior than the same animals after they had received experience (Lagoda et al., 2004). One possible explanation for the apparently similar effectiveness of the drug in the present experiments is the different numbers of animals in Experiments 1 and 2. There were 20 in Experiment 1, half in each group, whereas in Experiment 2 there were 30 that received counterbalanced treatments, resulting in greater

statistical power. In our previous experiment with L-NAME, the animals were first tested while sexually naïve; the same animals were then given sexual experience and tested with both drug and aCSF in counterbalanced order. Thus, unlike current Experiments 1 and 2, the naïve and experienced animals in the previous experiment were the same animals.

Therefore, Experiment 3 was designed to test the same animals when they were sexually naïve and again after attaining at least 3 ejaculations on weekly mating tests experiences. Indeed, there were greater drug effects when males were naïve than after they had gained experience. Fewer drug-treated naïve animals mounted, intromitted, or ejaculated, compared to controls. However, after the animals had gained sexual experience, there were no significant drug-related differences in numbers of animals performing those behaviors, although there were experience-related differences. Similarly, planned comparison statistical tests revealed significant drug-related decreases in intromissions and ejaculations only in naïve males. More sexually experienced males were able to mount, intromit, and ejaculate, regardless of drug treatment. On the other hand, there were both drug- and experience-related effects on numbers of mounts, intromissions, and ejaculations. MK-801 decreased IF and EF and lengthened EL, while sexual experience increased MF, IF, and EF and decreased IL. Thus, there were complex effects of both drug and experience.

In Experiment 4, microinjections of MK-801 before each of 7 daily noncopulatory female exposures significantly impaired animals on the drug-free test on day 8, compared with aCSF-treated exposed animals. Drug-treated males displayed fewer mounts, intromissions and ejaculations, compared with the aCSF-treated exposed group and fewer mounts than the unexposed control group. Also, the aCSF-treated exposed group displayed more mounts and intromissions than the unexposed control group. Indeed, these results indicate that NMDA receptors in the MPOA are important for the control of both copulation and stimulus sensitization.

The greater number of mounts by unexposed control animals, compared to MK-801 exposed males, was unexpected. One possible explanation is that there might be some deficit in the basal regulation of normal brain activity of animals repeatedly exposed to MK-801. In other words, basal levels of glutamatergic activity in the MPOA may play an important role in the facilitation of male sexual behavior. By microinjecting MK-801 for 7 consecutive days, glutamate could not stimulate NMDA receptors normally. Thus, repeated antagonism of NMDA receptors may inhibit a mechanism that up-regulates normal responsiveness to sexually relevant stimuli.

There is evidence that repeated systemic administration of MK-801 increased binding of low-affinity nicotinic acetylcholine receptors in the hypothalamus, hippocampus and colliculus of rats and also NMDA binding in the hippocampus (Levin et al., 2005). However, while muscarinic acetylcholine receptors in the MPOA have been shown to reduce the threshold for ejaculation (Hull et al., 1988a, 1988b), there is no evidence regarding nicotinic receptors in the MPOA affecting male sexual behavior. Also, 5 and 10 days of systemic MK-801 treatment increased the phosphorylation of enzymes in the frontal cortex of rats that regulate protein synthesis (Yoon et al., 2008) or pro-survival and anti-apoptotic activity (Seo et al., 2007). Again, whether local drug administration in the MPOA would have a similar effect is unknown, although it would seem unlikely that such changes would induce the sexual impairment observed in Experiment 4. Finally, in monkey prefrontal cortex there was decreased extracellular glutamate and dopamine after 13 days of i.m. injections of MK-801 (Tsukada et al., 2005). If a similar effect were produced in the MPOA with local microinjections, that could explain the deficits observed in the present experiments, since both dopamine and glutamate in the MPOA facilitate male sexual behavior (reviewed in Hull & Rodriguez-Manzo, 2009).

Sensitization is an increase in the strength of a specific stimulus following repeated presentations, in this case, the receptive female. Olfaction is critical for successful expression of male reproductive behavior among many rodent species. Olfactory bulbectomy or complete removal of the vomeronasal organ produces deficits in male sexual behavior in rats (Edwards, Griffs, & Tardival, 1990; reviewed in Hull & Rodriguez-Manzo, 2009). Projections from the main and accessory olfactory bulbs travel to the amygdala, which integrates sexually relevant sensory information and relays it to the MPOA via the stria terminalis and the ventral amygdalofugal pathway (Meisel, Lumia, & Sachs, 1980). Furthermore, gonadal steroid receptors in the amygdala, in its projection to the bed nucleus of the stria terminalis (BNST), and in the MPOA enable transmission of the relevant information (reviewed in Newman, 1999; Wood, 1997). A process of consolidation of such sensory input may explain the better sexual performance of animals that received aCSF before exposure to inaccessible receptive females, compared to the animals that received MK-801 before exposure and those that received aCSF but were not exposed to females.

Previous evidence for experience-induced sensitization in the MPOA is provided by Lumley & Hull (1999). Males that had 10 previous copulatory experiences had greater ejaculation-induced immunoreactivity (ir) to c-Fos, the protein product of the immediate-early gene *c-fos*, a measure of neural activity. This occurred despite the fact that the experienced males had somewhat fewer intromissions preceding the ejaculation. Thus, each intromission may have been more effective in the experienced males. Furthermore, activation of NMDA receptors is necessary for mating-induced activation of Fos-ir in the MPOA (Dominguez et al., 2007). The latter study also showed that virtually all Fos-ir neurons in the medial preoptic nucleus (MPN) contained NMDA receptors, and that mating increased phosphorylation of NMDA receptors in the MPN. Finally, it reported that microinjections of MK-801 (1.25 µg/0.25 µl) bilaterally into the MPOA decreased the number of ejaculations and increased ejaculation latency, postejaculatory interval, and mounts preceding ejaculation in sexually experienced males. Therefore, NMDA receptors are important both for performance of mating behavior and for stimulus sensitization.

There are several reports of facilitative effects of glutamate in the MPOA on male sexual behavior. In anesthetized male rats microinjection of glutamate into the MPOA elicited erection (Giuliano et al., 1996) or the urethrogenital reflex (a model of orgasm, Marson & McKenna, 1994). Glutamate is released in the MPOA during copulation and increases dramatically during ejaculation; and increased glutamate levels, resulting from reverse-dialysis of glutamate uptake inhibitors, facilitated copulation (Dominguez et al., 2006). The glutamate released in the MPOA may arise, at least in part, from glutamatergic projections from the posterodorsal medial amygdala, as has been demonstrated in male gerbils (Simmons & Yahr, 2003).

One way in which NMDA receptors may facilitate sensitization of sexual behavior is by increasing the production of nitric oxide (NO). Neuronal NO synthase (nNOS) is activated by Ca^{2+} , admitted via NMDA receptors (reviewed in Garthwaite, 1991; Luo & Zhu, 2011; Yun, Dawson, & Dawson, 1996). NO has been implicated in the control of male sexual behavior (reviewed in Hull & Rodriguez-Manzo, 2009; Simmons & Yahr, 2011; Succu et al., 2008). We have previously reported that NO is essential for sensitization in an experiment similar to the present one (Lagoda et al., 2004). In addition, glutamate, reverse-dialyzed into the MPOA elicits dopamine release via NO (Dominguez et al., 2004). We have previously reported that dopamine in the MPOA facilitates male sexual behavior (reviewed in Hull & Rodriguez-Manzo, 2009).

One possible explanation for the impairment of sexual behavior of MK-801-treated animals is a decrease in a process similar to LTP, in which rapid stimulation of afferent neurons

increases responsiveness of the recipient neurons. This change can last for hours, days, or weeks and may lay a foundation for more permanent changes, such as the construction of new synapses between the neurons (Bennett, 2000). LTP is associated with increased presynaptic neurotransmitter release and enhanced postsynaptic receptor activity, including insertion of AMPA receptors into the postsynaptic membrane (reviewed in Rao & Finkbeiner, 2007; Malenka & Bear, 2004). NMDA receptor channel opening and the resultant calcium influx are necessary for initiating LTP; inhibition of NMDA receptors may impede the cascade of events that would normally lead to consolidation of sexually relevant stimuli in male rats. We suggest that the presence of sexual stimuli, which elicit the release of glutamate in the MPOA (Dominguez et al., 2006), contribute to sensitization to those stimuli and also facilitate copulation.

In summary, acute microinjections of MK-801 into the MPOA decreased the numbers and increased the latencies of mounts, intromissions, and/or ejaculations in sexually naïve and experienced male rats. Repeated microinjections of MK-801 into the MPOA before each of 7 noncopulatory exposures to an estrous female also prevented the improvement that normally would have resulted from such exposures. These results suggest that glutamate in the MPOA facilitates the initiation of sexual behavior in sexually naïve male rats, the progression of copulation in sexually experienced males, and the enhancement of copulation that results from repeated non-copulatory exposures to a female.

Acknowledgments

This research was supported by National Institutes of Mental Health grants MH R01-40826 and MH K02-001714 to EMH. The authors thank Angela A. Loewke for assistance with behavioral testing and Christopher Robison for much assistance with figures.

References

- Barker GR, Warburton EC. NMDA receptor plasticity in the perirhinal and prefrontal cortices is crucial for the acquisition of long-term object-in-place associative memory. *The Journal of Neuroscience*. 2008; 28:2837–2844. [PubMed: 18337414]
- Bennett MR. The concept of long term potentiation of transmission at synapses. *Progress in Neurobiology*. 2000; 60:109–137. [PubMed: 10639051]
- Bialy M, Rydz M, Kaczmarek L. Precontact 50-kHz vocalizations in male rats during acquisition of sexual experience. *Behavioral Neuroscience*. 2000; 114:983–990. [PubMed: 11085613]
- Brigman JL, Wright T, Talani G, Prasad-Mulcare S, Jinde S, Seabold GK, Mathur P, Davis MI, Bock R, Gustin RM, Colbran RJ, Alvarez VA, Nakazawa K, Delpire E, Lovinger DM, Holmes A. Loss of GluN2B-containing NMDA receptors in CA1 hippocampus and cortex impairs long-term depression, reduces dendritic spine density, and disrupts learning. *The Journal of Neuroscience*. 2010; 30:4590–4600. [PubMed: 20357110]
- Debanne D, Daoudal G, Sourdet V, Russier M. Brain plasticity and ion channels. *Journal of Physiology Paris*. 2003; 97:403–414.
- Delint-Ramírez I, Salcedo-Tello P, Bermudez-Rattoni F. Spatial memory formation induces recruitment of NMDA receptor and PSD-95 to synaptic lipid rafts. *Journal of Neurochemistry*. 2008; 106:1658–1668. [PubMed: 18700282]
- de Jonge FH, Oldenburger WP, Louwerse AL, Van de Pol NE. Changes in male copulatory behavior after sexual exciting stimuli: Effects of medial amygdala lesions. *Physiology & Behavior*. 1992; 52:327–332. [PubMed: 1523261]
- Dewsbury DA. Copulatory behaviour of rats (*Rattus norvegicus*) as a function of prior copulatory experience. *Animal Behaviour*. 1969; 17:217–223. [PubMed: 5349578]
- Dominguez JM, Balfour ME, Lee HS, Brown JL, Davis BA, Coolen LM. Mating activates NMDA receptors in the medial preoptic area of male rats. *Behavioral Neuroscience*. 2007; 121:1023–1031. [PubMed: 17907833]

- Dominguez JM, Gil M, Hull EM. Preoptic glutamate facilitates male sexual behavior. *The Journal of Neuroscience*. 2006; 26:1699–1703. [PubMed: 16467517]
- Dominguez JM, Muschamp JW, Schmich JM, Hull EM. Glutamate-evoked dopamine release in the medial preoptic area is mediated by nitric oxide: Implications for male rat sexual behavior. *Neuroscience*. 2004; 125:203–210. [PubMed: 15051159]
- Edwards DA, Griffs KT, Tardival C. Olfactory bulb removal: Effects on sexual behavior and partner preference in male rats. *Physiology & Behavior*. 1990; 48:447–450. [PubMed: 2267253]
- Fleming AS, Kucera C. Sexual experience effects are blocked both by the protein-synthesis inhibitor, cycloheximide, and by the noncompetitive NMDA antagonist, MK-801. *Behavioral and Neural Biology*. 1991; 56:319–329. [PubMed: 1836946]
- Garthwaite J. Glutamate, nitric oxide and cell-cell signaling in the nervous system. *Trends in Neuroscience*. 1991; 14:60–67.
- Gilbert ME, Mack CM. The NMDA antagonist, MK-801, suppresses long-term potentiation, kindling, and kindling-induced potentiation in the perforant path of the unanesthetized rat. *Brain Research*. 1990; 519:89–96. [PubMed: 2204470]
- Giuliano F, Rampin O, Brown K, Courtois F, Benoit G, Jardin A. Stimulation of the medial preoptic area of the hypothalamus in the rat elicits increases in intracavernous pressure. *Neuroscience Letters*. 1996; 209:1–4. [PubMed: 8734895]
- Hernandez PJ, Andrzejewski ME, Sadeghian K, Panksepp JB, Kelley AE. AMPA/kainate, NMDA, and dopamine D1 receptor function in the nucleus accumbens core: a context-limited role in the encoding and consolidation of instrumental memory. *Learning and Memory*. 2005; 12:285–295. [PubMed: 15930507]
- Hull EM, Bitran D, Pehek EA, Holmes GM, Warner RK, Band LC, Clemens LG. Cholinergic regulation of male sexual behavior in rats: Effects of intra-cerebrally microinjected agonists. *Pharmacology, Biochemistry & Behavior*. 1988; 31:169–174.
- Hull EM, Pehek EA, Bitran D, Warner RK, Holmes GM, Band LC, Bazzett T, Clemens LG. Cholinergic regulation of male sexual behavior in rats: Effects of an intracerebrally microinjected antagonist. *Pharmacology, Biochemistry & Behavior*. 1988; 31:175–178.
- Hull, EM.; Rodriguez-Manzo, G. Male Sexual Behavior.. In: Pfaff, Donald, editor. *Hormones, Brain and Behavior*. Second Edition. Vol. 1. Elsevier Press; Amsterdam: 2009. p. 5-65.
- Lagoda G, Muschamp JW, Vigdorchik A, Hull, E. M. A nitric oxide synthesis inhibitor in the medial preoptic area inhibits copulation and stimulus sensitization in male rats. *Behavioral Neuroscience*. 2004; 118:1317–1323. [PubMed: 15598140]
- Larsson, K. Experiential factors in the development of sexual behavior.. In: Hutchison, JB., editor. *Biological determinants of sexual behaviour*. Wiley; New York: 1978. p. 55-86.
- Lee JL, Everitt BJ. Appetitive memory reconsolidation depends upon NMDA receptor-mediated neurotransmission. *Neurobiology, Learning and Memory*. 2008; 90:147–154.
- Lisk RD, Heimann J. The effects of sexual experience and frequency of testing on retention of copulatory behavior following castration in the male hamster. *Behavioral and Neural Biology*. 1980; 28:156–171. [PubMed: 7362582]
- Lumley LA, Hull EM. Effects of a D1 antagonist and of sexual experience on copulation-induced Fos-like immunoreactivity in the medial preoptic nucleus. *Brain Research*. 1999; 829:55–68. [PubMed: 10350530]
- Luo CX, Zhu DY. Research progress on neurobiology of neuronal nitric oxide synthase. *Neuroscience Bulletin*. 2011; 27:23–35. [PubMed: 21270901]
- Maliszewska-Scislo M, Trojnar W. Effect of unilateral injection of MK-801 into the area of A10 cells on feeding evoked by stimulation of homologous area in the contralateral hemisphere. *Acta Neurobiologiae Experimentalis*. 2000; 60:489–494. [PubMed: 11200176]
- Marson L, McKenna KE. Stimulation of the hypothalamus initiates the urethro-genital reflex in male rats. *Brain Research*. 1994; 638:103–108. [PubMed: 8199850]
- Meisel RL, Lumia AR, Sachs BD. Effects of olfactory bulb removal and flank shock on copulation in male rats. *Physiology & Behavior*. 1980; 25:383–387. [PubMed: 7443807]
- Newman SW. The medial extended amygdala: A node in the mammalian social behavior network. *Annals of the New York Academy of Sciences*. 1999; 877:242–257. [PubMed: 10415653]

- Paredes RG, Haller AE, Manero MC, Alvarado R, Ågmo A. Medial preoptic area kindling induces sexual behavior in sexually inactive male rats. *Brain Research*. 1990; 515:20–26. [PubMed: 2357558]
- Pellegrino, LJ.; Pellegrino, AS.; Cushman, AJ. *A Stereotaxic Atlas of the Rat Brain*. 2nd ed.. Academic Press; New York: 1979.
- Pfaus JG, Wilkins MF. A novel environment disrupts copulation in sexually naïve but not experienced male rats: Reversal with naloxone. *Physiology & Behavior*. 1995; 57:1045–1049. [PubMed: 7652023]
- Popescu AT, Saghyian AA, Paré D. NMDA-dependent facilitation of corticostriatal plasticity by the amygdala. *Proceedings of the National Academy of Sciences U S A*. 2007; 104:341–346.
- Powell WS, Dominguez JM, Hull EM. An NMDA antagonist impairs copulation and the experience-induced enhancement of male sexual behavior in the rat. *Behavioral Neuroscience*. 2003; 117:69–75. [PubMed: 12619909]
- Seo MS, Kim SH, Ahn YM, Kim Y, Jeon WJ, Yoon SC, Roh MS, Juhnn YS, Kim YS. The effects of repeated administrations of MK-801 on ERK and GSK-3 β signalling pathways in the rat frontal cortex. *International Journal of Neuropsychopharmacology*. 2007; 10:359–368. [PubMed: 16780607]
- Simerly RB, Swanson LW. The organization of neural inputs to the medial preoptic nucleus of the rat. *Journal of Comparative Neurology*. 1986; 246:312–342. [PubMed: 3517086]
- Simerly RB, Swanson LW. Projections of the medial preoptic nucleus: *APhaseolis vulgaris* leucoagglutinin anterograde tract-tracing study in the rat. *Journal of Comparative Neurology*. 1988; 270:209–242. [PubMed: 3259955]
- Simmons DA, Yahr P. Nitric oxidergic cells related to ejaculation in gerbil forebrain contain androgen receptor and respond to testosterone. *Journal of Comparative Neurology*. 2011; 519:900–915. [PubMed: 21280043]
- Simmons DA, Yahr P. GABA and glutamate in mating-activated cells in the preoptic area and medial amygdala of male gerbils. *Journal of Comparative Neurology*. 2003; 459:290–300. [PubMed: 12655511]
- Swanson, LW. *Brain maps III: Structure of the rat brain*. 3rd ed.. Elsevier; New York: 2004.
- Succu S, Sanna F, Cocco C, Melis T, Boi A, Ferri GL, Argiolas A, Melis MR. Oxytocin induces penile erection when injected into the ventral tegmental area of male rats: Role of nitric oxide and cyclic GMP. *European Journal of Neuroscience*. 2008; 28:813–821. [PubMed: 18671741]
- Tsukada, H.; Miyasato, K.; Nishiyama, S.; Fukumoto, D.; Kakiuchi, T.; Domino, EF. Nicotine normalizes increased prefrontal cortical dopamine D1 receptor binding and decreased working memory performance produced by repeated pretreatment with MK-801: a PET study in conscious monkeys. 2005. p. 2144–2153.
- Wong EHF, Kemp JA, Priestley T, Knight AR, Woodruff GN, Iversen LL. The anticonvulsant MK-801 is a potent *N*-methyl-D-aspartate antagonist. *Proceedings of the National Academy of Sciences, USA*. 1986; 83:7104–7108.
- Wood RI. Thinking about networks in the control of male hamster sexual behavior. *Hormones and Behavior*. 1997; 32:40–45. [PubMed: 9344690]
- Yoon SC, Seo MS, Kim SH, Jeon WJ, Ahn YM, Kang UG, Kim YS. The effect of MK-801 on mTOR/p70S6K and translation-related proteins in rat frontal cortex. *Neuroscience Letters*. 2008; 434:23–28. [PubMed: 18262357]
- Yun HY, Dawson VL, Dawson TM. Neurobiology of nitric oxide. *Critical Reviews in Neurobiology*. 1996; 10:291–316. [PubMed: 8978984]
- Zhang L, Meng K, Li YH, Han TZ. NR2A-containing NMDA receptors are required for L-LTP induction and depotentiation in CA1 region of hippocampal slices. *European Journal of Neuroscience*. 2009; 29:2137–2144. [PubMed: 19490087]

Figure 1. Location of MPOA injection sites in coronal sections based on the atlas by Swanson (2004) for animals in all experiments. *Closed circles* represent accurate MPOA placements. *Open circles* represent cannula placements outside of the MPOA.

Figure 2.

Effects of intra-MPOA administration of MK-801 on copulation in sexually naïve males, Experiment 1. A) Mean ($\pm$ SE) total mounts, intromissions, and ejaculations; and intromission ratio for the first ejaculatory series, in sexually naïve animals treated with MK-801 (2.5 μ g/ μ l) or aCSF in the MPOA. * $p < .05$, ** $p < .01$. B) Mean ($\pm$ SE) latencies for mounts, intromissions, and ejaculations in sexually naïve animals treated with MK-801 (2.5 μ g/ μ l) or aCSF in the MPOA. * $p < .05$.

Figure 3.

Effects of intra-MPOA administration of MK-801 on copulation in sexually experienced males, Experiment 2. A) Mean ($\pm$ SE) total mounts, intromissions, and ejaculations; and intromission ratio for the first ejaculatory series, in sexually experienced animals treated with MK-801 (2.5 μ g/ μ l) or aCSF in the MPOA. * $p < .05$. B) Mean ($\pm$ SE) latencies for mounts, intromissions, and ejaculations in sexually experienced animals treated with MK-801 (2.5 μ g/ μ l) or aCSF in the MPOA. * $p < .05$.

Figure 4.

Effects of intra-MPOA administration of MK-801 on copulation before and after repeated sexual experience, Experiment 3. Mean ($\pm$ SE) total: A) mounts, B) intromissions, and C) ejaculations in animals tested when sexually naïve and after three sexual experiences, after treatment with MK-801 (2.5 μ g/ μ l) or aCSF in the MPOA. For animals tested with aCSF when sexually naïve, only data from the aCSF test after experience were analyzed; similarly, for animals tested with drug while naïve, only the drug test after experience was analyzed. Data were analyzed with 2-way ANOVAs with planned comparisons between drug and aCSF in naïve and experienced males. * p < .05 for experienced vs. naïve animals and drug vs. aCSF for naïve animals; ** p < .01 for experienced vs. naïve animals and drug vs. aCSF for naïve animals; # p = .08 for drug vs. aCSF for experienced animals. D) Mean ($\pm$ SE) total mounts, intromissions, and ejaculations in sexually experienced animals treated with MK-801 (2.5 μ g/ μ l) or aCSF in counterbalanced order. Data were analyzed with a paired t-test. ** p < .01. E) Mean ($\pm$ SE) latencies for the first mount, intromission, and ejaculation in sexually experienced animals treated with MK-801 (2.5 μ g/ μ l) or aCSF. Data were analyzed with a paired t-test. * p < .05 for drug vs. aCSF for mounts; # p = .07 for drug vs. aCSF for intromission latency. Intro = intromission; Ejac = ejaculation.

Figure 5.

Effects of intra-MPOA administration of MK-801 on stimulus sensitization, Experiment 4. Mean ($\pm$ SE) total mounts, intromissions, and ejaculations; and intromission ratio for the first ejaculatory series, in sexually naïve animals treated with MK-801 (2.5 μ g/ μ l) or aCSF before each of 7 noncopulatory exposures to a receptive female and tested drug-free on Day 8, and in untreated naïve males not pre-exposed to a female. Open bars indicate animals that received a microinjection of aCSF before each exposure; black bars indicate animals that received an MK-801 microinjection before each exposure; and gray bars indicate animals that received 7 daily microinjections of aCSF, but did not receive any pre-exposure. * $p < .05$, ** $p < .01$.