Published in final edited form as:

J Neurovirol. 2010 October ; 16(5): 399–404. doi:10.3109/13550284.2010.504248.

Correlation of major depressive disorder symptoms with FKBP5 but not FKBP4 expression in human immunodeficiency virus–infected individuals

Erick T Tatro¹, Timothy B Nguyen², Chad A Bousman¹, Eliezer Masliah³, Igor Grant¹, J Hampton Atkinson¹, and Ian P Everall²

¹Department of Psychiatry, University of California San Diego, La Jolla, California, USA

²Department of Psychiatry, University of Melbourne, Melbourne, Australia

³Department of Neurosciences, University of California San Diego, La Jolla, California, USA

Abstract

Major depressive disorder (MDD) is a significant cause of morbidity in people living with the human immunodeficiency virus (HIV). FKBP5 is a candidate gene with single-nucleotide polymorphisms (SNPs) rs1360780 and rs3800373 associated with MDD. This gene product and its relative, FKBP4, physically associate with the glucocorticoid receptor whose function is implicated in MDD pathophysiology. Because these genes are expressed in blood and brain and elevated in HIV infection, we explored the relationship between gene expression, genotype, and MDD symptoms. Longitudinally followed subjects (N = 57) as part of the CNS HIV AntiRetroviral Effects Research study, with diagnosed MDD and who donated blood for genotyping and gene expression analysis, were assessed. Subjects donated blood on adjacent visits with and without meeting criteria for MDD episode. Changes in clinical parameters were compared changes in gene expression. Change in FKBP5 expression correlated with change in Beck Depression Inventory (BDI) for MDD → euthymic comparison in GG genotype of rs3800373 (P = .013) and TT carriers of rs1360780 (P = .02). In euthymic → MDD comparison, GG homozygous, FKBP5 expression correlated with more severe change in BDI. Change in FKBP4 expression did not correlate with changes in clinical or depression measurements. Higher FKBP5 expression correlated with greater symptom change for GG carriers of rs3800373.

Keywords

FKBP4; FKBP5; gene expression; genotyping; HIV; human immunodeficiency virus; major depression; major depressive disorder

© 2010 Journal of NeuroVirology

Address correspondence to Erick T. Tatro, Department of Psychiatry, University of California San Diego, 9700 Gilman Drive, La Jolla, CA 92093-0603, USA. etatro@ucsd.edu.

Erick T. Tatro and Timothy B. Nguyen are joint first authors.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Supplementary material available online

Figures S1, S2 and S3

Tables S1 and S2

Introduction

FKBP5, a candidate gene for major depressive disorder (MDD), is inducible by the stress-related hormones glucocorticoids (Binder et al, 2004). FKBP5 and its relative FKBP4 function to differentially regulate trafficking and signaling of hormone receptors (Wohnik et al, 2005). We previously showed that FKBP4 and FKBP5 were up-regulated in the frontal cortex of human immunodeficiency virus (HIV)-infected individuals (Tatro et al, 2009b). Taken together, along with the fact that the HIV-infected population is disproportionately affected by MDD (Bing et al, 2001), one mission of the neuroAIDS field is to understand whether and how HIV affects the brain that results in the frequency of MDD high in the era of antiretroviral (ARV) therapy and long-term survival with HIV infection. We reported some changes in gene expression in MDD of HIV subjects through microarray analysis in frontal cortex including somatostatin (Everall et al, 2006). We explored the relationship between FKBP gene expression in peripheral blood mononuclear cells and MDD symptoms in HIV-infected individuals on ARV. Here, we examined the longitudinal gene expression of both FKBP4 and FKBP5 in HIV-infected individuals diagnosed with MDD by comparing transcript levels during an episode of MDD to expression during the euthymic state as it relates to FKBP5 genotype.

Association between FKBP5 and MDD recurrence was originally discovered by genome-association study of three chromosome 6 genes related to glucocorticoid receptor (GR) function and the importance of FKBP5 for MDD were confirmed in the STAR*D cohort (Binder et al, 2004; Lekman et al, 2008). A report found increased risk of peritraumatic dissociation with rs1360780 minor allele (Koenen et al, 2005). Additionally, the minor allele, in which FKBP5 is hyperinduced by cortisol, is associated with the development of posttraumatic stress disorder (Binder et al, 20088). However, a more recent study reported no linkage between FKBP5 single-nucleotide polymorphisms (SNPs) and late-life MDD (Sarginson et al, 2010). The hyper-induction allele is thought to result in decreased negative feedback inhibition of cortisol release and therefore dysfunctionally prolonged physiologic stress response after stimuli (Binder, 2009). HIV-infected individuals are more likely to be hypercortisolemic and the infection is thought to, in part, disrupt the hypothalamic-pituitary-adrenal (HPA) axis (Chittiprol et al, 2007).

In this study, gene expression of FKBP5 correlated with change in depressive symptom severity in major depressive episode when HIV-positive subjects were homozygous for minor alleles of either FKBP5 SNPs rs1360780, an intronic SNP near enhancer and promoter elements, or rs3800373, in the 3' untranslated region (UTR). Our results provide evidence that FKBP5 transcriptional dysregulation together with FKBP4 as its functional antagonist are implicated in biological features of MDD symptoms.

Methods

Ascertainment of cases

Cases were ascertained as part of the CNS HIV Antiretroviral Therapy Effects Research (CHARTER) Study (https://www.charterresource.ucsd.edu/images/M_images/x-sect_cohort_rep.pdf). Subjects were longitudinally assessed semiannually and neuromedical, behavioral, and laboratory data were obtained, including psychiatric assessment using the Beck Depression Inventory (BDI) (Beck and Brown, 1996) and the Composite International Diagnostic Interview (CIDI) (Kessler et al, 2003), which yield, current and lifetime psychiatric diagnosis based on Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria. A summary of parameters measured and methods is available at the CHARTER resource site (CHARTER Group, 2009). We chose subjects who met criteria for major depressive

disorder in whom blood samples were available during both an episode of MDD and when euthymic at adjacent visit. All subjects included in this report tested negative for drugs of abuse and hepatitis C virus antibodies at visits. Seventy-four percent were male (40:14 M:F ratio). Summary of clinical parameters at the two time points for subjects meeting these criteria are in Table 1. Note that global deficit score is a measure of neurocognitive impairment standardized at the HIV Neurobehavioral Research Center and is described by Carey et al (2004). All subjects gave informed consent as approved by the University of California San Diego Institutional Review Board.

The depression-remission patterns differed temporally among the subjects and could be divided into two cohorts: Cohort 1: (N = 56) MDD at first visit → euthymic at second visit; and Cohort 2: (N = 30) euthymic at first visit → MDD at second visit.

Gene expression and genotyping

Approximately 10 ml blood was drawn directly into EDTA-coated glass tubes and peripheral blood mononuclear cell RNA was isolated according to manufacturer's protocols (Paxgene, Valencia, CA). After reverse transcription using Superscript III enzyme (Invitrogen, Carlsbad, CA), Taqman quantitative polymerase chain reaction (PCR) was performed in 384-well format. Taqman primers were obtained from Applied Biosystems (Foster City, CA): for FKBP4, assay ID Hs00427038; for FKBP5, assay ID Hs00188025; for GAPDH, assay ID Hs02758991. Samples were analyzed in triplicate on the same plate to avoid batch-effect. Gene expression was quantified using $\Delta\Delta C_T$ method with glutaraldehyde 3-phosphate dehydrogenase (GAPDH) as endogenous control. Taqman assays were run and analyzed at the University of California San Diego Center for AIDS Research Genomics Core. Considering that genotype of FKBP5 is known to affect its transcript levels, we genotyped the rs1360780 and rs3800373 SNPs using Taqman allelic discrimination assays (Applied Biosystems) in each subject. We used multivariate regression analysis grouping by genotype to compare change in gene expression from euthymic → MDD and MDD → euthymic with changes in BDI.

Results

Effect of depression status on FKBP4 and FKBP5 gene expression

We considered that there would be differences between the two patterns (first MDD, then euthymic at 6-month follow-up, and vice versa), as the changes may either correlate with onset of major depression episode or with recovery and medication effects. Separating the sample into these two cohorts, there was a small to medium effect of depression status on gene expression for FKBP5 (Cohort 1: Cohen's $d = 0.23$, $P = .12$, Cohort 2: $d = 0.31$, $P = .09$) but no effect for FKBP4 (Cohort 1, $d = 0.06$, $P = .66$, Cohort 2: $d = 0.07$, $P = .78$) regardless of direction of change. In both cohorts, expression of FKBP5 was the same under the euthymic state. However, FKBP5 was elevated during MDD in Cohort 1, and decreased during MDD in Cohort 2, indicating a potential temporal difference in whether changes were due to onset or recovery.

Peripheral blood mononuclear cell (PBMC) RNA was available for analysis of a smaller subset of the subjects at a third, euthymic time point. In most cases, FKBP4 (81%) and FKBP5 (78%) RNA quantities were stable at two euthymic time points compared to the MDD time point, even though blood was drawn with 12 intervening months (see Figure S1; online Supplementary Materials, available at ...). Further, other candidate genes were measured in a preliminary analysis, STAT1 and GADD45b, which had consistent levels at euthymic and MDD states (Figure S1) and so were not further analyzed. These observations help to rule out assay variability.

Effect of sex on FKBP4 and FKBP5 expression

To test for sex as a possible confounding variable, we segregated the subjects to compare change in gene expression and change in BDI by male versus female in both cohorts. In Cohort 2, (euthymic → MDD), the magnitude of change in FKBP4 in females was higher than males (difference = 0.35-fold, $P = .04$) and the magnitude of change in FKBP5 was higher in females than males (difference = 0.30-fold, $P = .02$). There was no difference with respect to BDI changes in Cohort 2. No statistically significant male/female differences were found in Cohort 1. These statistics are provided in Table S2 in the online Supplementary Materials.

Correlation of FKBP5 gene expression with BDI

Using a multivariate regression analysis comparing changes in euthymic and MDD gene expression versus changes in other clinical and laboratory parameters (listed in Table 1), separating by genotype, we found that change in FKBP5 expression correlated with change in BDI for MDD → euthymic comparison in GG genotype of rs3800373 ($P = .013$, Pearson's $R = .61$; Figure 1, left). Using this comparison, the same correlation was observed for rs1360780 ($P = .02$, $R = .56$; Figure S2). Because the SNPs were in linkage disequilibrium, we present the rs1360780 comparisons in Supplementary Materials. In euthymic → MDD comparison, GG homozygous FKBP5 expression correlated with more severe change in BDI (Figure 1, right; $P = .0499$, $R = .40$). In sum, higher gene expression correlated with greater symptom change (Figure 1). Change in FKBP4 gene expression did not significantly correlate with changes in clinical or MDD mood measurements.

Conclusions

In this study we have observed a correlation of change in the severity of depressive mood, as measured by the BDI, with FKBP5 gene expression in individuals homozygous for GG of the FKBP5 SNP rs3800373. We suggest that this link of FKBP5 to BDI improvement in Cohort 1 is a component of the pathophysiological substrate of MDD. In this cohort, whose first visit was while they had MDD and the second when they were euthymic, the gene expression observations were more robust ($P = .02$ versus $P = .049$) and significant at both SNPs. Although both of the FKBP5 SNPs are in noncoding regions of the gene, the variants may affect translation efficiency and therefore the regulatory effectiveness of immunophilins on GR signaling. Our clinical and genetic results are correlational and animal or in vitro models of HIV to manipulate these genes would help determine causative roles in neurobehavioral symptoms we measure. Future work to analyze protein levels of these genes as in Binder et al (2004) should be carried out in a longitudinally followed HIV-positive cohort to further our understanding of the possible roles these genes play in MDD symptomatology.

Susceptibility to mood disorders is determined by interactions between genes and environment, heritability estimates range from 30% to 80% (Kendler, 1995). An important, but not sole, system that mediates long-term effects of stress on psychiatric disorders is the HPA system. Stress activates corticotropin release hormone (CRH) (Vale et al, 1981). In the central nervous system (CNS), CRH neurons innervate the locus coeruleus and activate the noradrenergic and sympathetic systems (Valentino et al, 1983). CRH regulates adrenal cortisol release as well as norepinephrine and epinephrine secretion (Tsatsanis et al, 2007). Glucocorticoids such as cortisol promote the fight and flight response in end organs mediated through the GR, which is critical for the negative-feedback loop terminating the response (Pariante and Miller, 2001). Hypercortisolemia and evasion from dexamethasone-induced cortisol suppression are physiologic features of MDD (Heuser et al, 1994). Because of hypercortisolemia observed in a significant number HIV-infected individuals (Chittiprol

et al, 2007) and the increased risk of MDD in this population (Bing et al, 2001), we sought to determine if the GR-associated genes were implicated.

The proteins that are products of FKBP4 and FKBP5 associate physically with hormone receptors in many tissues, including peripheral blood mononuclear cells and the brain (Tatro et al, 2009b; Davies et al, 2002). FKBP4 gene product promotes fast nuclear translocation in neurons and FKBP5 gene product inhibits translocation of GR in the absence of hormone in neurons (Tatro et al, 2009a). Together, these genes regulate an ultrafast feedback loop within the cells. Our finding of a genotype-dependent correlation of FKBP5 mRNA expression with greater major depressive symptom recovery is in agreement with past studies (Kirchheiner et al, 2008), and we recapitulate this in HIV-infected individuals. FKBP5 may offer the future possibility of a bio-marker to aid as a diagnostic or prognostic indicator for recovery of MDD in this population.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding to perform this work was provided by National Institutes of Health Grant Numbers MH076681 and MH079881. This work was performed with the support of the Genomics Core at the University of California San Diego Center for AIDS Research (AI36214) and the San Diego Veterans Medical Research Foundation.

References

- Binder E. The role of FKBP5, a co-chaperone of the glucocorticoid receptor in the pathogenesis and therapy of affective and anxiety disorders. *Psycho-neuroendocrinology*. 2009; 34S:S186–S195.
- Beck, A.; Brown, G. BDI-II manual. San Antonio, TX: The Psychological Corporation; 1996.
- Binder EB, Bradley RG, Liu W, Epstein MP, Deveau TC, Mercer KB, Tang Y, Gillespie CF, Heim CM, Nemeroff CB, Schwartz AC, Cubells JF, Ressler KJ. Association of FKBP5 polymorphisms and childhood abuse with risk of posttraumatic stress disorder symptoms in adults. *JAMA*. 2008; 299:1291–1305. [PubMed: 18349090]
- Bing EG, Burnam MA, Longshore D, Fleishman JA, Sherbourne CD, London AS, Turner BJ, Eggen F, Beckman R, Vitiello B, Morton SC, Orlando M, Bozzette SA, Ortiz-Barron L, Shapiro M. Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Arch Gen Psychiatry*. 2001; 58:721–728. [PubMed: 11483137]
- Binder EB, Salyakina D, Lichtner P, Wochnik GM, Ising M, Putz B, Papiol S, Seaman S, Lucae S, Kohli MA, Nickel T, Kunzel HE, Fuchs B, Majer M, Pfennig A, Kern N, Brunner J, Modell S, Baghai T, Deiml T, Zill P, Bondy B, Rupprecht R, Messer T, Kohnlein O, Dabitz H, Bruckl T, Muller N, Pfister H, Lieb R, Mueller JC, Lohmussaar E, Strom TM, Bettecken T, Meitinger T, Uhr M, Rein T, Holsboer F, Muller-Myhsok B. Polymorphisms in FKBP5 are associated with increased recurrence of depressive episodes and rapid response to antidepressant treatment. *Nat Genet*. 2004; 36:1319–1325. [PubMed: 15565110]
- CHARTER Group. Technical report. La Jolla, CA: University of California San Diego; 2009. CHARTER variables of interest—cross sectional complete. Available at: https://www.charterresource.ucsd.edu/images/Mimages/x-sect_cohort_rep.pdf
- Chittiprol S, Shetty KT, Kumar AM, Bhimasenarao RS, Satishchandra P, Subbakrishna DK, Desai A, Ravi V, Satish KS, Gonzalez L, Kumar M. HPA axis activity and neuropathogenesis in HIV-1 clade C infection. *Front Biosci*. 2007; 12:1271–1277. [PubMed: 17127379]
- Carey CL, Woods SP, Gonzalez R, Conover E, Marcotte TD, Grant I, Heaton RK. Predictive validity of global deficit scores in detecting neuropsychological impairment in HIV infection. *J Clin Exp Neuropsychol*. 2004; 26:307–319. [PubMed: 15512922]

- Davies TH, Ning YM, Sanchez ER. A new first step in activation of steroid receptors: hormone-induced switching of FKBP51 and FKBP52 immunophilins. *J Biol Chem*. 2002; 277:4597–4600. [PubMed: 11751894]
- Everall IP, Salaria S, Atkinson JH, Young C, Corbeil J, Grant I, Masliah E. HNRC (HIV Neurobehavioral Research Center) . Diminished somatostatin gene expression in individuals with HIV and major depressive disorder. *Neurology*. 2006; 67:1867–1869. [PubMed: 17130427]
- Heuser I, Yassouridis A, Holsboer F. The combined dexamethasone/CRH test: a refined laboratory test for psychiatric disorders. *J Psychiatr Res*. 1994; 28:341–356. [PubMed: 7877114]
- Kendler K. Genetic epidemiology in psychiatry: taking both genes and environment seriously. *Arch Gen Psychiatry*. 1995; 15:895–899. [PubMed: 7487337]
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, Rush AJ, Walters EE, Wang PS. National Comorbidity Survey Replication . The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003; 289:3095–3105. [PubMed: 12813115]
- Kirchheiner J, Lorch R, Lebedeva E, Seeringer A, Roots I, Sasse J, Brockmüller J. Genetic variants in FKBP5 affecting response to antidepressant drug treatment. *Pharmacogenomics*. 2008; 9:841–846. [PubMed: 18597649]
- Koenen KC, Saxe G, Purcell S, Smoller JW, Bartholomew D, Miller A, Hall E, Kaplow J, Bosquet M, Moulton S, Baldwin C. Polymorphisms in FKBP5 are associated with peritraumatic dissociation in medically injured children. *Mol Psychiatry*. 2005; 10:1058–1059. [PubMed: 16088328]
- Lekman M, Laje G, Charney D, Rush AJ, Wilson AF, Sorant AJM, Lipsky R, Wisniewski SR, Manji H, McMahon FJ, Paddock S. The FKBP5-gene in depression and treatment response—an association study in the Sequenced Treatment Alternatives to Relieve Depression (STAR*D) cohort. *Biol Psychiatry*. 2008; 63:1103–1110. [PubMed: 18191112]
- Pariante CM, Miller AH. Glucocorticoid receptors in major depression: relevance to pathophysiology and treatment. *Biol Psychiatry*. 2001; 49:391–404. [PubMed: 11274650]
- Sarginson J, Lazzaroni L, Ryan H, Schatzberg A, Murphy G. FKBP5 polymorphisms and antidepressant response in geriatric depression. *Am J Med Genet B*. 2010; 153B:554–560.
- Tsatsanis C, Dermitzaki E, Venihaki M, Chatzaki E, Minas V, Gravanis A, Margioris AN. The corticotropin-releasing factor (CRF) family of peptides as local modulators of adrenal function. *Cell Mol Life Sci*. 2007; 64:1638–1655. [PubMed: 17453142]
- Tatro ET, Everall IP, Kaul M, Achim CL. Modulation of glucocorticoid receptor nuclear translocation in neurons by immunophilins FKBP51 and FKBP52: implications for major depressive disorder. *Brain Res*. 2009a; 1286:1–12. [PubMed: 19545546]
- Tatro E, Everall I, Masliah E, Hult B, Lucero G, Chana G, Soontornniyomkij V, Achim C. HNRC . Differential expression of immunophilins FKBP51 and FKBP52 in the frontal cortex of HIV-infected patients with major depressive disorder. *J Neuroimmune Pharmacol*. 2009b; 4:218–226. [PubMed: 19199039]
- Valentino RJ, Foote SL, Aston-Jones G. Corticotropin-releasing factor activates norad-renergic neurons of the locus coeruleus. *Brain Res*. 1983; 270:363–367. [PubMed: 6603889]
- Vale W, Spiess J, Rivier C, Rivier J. Characterization of a 41-residue bovine hypothalamic peptide that stimulates secretion of corticotropin and beta-endorphin. *Science*. 1981; 213:1394–1397. [PubMed: 6267699]
- Wochnik GM, Ruegg J, Abel GA, Schmidt U, Holsboer F, Rein T. FK506-binding proteins 51 and 52 differentially regulate dynein interaction and nuclear translocation of the glucocorticoid receptor in mammalian cells. *J Biol Chem*. 2005; 280:4609–4616. [PubMed: 15591061]

Appendix

The CNS HIV Antiretroviral Therapy Effects Research (CHARTER) is supported by award N01 MH22005 from the National Institutes of Health. The CHARTER group is affiliated with the Johns Hopkins University, Mount Sinai School of Medicine, University of California, San Diego, University of Texas, Galveston, University of Washington, Seattle, and Washington University, St Louis; is headquartered at the University of California, San

Diego; and includes the following individuals: Director: Igor Grant, MD; Codirectors: J. Allen McCutchan, MD, Ronald J. Ellis, MD, PhD, and Thomas D. Marcotte, PhD; Center Manager: Shondra Neumayer, RN, FNP; Neuromedical Component: Ronald J. Ellis, MD, PhD (principal investigator [PI]), J. Allen McCutchan, MD, and Terry Alexander, RN; Laboratory, Pharmacology, and Immunology Component: Scott Letendre, MD (PI), Edmund Capparelli, PharmD, and Janis Durelle, BS; Neurobehavioral Component: Robert K. Heaton, PhD (PI), J. Hampton Atkinson, MD, Steven Paul Woods, PsyD, and Donald Franklin, BS; Virology Component: Joseph K. Wong, MD (PI), and Caroline Ignacio, BA; Imaging Component: Terry Jernigan, PhD (PI), Michael J. Taylor, PhD, and Rebecca Theilmann, PhD; Data Management Unit: Anthony C. Gamst, PhD (PI), Clint Cushman, BA, and Michelle Frybarger, BA; Statistics Unit: Ian Abramson, PhD (PI), and Deborah Lazzaretto, MS; Protocol Coordinating Component: Thomas D. Marcotte, PhD (PI), and Rodney von Jaeger, MPH; Johns Hopkins University Site: Justin C. McArthur, MBBS (PI); Mount Sinai School of Medicine Site: Susan Morgello, MD (Co-PI), and David Simpson, MD (Co-PI); University of California, San Diego, Site: J. Allen McCutchan, MD (PI); University of Washington, Seattle, Site: Ann Collier, MD (Co-PI), and Christina Marra, MD (Co-PI); University of Texas, Galveston, Site: Benjamin Gelman, MD, PhD (PI); and Washington University, St Louis, Site: David Clifford, MD (PI), and Muhammad Al-Lozi, MD.

Figure 1.

Correlation of Beck's Depression Inventory with FKBP4 and FKBP5 gene expression by rs3800373 genotype. After separating the subjects by FKBP5 genotype, correlation coefficients between change in BDI with change in gene expression from euthymic versus MDD status were calculated. For the euthymic → MDD group: as change in FKBP5 increased, BDI worsened. Correspondingly, for the MDD → euthymic group, increasing FKBP5 expression correlated with greater BDI improvement.

Table 1

Summary statistics of clinical parameters of subjects

Parameter	Median	Range	SD
Euthymic			
BDI	19	53	11.24
Plasma HIV	1.69	3.52	1.09
Number ARVs	3	5	1.2
GDS	0.33	3.43	0.76
CD4	490	1398	292
MDD			
BDI	25	42	9.67
Plasma HIV	1.69	3.56	1.28
Number ARVs	3	4	1.22
GDS	0.17	3.6	0.85
CD4	473	1057	251

Note. BDI = Beck's Depression Inventory; plasma HIV = log (plasma HIV RNA); ARVs = antiretroviral medications; GDS = global deficit score; CD4 = CD4+ T-cell count.