Restriction endonuclease *EcaI* from *Enterobacter cloacae*

G.Hobom[†], E.Schwarz[†], M.Melzer^{*} and H.Mayer^{*}[†]Institut für Biologie III der Universität Freiburg, Schänzlestrasse 1, D-7800 Freiburg i.Br., and^{*}Gesellschaft für Biotechnologische Forschung mbH., Mascheroder Weg 1, D-3300 Braunschweig, GFR

Received 27 July 1981

ABSTRACT

Restriction endonuclease *EcaI* obtained from *Enterobacter cloacae* DSM30056 recognizes the group of heptanucleotide palindromes 5'-G⁺G-T-N-A-C-C-3', and on cleavage (arrow) produces fragments with 5'-terminal pentanucleotide extensions. It is identical in specificity with restriction endonuclease *BstEII* from *Bacillus stearothermophilus* ET.

INTRODUCTION

Restriction endonucleases of type II have been isolated in growing numbers from many bacterial species. Most of the DNA recognition sites reported for these endonucleases contain symmetric tetra- or hexanucleotide sequences (palindromes), and several have been shown to be extended by one or three basepairs of degenerate character in a central position, as is known for *HinfI* or *BglII* or several others (see reference 1). In these cases penta- or nonanucleotide sequences with external symmetry, i.e. hyphenated palindromes are recognized by such a group of enzymes.

From the large family of *enterobacteriaceae* only one restriction enzyme, *EclI* from *Enterobacter cloacae* has been obtained so far (2). We have isolated a restriction enzyme from *Enterobacter cloacae* DSM30056 (*EcaI*) which recognizes a symmetric hexanucleotide DNA sequence that is hyphenated by a single degenerate basepair: 5'-GGT-N-ACC-3'. In the course of this work it has been detected that restriction endonuclease *BstEII* from *Bacillus stearothermophilus* ET recognizes the same sequence, as has also been determined independently by Lautenberger et al. (3). An additional isoschizomer of both enzymes (*BstPI*) has been isolated from another strain of *B. stearothermophilus* (4). - This publication extends a previous note on the isolation and characterization of *EcaI* (5).

MATERIALS AND METHODS

Enterobacter cloacae strain DSM30056 was obtained from the German Col-

lection of Microorganisms, Göttingen. Restriction endonuclease *Bst*EII was purchased from BRL, Rockville (Md), bacterial alkaline phosphatase and DNaseI from Worthington, snake venom phosphodiesterase from Sigma, T4 polynucleotide kinase from Biogenics Research Corporation, and T4 DNA ligase from Miles. Plasmid DNAs were prepared following a cleared lysate procedure (6), λ DNAs were isolated from phage preparations obtained after induction of λ CI857S7 or λ CI857t68 lysogenic strains. SV40 DNA was purchased from MRE, England.

Assay for *Eca*I restriction endonuclease activity: 0.1 - 0.3 μ g DNA were incubated in 50 μ l with 5-10 μ l of activity containing fractions at 37°C for 60 min either in 10 mM Tris HCl pH 8, 10 mM MgCl₂, or in 6 mM Tris HCl pH 7.4, 6 mM MgCl₂, 50 mM NaCl, 6 mM 2-mercaptoethanol. The resulting fragment mixtures were separated by electrophoresis on agarose or polyacrylamide slab gels. Cleavages with endonuclease *Bst*EII were done in 100 mM Tris HCl pH 7.6, 5 mM MgCl₂, 6 mM 2-mercaptoethanol, 1 μ g/ml of bovine serum albumine at 37°C for 60 min.

Analysis of the *Eca*I recognition site: About 1 μ g DNA each of plasmids λ dv1, λ dvh93 (7), pHL81 (a 9 kb segment of phage 434 DNA cloned into pBR313, ref. 8) and pHL102 (a *lac*I gene fragment cloned into pBR322, G. Hobom, unpublished) were digested with *Eca*I, ethanol precipitated and resuspended in 20 μ l H₂O. Dephosphorylation was carried out in 50 μ l 10 mM Tris HCl pH 9.3 with 1 μ g bacterial alkaline phosphatase for 1 h at 37°C. The enzyme was inactivated by addition of EGTA (final conc. 6.7 mM) and subsequent boiling of the reaction mixture for 3 min. 5'-terminal labeling was achieved using T4 polynucleotide kinase and γ -[³²P]-ATP, following standard procedures (9). After ethanol precipitation, the labeled fragment mixtures were separated on 1% agarose slab gels in order to remove residual γ -[³²P]-ATP and ³²P-phosphate; fragments were isolated from the gels by electrophoretic elution (10).

To achieve complete digestion to 5'-mononucleotides isolated *Eca*I restriction fragments were incubated with 1 μ g of DNaseI and 1 μ g of snake venom phosphodiesterase in 30 μ l 10 mM Tris HCl pH 8.5, 10 mM MgCl₂ for 1 h at 37°C. The resulting nucleotides were separated by electrophoresis on Whatman 3MM paper at pH 3.5. The bands were visualized under UV light illumination and upon isolation measured for their ³²P-activities.

Partial digestions of ³²P-labeled fragments were carried out by incubation with 1 μ g DNaseI in 30 μ l for 30 min at 37°C, followed by the addition of 0.05 μ g snake venom phosphodiesterase and further incubation for 0.5, 1, and 2 minutes (one third volume each). The three fractions were recombined, and the products separated twodimensionally, according to Brownlee and Sanger

(11), using electrophoresis at pH 3.5 on cellulose acetate strips in the first dimension, and homochromatography on 20% DEAE-cellulose thin-layer plates at 65°C in the second dimension, with a 1:1 diluted 3% RNA homomix solution.

DNA sequencing was carried out following the method of Maxam and Gilbert (9).

RESULTS

Isolation of restriction endonuclease *EcaI*

Enterobacter cloacae DSM30056 was cultured in enriched nutrient broth (8 g nutrient broth (Difco), 5 g bacto-peptone (Difco), 5 g NaCl, 3.5 g Na₂HPO₄, 1.5 g KH₂PO₄, 1 g glucose, 5 mg thiamine HCl per 1000 ml) at 37°C to late logarithmic phase, and cells were collected. 4 g of frozen cells were thawed in 8 ml buffer A (10 mM potassium phosphate, 1 mM EDTA, 0.1% 2-mercaptoethanol, and 0.1% Triton X 100, pH 7.5), and disrupted at 4°C in a Branson sonifier for 15 min at 60 Watt. The homogenate was centrifuged at 45000 g for 20 min at 2°C and the supernatant adjusted to 1 M NaCl. DNA was precipitated by addition of a 10% streptomycin sulfate solution to give a final concentration of 1%. The precipitate was removed by centrifugation as above, and the supernatant was dialysed against buffer A. Endonuclease activity was detectable in this crude extract.

The dialysate was applied to a DE52-column (2.5 X 20 cm) equilibrated with buffer A. Elution was performed with a 200 ml (0 - 0.5 M NaCl) linear gradient in buffer A plus a final step of 100 ml 1 M NaCl in buffer A. Fractions of 4 ml were collected and assayed for the presence of endonuclease activity which eluted predominantly in the 1 M NaCl step. Active fractions were combined and dialysed against buffer A.

The active DE-cellulose-fractions were applied to a phospho-cellulose column 1 X 15 cm, equilibrated with buffer A.

The *EcaI* activity eluted between 0.2 - 0.3 M NaCl of a 50 ml (0 - 0.6 M NaCl) linear gradient in buffer A. The combined active fractions were stored at -20°C in 50% glycerol. The yield of *EcaI* activity was about 1000 U/4 g cells (wet weight). (1 U of activity is defined by digestion of 1 µg λDNA to its complete, final digestion pattern in 1 h at 37°C.) Only insignificant loss of activity could be detected over a period of 8 months. Exonuclease or phosphatase contaminations have been excluded from these *EcaI* preparations, since overnight incubation with 5'-[³²P]-labeled DNA fragments failed to yield non-precipitable ³²P-activity.

Determination of the recognition site of *EcaI*

The small λ -derived plasmids λ dvh93 und λ dvO21 (7) contain a single *EcaI* cleavage site which has been precisely mapped in a region of the *P* gene, and plasmid λ dv1 contains two such sites, of which the second one is located in the *l_{rex}* region. Similarly, in plasmid pHL102 DNA the single *EcaI* recognition site could be located through double digestions with other restriction endonucleases near the center of the *lacI* gene. In all three cases the DNA sequence of the respective region is known (12-15) and because of the location of the cleavage sites a common palindrome sequence GGTNACC could be pointed out at this stage as a possible candidate for the *EcaI* recognition site.

For a biochemical characterization of the *EcaI* recognition and cleavage site, restriction fragments obtained by cleaving plasmids pHL81 (5 cleavage sites), λ dv1 (2 sites) and pHL102 (1 site) with *EcaI*, and subsequently labeled at their 5'-ends with 32 P-phosphate were mixed in equivalent proportions, or individually subjected to a) complete digestion to 5'-mononucleotides for determination of the 5'-terminal nucleotide in paper electrophoresis, and b) partial digestion followed by two-dimensional separation of the products for determination of the 5'-oligonucleotide sequence (for experimental details see Methods). In all cases separation of the 5'- 32 P]-mononucleotides showed close to 98% of the radioactivity associated with guanylic acid. From the pattern obtained after two-dimensional separation of the partial DNaseI/snake venom phosphodiesterase digestion products a 5'-terminal nucleotide sequence 5'-G-T-N-A-C-C-N- could be deduced (Fig. 1). This confirms a 5'-G-T-N-A-C-C-3' recognition and cleavage site for *EcaI* in which the central position can be occupied by any of the four nucleotides.

A further analysis of the exact location of the cleavage site within the recognition sequence was made by sequencing a 38 basepair *HinfI:EcaI* fragment labeled at its *HinfI* site, which was obtained from a 145 bp λ dvh93 DNA:*HinfI* fragment by secondary *EcaI* cleavage and fragment separation. The autoradiogram of the sequencing gel clearly resolved a 3'-terminal G residue, below the band of unreacted material (see Fig. 2). Finally, a 32 P-*EcaI* labeled *HinfI-EcaI* fragment (129 bp) derived from the λ dv1-*rex* region was treated according to the Maxam-Gilbert sequencing reactions (9) and was run on a 20% polyacrylamide sequencing gel in between two other fragments of known DNA sequence which were to be used as a ladder for determining the equidistant bands from the 5'-end. From this experiment an extrapolation down to the position of the 5'-terminal nucleotide (against the known se-

4827

Figure 2. Autoradiogram of the upper part of a DNA sequencing gel obtained after *Eco*AI subfragmentation of a 5'-[³²P]-labeled *Hinf*I fragment which was derived from the *P*-region of λ dvh93 DNA. The 38 bp subfragment was isolated, treated according to the Maxam-Gilbert sequencing reactions (9), and the resulting products were fractionated on a 20% polyacrylamide/7 M urea gel. The sequence observed represents positions λ :40862 through λ :40881 from the *L*-strand of λ DNA, the product resulting from interaction with the 3'-terminal G-residue (arrow) is clearly separated from the final band of unreacted material.

conclude that no other related DNA sequences are recognized by this enzyme. In particular all types of degeneracies known for other type II restriction endonuclease recognition sites (such as the *Acc*II, *Ava*II, *Hae*I, *Hae*II and *Hind*II specific patterns) have been excluded in this way, and also GGT-N_x-ACC palindromes with an extended length of degenerate hyphenation ($x = 2$ to $x = 7$). All of the *Eco*AI variant sequences mentioned have been determined in this computer search at one or more positions in any of the DNAs which are not cleaved by this enzyme. On the other hand no nucleotide other than the hyphenated hexanucleotide palindrome sequence is common to all three *Eco*AI sites studied here in detail, in a symmetrical or asymmetrical location within the next seven basepairs on either side.

As is known from other type II restriction endonucleases cleavage by *Eco*AI appears to result in fragments with 5'-phosphate bearing ends. This conclusion was drawn from the ability to religate *Eco*AI linearised λ dvh93, λ dv021 and other DNAs back into circular molecules and linear concatemers, as is shown in Figure 3. T4 DNA ligase reactions are known to depend on 5'-phosphate ends of substrate DNA fragments. Due to their pentanucleotide single stranded ends *Eco*AI fragments are particularly good substrates for

Figure 3. Comparison of *EcaI* and *BstEII* cleavage reactions, and mixed ligation of *EcaI* and *BstEII* generated DNA fragments in an electrophoretic separation on a 0.7% agarose gel. Lanes (1) and (2): pBR313 DNA, digested by *BstEII* and *EcaI*, respectively. (3): λ dv1 DNA, digested by *HaeIII* (size marker: 1670 bp, 1310 bp). (4) and (5): λ cI857S7 DNA, digested by *BstEII* and *EcaI*, respectively. Both samples have been incubated at 65°C and quickly cooled prior to electrophoresis; the faint uppermost band originates from a partial addition of the right and left terminal fragments, bands 1 and 4. (6): octomer λ dvO21 plasmid DNA, partially digested by *EcoRI* (size marker for multimers of 3.153 kb). (7): λ dvh93 DNA, *BstEII* digested, and religated by T4 DNA ligase at 160 μ g DNA/ml. (8): mixture of λ dvh93 digested by *BstEII* and λ dvO21 digested by *EcaI*, ligated by T4 DNA ligase at 190 μ g DNA/ml (1:1). (9): λ dvO21 DNA *EcaI* digested, and religated by T4 DNA ligase at 225 μ g DNA/ml. Arrows at the right mark positions of the mixed linear dimer (1 λ dvh93+1 λ dvO21), and of mixed linear trimers (2 λ dvh93 + 1 λ dvO21, and 1 λ dvh93 + 2 λ dvO21, respectively) in lane 8, from bottom to top. The faint band in lane 7 at a position slightly below the mixed linear dimer band in lane 8 is caused by λ dvh93 dimer nicked circles, and the band slightly above the position of linear λ dvO21 monomers in lanes 8 and 9 originates from λ dvO21 monomer nicked circles. Other circular molecular species of both plasmids, supercoiled or relaxed, are also visible besides the dominant series of linear oligomers.

DNA ligase reactions, and *Eco*I linearised λ dvO21 has been used as a substrate for assaying ligase activity during a routine preparation of this enzyme.

Plasmid pMB1 contains a single *Eco*I recognition site outside the gene for the *Eco*RI endonuclease, and in pBR313 three cleavage sites have been observed (Fig. 3) which are located at 3.2 kb, 4.7 kb, and 6.3 kb relative to the *Eco*RI reference point. None of them is, however, retained in the smaller pBR322 plasmid.

The enzyme cuts λ DNA into 13 fragments, 11 of which are visible in Fig. 3. Among these fragments, the largest one of 8.7 kb represents the right, and the fourth largest of 5.6 kb the left terminal fragment of λ DNA which together give rise to an additional band of 14.3 kb due to hybridisation via their single-stranded cohesive ends. In the course of these experiments a partial mapping of the λ :*Eco*I fragments visible in Fig. 3 has been done, which was based on *Eco*I double digestions with enzymes for which the λ DNA map is known (*Eco*RI, *Hind*III, *Egl*II, *Bam*HI, *Sal*I, and *Kpn*I), on similar cleavages of λ DNA variants such as λ b2, λ nin5, λ imm434, λ imm21 and others, and on *Eco*I digestion of isolated or cloned individual λ DNA fragments (data not shown). These experiments resulted in the following order of λ DNA:*Eco*I fragments: 5.6 kb/4.3 kb/(0.8 kb, 1.3 kb, 1.45 kb, 2.35 kb)/1.9 kb/7.7 kb/4.8 kb/6.4 kb/3.7 kb/8.7 kb, with the final order of four fragments in the region of genes G-K (λ :10500- λ :16600) not yet determined. Besides the fragment of 0.8 kb listed above a very small fragment of approximately 60 bp has been observed on polyacrylamide gels, the position of which remains undetermined, but can be excluded from the two rightmost *Eco*I cuts which are sequenced and occur in λ genes *rex* and *P*, at λ :37206 and λ :40881 relative to the reference point at λ -*Eco*RI(*ori*):40000.

Isoschizomeric restriction by endonucleases *Eco*I and *Bst*EII

For all of the DNA substrates used the same restriction patterns were obtained for both the *Eco*I and *Bst*EII enzymes (see Fig. 3 for λ and pBR313 DNA), and a detailed mapping of the *Bst*EII versus the *Eco*I cut within λ gene *P* resulted in the same series of very small subfragmentations in various double digests. While this proves the recognition sequences to be identical for both enzymes, their cleavage positions might still be different, as for example in the pair of *Sma*I/*Xma*I endonucleases (1).

Because identical protruding single-stranded ends on *Bst*EII fragments would allow mixed ligations with *Eco*I fragments, and different ends would prohibit an interaction between these two types of fragments, we decided

to cleave circular λ dvh93 DNA with *Bst*EII, and circular λ dvO21 DNA with *Eca*I for subsequent ligation reactions. Both plasmids contain an identical λ gene *P* sequence including the same single *Eca*I/*Bst*EII recognition site (GGTTACC), they differ, however, in length: 2.324 kb versus 3.153 kb. The linearised λ dv plasmids were treated individually, and also in an equimolar mixture with T4 DNA ligase, and an approximately equivalent formation of the uneven dimer (5.48 kb) as compared to the sum of the even dimers (4.65 kb and 6.35 kb) indicated the ability of *Bst*EII ends to interligate with *Eca*I ends. This is supported by the results for the uneven trimers (2 λ dvh93 + 1 λ dvO21, and 1 λ dvh93 + 2 λ dvO21 units), and also the uneven tetramers (see Fig. 3). - The analysis appears to be slightly biased by the presence of a trace of exonuclease in the purchased *Bst*EII enzyme which was used for linearisation of the λ dvh93 DNA.

DISCUSSION

Similar to several other restriction endonucleases which cleave an odd numbered palindromic sequence, such as *Asu*I, *Fnu*HI and *Hin*FI (1), the central position of the heptanucleotide sequence recognised by *Eca*I and *Bst*EII is degenerate (N), and therefore the protruding pentanucleotide, single-stranded tails in a mixture of fragments produced by these enzymes will not be fully complementary to each other. While this situation appears to be disadvantageous with respect to cloning of *Eca*I/*Bst*EII fragments in general, it may be favorable for special applications in cases where the sequences to be joined would be known.

It is also worth mentioning that in a mixed religation experiment of the two fragments resulting from *Eca*I cleavage of λ dv1 DNA (which have been chosen because they terminate one each in the four different pentanucleotide ends possible) low yields of dimerized individual fragments have been observed in addition to the dominating ligation of different fragments presumably through their complementary ends. The former ligation products have to be attributed to reactions involving N-mismatched tail sequences. Besides allowing additional cloning reactions, ligated *Eca*I fragments of such constitution may be potentially useful for further studies of mismatch repair processes both *in vivo* and *in vitro*.

A 6.6 kb cosmid cloning vector pHC79 has been developed which carries a single *Eca*I (*Bst*EII) recognition site very close to its hybrid *cos* sequence, and is contained in the right end of ϕ 80 DNA fused onto the left end of λ (22). Also, the single cleavage site present in pMB1 or the single

site present in the λ -P gene in λ -pBR322 hybrid plasmids such as pHL95 (8) could be used for cloning DNA fragments generated by *Eco*I or isoschizomer endonucleolytic digestions.

ACKNOWLEDGEMENT

We thank Mrs. G. Schlingmann for expert technical assistance, H.J. Gross for computer programming, J. Collins for a critical reading of the manuscript, and H. Kössel for helpful suggestions. These investigations were supported by a grant from the Deutsche Forschungsgemeinschaft to G.H.

REFERENCES

1. Roberts, R.J. (1978) *Gene* 4, 183-193
2. Hartmann, H. and Goebel, W. (1977) *FEBS Letters* 80, 285-287
3. Lautenberger, J.A., Edgell, M.H. and Hutchison III, C.A. (1980) *Gene* 12, 171-174
4. Pugatsch, T. and Weber, H. (1979) *Nucl. Acids Res.* 7, 1429-1444
5. Mayer, H., Schwarz, E., Melzer, M., Grosschedl, R., Schütte, H. and Hobom, G. (1978) Abstracts XII Int. Congr. Microbiology, München, p. 106
6. Mayer, H., Luijbrand, G. and Goebel, W. (1977) *Molec. gen. Genet.* 152, 145-152
7. Streeck, R.E. and Hobom, G. (1975) *Eur. J. Biochem.* 57, 595-606
8. Lusky, M. and Hobom, G. (1979) *Gene* 6, 137-172; 173-197
9. Maxam, A. and Gilbert, W. (1977) *Proc. Natl. Acad. Sci. USA* 74, 560-564
10. Wienand, U., Schwarz, Z. and Feix, G. (1979) *FEBS Letters* 98, 319-323
11. Brownlee, G.G. and Sanger, F. (1969) *Eur. J. Biochem.* 11, 395-399
12. Schwarz, E., Scherer, G., Hobom, G. and Kössel, H. (1980) *Biochemistry International* 1, 386-394
13. Pirrotta, V., Ineichen, K. and Walz, A. (1980) *Molec. gen. Genet.* 180, 369-376
14. Landsmann, J. (1980) thesis, Univ. Köln
15. Farabaugh, P.J. (1978) *Nature* 274, 765-769
16. Beck, E., Sommer, R., Auerswald, E.A., Kurz, C., Zink, B., Osterburg, G. and Schaller, H. (1978) *Nucleic Acids Res.* 5, 4495-4503
17. Sanger, F., Coulson, A.R., Friedman, T., Air, G.M., Barrell, B.G., Brown, N.L., Fiddes, J.C., Hutchison III, C.A., Slocombe, P.M. and Smith, M. (1978) *J. Mol. Biol.* 125, 225-246
18. Fiers, W., Contreras, R., Haegeman, G., Rogiers, R., Van de Voorde, A., Van Heuverswyn, H., Van Herreweghe, J., Volckaert, G. and Ysebaert, M. (1978) *Nature* 273, 113-120
19. Reddy, V.B., Thimmappaya, B., Dhar, R., Subramanian, K.N., Zain, B.S., Pan, J., Ghosh, P.K., Celma, M.L. and Weissman, S.M. (1978) *Science* 200, 494-502
20. Sutcliffe, J.G. (1979) *Cold Spring Harbor Symp. Quant. Biol.* 43, 77-90
21. Grosschedl, R. (1978) thesis, Univ. Freiburg
22. Hohn, B. and Collins, J. (1980) *Gene* 11, 291-298