Purification and properties of a new DNase activity from KB cells

Gerald D.Frenkel, Karen Randles and Neil Berns

Division of Laboratories and Research, New York State Department of Health, Albany, NY 12201, USA

Received 14 August 1981

ABSTRACT

A deoxyribonuclease activity with specificity towards single-stranded DNA has been purified approximately four hundred-fold from KB cells, by chromatography on DEAE-cellulose, phosphocellulose and hydroxylapatite. The last step of the purification results in separation of the enzyme from a DNase activity which has been described previously (Wang, E.C., Furth, J.J. and Rose, J.A., (1978) *Biochemistry* 17: 544-549). The properties of the new DNase activity are significantly different from those of the enzymes which have previously been identified in these cells. The activity sediments at approximately 7.5S in a glycerol gradient. The DNase activity is optimal at pHs between 6.0 and 6.5. It cleaves DNA endonucleolytically and hydrolyzes single-stranded DNA at about 11 times the rate of double-stranded DNA and at twice the rate of Poly (dA). The activity is moderately sensitive to inhibition by N-ethylmaleimide and is inhibited 80% by 50 mM NaCl. It is stimulated twenty-fold by Mn^{++} at an optimal concentration of approximately 0.7 mM. It is stimulated to a lesser extent by Mg^{++} , but not by Ca^{++} .

INTRODUCTION

Deoxyribonucleases are hydrolytic enzymes which appear to be ubiquitous in biological systems (1). They display a wide variety of enzymatic properties, substrate specificities and modes of hydrolysis. They are clearly involved in a number of cellular functions; however, the definitive demonstration of the role(s) of specific DNases has proven to be a difficult task (2). Nevertheless, over the past few years several DNases have been shown to be involved in the processes of DNA replication, genetic recombination, DNA repair, and restriction of foreign DNA in procaryotic systems (3-5). Due to the lack of appropriate mutants, the function(s) of DNases in higher cells have not yet been demonstrated. It is nevertheless reasonable to suppose that these enzymes play correspondingly important roles in higher cells. For this reason it is important to characterize the types of DNase activities which are present in a eucaryotic cell. KB cells, a permanent line of human cells derived from a nasopharyngeal carcinoma (6), have been utilized in many studies on cellular and viral DNA

metabolism. There have been several reports of the isolation of DNases from these cells (7-11). During the course of studies on the effect on adenovirus infection on the level of cellular DNase (12) we discovered an activity which had not been previously described. In this communication we describe the properties of this new DNase activity.

EXPERIMENTAL PROCEDURES

Materials - ^3H -adenovirus DNA was purchased from Bethesda Research Laboratories. Single-stranded DNA was prepared by alkaline denaturation of native DNA. ^3H -poly(dA) was purchased from Miles. DEAE-cellulose (DE-52) and phosphocellulose were obtained from Whatman. Hydroxylapatite was obtained from Bio-Rad, N-ethylmaleimide and MES buffer from Calbiochem. The bovine serum albumin used in all experiments was crystallized grade from Miles.

Growth of Cells - KB cells were grown in suspension cultures in Minimal Essential Medium (Flow Labs) containing 10% heat-inactivated fetal calf serum.

Assay of DNase Activity - The assay measures the conversion of denatured ^3H -DNA to an acid-soluble form. The reaction mixture (0.3 ml) contained 0.4 nmole of denatured ^3H -adenovirus DNA (1.5×10^5 cpm/nmole), 33 mM MES buffer (pH 6.1), 0.67 mM MnCl_2 , 10 mM mercaptoethanol, 0.2 mg/ml bovine serum albumin, and enzyme diluted in 10 mM Tris-HCl buffer (pH 7.5) containing 1 mM mercaptoethanol and 0.5 mg/ml bovine serum albumin. After incubation for 30 min. at 37°C , 0.2 ml of salmon sperm DNA (2.5 mg/ml) and 0.5 ml of 10% trichloroacetic acid were added. After centrifugation at $10,000 \times g$ for 10 min., 0.5 ml of the supernatant fluid was removed and the radioactivity determined in a scintillation vial containing 4 ml of ACS scintillation fluid (Amersham). One unit of enzyme is defined as the amount catalyzing the conversion of 1 nmole of DNA-nucleotide to an acid-soluble form in 30 min. at 37°C .

RESULTS

Purification

All procedures were carried out at 0° - 4°C . A summary of a typical purification is given in Table I. The increase in total activity which occurs during the course of the purification is most likely due to removal of the DNase inhibitor protein which we have previously described (14).

Preparation of Extract - KB cells (5×10^9) were washed twice with 0.15 M sodium phosphate buffer (pH 7.6) containing 0.1 M NaCl and resuspended in 40 ml of 0.1 M Tris-HCl buffer (pH 8.1) containing 2 mM mercaptoethanol. The cells were lysed by sonic treatment for 3 min. (in 30 second pulses) at 4 amps in a

TABLE I
Purification of DNase activity^a

Fraction		Total Units	Specific Activity
			<u>units/mg</u>
I.	Extract	46	0.09
II.	Ammonium Sulfate	80	0.35
III.	DEAE-cellulose	97	3.5
IV.	Phosphocellulose	111	7.9
V.	Hydroxylapatite	21 ^b	40

^aDetails of the purification are described in the text. Protein was determined by the method of Lowry et al. (13)

^bTotal units for Fraction V is corrected for the percentage of Fraction IV which was utilized for the hydroxylapatite chromatography.

Branson sonifier. The extract was centrifuged for 90 min. at 16,000 rpm in the Sorvall centrifuge, and the supernatant fluid was recovered (Fraction I).

Ammonium Sulfate Precipitation - $(\text{NH}_4)_2\text{SO}_4$ was added to Fraction I to 30% saturation with stirring over 30 min. After centrifugation for 20 min. at 10,000 rpm, the supernatant fluid was recovered, made 50% in $(\text{NH}_4)_2\text{SO}_4$, and centrifuged as above. The precipitates were dissolved in 0.1 Tris-HCl buffer (pH 7.5) containing 2 mM mercaptoethanol and dialyzed against 10 mM Tris-HCl buffer (pH 7.5) containing 10 mM mercaptoethanol and 10% glycerol. The precipitate obtained at 50% saturation contained 90% of the DNase activity recovered (Fraction II).

DEAE-Cellulose Chromatography - A column of DEAE-cellulose (2.5 cm x 13 cm) was washed with 50 column volumes of 10 mM Tris-HCl buffer (pH 7.5) containing 10 mM mercaptoethanol and 10% glycerol, and Fraction II was applied to the column. The column was washed with 4 column volumes of the same buffer, and the DNase was eluted with 2 column volumes of the same buffer containing 0.15 M NaCl (Fraction III).

Phosphocellulose Chromatography - A column of phosphocellulose (2.5 cm x 10 cm) was washed with 50 column volumes of 10 mM Tris-HCl buffer (pH 7.5) containing 10 mM mercaptoethanol and 10% glycerol, and Fraction III was applied to the column. The column was washed successively with 2 column volumes each of the same buffer containing 0.15 and 0.3 M NaCl. The 0.3 M NaCl eluate was

dialyzed against 10 mM Tris-HCl buffer (pH 7.5) containing 10 mM mercaptoethanol and 10% glycerol (Fraction IV).

Hydroxylapatite Chromatography - A column of hydroxylapatite (1.0 cm x 10 cm) was washed with 20 column volumes of 10 mM potassium phosphate buffer (pH 7.6). Fraction IV (7 mg of protein) was loaded on the column, which was then washed with 2 column volumes of 10 mM potassium phosphate buffer (pH 7.6) containing 10 mM mercaptoethanol and 10% glycerol. The proteins were eluted with a continuous gradient of 10 mM to 300 mM potassium phosphate buffer (pH 7.6) containing 10 mM mercaptoethanol and 10% glycerol, and the fractions collected were assayed for DNase activity (Fig. 1). Fractions 32-35, eluting with a phosphate concentration of approximately 0.09 M, constitute Fraction V.

When the column fractions were assayed at alkaline pH in the presence of Mg^{++} , a second peak of DNase activity was evident eluting with a phosphate concentration of approximately 0.14 M in fractions 40-45. This activity had a pH optimum of 8.5 and is probably identical to the DNase described by Wang *et al* (10) (see Discussion). This enzyme has no detectable activity in the presence of Mn^{++} (10), so it does not contribute to the activity seen in Figure 1. Fraction V is not significantly contaminated with this enzyme since the former has no activity at pH 8.5 in the presence of Mg^{++} .

Fig. 1. Hydroxylapatite chromatography of DNase. Details of the chromatography are described in the text. Aliquots (10 μ l) of the fractions were assayed for DNase activity under standard conditions, as described under Experimental Procedures. Phosphate concentration was determined by measurement of refractive index. Protein concentration was determined by the method of Lowry *et al* (13).

Properties of the DNase activity

Stability - Fraction V DNase activity was rather unstable; loss of 50% of the activity occurred within two weeks at 0°C. Dialysis or dilution of Fraction V resulted in immediate loss of most of the activity.

Sedimentation Analysis - Sedimentation analysis of Fraction V in a 10-30% glycerol gradient yielded a single broad peak of DNase activity, sedimenting at approximately 7.5S relative to a bovine serum albumin marker. (Fig. 2)

pH Optimum - Optimal DNase activity is obtained at pHs between 6.0 and 6.5, with very little activity at more acid or alkaline pHs. This is in sharp contrast to the pH profile of the DNase activity which elutes from hydroxylapatite at 0.14 M phosphate (see above), which shows very little activity at pHs 6.0 to 6.5.

Requirements for Activity - In the absence of a divalent cation only 5% of maximal activity is obtained. Maximal activity is obtained in the presence of Mn^{++} , at approximately 0.7 mM (Fig. 3). A profile of enzyme activity in the presence of Mg^{++} (Fig. 3) indicates that Mg^{++} can substitute for Mn^{++} , but with a resulting relative optimal activity of only 30%. Addition of Ca^{++} does not increase the level of activity above that obtained in the absence of a divalent cation (Table II).

Other properties of the DNase activity are shown in Table II. Omission of mercaptoethanol from the reaction results in only a 13% decrease in enzyme

Fig. 2. Sedimentation analysis of DNase activity in a glycerol gradient. Fraction V (0.5 unit) was layered on a 10-30% glycerol gradient containing 10 mM Tris-HCl buffer (pH 8) and 10 mM mercaptoethanol. After centrifugation for 16 h at 38,000 rpm in the Beckman SW50.1 rotor, fractions were collected (total 29) and 50 μ l aliquots were assayed for DNase activity as described under Experimental Procedures. The arrow indicates the position to which bovine serum albumin sedimented in a parallel gradient, as determined by measurement of optical density at 260 nm. Direction of sedimentation is from right to left.

Fig. 3. DNase activity as a function of concentration of divalent cation. Fraction V was assayed for DNase activity under standard conditions (see Experimental Procedures) except that the concentration of divalent cation was as indicated. (O) MnCl_2 ; (Δ) MgCl_2 .

TABLE II

Reaction requirements of DNase activity

Reaction Mixture	Activity %
Complete ^a	100 ^b
- MnCl_2	5
- MnCl_2 + CaCl_2 (0.67 mM)	5
-Mercaptoethanol	87
-Mercaptoethanol + 5 mM N-ethylmaleimide	54
-Bovine Serum Albumin	47
+17 mM NaCl	81
+50 mM NaCl	19
+100 mM NaCl	5

^aThe complete reaction mixture is described under Experimental Procedures.

^b100% = 24 pmoles

activity. However, addition of the sulfhydryl reagent N-ethylmaleimide in the absence of mercaptoethanol results in a further decrease of about 30% in activity. The activity is sensitive to salt; the presence of 50 mM NaCl in the reaction inhibits the activity by 80%.

Substrate Specificity - A comparison of the rates of hydrolysis of single-stranded and double-stranded DNA is shown in Figure 4. Single-stranded DNA is clearly the preferred substrate; the rate of hydrolysis of single-stranded DNA is approximately 11-fold greater than that of double-stranded DNA. ^3H -poly(dA) is hydrolyzed at approximately 1/2 the rate of hydrolysis of single-stranded DNA.

Endonucleolytic Cleavage - In order to determine whether hydrolysis of the DNA is occurring endonucleolytically, denatured DNA was incubated with sufficient DNase to solubilize only a small percentage of the substrate. The size of the DNA was then determined by sedimentation analysis in an alkaline sucrose gradient. From a parallel reaction it was determined that 1% of the substrate had been made acid soluble in the reaction. The results (Fig. 5) showed that significant breakage of the DNA occurred even though only 1% of the DNA had been made acid-soluble. Since exonucleolytic hydrolysis of the DNA to 1% acid soluble would not produce a significant change in the sedimentation rate, it is clear that the enzyme is cleaving the DNA endonucleolytically.

Fig. 4. Hydrolysis of single-stranded and double-stranded DNA. The reaction mixture (2.4 ml) was as described under Experimental Procedures except that it contained 6.4 nmoles of either native or denatured ^3H -adenovirus DNA, which is twice the standard concentration. Fraction V (0.13 units) was added, and the reaction mixture was incubated at 37°C . At the indicated times 0.3 ml aliquots were removed and acid soluble nucleotide was determined as described under Experimental Procedures. The best fit lines were determined by least-squares analysis of the data. (O) single-stranded DNA; (●) double-stranded DNA.

Fig. 5. Sedimentation analysis of DNase-treated DNA. Denatured ^3H -adenovirus DNA (0.2 nmoles; 4.4×10^4 cpm/nmole) was incubated under standard conditions (see Experimental Procedures) either without enzyme or with sufficient Fraction V to make 1% of the substrate acid soluble (as determined from a parallel reaction). After the reaction EDTA was added to 6.7 mM and the reaction mixture was layered on a 5-20% alkaline sucrose gradient containing 0.2 N NaOH and 0.1 M NaCl. After centrifugation for 5 h at 41,000 rpm in the Beckman SW50.1 rotor at 4°C , fractions were collected into vials and the radioactivity determined in a liquid scintillation counter. (O) no DNase; (●) 0.009 units of DNase. Direction of sedimentation is from right to left.

DISCUSSION

During the past few years several deoxyribonucleases have been purified from KB cells. To facilitate discussion of these enzymes, as well as to clarify the distinction between them, we have listed them in Table III, and assigned them arbitrary designations. KB DNase α has maximal activity at acid pH and is active on native DNA (7,8,15). It has not been reported whether the enzyme has activity on denatured DNA. The presence of a DNase with activity on native DNA at neutral pH was also reported, but the activity was not characterized further (7). This enzyme may have been identical to KB DNase β which was subsequently described by Tsuruo *et al* (9). This enzyme shows maximal activity at alkaline pH, and has a clear preference for native DNA (9). Two activities have been described which clearly prefer single-stranded to double-stranded DNA (10,11). They have been referred to as KB₁ and KB₂ (10,11); we have designated them DNase γ and δ respectively.

The DNase activity which is the subject of this communication (which we designate DNase ϵ) is clearly distinct from these enzymes in both physical and enzymatic properties (Table III). DNase ϵ activity sediments at a rate which

TABLE III

DNase activities of KB cells

DNase Activity	Sedimentation Coefficient	pH Optimum	Preferred Divalent Cation	Preferred DNA	Reference
α	3.9	4.0	Mg^{++}	Native (?)	7, 8, 15
β	3.8	8.8	Mg^{++}	Native	9
γ	4.6	9.2	Mg^{++}	Denatured	10
δ	3.2	9.5	Mg^{++}	Denatured	11
ϵ	7.5	6.5	Mn^{++}	Denatured	This work

is significantly greater than the other enzymes. In addition to the obvious, striking differences in pH optimum and divalent cation preferences, DNase ϵ also differs in that it prefers denatured DNA over poly(dA), in contrast to DNases γ and δ which both hydrolyze poly(dA) more rapidly than denatured DNA (11,15).

It is not clear whether the DNase ϵ preparation contains only a single DNase. There are two aspects to this uncertainty. First, it is not clear whether or not the hydrolytic activity towards double-stranded DNA which is present in the preparation (see Fig. 4) is due to the same enzyme as the hydrolytic activity towards single-stranded DNA. Second, the rather broad peak of DNase ϵ activity which is obtained following sedimentation in a glycerol gradient (Fig. 2), as well as the incomplete inhibition of the activity by N-ethylmaleimide (Table II) raise the possibility that there is more than one enzyme contributing to the DNase ϵ activity. Because of the instability of the DNase activity (see Results) we have been unable as yet to resolve these uncertainties.

ACKNOWLEDGEMENTS

This work was supported in part by Grant CA28084 from the National Cancer Institute and Grant NP-198 from the American Cancer Society.

REFERENCES

1. Kowalski, D. and Laskowski, M. Sr. (1976) In: Handbook of Biochemistry and Molecular Biology; 3rd Ed., Nucleic Acids Vol. II. pp 491-531, CRC Press.

2. Kornberg, A. (1980) DNA Replication, W.H. Freeman and Co., San Francisco.
3. Grossman, L., Braun, A., Feldberg, R. and Mahler, I. (1975) Ann. Rev. Biochem. 44, 19-43.
4. Clark, A.J. (1973) Ann. Rev. Gen. 7, 67-86.
5. Roberts, R.J. (1976) CRC Critical Rev. Biochem. 4, 123-164.
6. Eagle, H. (1955) Proc. Soc. Exptl. Biol. Med. 89, 362-364.
7. Reif, U.M., Winterhoff, U., Lundholm, U., Philipson, L., and Doerfler, W. (1977) Eur. J. Biochem. 73, 313-325.
8. Reif, U., Winterhoff, U., Doerfler, W. (1977) Eur. J. Biochem. 73, 327-333.
9. Tsuruo, T., Arens, M., Padmanabhan, R. and Green, M. (1978) J. Biol. Chem. 253, 3400-3407.
10. Wang, E.C., Furth, J.J. and Rose, J.A. (1978) Biochemistry 17, 544-549.
11. Wang, E.C. and Rose, J.A. (1981) Biochemistry 20, 755-758.
12. Nass, K. and Frenkel, G.D. (1978) J. Virol. 26, 540-543.
13. Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951) J. Biol. Chem. 193, 265-275.
14. Nass, K. and Frenkel, G.D. (1979) J. Biol. Chem. 254, 3407-3410.
15. Padmanabhan, R., Stabel, S., Winterhoff, U. and Doerfler, W. (1979) Nucleic Acids Res. 6, 3161-3176.