Published in final edited form as:

ACS Comb Sci. 2012 February 13; 14(2): 85–88. doi:10.1021/co200199h.

Expedient Synthesis of a 72-Membered Isoxazolino- β -ketoamide Library by a 2•3-Component Reaction

John M. Knapp, Jie S. Zhu, Alex B. Wood, and Mark J. Kurth*

Department of Chemistry, University of California/Davis, One Shields Avenue, Davis, CA 95616

Abstract

An efficient 2•3-component reaction (2•3CR; a 2-component reaction followed, in one pot, by a 3-component reaction) is presented for the synthesis of isoxazolino- β -ketoamides. This 2•3CR proceeds by (i) a Meldrum's acid-generated acyl ketene, which is trapped by an amine to form a β -ketoamide intermediate in a 2CR followed, in one pot, by (ii) a Mannich reaction followed by elimination of dimethyl amine•HCl to generate an α,β -unsaturated β -ketoamide dipolarophile that reacts in a nitrile oxide 1,3-dipolar cycloaddition reaction. This one-pot 2•3CR process delivers the targeted isoxazolino- β -ketoamide product. A total of 72 compounds are presented, all of which have been submitted to the *NIH Molecular Libraries Small Molecule Repository* for high-throughput biological screening.

Introduction

Multicomponent reactions offer an expedient route to complex targets¹ since the combination of three or more reactants within a single reaction typically leads to a shorter reaction sequence and fewer purification steps.¹ Moreover, multicomponent products are comparatively far more elaborate than their starting material counterparts and, by altering a single reactive component, a wide range of diversity can be achieved. For these reasons, multicomponent reactions, such as the Biginelli,² Passerni,³ Hantzche,⁴ and Ugi⁵ reactions, have been widely used for the construction of combinatorial libraries.⁶ The development of new multicomponent reactions has broad implications in synthetic methodology, library production, and biological screening.

Isoxazoles have little precedence as pharmacophores. However, we have discovered that a class of isoxazoles function as a cystic fibrosis transmembrane regulator protein activator⁷ and others have previously shown this substructure to be bioactive in various other systems.⁸ As an under exploited heterocycle, isoxazoles could potentially find various new applications and serve as future synthetic targets. To address this possibility, expedient library synthesis is needed to enable the requisite random screening.

Isoxazoles are synthons for β -amino acids⁹ and γ -amino alcohols,¹⁰ which are abundant in natural products and medicinal compounds.¹¹ Consequently, the diversity oriented synthesis of isoxazoles can have considerable significance. To this end, we present here the synthesis of a seventy-two membered library of isoxazolino- β -ketoamides prepared by a modified multicomponent reaction with the ability for easy diversification.

* mjkurth@ucdavis.edu .

Supporting Information Available Detailed experimental procedures, characterization data, and ¹H and ¹³C NMR spectra of 20 representative compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Results and Discussion

We recently reported a 4-component reaction (4CR) for the synthesis of isoxazolino- β -ketoamides¹² as outlined in Scheme 1. This initial investigation was conducted as a proof-of-concept study for the development of a multicomponent macrocyclization reaction, which we have now evolved into a general protocol for the synthesis isoxazolino- β -ketoamides. This procedure incorporates an acyl ketene precursor (**1**) and an amine (**2**) to form the β -ketoamide moiety, which then undergoes a Mannich reaction with Eschenmoser's salt (**3**) followed by elimination of dimethyl ammonium chloride. The resulting doubly activated Michael acceptor then serves as a dipolarophile in a nitrile oxide cycloaddition ($\Rightarrow$ **4**) to, in one-pot, deliver the isoxazolino- β -ketoamide target (**5**). Our previously reported yields for this 4CR reaction ranged from 39 to 50%. Herein, we demonstrate modification and expansion of this versatile transformation.

The combination of four reactants can lead to undesirable side-product formation and, consequently, lower isoxazolino- β -ketoamide yields. Despite these inherent limitations, our 4CR produces modest yields (39–50%) of the targeted product because of the high reactivity of the acyl ketene with the amine nucleophile. This high acyl ketene/amine reactivity effectively suppresses side reactions that form isoxazol-5(2*H*)-ones (**6**) and *N*-hydroxyimidamides (**7**). That said, we report here a modified method that delivers isoxazolino- β -ketoamides in both higher average yields and cleaner reactions – advancements that enable efficient library synthesis.

As delineated in Scheme 2, performing the multicomponent reaction as a 2•3CR (a 2-component reaction followed by a 3-component reaction), instead of a 4CR, results in isoxazolino- β -ketoamide yields up to 88% as well as excellent product purity. This 2•3CR was accomplished as a one-pot procedure by heating a DMF solution of the acyl ketene precursor in the presence of only the amine nucleophile, thus pre-forming the β -ketoamide in a 2CR. While it is known that dioxolinones and Meldrum's acids fragment thermally to acyl ketenes, it must be said that an alternative mechanism where the amine attacks first, followed by loss of CO₂ and acetone, is possible. Subsequent addition of Eschenmoser's salt and chlorooxime reagents, to the same flask, results in a 3CR (collectively, a 2•3CR) that delivers the isoxazolino- β -ketoamide product. This 2•3CR modification essentially avoids both the chlorooxime + acyl ketene cycloaddition side reaction ($\rightarrow$ isoxazol-5(2*H*)-ones; **6**; Scheme 1) and the amine + chlorooxime side reaction ($\rightarrow$ *N*-hydroxyimidamides; **7**; Scheme 1), while proficiently effecting the desired transformation ($\rightarrow$ isoxazolino- β -ketoamide; **5**; Scheme 2).

In addition to the 2•3CR synthetic modification, we wanted to expand the scope of this reaction by utilizing an array of diverse reactants (Figure 1). A total of eight amines were used. Both aliphatic and aromatic amines worked well; however, yields dropped when aminothiazole, ethanolamine, or acetohydrazide were used. For the library reported here, three different acyl ketene precursors (*p*-methoxyphenyl,¹³ cyclopropyl,¹⁴ and methyl) as well as three different chlorooximes (*o*-pyridinyl,¹⁵ *o*-trifluoromethylphenyl,¹⁶ and carboethoxy¹⁷) were used. We also demonstrate that these substituted acyl ketenes and chlorooximes tolerate the 2•3CR reaction conditions. Isolated yields for all variations are shown in Table 1.

Considerable functional group diversity is represented in this library in which product molecular weights range from 272 to 496. Manipulation of the acyl ketene-, amine-, and chlorooxime-derived R-groups allow for dramatic alteration of the isoxazolino- β -ketoamide three-dimensional shape. Together, the ease of altering functional groups, molecular

weights, and molecular geometry with this 2•3CR-based methodology allows for the rapid assembly of a library of biological relevant isoxazolino- β -ketoamides.

Conclusion

We have successfully synthesized a library of seventy-two isoxazolino- β -ketoamides with novel 2•3CR-based methodology. This library effectively expands the scope of our multicomponent methodology and demonstrates its application in diversity-oriented synthesis. Compared to its 4CR variant, this one-pot 2•3CR method delivers the targeted scaffold with much improved yields – as much as 38% in side-by-side comparative reactions. The compounds reported here have been deposited in the *NIH Molecular Libraries Small Molecule Repository* for high-throughput biological screening and assay results for these molecules and similar analogs will likely follow.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank the National Institutes of Health (GM0891583 and RR1973) and the National Science Foundation [CHE-0910870; and CHE-0443516, CHE-0449845, CHE-9808183, and DBIO 722538 for NMR spectrometers] for their generous support.

References

- 1). (a) Trost BM. Atom economy – a challenge for organic synthesis: homogeneous catalysis leads the way. *Angew. Chem., Int. Ed.* 1995; 34:259–281. (b) Noyori R. synthesizing our future. *Nat. Chem.* 2009; 1:5–6. [PubMed: 21378778] (c) Bienayme H, Hulme C, Oddon G, Schmitt P. Maximizing synthetic efficiency: multi-component transformations lead the way. *Chem. Eur. J.* 2000; 6:3321–3329. [PubMed: 11039522] (d) Posner GH. Multicomponent one-pot annulations forming 3 to 6 bonds. *Chem. Rev.* 1986; 86:831–844. (e) Boger DL, Desharnais J, Capps K. Solution-phase combinatorial libraries: modulating cellular signaling by targeting protein-protein or protein-DNA Interactions. *Angew. Chem., Int. Ed.* 2003; 42:4138–4176.
- 2). (a) Biginelli P. The urea-aldehyde derivatives of acetoacetic esters. *Ber.* 1891; 24:2962–2967. (b) Biginelli P. The urea-aldehyde derivatives of acetoacetic esters. *Gazz. Chim. Ital.* 1893; 23:360–416.
- 3). Passerni M. Isonitriles. II. Compounds with aldehydes or with ketones and monobasic organic acids. *Gazz. Chim. Ital.* 1921; 51:181–189.
- 4). Hantzsch A. Synthesis of pyridine derivatives from acetoacetic esters and aldehydeammonia. *Liebigs Ann. Chem.* 1882; 215:1–82.
- 5). (a) Ugi I, Meyr R, Fetzer U, Steinbrückner C. Studies on isonitriles. *Angew. Chem.* 1959; 71:386. (b) Ugi I, Steinbrückner C. Concerning a new condensation principle. *Angew. Chem.* 1960; 72:267–268. (c) Ugi I. The α -addition of immonium ions and anions to isonitriles coupled with secondary reactions. *Angew. Chem.* 1962; 74:9–22.
- 6). (a) Ugi I, Domling A, Werner B. Since 1995 the new chemistry of multicomponent reactions and their libraries, including their heterocyclic chemistry. *J. Hetero. Chem.* 2000; 37:647–658. (b) Ugi I. The improved search of new products by multicomponent reactions and their libraries. *Lat. Kim. Z.* 1999; 1:47–58. (c) Ugi I. Fast and permanent changes in preparative and pharmaceutical chemistry through multicomponent reactions and their 'libraries.'. *Eesti. Tead. Akad., Keemia.* 1995; 44:237–273. (d) Bienayme H, Hulme C, Oddon G, Schmitt P. Maximizing synthetic efficiency: multi-component transformations lead the way. *Chem. Eur. J.* 2000; 6:3321–3329. [PubMed: 11039522]

- 7). Sammelson RE, Ma T, Galletta LJV, Verkman AS, Kurth MJ. 3-(2-Benzoyloxyphenyl)isoxazoles and isoxazolines: synthesis and evaluation as CFTR activators. *Bioorg. Med. Chem. Lett.* 2003; 13:2509–2512. [PubMed: 12852954]
- 8). (a) Del Corral JMM, Gordaliza M, Castro MA, Lopez-Vazquez ML, Garcia-Gravalo MD, Broughton HB, San Feliciano A. Bioactive isoxazoline and oxime derivatives from 7-Ketolignans. *Tetrahedron.* 1997; 53:6555–6564. (b) Cheng C, Yu P, Li Z, Shu J, Wang X. Synthesis of 2-methyl-3-(4-methoxyphenyl)-5-(3-methoxyphenyl)isoxazoline and its bioactivity. *Youji Huazue.* 2010; 30:833–836.
- 9). (a) Fuller AA, Chen B, Minter Aaron, R. Mapp AK. Succinct synthesis of β -amino acids via chiral isoxazolines. *J. Am. Chem. Soc.* 2005; 127:5376–5383. [PubMed: 15826175] (b) Sewald N. Synthetic Routes toward enantiomerically pure β -amino acids. *Angew. Chem. Int. Ed.* 2003; 42:5794–5795. (c) Minter AR, Fuller AA, Mapp AK. A Concise approach to structurally diverse β -amino acids. *J. Am. Chem. Soc.* 2003; 125:6846–6847. [PubMed: 12783519]
- 10). (a) Tang S, He J, Sun Y, He L, She X. Efficient and regioselective synthesis of 5-hydroxy-2-isoxazolines: versatile synthons for isoxazoles, β -lactams, and γ -amino alcohols. *J. Org. Chem.* 2010; 75:1961–1966. [PubMed: 20151710] (b) Jaeger V, Buss V, Schwab W. Syntheses via isoxazolines. III. Diastereoselective synthesis of γ -amino Alcohols with 2 and 3 chiral centers. *Tetrahedron Lett.* 1978; 34:3133–3136.
- 11). (a) Tirel PJ, Vaultier M, Carrie R. Ligand assisted hydride delivery: an expeditious stereoselective total synthesis of ($\pm$)-Norsedamine and its pyrrolidino analog. *Tetrahedron Lett.* 1989; 30:1947–1950. (b) Dondoni A, De Lathauwer G, Perrone D. A convergent synthesis of the renin inhibitor SPP-100 using a nitron intermediate. *Tetrahedron Lett.* 2001; 42:4819–4823. (c) Feske BD, Stewart JD. Chemoenzymatic formal total synthesis of (–)-Bestatin. *Tetrahedron Asym.* 2005; 16:3124–3127. (d) Aggarwal VK, Roseblade S, Alexander R. The use of enantiomerically pure ketene dithioacetal bis(sulfoxides) in highly diastereoselective intramolecular nitron cycloadditions. Application in the total synthesis of the p-amino acid (–)-Cispentacin and the First asymmetric synthesis of the cis-(3R,4R)-4-aminopyrrolidine-3-carboxylic acid. *Org. Biomol. Chem.* 2003; 1:684–691. [PubMed: 12929455] (e) Jiang J, Schumacher K, Joullie AA, Davis FA, Reddy RE. Approaches toward the total syntheses of isatins A, B, and C. *Tetrahedron Lett.* 1994; 35:2121–2124.
- 12). Knapp JM, Fetting JC, Kurth MJ. Multicomponent macrocyclization reactions (MCMRs) employing highly reactive acyl ketene and nitrile oxide intermediates. *Org. Lett.* 2011; 13:4732–4735. [PubMed: 21827181]
- 13). Yamamoto Y, Watanabe Y, Ohnishi S. 1,3-Oxazines and related compounds. XIII. Reactions of acyl Meldrum's acids with Schiff bases giving 2,3-disubstituted 5-acyl-3,4,5,6-tetrahydro-2H-1,3-oxazine-4,6-diones and 2,3,6-trisubstituted 2,3-dihydro-1,3-oxazin-4-ones. *Chem. Pharm. Bul.* 1987; 35:1860–1870.
- 14). Beyer WRC, Woithe K, Luke B, Schindler M, Antonicek H, Scherkenbeck J. Asymmetric total synthesis of the indole alkaloid cyclopiasonic acid and first structure-activity data. *Tetrahedron.* 2011; 67:3062–3070.
- 15). Grob JE, Nunez J, Dechantsreiter MA, Hamann LG. Regioselective synthesis and slow-release Suzuki-Miyaura cross-coupling of MIDA boronate-functionalized isoxazoles and triazoles. *J. Org. Chem.* 2011; 76:10241–10248. [PubMed: 22047083]
- 16). Guggenheim KG, Butler JD, Painter PP, Lorsch BA, Tantillo DJ, Kurth MJ. Synthesis of spiro-fused pyrazolidoylisoxazolines. *J. Org. Chem.* 2011; 76:5803–5812. [PubMed: 21650164]
- 17). Kozikowski AP, Tapadar S, Luchini DN, Kim KH, Billadeau DD. Use of the nitrile oxide cycloaddition (NOC) reaction for molecular probe generation: a new class of enzyme selective Histone Deacetylase inhibitors (HDACIs) showing picomolar activity at HDAC6. *J. Med. Chem.* 2008; 51:4370–4373. [PubMed: 18642892]

Scheme 1.
4CR route to isoxazolino- β -ketoamides.

Scheme 2.

Isoxazolino-β-ketoamides by a multicomponent (2•3CR) process.

Figure 1.
2•3CR diversity inputs.

Table 1

Library of isoxazolino- β -ketoamides.

Entry	Acyl Ketone	Chloro-Oxime	Amine	Product	Yield
1	{1}	{1}	{1}	5 {1,1,1}	80%
2	{1}	{1}	{2}	5 {1,1,2}	79%
3	{1}	{1}	{3}	5 {1,1,3}	30%
4	{1}	{1}	{4}	5 {1,1,4}	40%
5	{1}	{1}	{5}	5 {1,1,5}	40%
6	{1}	{1}	{6}	5 {1,1,6}	51%
7	{1}	{1}	{7}	5 {1,1,7}	76%
8	{1}	{1}	{8}	5 {1,1,8}	88%
9	{1}	{2}	{1}	5 {1,2,1}	75%
10	{1}	{2}	{2}	5 {1,2,2}	76%
11	{1}	{2}	{3}	5 {1,2,3}	31%
12	{1}	{2}	{4}	5 {1,2,4}	44%
13	{1}	{2}	{5}	5 {1,2,5}	31%
14	{1}	{2}	{6}	5 {1,2,6}	50%
15	{1}	{2}	{7}	5 {1,2,7}	78%
16	{1}	{2}	{8}	5 {1,2,8}	83%
17	{1}	{3}	{1}	5 {1,3,1}	81%
18	{1}	{3}	{2}	5 {1,3,2}	80%
19	{1}	{3}	{3}	5 {1,3,3}	19%
20	{1}	{3}	{4}	5 {1,3,4}	48%
21	{1}	{3}	{5}	5 {1,3,5}	41%
22	{1}	{3}	{6}	5 {1,3,6}	38%

Entry	Acyl Ketene	Chloro-Oxime	Amine	Product	Yield
23	{1}	{3}	{7}	5{1,3,7}	75%
24	{1}	{3}	{8}	5{1,3,8}	75%
25	{2}	{1}	{1}	5{2,1,1}	30%
26	{2}	{1}	{2}	5{2,1,2}	50%
27	{2}	{1}	{3}	5{2,1,3}	8%
28	{2}	{1}	{4}	5{2,1,4}	15%
29	{2}	{1}	{5}	5{2,1,5}	35%
30	{2}	{1}	{6}	5{2,1,6}	44%
31	{2}	{1}	{7}	5{2,1,7}	55%
32	{2}	{1}	{8}	5{2,1,8}	64%
33	{2}	{2}	{1}	5{2,2,1}	60%
34	{2}	{2}	{2}	5{2,2,2}	58%
35	{2}	{2}	{3}	5{2,2,3}	17%
36	{2}	{2}	{4}	5{2,2,4}	31%
37	{2}	{2}	{5}	5{2,2,5}	30%
38	{2}	{2}	{6}	5{2,2,6}	33%
39	{2}	{2}	{7}	5{2,2,7}	60%
40	{2}	{2}	{8}	5{2,2,8}	60%
41	{2}	{3}	{1}	5{2,3,1}	62%
42	{2}	{3}	{2}	5{2,3,2}	42%
43	{2}	{3}	{3}	5{2,3,3}	12%
44	{2}	{3}	{4}	5{2,3,4}	46%
45	{2}	{3}	{5}	5{2,3,5}	60%
46	{2}	{3}	{6}	5{2,3,6}	45%
47	{2}	{3}	{7}	5{2,3,7}	70%
48	{2}	{3}	{8}	5{2,3,8}	68%
49	{3}	{1}	{1}	5{3,1,1}	49%
50	{3}	{1}	{2}	5{3,1,2}	78%
51	{3}	{1}	{3}	5{3,1,3}	23
52	{3}	{1}	{4}	5{3,1,4}	31%
53	{3}	{1}	{5}	5{3,1,5}	37%

Entry	Acyl Ketene	Chloro-Oxime	Amine	Product	Yield
54	{3}	{1}	{6}	5{3,1,6}	44%
55	{3}	{1}	{7}	5{3,1,7}	52%
56	{3}	{1}	{8}	5{3,1,8}	62%
57	{3}	{2}	{1}	5{3,2,1}	77%
58	{3}	{2}	{2}	5{3,2,2}	70%
59	{3}	{2}	{3}	5{3,2,3}	18%
60	{3}	{2}	{4}	5{3,2,4}	39%
61	{3}	{2}	{5}	5{3,2,5}	35%
62	{3}	{2}	{6}	5{3,2,6}	43%
63	{3}	{2}	{7}	5{3,2,7}	54%
64	{3}	{2}	{8}	5{3,2,8}	62%
65	{3}	{3}	{1}	5{3,3,1}	76%
66	{3}	{3}	{2}	5{3,3,2}	62%
67	{3}	{3}	{3}	5{3,3,3}	9%
68	{3}	{3}	{4}	5{3,3,4}	34%
69	{3}	{3}	{5}	5{3,3,5}	69%
70	{3}	{3}	{6}	5{3,3,6}	32%
71	{3}	{3}	{7}	5{3,3,7}	63%
72	{3}	{3}	{8}	5{3,3,8}	40%