Bacterial Adaptation of Respiration from Oxidic to Microoxic and Anoxic Conditions: Redox Control

Emilio Bueno,¹ Socorro Mesa,¹ Eulogio J. Bedmar,¹ David J. Richardson,² and Maria J. Delgado¹

Abstract

Under a shortage of oxygen, bacterial growth can be faced mainly by two ATP-generating mechanisms: (i) by synthesis of specific high-affinity terminal oxidases that allow bacteria to use traces of oxygen or (ii) by utilizing other substrates as final electron acceptors such as nitrate, which can be reduced to dinitrogen gas through denitrification or to ammonium. This bacterial respiratory shift from oxidic to microoxic and anoxic conditions requires a regulatory strategy which ensures that cells can sense and respond to changes in oxygen tension and to the availability of other electron acceptors. Bacteria can sense oxygen by direct interaction of this molecule with a membrane protein receptor (*e.g.*, FixL) or by interaction with a cytoplasmic transcriptional factor (*e.g.*, Fnr). A third type of oxygen perception is based on sensing changes in redox state of molecules within the cell. Redox-responsive regulatory systems (*e.g.*, ArcBA, RegBA/PrrBA, RoxSR, RegSR, ActSR, ResDE, and Rex) integrate the response to multiple signals (*e.g.*, ubiquinone, menaquinone, redox active cysteine, electron transport to terminal oxidases, and NAD/NADH) and activate or repress target genes to coordinate the adaptation of bacterial respiration from oxidic to anoxic conditions. Here, we provide a compilation of the current knowledge about proteins and regulatory networks involved in the redox control of the respiratory adaptation of different bacterial species to microoxic and anoxic environments. *Antioxid. Redox Signal.* 16, 819–852.

I. Introduction	820
II. Aerobic Respiration Under Microoxic Conditions	820
A. Oxygen control of <i>cbh₃</i> oxidase expression	822
1. Involvement of Fnr	823
2. Involvement of FixLJ/K	824
III. Redox Control of the Adaptation of Aerobic Respiration from Oxidic to Microoxic Conditions	825
A. <i>E. coli</i> ArcBA	825
B. <i>Rhodobacter</i> species RegBA/PrrBA	827
C. <i>Pseudomonas</i> species RoxSR	829
D. <i>Br. japonicum</i> RegSR	831
E. <i>Bacillus subtilis</i> ResDE and Rex	833
IV. Nitrate Respiration: Denitrification and Ammonification	834
A. Respiratory nitrate reductases	835
1. Membrane-bound respiratory nitrate reductase	835
2. Periplasmic nitrate reductase	835
B. Other enzymes in bacterial denitrification	835
1. Nitrite reductases	835
2. Nitric oxide reductases	837
3. Nitrous oxide reductase	837
C. Oxygen and nitrogen oxides control of nitrate respiration	837
1. Oxygen sensors	837
2. Nitrate and nitrite sensors	837
3. NO sensors	838
4. Other regulators in denitrifying bacteria	838

Reviewing Editors: Timothy Donahue, Lars Hederstedt, Woojun Park, and James Shapleigh

¹Estación Experimental del Zaidín, CSIC, Granada, Spain.

²School of Biological Sciences, University of East Anglia, Norwich, United Kingdom.

V. Redox Control of Nitrate Respiration	838
A. <i>Rho. sphaeroides</i>	839
B. <i>Ag. tumefaciens</i>	840
C. <i>Br. japonicum</i>	841
D. <i>Ba. subtilis</i>	842
VI. Concluding Remarks	843

I. Introduction

RESPIRATION is a fundamental process to all living cells in which electrons produced from oxidation of low-redox-potential electron donors such as NADH are transferred sequentially through a series of membrane-bound or membrane-associated protein carriers, the electron transport chain (ETC), which terminates in the reduction of the high-redox-potential electron acceptor, oxygen (Fig. 1a). The free energy released during this electron transfer process is used to drive the translocation of protons across the membrane to generate an electrochemical gradient that can be used for a variety of purposes, such as ATP synthesis and active transport (Fig. 1a). In contrast to the respiratory systems found in the mitochondria of many higher eukaryotic organisms, prokaryotic cells can induce branched-respiratory chains terminating in multiple oxidases with different affinities for the oxygen or use alternative electron acceptors, which contributes to their ability to colonize many microoxic and anoxic environments [reviewed in (61, 183, 186, 197)] (Fig. 1b).

The oxidation of organic molecules in cell respiration releases electrons that are transferred to membrane mobile quinones which constitute the link between the electron donating enzymes and the electron accepting enzymes (Fig. 1b). Quinones are small, freely diffusible, lipophilic, membrane-entrapped organic molecules that can carry two electrons and two protons when fully reduced, that is, in the quinol state [for a review, see (221)]. Different kinds of quinones have different electrochemical potentials, and many bacteria can synthesize more than one type of quinone. *Escherichia coli* synthesizes three types of quinones, a benzoquinone (UQ-8), and two naphthoquinones, menaquinone (MK) and demethylmenaquinone. Ubiquinone (UQ) predominates under aerobic conditions, and MK predominates under anaerobic conditions when the cellular state is more reduced (221). From the lipophilic hydroquinones, electrons can be carried to two different types of terminal oxidases: cytochrome *c* oxidases or quinol oxidases, where dioxygen is reduced to water (Fig. 1b). When oxygen is not present in the medium, electrons can be transmitted to alternative reductases that reduce substrates such as nitrate (NO_3^-), nitrite (NO_2^-), nitric oxide (NO), nitrous oxide (N_2O), dimethyl-sulphoxide (DMSO), trimethylamine N-oxide, sulfate, sulfite, and fumarate as final electron acceptors. Among them, nitrate is one of the essential environmental components in the biosphere. It serves as a nutrient for plants and microorganisms, and is used as an electron acceptor by many bacteria, archaea, and also by several eukaryotes (109, 125, 266). During anaerobic respiration, nitrate can be reduced to dinitrogen gas (N_2) or to ammonium. The anaerobic reduction of nitrate to N_2 gas is called denitrification, and it constitutes one of the more important processes in the N-cycle (141). This reductive process occurs in four stages, reduction

of NO_3^- to NO_2^- and *via* the gaseous intermediates NO and N_2O to N_2 . The enzymes involved in denitrification are nitrate-, nitrite-, nitric oxide-, and nitrous oxide reductase, encoded by *nar/nap*, *nir*, *nor*, and *nos* genes, respectively [reviewed in (125, 247, 248)]. In agriculture and wastewater treatment, denitrification by microorganisms is an important issue due to the economical, environmental, and public health implications of this process (10, 105).

The bacterial respiratory flexibility requires a regulatory strategy which ensures that prokaryotic cells sustain life under different environments in response to changing oxygen tension. To date, three main modes to sense O_2 have been described in bacteria: two types by direct interaction of O_2 with a membrane protein receptor (as in the heme-based sensor kinase FixL in rhizobia [reviewed in (83, 101, 205)]) or by interaction with a regulatory protein such as the Fe-S-based fumarate and nitrate reductase regulatory protein (Fnr) in *E. coli* [reviewed in (65, 68, 84, 53, 101, 261)]. In addition, a third type of O_2 perception is based on monitoring environmental oxygen concentration by sensing changes in the redox state of molecules or pools of molecules within the cell. These changes are detected by various protein sensors that convert the redox signals into regulatory output at the transcriptional or post-transcriptional level. Redox signals are many and diverse, and can involve redox-active cofactors such as heme, flavins, pyridine nucleotides, iron-sulfur clusters, and redox-sensitive amino-acids chains such as cysteine thiols among others [reviewed in (3, 14, 102)]. This article focuses on a comparative study about the regulatory mechanisms used by different bacterial groups to undergo respiratory shifts from oxic to microoxic and anoxic environments.

II. Aerobic Respiration Under Microoxic Conditions

The branched ETC present in many bacteria often contains several terminal oxidases (Fig. 1b), which can be grouped into two main types based on the substrates used as electron donors [reviewed in (61, 177, 183)]. Reduced *c*-type cytochromes donate electrons to cytochrome oxidases, whereas hydroquinones deliver electrons to quinol oxidases (Fig. 1b). Both types of oxidases belong to the extensively studied family of heme-copper oxidases (HCOs). The common denominator in HCOs is a membrane-integral subunit I that carries as cofactors a low-spin heme and a high-spin heme-copper binuclear center (Cu_B site) where reduction of O_2 to H_2O takes place (Fig. 2). Among this family, quinol oxidases possess a cofactor-free subunit II, whereas cytochrome *c* oxidases have cofactors bound to subunit II. In most cases, this is a binuclear Cu-Cu center (Cu_A site) that is liganded by six highly conserved amino acids (87, 176). The HCO superfamily is very versatile due to the wide range of electron donors they can use, their subunit composition, and the types of heme groups that they

FIG. 1. Respiratory chain(s) in mammalian mitochondrion and bacteria. (a) A summary of the topology and bioenergetics of a basic aerobic respiratory electron transport system of a mammalian mitochondrion is shown. This figure is adapted from ref. (197). **(b)** Schematic representation of aerobic and anaerobic nitrate respiration pathways in bacteria. MK, menaquinone; UQ, ubiquinone; Cyt, cytochrome; SDH, succinate dehydrogenase; NDH, NADH dehydrogenase; *cd*₁Nir, *cd*₁-type nitrite reductase; CuNir, Cu-type nitrite reductase; Nor, nitric oxide reductase; Nos, nitrous oxide reductase; Nar, membrane-bound nitrate reductase; Nap, periplasmic nitrate reductase; NrfA, cytochrome *c* nitrite reductase; UQH₂, ubihydroquinone; MKH₂, menahydroquinone.

contain. Studies of evolutionary relationships led to classifying the HCO enzymes in three different types (176, 177) (Fig. 2): (i) type A oxidases (oxidases *aa*₃), which are structurally and functionally close to mitochondrial oxidases, (ii) type B oxidases, grouped as cytochrome oxidases *bo*₃, and (iii) type C

oxidases, which include the cytochrome *c* oxidases *cbb*₃. To gain new insights into the oxygen respiration process, C-family HCOs from pathogenic bacteria have been recently structurally characterized, and it has been shown that C-HCOs differs from the two other HCO families, A and B (40).

FIG. 2. Summary of the main types of heme-copper oxygen reductases belonging to the A, B, and C families and the bd-type quinol oxidase.

The X-ray structure of the C-family *cbb₃* oxidase from *Pseudomonas stutzeri* shows an electron supply system different from families A and B.

Under O₂ limitation (microoxic conditions), many bacteria induce high-affinity oxidases to respire traces of molecular oxygen. The *cbb₃*-type cytochrome oxidase is known to have a high affinity for oxygen. Thus far, the enzyme from *Bradyrhizobium japonicum* is the only cytochrome *cbb₃* oxidase in which substrate affinity has been measured showing a K_m for dioxygen in the order of 7 nM (185). As a consequence, it allows respiration under microoxic conditions, such as those encountered by rhizobia in the legume nodule ([O₂] estimated to be 30 nM) (61). This oxidase has a subunit II (CcoO or FixO) that is a monoheme *c*-type cytochrome instead of the Cu_A-containing protein (67, 176). Subunit III in *cbb₃*-type oxidases, which is a diheme cytochrome *c* (CcoP or FixP), is thought to relay the electrons from the cytochrome *bc₁* complex via CcoO to the redox centers of subunit I (126, 265). Subunit III of all other HCOs is cofactor-free, just like the nonconserved small subunit IV (176). Homology modelling and recent studies on the thermodynamic properties of the redox centers in *cbb₃* oxidase from *Rhodobacter sphaeroides* using a combination of optical and electron paramagnetic resonance redox titrations have allowed the characterization of the *cbb₃* active site (191). Furthermore, it has been recently shown that the redox-coupled proton movement in the proximal cavity of *cbb₃*-enzymes contributes to the low redox potential of heme *b₃*, and suggests its potential implications for the high oxygen affinity of these enzymes (219).

In addition to C-family *cbb₃* oxidases, another family of high-affinity oxidases comprises cytochrome *bd*-type oxidases (29). Among them, the *bd* oxidase of *E. coli* is known to have a high affinity for oxygen showing the K_d(O₂) value of 0.28 μM (19). The *bd*-type oxidases are quinol oxidases that show no sequence homology to any subunit of heme-cooper family members. They do not contain any copper atoms in the catalytic center, but probably replace the Cu_B atom with a second heme (Fig. 2). In contrast to the heme-copper enzymes that are found at most branches of the tree of life, *bd*-type oxidases are only found in bacteria and archaea (183). The *bd*-type cytochromes do not pump protons and, therefore, have a lower

total energetic efficiency compared with the heme-copper type oxidases (29).

A. Oxygen control of *cbb₃* oxidase expression

As just mentioned, *cbb₃* oxidase is quite distinct from other bacterial HCOs in terms of its strategy for receiving electrons, the heme prosthetic group present in the active site, and its affinity for oxygen. Cytochrome *cbb₃* oxidases have been purified from several organisms, including *Paracoccus denitrificans*, *Rho. sphaeroides*, *Rhodobacter capsulatus*, and *Br. japonicum* [reviewed in (179)]. The biogenesis of this multisubunit enzyme, encoded by the *ccoNOQP* operon, depends on the *ccoGHIS* gene products, which are proposed to be specifically required for cofactor insertion and maturation of *cbb₃*-type cytochrome *c* oxidases (185). In the facultative photosynthetic model organism *Rho. capsulatus*, CcoN, CcoO, and CcoQ assemble first into an inactive 210 kDa sub-complex, which is stabilized via its interaction with CcoH and CcoS. Binding of CcoP, and probably subsequent dissociation of CcoH and CcoS, generates the active 230 kDa complex (126). Recent results have proposed that CcoH behaves more similar to a subunit of the *cbb₃* oxidase rather than a transient assembly factor *per se* (173). The insertion of the heme cofactors into the *c*-type cytochromes CcoP and CcoO precedes sub-complex formation, while the cofactor insertion into CcoN could occur either before or after the 210 kDa sub-complex formation during the assembly of the *cbb₃*-type oxidase (126). CcoQ is required for optimal *cbb₃*-type oxidase activity, because it stabilizes the interaction of CcoP with the CcoNO core complex, leading subsequently to the formation of the active 230-kDa *cbb₃*-type oxidase complex (178).

Genes encoding the *cbb₃* complex were initially isolated from rhizobia and named *fixNOQP* due to its requirement for symbiotic nitrogen fixation (185). Since then, orthologous genes called *ccoNOQP* were identified in other Proteobacteria including photosynthetic and pathogenic bacteria, suggesting that this oxidase is not only required for diazotrophs to sustain N₂ fixation, but also for the successful colonization of anoxic tissues by human pathogens [reviewed in (50, 61, 179)]. Regulation of *ccoNOQP* expression has been investigated in

many bacteria with oxygen being the key effector that controls its expression. To date, four different strategies that activate the *ccoNOQP* operon have been described (Fig. 3): (a) in *Rho. sphaeroides*, *Pa. denitrificans*, *Pseudomonas aeruginosa*, *Rhizobium leguminosarum* bv. *viciae* (UPM791), and *Rho. capsulatus*, Fnr acts as oxygen sensor and transcriptional factor; (b) in rhizobial species, such as *Sinorhizobium meliloti*, *Br. japonicum*, and *Azorhizobium caulinodans*, FixLJ/FixK is the system implicated in the activation of *ccoNOQP* under oxygen deprivation; (c) in *Rhi. leguminosarum* (VF39) and *Rhizobium etli* CFN42, a mixed regulation strategy operates that involves both Fnr and FixK proteins; (d) and finally, in the human pathogenic bacteria *Helicobacter pylori* and *Campylobacter jejuni* (ϵ -Proteobacteria), which lack *fnr* and *fixLJ-fixK* genes, it has been proposed that *cbb₃* may be constitutively expressed allowing the growth under microoxic conditions in its hosts. In addition to the positive control of Fnr on *fixNOQP* expression, it has been shown in *Rhi. etli* that two novel Fnr-type transcriptional regulators (StoRd and StoRf, symbiotic terminal oxidase regulators) have a negative control on microoxic expression of *fixNOQP* genes (100).

1. Involvement of Fnr. The O₂-sensitive protein Fnr belongs to the cyclic AMP (cAMP) receptor protein (Crp) superfamily of transcription factors that trigger physiological changes in response to a variety of metabolic and environmental cues [reviewed in (53, 65, 68, 84, 101, 103, 124, 261)]. All members of this family are predicted to be structurally related to Crp. They consist of four functionally distinct domains: (i) a sensor domain, (ii) a series of β -strands (β -roll structure) that form a loop-like structure making contact with the RNA polymerase holoenzyme (RNAP), (iii) a long α -helix acting as a dimerization interface, and (iv) a C-terminal helix-turn-helix motif (H-T-H) involved in DNA binding (Fig. 4a). The best-characterized Fnr protein is that of *E. coli*. In *E. coli*, Fnr acts as a direct oxygen sensor and is the primary transcriptional regulator of the switch between aerobic and anaerobic growth. Thus, under anoxic condi-

tions, Fnr is in its active state and is able to bind specific palindromic sequences of DNA (Fnr box). Once bound to DNA, Fnr activates target gene expression by recruiting RNAP or represses transcription by preventing the formation of productive RNAP:DNA complexes (53). Active Fnr regulates genes involved in the adaptation of *E. coli* for anaerobic growth and controls one of the best-studied gene regulatory networks in the cell. Very recently, the Fnr regulatory network has been reviewed by Tolla and Savageau (238). Generally, Fnr activates genes encoding products involved in anaerobic metabolism, such as the *nar* operon (nitrate reductase), the *dms* operon (DMSO reductase), and the *frd* operon (fumarate reductase), and represses genes encoding products involved in aerobic metabolism, such as the *sdh* operon (succinate dehydrogenase) and *ndh* (NADH dehydrogenase II). Fnr is activated under anoxic conditions by the acquisition of one [4Fe-4S]²⁺ cluster per monomer (designated 4Fe-Fnr). Each [4Fe-4S]²⁺ cluster is ligated by four cysteine residues (Cys²⁰, Cys²³, Cys²⁹, and Cys¹²²), and the presence of the cluster promotes Fnr dimerization, increasing its capacity to bind specifically to DNA (52, 54, 132) (Fig. 4a). The [4Fe-4S]²⁺ cluster is susceptible to be attacked by O₂, resulting in rapid and reversible conversion to a [2Fe-2S]²⁺ and finally to apoFnr on exposure to high oxygen concentrations. In addition to its reaction with oxygen, the Fnr [4Fe-4S]²⁺ cluster is also sensitive to NO. On exposure to NO, [4Fe-4S]²⁺ becomes nitrosylated, forming a combination of monomeric and dimeric dinitrosyl-iron-cysteine complexes, again abolishing its ability to bind DNA (53, 57). It has been reported that Fnr represses genes involved in NO detoxification (e.g., *hmp*, which encodes the flavohemoglobin) (182). When cultures of *E. coli* are exposed to NO, Fnr repression of *hmp* is relieved, suggesting that the reaction between Fnr and NO is physiologically significant. This suggestion is supported by transcript profiling experiments which reveal that the abundances of many Fnr-activated genes are lower, and many Fnr-repressed genes are higher, when *E. coli* is exposed to NO (119, 187).

FIG. 3. Regulation strategies of *ccoNOQP* gene transcription. Four regulation strategies have been described in several bacterial groups (see text for details) that involve (a) Fnr, (b) FixLJ-FixK, (c) Fnr and FixLJ-FixK, or (d) an unknown regulator. Gene transcriptional activation is indicated by arrows. Protein inactivation by O₂ is indicated by perpendicular lines. This figure is adapted from ref. (50).

FIG. 4. Functional domains and model of Fnr and FixLJ mediated oxygen activation. (a) Fnr senses oxygen *via* an iron-sulfur center, which is coordinated with four cysteine residues. Under anoxic conditions, Fnr is in its active state and it promotes DNA-binding by formation of a homodimer leading to activation of gene transcription. **(b)** FixL is an oxygen receptor in which oxygen binds directly to a heme group that is coordinated to a histidine residue within a PAS domain. Detection of low oxygen tension changes the conformation of the input PAS domain, rendering increased autophosphorylation activity of the transmitter domain in His residue and repressing the phosphatase activity of FixL. FixL activates FixJ transcriptional activity by transferring the phosphoryl group to the N-terminal domain of FixJ (+) gene transcription activation.

2. Involvement of FixLJ/K. In some rhizobial species, the *fixNOQP* operon is under the control of one Fnr-type transcriptional regulator named FixK, which is also able to recognize Fnr boxes but lacks $[4Fe-4S]^{2+}$ clusters. FixK is thereby unable to sense oxygen directly. In this case, it is the two-component regulatory system FixLJ that ensures oxygen sensing under low-oxygen conditions and activates the transcription of the *fixK* gene. This regulatory system FixLJ/FixK has been studied in rhizobial species where it controls the expression of genes for microoxic respiration, as well as various other functions, in response to low oxygen (28, 56, 83, 145, 160). Interestingly, it has been shown that *Rhodospseudomonas palustris* FixK is not only required for optimal microaerobic growth, but is also required for phototrophic and autotrophic growth, and for growth on aromatic compounds (194). These authors also propose that FixK helps the control of the transition of *R. palustris* to anaerobic growth by activating the expression of the oxygen-sensing Fnr homolog, AadR.

The biochemistry of the FixLJ two-component system has been characterized in *Si. meliloti* and *Br. japonicum* (96, 211, 223). FixL is a histidine kinase that binds heme at an amino-terminal PAS domain (Fig. 4b). PAS domains are present in many proteins involved in the sensing of oxygen, redox, or light. They were first found in eukaryotes, and were named after homology to the *Drosophila* period protein (PER), the aryl hydrocarbon receptor nuclear translocator protein (ARNT), and the *Drosophila* single-minded protein (SIM). The binding of oxygen to heme inactivates FixL kinase activity. FixL, and the response regulator and transcription factor FixJ form a complex in which FixJ allosterically stimulates autophosphorylation of FixL. At very low oxygen concentrations, FixL

transphosphorylates FixJ, which, in turn, activates expression of the transcription factor gene *fixK* (Fig. 4b). In *Br. japonicum*, the nitrogen-fixing root-nodule endosymbiont of soybean (*Glycine max*), a FixK-like protein called FixK₂, acts as the key distributor of the low-oxygen signal perceived at the level of the FixLJ two-component regulatory system. FixK₂ is one of the 16 (Crp/Fnr)-type transcriptional regulators whose genes are present in the *Br. japonicum* genome [reviewed in (146)]. FixK₂ recognizes a palindromic sequence motif (TTG-N₈-CAA, termed the FixK₂ box) that is located around position-41 upstream of the transcription start site in the regulated promoters. Microoxically induced targets of FixK₂ include the operons *fixNOQP* and *fixGHIS*, both essential for microoxic respiration, several heme biosynthesis genes (*hemA*, *hemB*, *hemN₁*, and *hemN₂*), denitrification genes, and some hydrogen oxidation genes (*hup* genes) (16, 160). Recently, the *Br. japonicum* FixK₂ regulon was unraveled by using a transcriptomics approach (145). DNA binding site predictions, together with a FixK₂-dependent *in vitro* transcription assay, demonstrated that the *fixNOQP* operon is a direct target for FixK₂. The latter studies, carried out with purified FixK₂ protein, showed that FixK₂ is sufficient to activate transcription *in vitro* without any identified effector (148). This is puzzling in view of the fact that all Crp/Fnr-like proteins can be positively or negatively modulated in their activity through bound cofactors or intrinsic, reactive amino acids (101). However, recent findings have reported that post-translational control occurs at FixK₂, whereby a critical cysteine at position 183 in the polypeptide chain is a target for oxidation. This provides a second, important means of affecting FixK₂ activity, in addition to the regulation of its expression by FixLJ (147).

III. Redox Control of the Adaptation of Aerobic Respiration from Oxic to Microoxic Conditions

A. *E. coli* ArcBA

E. coli has three terminal quinol oxidases, of which one is a heme-copper bo_3 oxidase and two are *bd*-type oxidases [reviewed in (183, 186)]. The quinol bo_3 oxidase contains a heme-Cu cofactor (Cu_B) and is essential when oxygen is present at high concentrations. The protein complex is encoded by the *cyoABCDE* operon (186). Under conditions of low aeration, the high-affinity *bd* oxidase is the preferred terminal reductase in *E. coli* (29). The structural components are encoded by the genes *cydAB*, which form a heterodimer. CydC and D are encoded in the same operon as CydA and B and thought to form a heme transporter necessary for cytochrome *bd* oxidase biogenesis (29). There is a third terminal oxidase present in *E. coli* encoded by the genes *cyxAB*. It is a *bd* type oxidase but its function is unclear. It appears to be expressed under microoxic conditions (29).

Cytochrome *bd* oxidase prevails under oxygen-limiting conditions. Consistent with these properties, transcription of the *cydAB* operon is activated when oxygen becomes limiting. Under fully anoxic conditions, *cydAB* expression is subsequently repressed to an intermediate level relative to microoxic and oxygen-rich conditions (241). It is now solidly established that this oxygen control is achieved through the combined action of the oxygen sensing transcription factor Fnr and the aerobic respiration control (Arc)B/ArcA two-component system (217). Under anoxic conditions, Fnr acts as a repressor of the *cydAB* operon (Fig. 5). ArcBA is a two-component regulatory system involved in the transcriptional regulatory network that allows facultative aerobic bacteria to sense and respond to various respiratory conditions. While the Fnr protein is directly implicated in sensing oxygen, the ArcBA system activity is proposed to be modulated by the redox state of the quinones in the membrane (137, 138). The ArcBA system comprises the ArcB protein, a tripartite membrane-bound sensor kinase, and the cognate response regulator ArcA (Fig. 6a). ArcB belongs to the subclass of hybrid sensor kinases that also includes BarA, BvgS, EvgS, LemA (GacS), RteA, and TorS (2, 138). Under limiting oxygen conditions, ArcB autophosphorylates residue His²⁹² and, through a phosphorelay system involving Asp⁵⁷⁶ and His⁷¹⁷,

transphosphorylates ArcA (Fig. 6a). It has been previously reported that histidine kinases act as homodimers and that they autophosphorylate by an intermolecular reaction (229). That is, the gamma-phosphoryl group of ATP, which is bound to one monomer in the homodimer, is transferred to the other monomer. On the contrary, it has been recently proposed that ArcB autophosphorylates through an intramolecular and not through an intermolecular reaction as previously proposed (175).

On a shift to aerobic conditions, ArcB can form intermolecular disulphide bonds *via* Cys¹⁸⁰ and Cys²⁴¹ located in the PAS domain (Fig. 6a). The kinase activity of ArcB is highly dependent on this covalent linkage (137). The regulatory mechanisms that underlie the function of the ArcBA two-component system have been the subject of numerous studies. ArcB does not sense oxygen directly but is thought to sense the redox state of the UQ-ubiquinol pool in the aerobic respiratory chain, because UQ inhibits ArcB autokinase activity (137). However, this regulatory network becomes complex, as recent observations could not demonstrate a correlation between ArcB activation and oxygen availability (18). Bekker and colleagues (18) proposed that the *in vivo* activity of ArcB in *E. coli* is also modulated by the redox state of the MK pool and that the UQ/ubiquinol ratio is not the only determinant of ArcB activity (Fig. 6b). This combined regulation by the redox state of the UQ pool and MK pool provides a mechanism to explain the observed complex regulation of ArcB activation at variable rates of oxygen supply (Fig. 6b). Menaquinols would be the dominant activators under anoxic conditions, where the size of the UQ pool is approximately five times less than the size of the MK pool (18). Recent work has used controlled chemostat cultures subject to gradually decrease oxygen concentration to study the effects of oxygen availability on the transcriptome, redox state of the UQ-ubiquinol, outer membrane protein profiles, and terminal oxidase proteins of *E. coli* K-12 (208). A probabilistic model was used to predict the activity of ArcBA across the aerobiosis range. The model implied that the activity of the regulator ArcA correlated with aerobiosis, but not with the redox state of the UQ pool. Rolfe and colleagues (208) have proposed that, at least under the conditions used in this work, the rate of fermentation product synthesis (*q*acetate) exerts a greater influence on ArcA activity than the redox state of the quinone

FIG. 5. Regulation of *bd* and bo_3 terminal oxidase expression in *Escherichia coli*. Fnr and ArcB provide negative or positive transcriptional control in response to oxygen availability. The genes *cydAB* and *cyoABCDE* encode *bd* and bo_3 oxidase polypeptides, respectively. Positive regulation (+) is denoted by arrows, and negative regulation (–) is indicated by perpendicular lines.

FIG. 6. Functional domains and model of ArcBA-mediated redox control. (a) ArcB is attached to the membrane by two transmembrane regions (TM). A linker region containing a putative leucine-zipper and a PAS domain connects TM2 with the catalytic domains. The two cysteine-residues (Cys¹⁸⁰ and Cys²⁴¹) in the linker region are able to undergo oxidation and to form intermolecular disulfide bonds between both ArcB monomers. The transmitter domain (H1) contains the conserved His²⁹² together with the G1 and G2 nucleotide-binding motifs. The receiver domain (D1) harbors the conserved Asp⁵⁷⁶, and the histidine phosphotransfer domain (Hpt/H2) contains the conserved His⁷¹⁷. The ArcA component is represented with its N-terminal receiver domain carrying the conserved Asp⁵⁴ and its C-terminal H-T-H domain. This figure is adapted from ref. (138). (b) MKH₂, which predominates under anaerobic conditions, would be oxidized in the presence of low levels of oxygen (~20% aerobiosis), leading to increased levels of MKs and inactivation of ArcB kinase activity. UQH₂ predominate under microoxic conditions (~80% aerobiosis), giving rise to activation of ArcB kinase activity. In fully aerobic conditions (100%), the UQH₂ pool is subjected to oxidation, which results in increasing levels of UQs and oxidation of the key cysteines, leading to inactivation of the ArcB kinase activity. This figure is adapted from ref. (18).

pools. It is perhaps not surprising that, as a regulator of genes that have fundamental roles in different processes of the bacterial metabolism (aerobic respiration, anaerobic respiration, and fermentation), ArcB should integrate the response to multiple signals (e.g., UQ, MK, acetate, and other fermentation products) in response to different growth conditions.

Phosphorylated ArcA activates genes or operons needed to use traces of oxygen in the medium, for example, those for terminal oxidase *bd*, and represses expression of the aerobic cytochrome *bo*₃ terminal oxidase (136). Regulation of *bd* and cytochrome *bo*₃ terminal oxidases in response to oxygen concentration variation is also controlled by Fnr as just described (Fig. 5b). Fnr inhibits *cyd* and *cyo* gene activation from 0% to 10% oxygen tension. However, ArcA represses *cyo* expression and activates *cyd* transcription when O₂ concentration is between 10% and 20% (241) (Fig. 5). These authors proposed that when the oxygen tension drops, ArcA is phosphorylated by ArcB and activates *cydAB* transcription reaching a maximum at <2% O₂ tension. When oxygen is decreased further, Fnr becomes active and represses *cydAB* transcription. Fnr repression of *cydAB* requires the presence of a functional ArcA protein (51, 99). According to this observation, Fnr has been proposed to serve as an antiactivator by counteracting ArcA-mediated activation, rather than directly repressing transcription. This indirect regulation of *cydAB* repression by Fnr through ArcA is supported by gene expression profiling analyses where nearly two-thirds of the genes for which expression is affected by Fnr are also affected by ArcA (209, 210). In addition to controlling genes encoding respiratory oxidases, ArcA~P represses the expression of genes encoding dehydrogenases of the tricarboxylic acid cycle and enzymes of the glyoxylate shunt. Recent studies have demonstrated that ArcAB exerts an important influence on carbon and redox balances in *E. coli* growing under carbon-limited conditions with a restricted oxygen supply (163).

B. *Rhodobacter* species RegBA/PrrBA

Photosynthetic bacteria such as *Rho. Capsulatus* and *Rho. sphaeroides* conserve energy under oxic or microoxic condi-

tions by driving electrons through branched electron transfer chains to terminal oxidases with different affinity for oxygen (Fig. 7). One ubiquinol oxidase is found in *Rho. capsulatus* (*cydAB*), while two are found in *Rho. sphaeroides* (*qoxBA* and *qxtAB*). The cytochrome *c* oxidases depend on the cytochrome *bc*₁ complex (*petABC* in *Rho. capsulatus* and *fbfABC* in *Rho. sphaeroides*), and cytochromes *c*₂ (*cycA*) or *c*_y (*cycY*). While *Rho. capsulatus* contains only one *cbb*₃-type oxidase (*ccoNOQP*), an *aa*₃- (*ctaDC*) and a *cbb*₃-type oxidase (*ccoNOQP*) are found in *Rho. sphaeroides* [reviewed in (72, 120, 166, 233, 258)]. Gene expression experiments using the *lacZ* as reporter gene revealed that the *cbb*₃ oxidase from *Rho. capsulatus* and *aa*₃ oxidase from *Rho. sphaeroides* are considered low-oxygen-affinity oxidases which are induced under oxic conditions (233). However, the *Rho. capsulatus* ubiquinol *bd* oxidase and *Rho. sphaeroides* *cbb*₃ oxidase are classified as high-affinity terminal oxidases and are highly induced under oxygen-limiting conditions (233).

RegBA/PrrBA are members of the family of the two-component regulatory systems present in a large number of Proteobacteria. These proteins are named RegBA in *Rho. capsulatus*, *Rhodovulum sulfidophilum*, and *Roseobacter denitrificans* [reviewed in (72, 258)], PrrBA in *Rho. sphaeroides* (106, 167), ActSR in *Si. meliloti* (73) and *Agrobacterium tumefaciens* (8), RegSR in *Br. japonicum* (15), and RoxSR in *Ps. aeruginosa* (46). In *Rhodobacter* species, the RegBA/PrrBA regulon encodes proteins involved in numerous energy-generating and energy-utilizing processes such as photosynthesis, carbon fixation, nitrogen fixation, hydrogen utilization, aerobic and anaerobic respiration, and aerotaxis [reviewed in (72, 233, 258)]. Recent analyses of the transcriptome and the proteome of *Rho. sphaeroides* *prrA* mutant revealed that, in addition to the numerous PrrA gene targets already known, genes encoding proteins whose functions are involved in intermediary metabolism, repair of DNA and protein damage, cell motility and secretion, and translation constituted new targets for PrrA regulation (77).

The RegBA/PrrBA two-component systems comprise the membrane-associated RegB/PrrB histidine protein kinase, sensing changes in redox state and its cognate PrrA/RegA

response regulator. Under conditions where the redox state of the cell is altered due to generation of an excess of reducing potential, produced by either an increase in the input of reductants into the system (*e.g.*, presence of reduced carbon source) or a shortage of the terminal respiratory electron acceptor (*e.g.*, oxygen deprivation), the kinase activity of RegB/PrrB is stimulated relative to its phosphatase activity (72). This increases phosphorylation of the partner response regulators RegA/PrrA, which are transcription factors that bind DNA and activate or repress gene expression. The membrane-bound sensor kinase proteins RegB/PrrB contain an H-box site of autophosphorylation (His²²⁵), a highly conserved quinone binding site (the heptapeptide consensus sequence GGXXNPF, which is totally conserved among all known RegB homologues), and a conserved redox-active cysteine (Cys²⁶⁵, located in a “redox box”). The mechanism by which RegB controls kinase activity in response to redox changes has been an active area of investigation. A previous study demonstrated that RegB Cys²⁶⁵ is partially responsible for redox control of kinase activity. Under oxidizing growth conditions, Cys²⁶⁵ can form an intermolecular disulfide bond to convert active RegB dimers into inactive tetramers (235) (Fig. 8a). The

highly conserved sequence, GGXXNPF, located in a short periplasmic loop of the RegB transmembrane domain has also been implicated in redox sensing by interacting with the UQ pool [Swem *et al.* (234)] (Fig. 8a). Recently, kinase activity assays together with isothermal titration calorimetry (ITC) measurements indicated that RegB with a substitution in the cytosolic cysteine by serine in position 265 (RegB C²⁶⁵S) binds both oxidized and reduced UQ with almost equal affinity. However, only the oxidized UQ inhibits RegB kinase activity (259). The observation that the RegB C²⁶⁵S mutant is still redox responsive suggests that UQ binding is a signal input able of functioning independently from Cys²⁶⁵. However, the contribution of each redox sensing inputs is unknown.

In *Rho. sphaeroides*, the PrrB histidine kinase is a bifunctional enzyme that possesses both kinase and phosphatase activities (168). Several reports proposed that the *cbb*₃ oxidase transduced an inhibitory signal to the PrrBA under oxic conditions to prevent gene expression (165, 166). The dual function of the *cbb*₃ oxidase as both terminal oxidase and O₂/redox sensor and modulator of PrrB kinase/phosphatase activity represents a new model of redox sensing. In this model (Fig. 8b), the UQ binding site within the PrrB

a *Rhodobacter capsulatus*

b *Rhodobacter sphaeroides*

FIG. 8. Proposed models for redox-mediated sensing by RegB/PrrB. (a) In *Rho. capsulatus*, the conserved cysteine C²⁶⁵ of RegB functions as a redox switch that controls RegB kinase activity through a metal-dependent formation of a disulfide bond in response to redox changes (235). Binding of UQ to the RegB transmembrane domain has also been proposed to inhibit RegB kinase activity, whereas UQH₂ does not affect RegB kinase activity (234). (b) In *Rho. sphaeroides*, the *cbb*₃ oxidase generates an “inhibitory” signal that can directly stimulates the phosphatase activity of PrrB. Under low oxygen conditions when the electron flow is reduced, the inhibitory signal from *cbb*₃ is weakened, and PrrB would retain its kinase activity (165). PrrC has been proposed as a signal mediator between the *cbb*₃ oxidase and the sensor kinases PrrB (7).

transmembrane domain is not required for monitoring the PrrB kinase activity. Instead, a control based in direct interaction between components of the terminal oxidase *cbb₃* and PrrB is strengthened (122).

The photosynthetic regulatory response protein (PrrC) is a Sco homolog present in *Rho. sphaeroides* (75). Sco is thought to be involved in donating copper to the Cu_A center and, thus, it has a central role in cytochrome oxidase synthesis (12). *Rho. sphaeroides* PrrC, which reduces Cu²⁺ to Cu⁺, and possesses disulfide reductase activity, is required for the correct functioning of the sensor kinase/phosphatase PrrB (7). Similarly, the *Rho. capsulatus* SenC protein, homologous to PrrC, which is required for synthesis of a functional cytochrome *c* oxidase (232), might act as a signal mediator between the Q-pool and the sensor kinase RegB. However, at present, there is no direct evidence that SenC or *cbb₃* oxidase directly modulates the activity of the RegBA regulatory system.

RegA/PrrA contain conserved domains that are typical in two component response regulators such as a phosphate accepting aspartate, an "acid box" containing two highly conserved aspartate residues and a H-T-H DNA-binding motif (72). The phosphorylated form of RegA/PrrA has increased DNA binding capacity (129, 189). Under oxidizing conditions, RegB/PrrB shifts the relative equilibrium from the kinase to the phosphatase mode resulting in a dephosphorylated inactive RegA/PrrA form. Despite this evidence, it has been reported that inactivation of the *regA* gene affects expression of many different genes under oxidizing (aerobic) conditions, suggesting that both phosphorylated and unphosphorylated RegA/PrrA may be active transcriptional regulators (72, 233). In this context, it has been shown that both phosphorylated and unphosphorylated forms of RegA/PrrA are capable of binding DNA *in vitro* and activating transcription (189). Based on recent findings, there are two types of PrrA binding sites within the promoter of *Rho. sphaeroides hemA*, which codes for one of two isoenzymes catalyzing 5-aminolevulinate synthesis: (i) one site for which unphosphorylated PrrA has greater affinity, and (ii) another for which phosphorylated PrrA has greater affinity (190). Several consensus DNA binding sequences have been previously reported for PrrA (128, 139) and its homologs RegA in *Rho. capsulatus* (233, 253) and RegR in *Br. japonicum* (73, 134). While there is a measure of agreement in terms of the assignment of two imperfect quasi-symmetrical inverted repeats, or half-sites separated by a spacer region possessing variable length, the DNA sequence *per se* shows a high degree of degeneracy. Very recently, it has been demonstrated that PrrA can bind *in vitro* to DNA sequences with different lengths in the spacer regions between the half sites (76).

In *Rho. capsulatus*, RegA has been reported to activate the expression of both the *bd* and cytochrome *cbb₃* oxidases (Fig. 9a). An interesting paradox is that RegA activates expression of cytochrome *cbb₃* oxidase under oxic conditions, but represses expression of this oxidase under anoxic conditions (233). In many respects, this is similar to what occurs in *E. coli* where ArcA acts as an oxic activator, as well as an anoxic repressor of cytochrome *bo₃* oxidase (see section III.A; Fig. 5b).

Terminal oxidase expression in *Rho. capsulatus* involves a complex set of regulators beyond that of RegBA. In response to oxygen availability, these regulators provide negative or positive transcriptional control to coordinate enzyme synthesis for optimal growth. A regulatory scheme for control of

terminal oxidases in response to oxygen availability is shown in Figure 9a. Specifically, in *Rho. capsulatus* cytochrome *cbb₃*, oxidase expression is regulated by RegA and FnrL, as well as moderately regulated by HvrA, an activator of various photosynthetic components under low-light conditions [(Swem *et al.* (231)]. Ubiquinol *bd* oxidase expression was found to be regulated by RegA and HvrA as well as by AerR and CrtJ, which are aerobic repressors of photosystem gene expression (231). In *Rho. sphaeroides*, microarray analyses and quantitative reverse transcriptase-polymerase chain reaction showed that both PrrA and the repressor of genes involved in photosynthesis, PspR, affect the transcription of the *ccoNOQP* operon (Fig. 9b). The absence of PrrA prevented the expression of the *cco* genes, and a *prrA-pspR* double mutant leads to derepression of *ccoNOQP* operon (34). The fact that PspR binds directly the *cco* promoter indicates that PrrA/PspR could act as an antirepressor/repressor system in which PspR acts as the direct repressor of the *cco* operon and PrrA is the PspR antirepressor regulator (34). Thus, RegBA/PrrBA appears to be just one component of a more complex regulatory network that controls many cellular processes. In this context, recent comparative genomics analyses and characterization of *Rho. sphaeroides* FnrL regulon have revealed that *ccoNOQP* is also positively controlled by this protein (68).

C. *Pseudomonas* species *RoxSR*

Pseudomonads are opportunistic pathogens that inhabit a wide variety of environments, including soil, water, and animal, human, and plant roots. They are endowed with versatile respiratory chains that can be adapted to changing oxygen concentrations. Although they preferentially obtain energy from aerobic respiration, some of them, such as *Ps. aeruginosa*, can also grow anaerobically using nitrate as a final electron acceptor, or can even ferment arginine and pyruvate (254). The genome sequence and some biochemical studies have revealed that *Pseudomonads* have at least five terminal oxidases for aerobic respiration [reviewed in (183)] (Fig. 10a). In *Ps. aeruginosa*, UQs can be oxidized by either a *bo₃*-type oxidase, a cyanide-insensitive oxidase (named CIO), or a *bc₁* complex that transfers electrons to a cytochrome *c*, which, in turn, can be oxidized by an *aa₃* oxidase or by two related *cbb₃* oxidases, named *cbb₃-1* and *cbb₃-2* (46, 47, 49). The *Pseudomonas putida* genome contains genes coding for all these terminal oxidases (161). Each oxidase is expected to have a specific affinity for oxygen, efficiency of proton translocation, and redox potential. *Ps. aeruginosa* has three predicted high-affinity oxidases; CIO, *cbb₃-1*, and *cbb₃-2*, which are believed to be important in the adaptation to oxygen-limiting conditions during growth under microoxic conditions. In this context, Alvarez-Ortega and Harwood (1) reported that any one of CIO, *cbb₃-1* and *cbb₃-2*, could support growth under 2% oxygen concentrations and that either of *cbb₃-1* and *cbb₃-2* was indispensable for the growth under 0.4% oxygen.

In *Ps. aeruginosa*, the Anr global transcriptional regulator plays a pivotal role in adaptation to microoxic or anoxic conditions. Anr is a homolog of *E. coli* Fnr and regulates expression of the enzymes required for nitrate respiration (4, 213), and of at least some of the aerobic terminal oxidases, such as the CIO oxidase and cytochrome *cbb₃-2* oxidase (47, 49). Recently, it has been shown that under low-oxygen conditions, Anr is involved in the repression of the CIO genes and

a *Rhodobacter capsulatus*

b *Rhodobacter sphaeroides*

FIG. 9. Regulation of *bd* and *cbb₃* terminal oxidases expression in *Rho. capsulatus* (a) and *Rho. sphaeroides* (b). In *Rho. capsulatus*, RegA regulates gene transcription in response to both oxic and anoxic conditions, FnrL, HvrA, in response to anaerobiosis, and AerR and CrtJ in response to aerobiosis. In *Rho. sphaeroides*, FnrL, PrrA, and PspR regulate gene transcription in response to anoxic conditions. Genes *cydAB* and *ccoNOQP* encode *bd* and cytochrome *cbb₃* oxidase polypeptides, respectively. Positive regulation (+) is denoted by arrows, and negative regulation (–) is indicated by perpendicular lines.

induction of the *cbb₃-2* genes, while the other three terminal oxidases are not significantly regulated by Anr (Fig. 10b) (121). Similarly, in *Ps. putida*, it has been shown that, under oxygen limitation, inactivation of *anr* led to an increased expression of the *bo₃*-type and the CIO terminal oxidases, and to a much lower expression of *cbb₃-1* (equivalent to *Ps. aeruginosa cbb₃-2*), suggesting that Anr is a transcriptional activator of *cbb₃-1* and a repressor of CIO genes (244).

In addition to Anr, another redox-responsive regulatory system has been proposed to be important for the regulation of terminal oxidases in *Pseudomonas* species, the RoxSR system. RoxSR belongs to the family of RegBA and PrrBA regulatory systems. It has been postulated in *Ps. aeruginosa* that the activation signal perceived by RoxS might depend on electron transport signals emerging from the oxidase *cbb₃-1* (47). Microarray analyses performed in *Ps. putida* (152) showed that a *cyo* mutation leads to a significant change in the transcriptome profile. They showed that the absence of *bo₃*-type oxidase was compensated by upregulation of CIO and *cbb₃-1* (corresponding to *cbb₃-2* of *Ps. aeruginosa*). A regulatory

system monitoring the electron flow through the *bo₃*-type oxidase, which is similar to PrrBA activity modulation by the *cbb₃* oxidase, might be operative in *Ps. putida*.

Microarray studies have reported that RoxSR is involved in the regulation of the terminal oxidases genes in *Ps. putida* (78) and *Ps. aeruginosa* (121). In *Ps. aeruginosa*, RoxSR functions as a positive regulator of the putative high-affinity oxidases CIO, *cbb₃-1* and *cbb₃-2*, and at the stationary phase where the dissolved oxygen concentration is low due to the high cell density, as a negative regulator of the putative low-affinity oxidase *aa₃* (121) (Fig. 10b). The RoxSR regulon of *Ps. putida* includes genes for respiratory function and maintenance of the redox balance, genes involved in sugar and amino acid metabolism, and the sulfur starvation response (78). These authors also showed that RoxSR participates in cell density signal transduction in *Ps. putida*. In *Ps. aeruginosa*, however, only a few genes related to sugar and amino-acid metabolism were identified as the members of the RoxSR regulon (121). The number of the putative RoxSR-regulated genes in *Ps. aeruginosa* seemed to be low compared with that in *Ps. putida*.

FIG. 10. Aerobic respiratory chains (a) and regulation of CIO, *cbb*₃, and *aa*₃ terminal oxidase expression (b) in *Pseudomonas aeruginosa*. In (a), the quinone-reducing (left) and quinol-oxidizing (right) branches with terminal oxidases induced under oxic conditions (gray boxes), and microoxic conditions (white boxes), are shown. In (b), Anr and RoxSR regulators provide negative or positive transcriptional control, in response to oxygen availability. Genes *cox*, *cioAB*, and *ccoNOPQ* encode *aa*₃, CIO, and *cbb*₃ oxidase polypeptides, respectively. Positive regulation (+) is denoted by arrows, and negative regulation (−) is indicated by perpendicular lines. CIO, cyanide insensitive oxidase.

These differences are presumably due to the different growth conditions used for the microarray analysis in the two *Pseudomonas* species.

D. *Br. japonicum* *RegSR*

The genera *Allorhizobium*, *Azorhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Rhizobium*, and *Sinorhizobium* (*Ensifer*), collectively referred to as rhizobia, are members, along with other genera, of the bacterial order *Rhizobiales* of the α -Proteobacteria. Rhizobia are soil bacteria with the unique ability to establish a N₂-fixing symbiosis on legume roots and on the stems of some aquatic legumes. During this interaction, bacteroids, as rhizobia are called in the symbiotic state, are contained in intracellular compartments called symbiosomes within a specialized organ, the nodule, where they fix N₂ [reviewed in (116, 169)]. In the nodule, maintenance of nitrogenase activity is subject to a delicate equilibrium. A high rate of O₂ respiration is necessary to supply the energy demands of the N₂-reduction process; but, on the other hand, O₂ also irreversibly inactivates the nitrogenase complex. In order

to keep the steady-state concentration of free-O₂ low, the cortex of nodules acts as a diffusion barrier, which greatly limits permeability to O₂ [reviewed in (150)]. Oxygen is delivered to the symbiosomes by the plant O₂-carrier leghemoglobin (Lb), which transports O₂ at a low, but stable, concentration, allowing for the simultaneous operation of nitrogenase activity and bacteroid respiration [(66) and references therein]. N₂-fixing bacteroids deal with the low levels of free O₂ by inducing a high-affinity cytochrome *cbb*₃-type oxidase [reviewed in (61)].

Similar to many other anaerobic facultative bacteria, *Br. japonicum* adapts to different environmental O₂ concentrations by inducing multiple terminal oxidases with different affinities for O₂. *Br. japonicum* has eight terminal oxidases, of which two are *bd*-type oxidases, and six are heme-copper oxidases, the latter being further divided into two quinol oxidases and four cytochrome oxidases [reviewed in (38)] (Fig. 11). Of particular interest is the *fixNOPQ*-encoded *cbb*₃-type oxidase, because it supports *Br. japonicum* aerobic respiration under free-living microoxic conditions and enables endosymbiotic *Br. japonicum* cells (bacteroids) to support the

FIG. 11. Aerobic respiratory chains in *Bradyrhizobium japonicum*. The quinone-reducing (left) and quinol-oxidizing (right) branches with terminal oxidases are shown. The *coxBACF*-encoded cytochrome *aa*₃ is the predominant heme-copper oxidase for aerobic growth (gray box), and the *fixNOQP*-encoded *cbb*₃-type oxidase (white box) supports respiration under free-living microoxic and under symbiotic conditions. ^aGene number according to RhizoBase (<http://genome.kazusa.or.jp/rhizobase>).

nitrogen fixation process. Cytochrome *cbb*₃ oxidase has been purified from *Br. japonicum* (185), where the experimentally determined *K_m* for O₂ is in the order of 7 nM, a value that is consistent with its function in the bacteroid.

The contribution of *Br. japonicum* CoxG and ScoI to the biogenesis of the *cbb*₃ oxidase has been recently investigated (38). These proteins are the respective paralogs of the mitochondrial chaperones Cox11 and Sco1 involved in the formation of Cu_B and Cu_A centers of cytochrome *aa*₃ (12). Analyses of *Br. japonicum* *coxG* and *scoI* mutants revealed that disparate pathways are used for *aa*₃- and *cbb*₃-type oxidases biogenesis [(Bühler *et al.* (38))].

Oxygen concentration is the primary effector of *fixNOQP* expression in *Br. japonicum*. Under O₂ limitation, the FixJ protein of the FixLJ system is phosphorylated by the oxygen-inhibitable, heme-based sensor kinase FixL (see section II.A Fig. 4b). The only known target of FixJ in *Br. japonicum* is *fixK₂*, whose product, FixK₂, has been shown to activate genes involved in anoxic or microoxic energy metabolism, including the *fixNOQP* operon (161). By using a transcriptomics approach (145), and FixK₂-dependent *in vitro* transcription assay (148), the involvement of FixK₂ as a direct transcriptional regulator of *cbb*₃ has been demonstrated.

RegSR are members of the family of two-component regulatory redox-responsive proteins. In *Br. japonicum*, RegSR induces expression of the *fixR-nifA* operon, which encodes the NifA regulatory protein (15) (see Fig. 15). Targets of NifA include *nif* and *fix* genes, which are directly involved in nitrogen fixation, and also genes that are indirectly related to nitrogen fixation or have a unknown function in this process (83, 108, 162). Although *fixNOQP* genes are involved in nitrogen fixation, these genes were not identified as targets of NifA in microarray-based experiments (108). Transcription of *fixR-nifA* is dependent on NifA and on RegSR, respectively (Fig. 15). RegR induces expression of the *fixR-nifA* operon in both oxic and anoxic conditions, suggesting that RegSR responds to the overall redox state of the cell rather than to oxygen directly (15). *Br. japonicum* RegS possesses a highly conserved quinone binding site and a conserved redox-active cysteine, which suggests that the redox state of the membrane-localized quinone pool or the redox-active cysteine might be involved in redox sensing in *Br. japonicum*. However, the precise nature of the signal that is transduced by the *Br. japonicum* RegSR is unknown.

Null mutations in the *regR* gene led to a dramatically decreased nitrogen fixation activity (15). Surprisingly, the phenotypic properties of *regS* mutants were largely indistinguishable from the wild type. Although it has been demonstrated, *in vitro*, that RegS is both a functional autokinase and a kinase (74), the *in vivo* role of RegS in RegR-mediated transcriptional activation is still unclear. In addition to *fixR-nifA*, a large number of novel members of the RegR regulon have been identified by transcriptomic analyses (134). Among them, a putative operon that encodes a predicted multidrug efflux system required for an efficient symbiosis specifically with soybean has been recently characterized (133).

In addition to oxygen respiration, *Br. japonicum* is able to obtain energy and grow from nitrate reduction to N₂ through denitrification when cultured under oxygen-limiting conditions with nitrate as the terminal electron acceptor. In *Br. japonicum*, denitrification reactions depend on *napEDABC*, *nirK*, *norCBQD*, and *nosRZDYFLX* genes encoding the nitrate-, nitrite-, nitric oxide-, and nitrous oxide reductases, respectively (see section V.C). Inactivation of *Br. japonicum* *cycA* or *napC*, which encode cytochromes *c* involved in the electron transfer through denitrification pathway, decreases the expression of *Br. japonicum* *cbb*₃ oxidase during nitrate-dependent anaerobic growth (35, 37). These results suggest that a change in the electron flow through the denitrification pathway may affect the quinone redox state, leading to alterations in *cbb*₃ expression. To further explore the possibility of redox-dependent regulation of *Br. japonicum* *cbb*₃ oxidase, the effect of reduced and oxidized carbon substrates on β -galactosidase activity from a *fixP'*-*lacZ* fusion was investigated. Levels of *fixP'*-*lacZ* expression were largely dependent on the oxidized or reduced nature of the carbon source in the medium (37). In order to study the involvement of *Br. japonicum* RegR in the redox-dependent regulation of *fixNOQP* genes, Bueno and colleagues (37) analyzed the expression of the *fixP'*-*lacZ* fusion in a *Br. japonicum* *regR* mutant. When cells were grown under denitrifying conditions with an oxidized carbon source, there was a significant decrease in β -galactosidase activity in the *regR* mutant compared with the expression levels detected in the wild-type strain. These observations suggest that *Br. japonicum* *fixNOQP* operon might be under the control of RegR. However, a whole-genome transcription-profiling analysis of the *Br. japonicum* *regR* mutant grown under microoxic

conditions demonstrated that transcript levels of *fixNOQP* operon were not RegR-dependent (134). In recent studies from our group (Maria J. Torres *et al.*, unpublished results), *fixNOQP* genes were not identified as RegR targets after transcriptomic analyses of the *regR* mutant grown under denitrifying conditions with succinate (an oxidized carbon source).

E. *Bacillus subtilis* *ResDE* and *Rex*

The endospore-forming Gram-positive bacterium *Bacillus subtilis* utilizes a branched ETC under aerobic conditions (250). To date, three terminal oxidases have been identified in *Ba. subtilis* (Fig. 12a). Both cytochromes *aa*₃ and *caa*₃ have been identified as heme-copper oxidases. The third oxidase has been shown to be a member of the cytochrome *bd* family encoded by the *cydABCD* operon (255). In addition, the presence of a fourth terminal oxidase of *bd* type, YthAB, can be predicted from the genome sequence (6).

Cytochrome *bd* is produced under conditions of low-oxygen tension and in cells grown in the presence of glucose. A perfect 16-bp palindromic sequence, upstream of the translation start site for *cydA*, was proposed as a potential operator binding site for a regulatory protein (255). Gel-shift

and DNase I footprinting analyses identified YdiH as a negative regulator of *cyd* genes (212). YdiH is a redox sensor protein whose activity is regulated by the levels of NAD⁺ and NADH in the cell (104). YdiH has been renamed Rex, as it is an ortholog of Rex in *Streptomyces coelicolor* (31) where this redox-sensitive transcription regulator was first described. In *St. coelicolor*, the Rex-DNA target region (Rex operator, ROP) is an 8-bp inverted repeat located upstream of genes encoding respiratory proteins, such as heme biosynthetic enzymes, NADH dehydrogenase, Rex itself, and cytochrome *bd* oxidase. In *Staphylococcus aureus*, it has been recently demonstrated, by combining various protein-DNA interaction studies with transcriptional analyses, that Rex acts on a multitude of anaerobically induced genes (172).

Brekasis and Paget (31) showed by electrophoretic mobility shift assay (EMSA) experiments that *St. coelicolor* Rex DNA-binding properties are influenced by the NADH/NAD⁺ ratio. Under aerobic conditions, raised NAD⁺/NADH ratio results in an increased binding of NAD⁺ to Rex, leading to its activation (Fig. 12b). Activated Rex binds to ROP, and inhibits the expression of several respiratory genes. However, under oxygen-limiting conditions, NADH replaces NAD⁺ bound to Rex, triggering a conformational change that dissociates Rex from the *cyd*-ROP sequence. Derepression of the *cyd* operon

FIG. 12. Aerobic respiratory chains (a) and regulation of *bd*, *aa*₃, and *caa*₃ terminal oxidase expression (b) in *Bacillus subtilis*. In (a), the quinone-reducing and quinol-oxidizing branches with terminal oxidases induced under oxic conditions (gray boxes), and microoxic conditions (white boxes), are shown. In (b), ResDE and Fnr provide positive transcriptional control in response to oxygen availability. ResD, but not ResE, is required for transcription of *cyd* operon (this complex regulation is indicated with a question mark in the figure). Redox-dependent repression of *cydABCD* (*bd* oxidase) by Rex is controlled by NAD⁺/NADH ratio in the cells. Operons *qox* and *cta* encode the aerobic terminal oxidases *aa*₃ and *caa*₃. Positive regulation (+) is denoted by arrows, and negative regulation (−) is indicated by perpendicular lines.

enables the cell to respire micromolar concentration of oxygen (31) (Fig. 12b). DNA binding studies of *Ba. subtilis* Rex with NADH or NAD⁺ showed that NAD⁺ boosted the binding activity of Rex, but that NADH seemed to have a negligible effect or a partial negative effect on DNA-binding activity (104). By contrast, in *St. coelicolor*, NADH completely inhibits DNA-binding activity. These data suggest that DNA-binding determinants of *Ba. subtilis* Rex are distinct from those of *St. coelicolor* Rex (31). Gyan and colleagues (104) proposed that Rex and NADH dehydrogenase together form a regulatory loop which functions to prevent a large fluctuation in the NADH/NAD⁺ ratio in *Ba. subtilis*.

In order to understand the conformational modifications involved in the redox sensing by Rex, Sickmier and colleagues (220) performed X-ray studies of the Rex homolog in the thermophile *Thermus aquaticus* (T-Rex). Its structure comprises two main domains, an N-terminal domain that adopts a winged H-T-H fold and which most likely is the region that interacts with DNA and a C-terminal NADH binding Rossmann fold domain (220). These studies revealed that T-Rex possesses the same functional characteristics as the *St. coelicolor* homolog Rex, which lead to speculation that the mechanism of repression of gene activation in response to the NADH/NAD⁺ redox pair is conserved among Rex family members. Small-angle X-ray scattering has been recently applied to obtain solution structures for *Ba. subtilis* apo B-Rex, B-Rex:NADH, and B-Rex:NAD⁺ showing that B-Rex presents rigid conformations for the complexes with NADH and NAD⁺ (251). Results from these studies indicated that the domain movements of B-Rex from the DNA bound form to the NADH (DNA-free) form could be different from the reported mechanism for T-Rex which is based on crystal structures (220).

In addition to Rex, transcription of the *Ba. subtilis* cytochrome *bd* oxidase during the transition between aerobic to anaerobic lifestyle is subjected to the action of other regulatory proteins such as Fnr, CcpA (carbon catabolite regulator protein), and the two-component system ResDE (188).

The ResD/ResE two-component signal transduction system has an important role in the regulation of both aerobic and anaerobic respiration [reviewed in (92)]. ResE is a membrane-bound sensor histidine kinase that, by responding to a redox signal(s), which have been postulated to be the redox state of intracellular MK, autophosphorylates and subsequently donates a phosphate to its cognate response regulator, ResD. It has been demonstrated that ResD activates transcription by interaction with the C-terminal domain of the alpha subunit of RNAP (91). The ResD/ResE system activates transcription of genes involved in the respiratory pathway that transfers electrons to oxygen (under oxic conditions) or to nitrate (under anoxic conditions) as well as genes encoding transcriptional factors such as Fnr (91, 92) (Fig. 12b). Since ResDE is essential for both *ctaA* and *ctaB* expression, which are required for heme A biosynthesis, *resDE* mutant strains lack cytochromes *aa₃* and *caa₃*. ResD also activates the genes *ctaCDEF*, encoding the structural proteins for cytochrome *caa₃* (264). By using a *cydA::lacZ* transcriptional fusion in *resD* and *resE* mutant backgrounds, it was shown that ResD, but not ResE, is required for the transcription of the *cydA* promoter, suggesting that another sensor might be involved in *cydA* activation by ResD. The direct interaction of ResD with the *cydA* promoter was found around -58 and -108 nucleotides

upstream from the transcription start site by using a DNase I footprinting assay (188).

IV. Nitrate Respiration: Denitrification and Ammonification

When faced with a shortage of oxygen, many bacterial species are able to use nitrate as an electron acceptor of the respiratory chain. This switch from oxygen to nitrate respiration leads to a reduction in the ATP yield rates, but allows bacteria to survive and multiply (221). Denitrification has been defined as the dissimilatory reduction of nitrate or nitrite to a gaseous N-oxide concomitant with free energy transduction (266). The process requires four separate enzymatically catalyzed reactions:

For many years, it was believed that denitrification is performed exclusively by eubacteria. However, there are indications that some fungi (e.g., the pathogenic species *Fusarium oxysporum*) and archaea are also able to denitrify (236, 266). It has also been shown that nitrifiers also have genes involved in denitrification (44). Some bacteria such as *E. coli* or *Ba. subtilis* are able to perform nitrate respiration and release gaseous nitrogen oxides, but they do not denitrify with dinitrogen as a product. Instead, they reduce nitrate to ammonium, so-called nitrate-ammonification (45). In many species of nitrate-ammonifying bacteria, there are two biochemically distinct nitrate reductases, one membrane-bound with the active site located in the cytoplasm and the other in the periplasm. These are coupled to two nitrite reductases (Nir) that provide independent pathways for nitrate reduction to ammonium (nitrate-ammonification) in the two cellular compartments. In the cytoplasm, nitrate is reduced to nitrite by a membrane-bound respiratory nitrate reductase system (NarGHI):

The nitrite produced can then be further reduced to ammonium by a siroheme-containing Nir (NirB):

In the periplasm, the process involves two different enzymes, a periplasmic nitrate reductase (NapA) that reduces nitrate to nitrite and a periplasmic cytochrome *c* Nir that further reduces nitrite to ammonium. This process of nitrate ammonification produces nitric oxide and nitrous oxide as side-products (95, 149, 222, 225).

Products of denitrification and nitrate ammonification have manifold, mainly adverse, effects on the atmosphere, soils, and waters and thus, have both an agronomic and environmental impact (10, 105). When nitrate is converted to gaseous nitrogen by denitrifying bacteria in agricultural soils, nitrogen is lost as an essential nutrient for the growth of plants. In contrast to ammonium, which is tightly bound in soil, nitrate is easily washed out and flows to the groundwater where it (and its reduction product nitrite) adversely affects water

quality. In addition, nitrogenous oxides released from soils and waters are in part responsible for the depletion of the ozone layer above the Antarctic, and in part for the initiation of acid rain and global warming (192). Among them, N_2O , commonly known as laughing gas, has received special attention in the last years, because it is a powerful greenhouse gas that can persist for up to 150 years while it is slowly broken down in the stratosphere. Although N_2O only accounts for around 0.03% of total greenhouse gas emissions, it has a 300-fold greater potential for global warming effects, based on its radiative capacity compared with that of carbon dioxide (10, 13, 192, 195).

A. Respiratory nitrate reductases

The first reaction of denitrification and ammonification (reactions 1 and 5), the conversion of nitrate to nitrite, is catalyzed by a membrane-bound nitrate reductase (Nar), or a Nap [reviewed in (98, 153, 154, 184, 198, 199, 201, 247)].

The Nar enzyme employs a redox loop to couple quinol oxidation with proton translocation and energy conservation, which permits cell growth under oxygen-limiting conditions (23, 221). Nap is also linked to quinol oxidation, but does not transduce the free energy in the $\text{QH}_2\text{-NO}_3^-$ couple into proton motive force (PMF) to synthesize ATP (221). NO_3^- reduction *via* Nap can only be coupled to free-energy transduction if the primary quinone reductase, for example NADH dehydrogenase or formate dehydrogenase, generates an H^+ -electrochemical gradient (221). In contrast to Nar, which has a respiratory function, Nap systems have a range of physiological functions that include the disposal of reducing equivalents during aerobic growth on reduced carbon substrates and anaerobic nitrate respiration as a part of bacterial ammonification or denitrification pathways (184).

1. Membrane-bound respiratory nitrate reductase. The *E. coli* and *Paracoccus* Nar enzymes have been the focus of the most biochemical and genetic studies (reviewed in references just mentioned). Nar is a structurally defined 3-subunit enzyme composed of NarGHI (Fig. 13a) (24, 118). NarG is the catalytic subunit of about 140 kDa that contains a bis-molybdopterin guanine dinucleotide (*bis*-MGD) cofactor and a [4Fe-4S] cluster. NarH, of about 60 kDa, contains four additional iron-sulfur centers: one [3Fe-4S] and three [4Fe-4S]. NarG and NarH are located in the cytoplasm and associate with NarI. NarI is an integral membrane protein of about 25 kDa with five transmembrane helices and the N-terminus facing the periplasm (Fig. 13a). Nar proteins are encoded by genes of a *narGHJI* operon. The organization of this operon is conserved in most species that express Nar. The *narGHI* genes encode the structural subunits, and *narJ* encodes a dedicated chaperone required for the proper maturation and membrane insertion of Nar (27). *E. coli* has a functional duplicate of the *narGHJI* operon named *narZYWV*. The subunits of the two enzymes encoded by both operons are interexchangeable, but physiologically, NarZYWV has a function during stress response rather than anaerobic respiration *per se* (26, 224). A number of *nar* gene clusters in denitrifying bacteria also have a gene encoding a nitrogen ox anion transporter NarK that can transport nitrate into the cell and nitrite out of the cell (97, 257), a homolog of which is also found in many nitrate-

ammonifying bacteria (45). A variation of the Nar system occurs in some archaea and bacteria where the NarGH subunits are on the outside, rather than the inside of the cytoplasmic membrane. These systems have thus far been less well studied biochemically, but a nitrate transporter is not needed for these systems (142).

2. Periplasmic nitrate reductase. Nap is widespread in all classes of denitrifying and non-denitrifying proteobacteria [reviewed in (98, 153, 154, 184, 198, 199, 201, 247)]. The best-studied Nap enzymes were isolated from *Paracoccus pantotrophus* *E. coli*, *Rho. sphaeroides*, and *Desulfovibrio desulfuromonas* (5, 64, 115). Nap is commonly a 2-subunit enzyme composed of the NapAB complex located in the periplasm and associates with a transmembrane NapC component (Fig. 13a). The catalytic subunit NapA contains the *bis*-MGD cofactor at its active site and an FeS center. NapB is di-heme cytochrome c_{552} , and NapC is a *c*-type tetra-heme membrane-anchored protein that is involved in the electron transfer from the quinol pool to NapAB (43, 207). Mono-subunit forms of NapA also occur, which may interact with small FeS proteins as well as cytochromes (114, 115, 140).

In contrast to the *nar* operon, the *nap* operons present considerable heterogeneity in gene composition as well as in ordering. Eight different genes have been identified as components for operons that encode Naps in different organisms (199). Except for *Shewanella oneidensis*, *Wollinella succinogenes*, *Desulfobacterium hafniense*, and *C. jejuni*, which lack *napB* or *napC*, all operons studied thus far have the *napABC* genes in common. The remaining *napDEFKL* genes encode for different proteins that are not directly involved in the nitrate reduction. NapD is a cytoplasmic protein that acts as a chaperone. NapF is a cytoplasmic iron-sulfur containing protein with four loosely bound [4Fe-4S] clusters, and is thought to participate in the assembling of the iron-sulfur cluster of NapA (164, 170). The *napEKL* genes encode for proteins with so far unknown functions. In *E. coli*, the *nap* operon includes *napGH* genes encoding a periplasmic and an integral membrane protein with [4Fe-4S] clusters. NapH and NapG interact, making an electron transfer supercomplex that can channel electrons from both menaquinol and ubiquinol to NapA (32, 33).

B. Other enzymes in bacterial denitrification

The enzymes involved in denitrification are nitrate-, nitrite-, nitric oxide-, and nitrous oxide reductase, encoded by *nar/nap*, *nir*, *nor*, and *nos* genes, respectively. A scheme of the denitrification reactions is shown in Figure 13a. Comprehensive reviews covering the physiology, biochemistry, and molecular genetics of denitrification have been published elsewhere (11, 16, 110, 125, 199, 247, 248, 266).

1. Nitrite reductases. In denitrifying bacteria, two types of respiratory Nir have been described: the NirS *cd*₁ nitrite reductase, a homodimeric enzyme with hemes *c* and *d*₁, and the NirK, a copper-containing Nir [reviewed in (202, 203, 246, 247)]. Both are located in the periplasmic space, and receive electrons from cytochrome *c* and/or a blue copper protein, pseudoazurin *via* the cytochrome *bc*₁ complex (Fig. 13a). They catalyze the one-electron reduction of nitrite to nitric oxide. Neither of the enzymes is electrogenic.

FIG. 13. Denitrification pathway (a) and regulation (b) in bacteria. In (a), the topological organization of denitrification enzymes is shown. The membrane-bound (NarGHI), and periplasmic, (NapABC) nitrate reductases as well as the nitrite reductases (Cu-type or cd_1 -type), nitric oxide reductases (cNor, qNor, and qCu_ANor), and nitrous oxide reductase (NosZ) are shown. (b) Regulatory network of denitrification genes in response to O₂ concentration, nitrate/nitrite, and nitric oxide (NO). Positive regulation is denoted by arrows, and negative regulation is indicated by perpendicular lines. Further details are given in the text.

The cd_1 -type nitrite reductase (cd_1 Nir) is a homodimer, and each monomer carries a small heme *c* binding domain and a larger heme d_1 -binding domain. Electrons are transferred from the electron donor via the heme *c* of NirS to heme d_1 , where nitrite binds and is reduced to nitric oxide (202). The *nirS* gene encoding the cd_1 -Nir polypeptide is part of a

nir gene cluster. The best-characterized clusters are those from the denitrifying species *Ps. aeruginosa* (*nirSMCFDLGH-JEN*), *Pa. denitrificans* (*nirXISECFDLGHJN*), and *Ps. stutzeri* (*nirSTBMCFDLGH* and *nirJEN*, the two clusters being separated by a part of the *nor* gene cluster encoding nitric oxide reductase [Nor]).

Cu-type nitrite reductases (CuNir) are homotrimeric complexes harboring three type I copper centers, and three type II copper centers, which form the active site. Nitrite binds to the copper ion in the type II center, replacing an exogenous ligand (water or chloride), and by electron transfer from the type I copper site, nitrite is reduced to nitric oxide. The *nirK* gene encodes the CuNir (203).

2. Nitric oxide reductases. To date, three types of Nor have been characterized, and they are classified according to the nature of their electron donor as *c*-type nitric oxide-reductase (cNor), qNor, and qCu_ANor [reviewed in (59, 247, 248, 267)]. As a unique case, the Nor of *Ros. denitrificans* is similar to cNor, but differs in that it contains copper (143). The best-studied Nor is the cNor that uses membrane or soluble *c*-type cytochromes or small soluble blue copper proteins (azurin, pseudoazurin) as physiological electron donors. The qNor uses quinol or menaquinol as electron donors and it is found not only in denitrifying archaea and soil bacteria, but also in pathogenic microorganisms that do not denitrify (60). The qCu_ANor has thus far been found only in the Gram-positive bacterium *Bacillus azotoformans* (230). This enzyme is bifunctional using both menahydroquinone (MKH₂) and a specific *c*-type cytochrome *c*₅₅₁ as electron donor. It was suggested that the MKH₂-linked activity of qCu_ANor serves detoxification and the *c*₅₅₁ pathway has a bioenergetic function.

The cNor is an integral membrane enzyme harboring two subunits, NorC with a heme *c* group and NorB containing hemes *b* and a non-heme iron (Fig. 13a). Electron transfer from donor molecules to cNor is mediated by the cytochrome *bc*₁ complex and a soluble cytochrome *c* or pseudoazurin (174). Electrons are transferred to the heme *c* in subunit II and then *via* the heme *b* to the active site. There, two molecules of nitric oxide are reduced. The molecular mechanism of the NO reduction by Nor has been extensively studied through chemical, biochemical, and physicochemical techniques (80, 81, 252). The crystal structure of Nor from *Ps. aeruginosa* has been recently solved (112). Although the overall structure of Nor is closely related to the heme copper oxidases, neither the D- nor K-proton pathway, which connect the HCO active center to the intracellular space, was observed. This confirmed early studies with NorCB in intact cells of *Rho. capsulatus* which showed that it was not a proton pump or electrogenic (196). Site-specific mutagenesis has suggested that protons required for the Nor reaction are probably provided from the periplasmic side of the membrane *via* a channel that involves conserved glutamate residues (41, 85, 86, 237).

cNOR is encoded by the *norCBQD* operon. The *norC* and *norB* genes encode subunit II and subunit I, respectively. The *norQ* and *norD* genes encode proteins essential for activation of cNor. Some more specialized denitrifiers have additional *norEF* genes, the products of which are involved in maturation and/or stability of Nor activity (107).

3. Nitrous oxide reductase. The final step in denitrification consists of the two-electron reduction of nitrous oxide to dinitrogen gas. This reaction is performed by a copper containing homodimeric soluble protein located in the periplasmic space, the nitrous oxide reductase (Nos) [reviewed in (246–248, 268)] (Fig. 13a). The enzyme has been purified from a large number of denitrifying strains, including *Pa. deni-*

trificans *Pa. Pantotrophus*, and *Ps. stutzeri*. Nos is a homo-dimer of a 65 kDa copper-containing subunit. Each monomer is made up of two domains: the “Cu_A domain” and the “Cu_Z domain.” The recently reported structure of purple N₂O_R from *Ps. stutzeri* has revealed that N₂O bound side-on at Cu_Z, in close proximity to Cu_A (181). Electron input into Cu_A is usually *via c*-type cytochromes or cupredoxins (20, 22). The *nos* gene clusters often comprise the *nosRZDFYLX* genes. The *nosZ* gene encodes the monomers of Nos. The *nosDFYL* genes encode proteins that are apparently required for copper assemblage into Nos, although their specific role still remains unknown. The NosRX proteins have roles in transcription regulation, activation, and Cu assemblage of Nos (268).

C. Oxygen and nitrogen oxides control of nitrate respiration

The key molecules that act as signals for the regulation of nitrate respiration through denitrification or ammonification pathways are oxygen, nitrate, nitrite, and NO. Sensing of each of these molecules involves several types of sensor proteins (see Fig. 13b).

1. Oxygen sensors. There are two types of oxygen sensors involved in regulation of nitrate respiration, Fnr and FixL (reviewed in section II.A; see Fig. 4). Fnr controls the expression of genes involved in anaerobic respiration when O₂ is absent. For example, the *nar* and *nap* operons in *E. coli* and *Ba. subtilis* (193, 228, 238, 241) are activated by Fnr under anaerobic conditions (Fig. 13b). *Pa. denitrificans* FnrP controls expression of the *nar* gene cluster and the *cco*-gene cluster encoding the *ccb*₃-type oxidase (30, 249). Oxygen tension is sensed in *Ps. aeruginosa* by the Anr regulator, which carries a [4Fe-4S]²⁺ cluster (262).

2. Nitrate and nitrite sensors. In denitrifying species and species that anaerobically reduce nitrate to ammonium, there are three types of nitrate/nitrite sensing systems: NarXL, NarQP, and NarR. NarXL and NarQP are members of two-component regulatory systems (Fig. 13b). The NarX and NarQ proteins are the signal sensors, and NarL and NarP proteins are their cognate response regulators, respectively (227). In *E. coli* NarL and NarP bind DNA to control induction of the *nar* and *nap* operons and repression of genes encoding alternate anaerobic respiratory enzymes (58, 227, 228). The effects of nitrate and nitrite on the *E. coli* transcriptome during anaerobic growth have been investigated, resulting in a new list of operons that are regulated by Fnr, NarL, and NarP (48). To date, *narXL* and *narQP* genes are confined to species classified in the γ and β subdivisions of the proteobacteria such as *Escherichia*, *Salmonella*, *Klebsiella*, *Yersinia*, *Burkholderia*, *Ralstonia*, *Neisseria*, and *Pseudomonas* species among others. In *Ps. aeruginosa*, the regulators Anr, Dnr, and NarL in concert with integration host factor (IHF) activate transcription of the *narK1K2GHJI* operon encoding nitrate reductase and two transporters in response to oxygen limitation, nitrate, and N-oxides (213). Recently, it has been shown that during anaerobic growth of *Ps. aeruginosa* PAO1, NarL directly represses expression of *Nap*, while inducing maximal expression of membrane nitrate reductase (245).

NarR is a member of the Fnr family of transcription activators, but it lacks a [4Fe-4S] cluster. NarR of *Pa. pantotrophus*

and *Pa. denitrificans* is specifically required for transcription of the *narKGHJL* genes in the presence of nitrite (249, 256) (Fig. 13b). Genes encoding NarR are found in the α -proteobacteria *Brucella suis*, *Brucella melitensis*, *Pa. Denitrificans*, and *Pa. pantotrophus*. There are no indications that they have counterparts of *narXL*. It, therefore, seems that NarR substitutes the NarXL system in the α -proteobacteria. A recent proteomic analysis of *Pa. denitrificans* *fnrP*, *nnr*, and *narR* mutant strains grown oxidically, microoxidically, and microoxidically in the presence of nitrate has revealed new proteins involved in the FnrP, Nnr, and NarR regulons (30). Data from this work are in full agreement with the current view on the FnrP protein being a sensor for oxygen and on the Nnr and NarR proteins being sensors for intermediates of nitrate respiration. This study has also revealed that FnrP, Nnr, and NarR control transcription of target genes having both activating and repressing functions (30).

3. NO sensors. NO is a toxic, free radical gas with diverse biological roles in eukaryotes and bacteria. Several bacterial transcriptional regulators sense this molecule and regulate the expression of genes involved in both denitrification and NO detoxification [reviewed in (206, 226, 243, 267)]. Bacterial NO regulatory proteins can be roughly divided into (i) proteins whose primary role is not to sense NO, but which contain metal centers or cysteines that are nitrosylated by NO and (ii) those that appear to be solely dedicated to sensing NO (*i.e.*, they regulate genes involved in NO detoxification). An example of the first class is Fnr, whose primary role is to sense oxygen but that has also been shown to sense and respond to NO in *E. coli* (see section II.A).

The second class of NO sensors includes nitrite and nitric oxide reductase regulator (NnrR), NorR, and NsrR. NnrR also belongs to the Fnr family of transcription activators, but, similar to NarR, it lacks the cysteines to incorporate a [4Fe-4S] cluster. NnrR orthologs, sometimes named as Nnr, DnrR, or FnrD, are found in nearly all species containing the reductases Nir and Nor. In the presence of NO, these regulators orchestrate the expression of the *nir* and *nor* gene clusters. In many species, *nnrR* gene and its target genes are situated in close proximity on the chromosome, suggesting a local specificity. The promoters of these operons contain NnrR binding sites that resemble the consensus Fnr-box to a large extent. Nnr homologs, including NnrR in *Rho. sphaeroides* (131) and Dnr in *Ps. aeruginosa*, most likely sense NO *via* a ligated heme (93, 94). In denitrifying bacteria, knockout mutagenesis has shown that the Dnr and NnrR regulators are needed for the transcriptional control of the *nir*, *nor*, and *nos* operons.

NorR is another NO-responsive protein that was first identified in *Ralstonia eutropha* (180). This bacterium has two copies of the *norR* gene, both of which are located upstream of their *norAB* gene clusters where *norB* encodes a single-subunit Nor of the qNor type. NorR is a member of the NtrC family of response regulators. The absence of possible phosphorylation sites as well as the presence of a conserved GAF domain indicative for signal perception suggest that the protein belongs to a sub-family of response regulators which sense their signal themselves rather than *via* a cognate signal sensor (55). In response to anaerobiosis and the presence of NO, the NorR proteins specifically activate transcription of the σ^{54} -dependent *norAB* promoters and repress their own synthesis *via* negative autoregulation (39). NorR has also been found in *E. coli*, which senses NO directly through a mononuclear non-

heme iron center and responds by switching on expression of the flavorubredoxin NorVW to detoxify NO (242).

NsrR is an iron-sulfur-containing protein that senses NO directly *via* a [2Fe-2S] cluster. Nitrosylation of this cluster leads to a loss of DNA binding activity and, hence, derepression of NsrR target genes. In *E. coli*, this transcription repressor was shown to sense reactive nitrogen species and to switch on a regulon of at least 60 genes, including genes involved in nitrate respiration (82, 226, 243). In denitrifying bacteria, NsrR appears to have a specific role in coordinating production of the nitrite and NO reductase enzymes to prevent the build up of NO (243). Intriguingly, the same role is performed by Nnr homologs in denitrifying bacteria that do not contain NsrR. Two components of the denitrification pathway are transcriptionally regulated by NsrR in *Neisseria meningitidis*: the membrane-bound nitrite reductase AniA and the respiratory NO reductase NorB (111). In the closely related *Neisseria gonorrhoeae*, NsrR represses both *norB* and *aniA* expression in the absence of NO and, most notably, also exerts an autoregulatory effect on its own expression (113, 171).

4. Other regulators in denitrifying bacteria. In addition to the regulatory proteins that can monitor oxygen, nitrate, nitrite, and NO, regulation of the on-set and fine-tuning of nitrate respiration in some bacteria involves copper responsive regulators, redox sensing mechanisms and the NosR and NirI proteins (Van Spanning, 2011). The NosR is a membrane-bound protein containing six transmembrane helices, a large periplasmic domain, and cysteine-rich cytoplasmic domains that resemble the binding sites of [4Fe-4S] clusters. The periplasmic domain of NosR has a structural similarity with the FMN-binding domain of the NqrC subunit of the Na⁺-translocating NADH:quinone oxidoreductase and may play a role in activating or repairing oxygen damaged Nos (260). Although NosR proteins do not have properties of transcription factors, they are required for transcription activation of the *nos* genes in denitrifying bacteria. However, the regulatory function of NosR is not well understood (268). In *Pa. Denitrificans*, a homolog of *nosR*, *nirI*, representing thus far a singular case among the denitrifying bacteria, has been studied for its role in the transcription of *nirS*, encoding the cytochrome *cd₁*Nir.

V. Redox Control of Nitrate Respiration

Nitrate reduction by the membrane-bound Nar generates an H⁺-motive force (PMF) across the cytoplasmic membrane of the cell, allowing ATP synthesis (197, 221). During denitrification, the periplasmic Nir and Nos cannot themselves be involved in H⁺-movement across the membrane. Although Nor is an integral membrane protein, several lines of evidence show that H⁺ required for reduction of NO are taken from the periplasm and thus, no net charge movement occurs (see section IV.B). However, e-transfer from ubiquinol to NO₂⁻, NO or N₂O is coupled to H⁺-translocation across the membrane, because this involves the cytochrome *bc₁* complex (197). Quinol oxidation by the periplasmic Nap is not directly coupled to the generation of a PMF and is independent of the cytochrome *bc₁* complex. In many bacteria, Nap has a role in redox balancing under aerobic conditions (20, 42); indeed it seems to be biochemically tuned for this function (88). The need to dissipate reductants is likely to be most acute during

the metabolism of a reduced carbon substrate under conditions that are both oxygen and energy sufficient. Accordingly, expression of the Nap is significantly higher in *Paracoccus* species grown aerobically with butyrate or caproate to that when grown with malate or succinate (69, 71, 196, 215, 216). Taken together, these results suggest that Nap constitutes an example of redox-dependent regulation of the respiratory network in *Paracoccus* species. Two crucial regulatory sequences that control expression of the *nap* operon in response to the redox status of the cell were identified in *Pa. pantotrophus*, but the redox responsive transcriptional regulator has yet to be described (70). The use of Nap in redox homeostasis has been also shown in photosynthetic bacteria where it regulates the redox poise of the cyclic photosynthetic electron transport system (117), particularly during photoheterotrophic growth on reduced carbon substrates (89, 200).

Many bacteria can also recruit Nap for anaerobic nitrate respiration as a part of bacterial ammonification (*Enterobacteriaceae*) or to promote denitrification (*Rhizobiaceae*). In fact, some rhizobia species (e.g., *Pseudomonas* sp. G179 [actually *Rhizobium galegae*], *Br. japonicum*) can express *nap* genes under anaerobic conditions, and disruption of the *nap* genes is lethal for growth under denitrifying conditions (17, 62). In addition, in *Rho. sphaeroides* f. sp. *denitrificans*, which can express both Nar and Nap, inactivation of Nap is lethal for anaerobic denitrification (135). Thus, in these organisms, the physiological role of Nap is in anaerobic denitrification. When considered as single enzymes, the energy coupling of Nar and Nap appear markedly different: $q + 1/2e^- = 6$ (Nar) and 4 (Nap) with NADH as electron donor and 2 (Nar) and 0 (Nap) with succinate as electron donor. However, when considered in the context of the entire denitrification pathway, the $q + 1/2e^-$ ratio is 24 (Nar) or 22 (Nap) with NADH and 8 (Nar) or 6 (Nap) with succinate. Thus, the energetic loss of using Nap rather than Nar is only 8% when NADH is the electron donor to the respiratory system (197).

To date, the knowledge about the redox-dependent control of anaerobic nitrate respiration as well as the identification of the regulators and the signal transduction mechanism involved in this response is limited. In this section, we will focus on recent advances in understanding nitrate respiration redox-dependent control in *Rho. sphaeroides*, *Ag. tumefaciens*, *Br. japonicum*, and *Ba. subtilis*.

A. *Rho. sphaeroides*

Rho. sphaeroides strains are photosynthetic bacteria able to induce the denitrification pathway by expressing the enzymes Nap, Nir, and cNor, which are encoded by the *nap*, *nirK*, and *nor* genes, respectively, when oxygen concentration is limited and nitrate or nitrite is present in the medium. In *Rho. sphaeroides* 2.4.3., Nap is the enzyme responsible for denitrification as judged by the fact that strains with mutations in the *nap* gene cluster lost nitrate reductase activity as well as the ability to grow with nitrate under anaerobic-dark conditions (135). In the strain IL106 of *Rho. sphaeroides*, *nap* genes showed the same level of expression under both oxic and denitrifying conditions with nitrate. Expression increases in the presence of the reduced carbon compound butyrate, indicating that this protein probably plays a role in redox homeostasis [reviewed in (218)]. A slightly different *nap* expression pattern was observed in *Rho. sphaeroides* strain 2.4.1, where expression of the

nap operon increases significantly when cells are shifted from anoxic photosynthetic conditions to oxic conditions and do not require nitrate [reviewed in (218)].

Although the expression of *nap* has been shown to be variable, the expression of the genes encoding Nir and Nor only occurs under microoxic conditions. In *Rho. sphaeroides* 2.4.3, *nir* and *nor* expression requires low oxygen and also requires the presence of a nitrogen oxide. It has been shown that NnrR is necessary for the expression of *nirK* and *nor* during denitrification (240). Studies on regulation by NnrR demonstrated that the presence of NO is required for transcriptional activation of *nirK* and *nor*, yet the mechanism whereby NnrR senses NO is not known. Analysis of the *Rho. sphaeroides* 2.4.3 genome and expression studies demonstrated that genes designated *paz* and *norEF* are also members of the NnrR regulon involved in denitrification (107).

Genes for anaerobic growth in many photosynthetic bacteria are also regulated by the two-component PrrBA system. In *Rho. sphaeroides* strain 2.4.1 where *nap* genes are maximally expressed under oxic conditions, the phosphorylated form of the response regulator PrrA is responsible for repressing *nap* expression under low oxygen [reviewed in (218)], see Fig. 14a). By contrast, in *Rho. sphaeroides* 2.4.3, inactivation of *prrA* eliminated the ability to grow both photosynthetically and anaerobically in the dark on nitrite-amended medium (130). The PrrA-deficient strain exhibited a severe decrease in both Nir activity and expression of a *nirK-lacZ* fusion (130) when environmental oxygen tension was limited. This regulation is not mediated by NnrR, as *nnrR* is fully expressed in a PrrA mutant background (Fig. 14a). By using truncated versions of the *nirK* promoter fused to the *lacZ* marker gene, Laratta and colleagues (130) revealed that the DNA region expanded upstream to the putative NnrR binding site is required for full activation of *nirK*. These results suggested that this DNA region could act as a binding site of PrrA or additional regulatory proteins in order to control *nirK* expression. In fact, the presence upstream of the *nirK* start site of 4 imperfect repeats proposed as binding sites for RegR (a PrrA homolog in *Br. japonicum*) suggests again that PrrA could bind directly to such sites. However, binding assays using purified PrrA would be required to confirm this. Unlike the *nirK* promoter region, the *nor* promoter has no sites with similarity to the imperfect repeats proposed as binding sites for PrrA homologs, suggesting that PrrA may not directly regulate *norB* expression (130). The model proposed by Laratta and colleagues (130) to understand the mechanism of *nirK* control by PrrBA is that under low oxygen tensions, the kinase activity of PrrB increases relative to its phosphatase activity, resulting in an increase in the concentration of PrrA-P. Under microoxic conditions and in the presence of NO, NnrR together with PrrA-P activates transcription of *nirK* (Fig. 14a). As mentioned in section III.B (Fig. 8c), inactivation of genes encoding the *cbb₃* oxidase in *Rho. sphaeroides* has been shown to lead to high levels of PrrA-P under both oxic and microoxic conditions (190). Unexpectedly, loss of the *cbb₃* oxidase resulted in a significant decrease in *nirK* expression and Nir activity (130). Recent results have demonstrated that the *cbb₃* mutant retains Nir activity at very low oxygen concentrations (107). The simplest explanation for this result is that, under transition conditions, loss of the *cbb₃* oxidase results in high residual oxygen levels in the culture, which prevents expression of *nirK* and *nor*.

FIG. 14. Schematic representation of denitrification control in *Rho. sphaeroides* (a) and *Agrobacterium tumefaciens* (b). (a) In *Rho. sphaeroides*, PrrBA provides control of *nirK* expression in response to external redox variations, and NnrR provides control of expression of *nirK* and *nor* genes in response to NO. This figure is adapted from ref. (218). (b) In *Ag. tumefaciens*, ActSR provides control of *nirK* expression presumably in response to external redox variations, and FnrN and NnrR provide control of *nirK* and *nor* expression in response to low oxygen tension and presence of NO, respectively. Positive regulation is denoted by arrows, and negative regulation is indicated by perpendicular lines.

B. *Ag. tumefaciens*

Ag. tumefaciens is a plant pathogen that can grow under free-living conditions in the absence of oxygen *via* denitrification. The ability to respire nitrogen oxides could be significant for the survival and growth of free-living and plant-associated *Ag. tumefaciens* cells. The ability to respire nitrate has been shown to be advantageous to bacteria in the rhizosphere, as nitrate provides an alternative oxidant when oxygen concentrations are low (25). In addition, the ability to reduce nitrogen oxides may be useful for countering some of the plant defenses against plant pathogens (63, 79). *Ag. tumefaciens* is considered a partial denitrifier, as it lacks the genes encoding Nos (9). Consequently, nitrous oxide is the final product of denitrification.

Ag. tumefaciens has a *nap* cluster that is maximally expressed under denitrifying conditions, but is not influenced by nitrate (218). The regulator(s) responsible for expression of *nap* in *Ag. tumefaciens* are uncertain, as there are no conserved sequence motifs in the promoter region that might indicate the involvement of a particular family of regulators. Similar to *Rho. sphaeroides*, activation of *nirK* and *norCBQD* expression requires low oxygen concentration and the presence of NO (8, 9, 21). By using the *lacZ* marker gene fused to the *nirK* and *nor*

promoter and by *in vitro* binding assay experiments, it has been demonstrated that activation of *nirK* and *nor* requires NnrR. This protein is directly activated by FnrN when oxygen tension in the environment becomes depleted (Fig. 14b). However, activation of *nirK* and *nor* is raised when NO, as a reduction product of nitrate or nitrite, is present in the medium, due to the maximal NO-dependent activation of NnrR. In *Ag. tumefaciens*, electron transfer through the denitrification pathway is facilitated by soluble small cytochromes such as the copper-containing pseudoazurin, encoded by the *paz* gene, whose expression also depends on FnrN and NnrR. In addition to FnrN and NnrR, the ActRS system is involved in denitrification (Fig. 14b). As observed in *Rho. sphaeroides*, insertional inactivation of the response regulator ActR significantly reduced *nirK* expression and Nir activity but not *nnrR* expression. In *Ag. tumefaciens*, a putative ActR binding site was identified in the *nirK* promoter region using mutational analysis and an *in vitro* binding assay (8). These studies also show that purified ActR bound to the *nirK* promoter but not to the *nor* or *nnrR* promoter. Additional experiments revealed that expression of *paz* was highly reduced in the *actR* or *fnrN* mutants and that ActR binds to the *paz* promoter (8). Due to the similarity to the PrrBA/RegBA systems, the ActSR system is believed to mediate gene activation on sensing redox

FIG. 15. Schematic representation of the regulatory network of denitrification in *Br. japonicum*. FixLJ regulates *fixK₂* gene transcription in response to a shortage of oxygen, and FixK₂ provides control of transcription on *nap* and *nirK* denitrification genes as well as the *nnrR* gene, whose product, in turn, activates transcription of *nor* genes. RegSR activates expression of *fixRnifA* operon presumably by sensing changes in the redox state of the cell. The NifA regulatory protein activates gene expression at very low oxygen concentrations. Control of denitrification genes by NifA has been demonstrated by our group (36). Furthermore, recent results have shown that NorC expression is significantly reduced in a *Br. japonicum* *regR* mutant (239). Sigma⁵⁴ links both regulatory cascades. Positive regulation is denoted by arrows, and negative regulation is indicated by perpendicular lines.

changes originated from the ETC through *cbb₃*-oxidase (PrrB) or through the redox state of the UQ pool (RegB). However, there is no experimental evidence as yet to confirm these hypotheses.

With all of this experimental evidence, we can see that activation of denitrification in *Ag. tumefaciens* depends on regulatory proteins involved in sensing different environmental signals. While the Fe-S-containing FnrN protein is an oxygen sensor, NnrR activates denitrification genes in response to NO. In this complex regulatory process, the Prr/Reg orthologs protein ActSR would adjust the denitrification machinery in response to redox status as an indirect measure of external oxygen tension (Fig. 14b).

C. *Br. japonicum*

Similar to *Ag. tumefaciens*, the nitrogen-fixing root nodule symbiont of soybean *Br. japonicum* has the ability to sustain cell growth by using nitrate as respiratory substrate when oxygen concentrations are very low. In contrast to *Ag. tumefaciens*, *Br. japonicum* contains all the denitrification enzymatic machinery necessary to reduce sequentially nitrate to molecular nitrogen.

In *Br. japonicum*, denitrification is dependent on the *napEDABC*, *nirK*, *norCBQD*, and *nosRZDYFLX* genes that encode an Nap, a Cu-containing Nir (NirK), a cNor, and an Nos, respectively (16). In addition, accessory cytochromes are

necessary to support electron transport during denitrification. For example, the cytochrome *c₅₅₀*, encoded by *cycA*, is essential for the electron delivery to the NirK reductase (35). Very recently, it has also been found that the cytochrome *c₆*, encoded by the *cycS* gene, has a role in denitrification, although its precise biochemical function still remains unknown (145). Neither azurin- nor pseudoazurin-like copper proteins has been annotated in the genome sequence of *Br. japonicum* (www.kazusa.jp/rhizobase/).

In *Br. japonicum*, maximal expression of denitrification genes requires both the absence of oxygen and the presence of nitrate or a derived N-oxide. Microoxic induction of transcriptional fusions from the *nap*, *nir*, *nor*, and *nos* promoter regions to the *lacZ* reporter gene depends on the FixLJ/FixK₂ regulatory cascade (Fig. 15) (16, 204). Sequences homologous to the consensus DNA binding sites of the Fnr and FixK proteins are present in the promoter region of each denitrification gene. Regulatory studies indicated that, in addition to low oxygen, nitrate, or an N-oxide derived from it, presumably either NO₂⁻ or NO or both, are required for maximal induction of *Br. japonicum* denitrification genes. This N oxide-mediated induction of *nap*, *nir*, and *nor* genes depended on NnrR (144, 204). Thus, NnrR expands the FixLJ/FixK₂ regulatory cascade by an additional control level that integrates the N oxide signal required for maximal induction of the denitrification genes (Fig. 15). Induction of the *norCBQD* promoter is completely abolished in the absence of a functional

nrrR gene. By contrast, microoxic induction of the *nap* or *nirK* promoters is retained in a *nrrR* mutant background, implying that the *nap* or *nirK* and the *norCBQD* promoters exhibit slight differences with regard to their dependence on FixK₂. In this context, recent results from our group have demonstrated that purified FixK₂ activates transcription from *nap*- or *nirK*-dependent promoters but not from *nor*-dependent promoter (E. Bueno, unpublished work). By contrast, ITC allowed us to demonstrate that NnrR bound to a specific DNA fragment from the promoter region of the *nor* genes, but not to those from the *nap* and *nirK* genes and that this interaction requires anoxic conditions but not the presence of an N oxide (unpublished results from our laboratory). Supporting these observations, a genome-wide transcription profiling of *Br. japonicum* *fixJ* and *fixK₂* mutant strains grown in free-living microoxic conditions have shown that *nap*, *nirK*, and *nrrR* but not *nor* are targets of *fixK₂* regulatory gene (145).

In addition to FixLJ/FixK₂-NnrR, in *Br. japonicum*, a second oxygen responsive regulatory cascade has been described, the RegSR/NifA cascade (Fig. 15). NifA activates gene expression in concert with the RNAP containing the alternative sigma factor (σ^{54}), which, in *Br. japonicum*, is encoded by the two highly similar and functionally equivalent genes (*rpoN₁* and *rpoN₂*) (127). Since *rpoN₁* is under the control of FixK₂, this gene represents the link between the two regulatory cascades.

While the FixLJ/FixK₂-NnrR cascade activates gene transcription at a moderately low oxygen concentration ($\leq 5\%$), activation of the RegSR/NifA system requires oxygen concentration $\leq 0.5\%$ (214). Recent results from our group showed that NifA is required for maximal expression of *nap*, *nirK*, and *nor* genes, suggesting a role for the RegSR/NifA regulatory cascade in the control of the denitrification process in *Br. japonicum* (36). In that study, it was shown that disruption of *nifA* caused a growth defect in *Br. japonicum* cells when grown under denitrifying conditions, as well as decreased activity of Nap and Nir enzymes. Furthermore, expression of *napE-lacZ*, *nirK-lacZ*, or *norC-lacZ* transcriptional fusions, as well as levels of *nirK* transcripts, were significantly reduced in the *nifA* mutant after incubation under nitrate-respiring conditions. These results suggest that maximal induction of denitrification genes is only triggered once the cellular oxygen falls to a sufficiently low level, that is, within the sensory range of the NifA-dependent regulatory cascade. Similarly, in *Ag. tumefaciens*, a study of expression and activity of Nir and Nor as a function of oxygen has shown that *nirK* and *nor* are expressed once oxygen reaches $< 0.5 \mu\text{M}$ or approximately 0.2% of air-saturating oxygen (21). These oxygen concentrations are low enough to allow formation of ActR-P necessary for *nirK* activation in *Ag. tumefaciens*.

Concerning the involvement of *Br. japonicum* RegSR in denitrification, it has been shown that free-living growth of *regR* mutants under anoxic conditions with nitrate as the terminal electron acceptor was severely impaired [Bauer *et al.* (15)]. Furthermore, expression of NorC is significantly lower in membranes from the *regR* mutant (239). As just mentioned, it has been shown that disruption of *Rho. sphaeroides* *prfA* or *prfB* causes a significant decrease in both *nirK* expression and Nir activity. Similarly, in *Ag. tumefaciens*, it has been shown that purified ActR binds to the *nirK* promoter, but not to the *nor* or *nrrR* promoters. In *Br. japonicum*, the involvement of RegR in *nirK* expression is at the moment unknown, and further investigations are needed to demonstrate the in-

volvement of RegR in *norCBQD* gene expression and to establish whether these genes are direct, or indirect, targets of RegR.

D. *Ba. subtilis*

Once the oxygen present in the medium has been depleted by the *bd* oxidase (see section III.E; Fig. 12a), *Ba. subtilis* has the ability to exploit the use of an alternative final electron acceptor such as nitrate (158). Two distinct operons specifying two nitrate reductases have been characterized: (i) *narGHJ* encoding a respiratory nitrate reductase, which is required for nitrate respiration and is dependent on Fnr, and (ii) *nasBC* encoding a assimilatory nitrate reductase involved in the assimilation of nitrogen from nitrate under oxic conditions (159) and references therein) (Fig. 16). *Ba. subtilis* does not contain *nirK*, *nor*, or *nos* denitrification genes. Under anoxic conditions with nitrate or nitrite, this bacterium reduces nitrite to ammonium by the Nir NasDE encoded by *nasD* and *nasE* (Fig. 16) (*i.e.*, reactions 5 and 6). The NasDE Nir was previously identified as an assimilatory enzyme [(157) and references therein]. Under aerobic conditions of nitrogen limitation, this enzyme catalyzes the reduction of nitrite to ammonia for the anabolic incorporation of nitrogen into biomolecules. It also functions catabolically in anaerobic respiration, which involves the use of nitrite as a terminal electron acceptor (157). Nitrite reduction catalyzed by *Ba. subtilis* Nir does not result in a proton gradient and coupled ATP generation. Instead, nitrite enhances anaerobic growth by serving as an electron sink to recycle the cellular pyridine nucleotide pool (157).

The *nasDE* genes, together with *nasBC* (encoding assimilatory nitrate reductase) and *nasF* (required for Nir siroheme cofactor formation), constitute the *nas* operon (157). Transcription of *nasDEF* is driven not only by the *nasBCDEF* operon promoter, but also from an internal promoter residing between the *nasC* and *nasD* genes. Transcription from both promoters is activated by nitrogen limitation during aerobic growth by the nitrogen regulator, TnrA. However, under conditions of oxygen limitation, *nasDEF* expression and Nir activity were significantly induced (157, 158).

The expression of genes involved in nitrate respiration in *Ba. subtilis* is also regulated by the ResDE two-component signal transduction system (Fig. 16). To successfully switch from aerobic growth to nitrate respiration, the ResDE two-component signal transduction system should be activated, which allows induction of genes that function in anaerobic metabolism including *fnr*, *nasDEF*, and *hmp* (encoding flavohemoglobin) [(90) and references therein]. Footprinting analysis revealed that ResD tandemly binds to the -41 to -83 region of *hmp* and the -46 to -92 region of *nasD* (90). *In vitro* run-off transcription experiments showed that ResD is necessary and sufficient to activate transcription of the ResDE regulon. Although phosphorylation of ResD by ResE kinase greatly stimulated transcription, unphosphorylated ResD, and ResD with a phosphorylation site (Asp⁵⁷) mutation, it was able to activate transcription at a low level (90). These results suggest that ResD itself, in addition to its activation through phosphorylation-mediated conformation change, senses oxygen limitation *via* an unknown mechanism leading to anoxic gene activation. Although ResDE was initially believed to sense anoxic conditions, subsequent studies have revealed that NO is also required (151, 155). It has been

FIG. 16. Schematic representation of denitrification control in *Ba. subtilis*. ResDE regulates gene transcription in response to both redox/oxygen changes and presence of NO, Fnr provides control in response to low oxygen concentration, NsrR in response to NO, and TnrA in response to high oxygen concentration and limited nitrogen availability. *narGHJI* encodes a respiratory nitrate reductase. *nasBC* encodes an assimilatory nitrate reductase; *nasDEF*, a nitrite reductase; and *hmp*, a flavohemoglobin. Positive regulation is denoted by arrows, and negative regulation is indicated by perpendicular lines.

proposed that *Ba. subtilis* senses NO and responds through activation of ResDE-dependent transcription (156). In this context, it has been recently discovered that the NO-sensitive transcription factor NsrR is involved in ResDE-dependent gene regulation (156). In *Ba. subtilis*, NsrR represses transcription of the Nir (*nasDEF*) genes that are under positive control of ResDE (Fig. 16). Derepression is achieved by reaction of NO with NsrR. Anaerobically purified *Ba. subtilis* NsrR (BsNsrR) bears a [4Fe-4S] cluster that, on exposure to NO, forms dinitrosyl iron complexes (263). [4Fe-4S]-NsrR binds around the -35 element of the *nasD* promoter with much higher affinity than apo-NsrR, and this binding is sensitive to NO. RNAP and phosphorylated ResD make a ternary complex at the *nasD* promoter, and NsrR dissociates the preformed ternary complex (123).

VI. Concluding Remarks

Genetic, biochemical, and transcriptomic analyses have revealed that in many bacterial species, the adaptation of respiratory metabolism to changing environments with different oxygen conditions is controlled by Fnr-type transcriptional regulators that directly sense O₂ or by the Fnr paralog, FixK; which senses O₂ through the FixLJ two-component system. In addition to external oxygen concentration, other signals such as redox changes can regulate the expression of genes involved in respiration. Two-component regulatory systems (e.g., ArcBA, RegBA/PrrBA, RoxSR, RegSR, ResDE, and ActSR) from different bacterial species integrate the response to multiple redox signals. The redox state of the membrane-localized UQ/ubiquinol pool as well as the redox state of the conserved cysteine²⁶⁵ were considered as major indicators of redox variations detected by these regulatory systems. Recently, it has been proposed that both signals are able of functioning independently (259). In addition, a third model has been proposed in *Rho. sphaeroides* and *Pseudomonas*, where redox changes are sensed by the terminal oxidase *cbb*₃,

which modulates PrrB and RoxS kinase/phosphatase activity (47, 122, 152). Those redox-dependent responses are complex, as recent reports have proposed that the activity of the sensor kinase ArcA from *E. coli* can also be modulated by the redox state of the MK pool (18) or the presence of fermentation products (208). Particularly interesting is the case for the Gram-positive bacteria such as *Ba. subtilis*, where, in addition to the Fnr-type and the ResDE two-component system, the respiratory shift from oxic to microoxic conditions is also controlled by the repressor Rex, whose activity is influenced by the balance of the NADH/NAD⁺ pool in the cell (31, 92).

The respiratory shift in response to changes in oxygen concentration becomes even more sophisticated in bacterial species that use nitrate as respiratory substrate through denitrification or ammonification pathways. This shift is controlled not only by oxygen limitation (FixL and Fnr-type), but also by the presence of nitrate/nitrite (NarXL, NarQP, and NarR) or/and NO (NnrR, NorR, and NsrR). Furthermore, recent findings have shown that redox-responsive two-component regulatory systems (e.g., PrrBA, ActSR, RegSR, and ResDE) are also involved in the regulation of genes needed for anaerobic nitrate respiration. Intriguingly, *Ba. subtilis* ResDE requires anaerobiosis and the presence of NO to activate transcription of nitrate respiration genes (151, 155, 156). In this context, it has been recently discovered that the NO-sensitive transcription factor NsrR plays a role in ResDE-dependent gene regulation (123, 156). However, in Gram negative bacteria, the involvement of such NO-sensitive repressor in regulation of denitrification genes by others redox-responsive regulators (PrrBA, ActSR, RegSR, etc) is at the moment unknown.

In summary, it is clear that bacteria adopt a range of strategies to monitor the redox cellular state to coordinate the respiratory shift from oxic to anoxic environments. This control is based on a complex regulatory network where different regulators responding to diverse redox signals interact and control expression of target genes. The study and comparison

of regulatory pathways in different bacterial groups is very useful to better understand how bacteria integrate different signals and how the regulatory systems respond to these signals. In spite of their diversity, regulatory networks from different bacteria share common proteins as well as their complexity. With the advent of the post-genomic era, and the move toward more predictive biology, a better understanding of how these complex regulatory circuits interact to integrate transcriptional responses to multiple environmental cues will be achieved.

Acknowledgments

The authors would like to thank all of their colleagues who have contributed to the research arising from their laboratories that is discussed in this article and also Junta de Andalucía (grant RMN-4746 partially funded by FEDER), CYTED (grant 107PICO312), and Ministerio de Ciencia e Innovación (grants AGL2010-18607 and 200940I116 partially funded by FEDER) for financial support. Support from the Junta de Andalucía (BIO-275) is also acknowledged. D.J.R. thanks the Royal Society and Wolfson Foundation for a Merit Award Fellowship and the Biotechnology and Biological Research Council for financial support for N-cycle research.

References

1. Alvarez-Ortega C and Harwood CS. Responses of *Pseudomonas aeruginosa* to low oxygen indicate that growth in the cystic fibrosis lung is by aerobic respiration. *Mol Microbiol* 65: 153–165, 2007.
2. Alvarez AF and Georgellis D. *In vitro* and *in vivo* analysis of the ArcB/A redox signaling pathway. *Methods Enzymol* 471: 205–228, 2010.
3. Antelmann H and Helmann JD. Thiol-based redox switches and gene regulation. *Antioxid Redox Signal* 14: 1049–1063, 2011.
4. Arai H, Kodama T, and Igarashi Y. Cascade regulation of the two CRP/FNR-related transcriptional regulators (ANR and DNR) and the denitrification enzymes in *Pseudomonas aeruginosa*. *Mol Microbiol* 25: 1141–1148, 1997.
5. Arnoux P, Sabaty M, Alric J, Frangioni B, Guigliarelli B, Adriano JM, and Pignol D. Structural and redox plasticity in the heterodimeric periplasmic nitrate reductase. *Nat Struct Biol* 10: 928–934, 2003.
6. Azarkina N, Siletsky S, Borisov V, von Wachenfeldt C, Hederstedt L, and Konstantinov AA. A cytochrome *bb'*-type quinol oxidase in *Bacillus subtilis* strain 168. *J Biol Chem* 274: 32810–32817, 1999.
7. Badrick AC, Hamilton AJ, Bernhardt PV, Jones CE, Kappler U, Jennings MP, and McEwan AG. PrrC, a Sco homologue from *Rhodobacter sphaeroides*, possesses thiol-disulfide oxidoreductase activity. *FEBS Lett* 581: 4663–4667, 2007.
8. Baek SH, Hartsock A, and Shapleigh JP. *Agrobacterium tumefaciens* C58 uses ActR and FnrN to control *nirK* and *nor* expression. *J Bacteriol* 190: 78–86, 2008.
9. Baek SH and Shapleigh JP. Expression of nitrite and nitric oxide reductases in free-living and plant-associated *Agrobacterium tumefaciens* C58 cells. *Appl Environ Microbiol* 71: 4427–4436, 2005.
10. Baggs EM and Philippot L. Nitrous oxide production in the terrestrial environment. In: *Nitrogen Cycling in Bacteria*, edited by Moir JWB. Norkfolk, UK: Caister Academic Press, 2011.
11. Baker SC, Ferguson SJ, Ludwig B, Page MD, Richter OM, and van Spanning RJ. Molecular genetics of the genus *Paracoccus*: metabolically versatile bacteria with bioenergetic flexibility. *Microbiol Mol Biol Rev* 62: 1046–1078, 1998.
12. Balatri E, Banci L, Bertini I, Cantini F, and Ciofi-Baffoni S. Solution structure of Sco1: a thioredoxin-like protein involved in cytochrome *c* oxidase assembly. *Structure* 11: 1431–1443, 2003.
13. Bates BC. *Climate Change and Water*, edited by IPCC Secretariat 2008. <http://www.ipcc.ch/meetings/session28/doc13.pdf>
14. Bauer CE, Elsen S, and Bird TH. Mechanisms for redox control of gene expression. *Annu Rev Microbiol* 53: 495–523, 1999.
15. Bauer E, Kaspar T, Fischer HM, and Hennecke H. Expression of the *fixR-nifA* operon in *Bradyrhizobium japonicum* depends on a new response regulator, RegR. *J Bacteriol* 180: 3853–3863, 1998.
16. Bedmar EJ, Mesa S, Velasco L, Robles EF, Bonnard N, and Delgado MJ. Desnitrificación en *Bradyrhizobium japonicum*: Un misterio resuelto. In: *Avances en el Metabolismo del Nitrógeno: De los Microorganismos a las Plantas*, edited by Vega JM, Márquez AJ, Vígara AJ, Vélchez C, and Garbayo I. Huelva, Spain: Servicio de Publicaciones de la Universidad de Huelva, 2005, pp. 35–44.
17. Bedzyk L, Wang T, and Ye RW. The periplasmic nitrate reductase in *Pseudomonas* sp. strain G-179 catalyzes the first step of denitrification. *J Bacteriol* 181: 2802–2806, 1999.
18. Bekker M, Alexeeva S, Laan W, Sawers G, Teixeira de Mattos J, and Hellingwerf K. The ArcBA two-component system of *Escherichia coli* is regulated by the redox state of both the ubiquinone and the menaquinone pool. *J Bacteriol* 192: 746–754, 2010.
19. Belevich I, Borisov VB, Konstantinov AA, and Verkhovsky MI. Oxygenated complex of cytochrome *bd* from *Escherichia coli*: stability and photolability. *FEBS Lett* 579: 4567–4570, 2005.
20. Bell LC, Richardson DJ, and Ferguson SJ. Periplasmic and membrane-bound respiratory nitrate reductases in *Thiosphaera pantotropha*. The periplasmic enzyme catalyzes the first step in aerobic denitrification. *FEBS Lett* 265: 85–87, 1990.
21. Bergaust L, Shapleigh J, Frostegard A, and Bakken L. Transcription and activities of NO_x reductases in *Agrobacterium tumefaciens*: the influence of nitrate, nitrite and oxygen availability. *Environ Microbiol* 10: 3070–3081, 2008.
22. Berks BC, Baratta D, Richardson J, and Ferguson SJ. Purification and characterization of a nitrous oxide reductase from *Thiosphaera pantotropha*. Implications for the mechanism of aerobic nitrous oxide reduction. *Eur J Biochem* 212: 467–476, 1993.
23. Berks BC, Page MD, Richardson DJ, Reilly A, Cavill A, Outen F, and Ferguson SJ. Sequence analysis of subunits of the membrane-bound nitrate reductase from a denitrifying bacterium: the integral membrane subunit provides a prototype for the dihaem electron-carrying arm of a redox loop. *Mol Microbiol* 15: 319–331, 1995.
24. Bertero MG, Rothery RA, Palak M, Hou C, Lim D, Blasco F, Weiner JH, and Strynadka NC. Insights into the respiratory electron transfer pathway from the structure of nitrate reductase A. *Nat Struct Biol* 10: 681–687, 2003.
25. Binnerup SJ and Sorensen J. Nitrate and nitrite microgradients in barley rhizosphere as detected by a highly sensitive denitrification bioassay. *Appl Environ Microbiol* 58: 2375–2380, 1992.

26. Blasco F, Iobbi C, Ratouchniak J, Bonnefoy V, and Chippaux M. Nitrate reductases of *Escherichia coli*: sequence of the second nitrate reductase and comparison with that encoded by the *narGHJI* operon. *Mol Gen Genet* 222: 104–111, 1990.
27. Blasco F, Pommier J, Augier V, Chippaux M, and Giordano G. Involvement of the *narJ* or *narW* gene product in the formation of active nitrate reductase in *Escherichia coli*. *Mol Microbiol* 6: 221–230, 1992.
28. Bobik C, Meilhoc E, and Batut J. FixJ: a major regulator of the oxygen limitation response and late symbiotic functions of *Sinorhizobium meliloti*. *J Bacteriol* 188: 4890–4902, 2006.
29. Borisov VB, Gennis RB, Hemp J, and Verkhovsky MI. The cytochrome *bd* respiratory oxygen reductases. *Biochim Biophys Acta* 1807: 1398–1413, 2011.
30. Bouchal P, Struharova I, Budinska E, Sedo O, Vyhldalova T, Zdrahal Z, van Spanning R, and Kucera I. Unraveling an FNR based regulatory circuit in *Paracoccus denitrificans* using a proteomics-based approach. *Biochim Biophys Acta* 1804: 1350–1358, 2010.
31. Brekasis D and Paget MS. A novel sensor of NADH/NAD⁺ redox poise in *Streptomyces coelicolor* A3(2). *EMBO J* 22: 4856–4865, 2003.
32. Brondijk TH, Fiegen D, Richardson DJ, and Cole JA. Roles of NapF, NapG and NapH, subunits of the *Escherichia coli* periplasmic nitrate reductase, in ubiquinol oxidation. *Mol Microbiol* 44: 245–255, 2002.
33. Brondijk TH, Nilavongse A, Filenko N, Richardson DJ, and Cole JA. NapGH components of the periplasmic nitrate reductase of *Escherichia coli* K-12: location, topology and physiological roles in quinol oxidation and redox balancing. *Biochem J* 379: 47–55, 2004.
34. Bruscella P, Eraso JM, Roh JH, and Kaplan S. The use of chromatin immunoprecipitation to define PpsR binding activity in *Rhodobacter sphaeroides* 2.4.1. *J Bacteriol* 190: 6817–6828, 2008.
35. Bueno E, Bedmar EJ, Richardson DJ, and Delgado MJ. Role of *Bradyrhizobium japonicum* cytochrome *c₅₅₀* in nitrite and nitrate respiration. *FEMS Microbiol Lett* 279: 188–194, 2008.
36. Bueno E, Mesa S, Sanchez C, Bedmar EJ, and Delgado MJ. NifA is required for maximal expression of denitrification genes in *Bradyrhizobium japonicum*. *Environ Microbiol* 12: 393–400, 2010.
37. Bueno E, Richardson DJ, Bedmar EJ, and Delgado MJ. Expression of *Bradyrhizobium japonicum cbb₃* terminal oxidase under denitrifying conditions is subjected to redox control. *FEMS Microbiol Lett* 298: 20–28, 2009.
38. Bühler D, Rossmann R, Landolt S, Balsiger S, Fischer HM, and Hennecke H. Disparate pathways for the biogenesis of cytochrome oxidases in *Bradyrhizobium japonicum*. *J Biol Chem* 285: 15704–15713, 2010.
39. Büsch A, Strube K, Friedrich B, and Cramm R. Transcriptional regulation of nitric oxide reduction in *Ralstonia eutropha* H16. *Biochem Soc Trans* 33: 193–194, 2005.
40. Buschmann S, Warkentin E, Xie H, Langer JD, Ermler U, and Michel H. The structure of *cbb₃* cytochrome oxidase provides insights into proton pumping. *Science* 329: 327–330, 2010.
41. Butland G, Spiro S, Watmough NJ, and Richardson DJ. Two conserved glutamates in the bacterial nitric oxide reductase are essential for activity but not assembly of the enzyme. *J Bacteriol* 183: 189–199, 2001.
42. Carter JP, Hsaio YH, Spiro S, and Richardson DJ. Soil and sediment bacteria capable of aerobic nitrate respiration. *Appl Environ Microbiol* 61: 2852–2858, 1995.
43. Cartron ML, Roldan MD, Ferguson SJ, Berks BC, and Richardson DJ. Identification of two domains and distal histidine ligands to the four haems in the bacterial *c*-type cytochrome NapC; the prototype connector between quinol/quinone and periplasmic oxido-reductases. *Biochem J* 368: 425–432, 2002.
44. Cebren A and Garnier J. *Nitrobacter* and *Nitrospira* genera as representatives of nitrite-oxidizing bacteria: detection, quantification and growth along the lower Seine River (France). *Water Res* 39: 4979–4992, 2005.
45. Cole JA and Richardson DJ. Respiration of nitrate and nitrite. In: *EcoSal-Escherichia coli and Salmonella*, edited by Böck A, Curtiss III R, Kaper JB, Karp PD, Neidhardt FC, Nyström T, Slauch JM, and Squires CL. Washington DC: ASM Press, 2008.
46. Comolli JC and Donohue TJ. *Pseudomonas aeruginosa* RoxR, a response regulator related to *Rhodobacter sphaeroides* PrrA, activates expression of the cyanide-insensitive terminal oxidase. *Mol Microbiol* 45: 755–768, 2002.
47. Comolli JC and Donohue TJ. Differences in two *Pseudomonas aeruginosa cbb₃* cytochrome oxidases. *Mol Microbiol* 51: 1193–1203, 2004.
48. Constantinidou C, Hobman JL, Griffiths L, Patel MD, Penn CW, Cole JA, and Overton TW. A reassessment of the FNR regulon and transcriptomic analysis of the effects of nitrate, nitrite, NarXL, and NarQP as *Escherichia coli* K12 adapts from aerobic to anaerobic growth. *J Biol Chem* 281: 4802–4815, 2006.
49. Cooper M, Tavankar GR, and Williams HD. Regulation of expression of the cyanide-insensitive terminal oxidase in *Pseudomonas aeruginosa*. *Microbiology* 149: 1275–1284, 2003.
50. Cosseau C and Batut J. Genomics of the *ccoNOQP*-encoded *cbb₃* oxidase complex in bacteria. *Arch Microbiol* 181: 89–96, 2004.
51. Cotter PA, Melville SB, Albrecht JA, and Gunsalus RP. Aerobic regulation of cytochrome *d* oxidase (*cydAB*) operon expression in *Escherichia coli*: roles of Fnr and ArcA in repression and activation. *Mol Microbiol* 25: 605–615, 1997.
52. Crack J, Green J, and Thomson AJ. Mechanism of oxygen sensing by the bacterial transcription factor fumarate-nitrate reduction (FNR). *J Biol Chem* 279: 9278–9286, 2004.
53. Crack JC, Gaskell AA, Green J, Cheesman MR, Le Brun NE, and Thomson AJ. Influence of the environment on the [4Fe-4S]²⁺ to [2Fe-2S]²⁺ cluster switch in the transcriptional regulator FNR. *J Am Chem Soc* 130: 1749–1758, 2008.
54. Crack JC, Green J, Cheesman MR, Le Brun NE, and Thomson AJ. Superoxide-mediated amplification of the oxygen-induced switch from [4Fe-4S] to [2Fe-2S] clusters in the transcriptional regulator FNR. *Proc Natl Acad Sci U S A* 104: 2092–2097, 2007.
55. Cramm R, Busch A, and Strube K. NO-dependent transcriptional activation of gene expression in *Ralstonia eutropha* H16. *Biochem Soc Trans* 34: 182–184, 2006.
56. Crosson S, McGrath PT, Stephens C, McAdams HH, and Shapiro L. Conserved modular design of an oxygen sensory/signaling network with species-specific output. *Proc Natl Acad Sci U S A* 102: 8018–8123, 2005.
57. Cruz-Ramos H, Crack J, Wu G, Hughes MN, Scott C, Thomson AJ, Green J, and Poole RK. NO sensing by FNR: regulation of the *Escherichia coli* NO-detoxifying flavohaemoglobin, Hmp. *EMBO J* 21: 3235–3244, 2002.
58. Darwin AJ, Ziegelhoffer EC, Kiley PJ, and Stewart V. Fnr, NarP, and NarL regulation of *Escherichia coli* K-12 *napF*

- (periplasmic nitrate reductase) operon transcription *in vitro*. *J Bacteriol* 180: 4192–4198, 1998.
59. de Vries R, Suharti S, and Pouvreau LAM. Nitric oxide reductase: structural variations and catalytic mechanism. In: *Biology of the Nitrogen Cycle*, edited by Bothe H, Ferguson SJ, and Newton WE. The Netherlands: Elsevier, 2007.
 60. de Vries S, Strampraad MJ, Lu S, Moenne-Loccoz P, and Schroder I. Purification and characterization of the $\text{MQH}_2\text{:NO}$ oxidoreductase from the hyperthermophilic archaeon *Pyrobaculum aerophilum*. *J Biol Chem* 278: 35861–35868, 2003.
 61. Delgado MJ, Bedmar EJ, and Downie JA. Genes involved in the formation and assembly of rhizobial cytochromes and their role in symbiotic nitrogen fixation. *Adv Microb Physiol* 40: 191–231, 1998.
 62. Delgado MJ, Bonnard N, Tresierra-Ayala A, Bedmar EJ, and Muller P. The *Bradyrhizobium japonicum* *napEDABC* genes encoding the periplasmic nitrate reductase are essential for nitrate respiration. *Microbiology* 149: 3395–3403, 2003.
 63. Delledonne M, Zeier J, Marocco A, and Lamb C. Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease resistance response. *Proc Natl Acad Sci U S A* 98: 13454–13459, 2001.
 64. Dias JM, Than ME, Humm A, Huber R, Bourenkov GP, Bartunik HD, Bursakov S, Calvete J, Caldeira J, Carneiro C, Moura JJ, Moura I, and Romao MJ. Crystal structure of the first dissimilatory nitrate reductase at 1.9 Å solved by MAD methods. *Structure* 7: 65–79, 1999.
 65. Donohue TJ and Kiley PJ. Bacterial responses to O_2 limitation. In: *Bacterial Stress Responses*, edited by Storz G and Hengge D. Washington, DC: American Society for Microbiology, 2010, pp. 175–189.
 66. Downie JA. Legume haemoglobins: symbiotic nitrogen fixation needs bloody nodules. *Curr Biol* 15: R196–R198, 2005.
 67. Ducluzeau AL, Ouchane S, and Nitschke W. The *cbb₃* oxidases are an ancient innovation of the domain bacteria. *Mol Biol Evol* 25: 1158–1166, 2008.
 68. Dufour YS, Kiley PJ, and Donohue TJ. Reconstruction of the core and extended regulons of global transcription factors. *PLoS Genet* 6: e1001027, 2010.
 69. Ellington MJ, Bhakoo KK, Sawers G, Richardson DJ, and Ferguson SJ. Hierarchy of carbon source selection in *Paracoccus pantotrophus*: strict correlation between reduction state of the carbon substrate and aerobic expression of the *nap* operon. *J Bacteriol* 184: 4767–4774, 2002.
 70. Ellington MJ, Fosdike WL, Sawers RG, Richardson DJ, and Ferguson SJ. Regulation of the *nap* operon encoding the periplasmic nitrate reductase of *Paracoccus pantotrophus*: delineation of DNA sequences required for redox control. *Arch Microbiol* 184: 298–304, 2006.
 71. Ellington MJ, Sawers G, Sears HJ, Spiro S, Richardson DJ, and Ferguson SJ. Characterization of the expression and activity of the periplasmic nitrate reductase of *Paracoccus pantotrophus* in chemostat cultures. *Microbiology* 149: 1533–1540, 2003.
 72. Elsen S, Swem LR, Swem DL, and Bauer CE. RegB/RegA, a highly conserved redox-responding global two-component regulatory system. *Microbiol Mol Biol Rev* 68: 263–279, 2004.
 73. Emmerich R, Hennecke H, and Fischer HM. Evidence for a functional similarity between the two-component regulatory systems RegSR, ActSR, and RegBA (PrrBA) in α -Proteobacteria. *Arch Microbiol* 174: 307–313, 2000.
 74. Emmerich R, Panglungtshang K, Strehler P, Hennecke H, and Fischer HM. Phosphorylation, dephosphorylation and DNA-binding of the *Bradyrhizobium japonicum* RegSR two-component regulatory proteins. *Eur J Biochem* 263: 455–463, 1999.
 75. Eraso JM and Kaplan S. From redox flow to gene regulation: role of the PrrC protein of *Rhodobacter sphaeroides* 2.4.1. *Biochemistry* 39: 2052–2062, 2000.
 76. Eraso JM and Kaplan S. Half-Site DNA sequence and spacing length contributions to PrrA binding to PrrA site 2 of RSP3361 in *Rhodobacter sphaeroides* 2.4.1. *J Bacteriol* 191: 4353–4364, 2009.
 77. Eraso JM, Roh JH, Zeng X, Callister SJ, Lipton MS, and Kaplan S. Role of the global transcriptional regulator PrrA in *Rhodobacter sphaeroides* 2.4.1: combined transcriptome and proteome analysis. *J Bacteriol* 190: 4831–4848, 2008.
 78. Fernandez-Piñar R, Ramos JL, Rodríguez-Hervá JJ, and Espinosa-Urgel M. A two-component regulatory system integrates redox state and population density sensing in *Pseudomonas putida*. *J Bacteriol* 190: 7666–7674, 2008.
 79. Ferrarini A, De Stefano M, Baudouin E, Pucciariello C, Polverari A, Puppo A, and Delledonne M. Expression of *Medicago truncatula* genes responsive to nitric oxide in pathogenic and symbiotic conditions. *Mol Plant-Microbe Interact* 21: 781–790, 2008.
 80. Field SJ, Roldan MD, Marritt SJ, Butt JN, Richardson DJ, and Watmough NJ. Electron transfer to the active site of the bacterial nitric oxide reductase is controlled by ligand binding to heme *b₃*. *Biochim Biophys Acta* 1807: 451–457, 2011.
 81. Field SJ, Thorndycroft FH, Matorin AD, Richardson DJ, and Watmough NJ. The respiratory nitric oxide reductase (NorBC) from *Paracoccus denitrificans*. *Methods Enzymol* 437: 79–101, 2008.
 82. Filenko N, Spiro S, Browning DF, Squire D, Overton TW, Cole J, and Constantinidou C. The NsrR regulon of *Escherichia coli* K-12 includes genes encoding the hybrid cluster protein and the periplasmic, respiratory nitrite reductase. *J Bacteriol* 189: 4410–4417, 2007.
 83. Fischer HM. Genetic regulation of nitrogen fixation in rhizobia. *Microbiol Rev* 58: 352–386, 1994.
 84. Fleischhacker AS and Kiley PJ. Iron-containing transcription factors and their roles as sensors. *Curr Opin Chem Biol* 15: 335–341, 2011.
 85. Flock U, Lachmann P, Reimann J, Watmough NJ, and Adelroth P. Exploring the terminal region of the proton pathway in the bacterial nitric oxide reductase. *J Inorg Biochem* 103: 845–850, 2009.
 86. Flock U, Thorndycroft FH, Matorin AD, Richardson DJ, Watmough NJ, and Adelroth P. Defining the proton entry point in the bacterial respiratory nitric-oxide reductase. *J Biol Chem* 283: 3839–3845, 2008.
 87. Garcia-Horsman JA, Barquera B, Rumbley J, Ma J, and Gennis RB. The superfamily of heme-copper respiratory oxidases. *J Bacteriol* 176: 5587–5600, 1994.
 88. Gates AJ, Richardson DJ, and Butt JN. Voltammetric characterization of the aerobic energy-dissipating nitrate reductase of *Paracoccus pantotrophus*: exploring the activity of a redox-balancing enzyme as a function of electrochemical potential. *Biochem J* 409: 159–168, 2008.
 89. Gavira M, Roldan MD, Castillo F, and Moreno-Vivian C. Regulation of *nap* gene expression and periplasmic nitrate reductase activity in the phototrophic bacterium *Rhodobacter sphaeroides* DSM158. *J Bacteriol* 184: 1693–1702, 2002.

90. Geng H, Nakano S, and Nakano MM. Transcriptional activation by *Bacillus subtilis* ResD: tandem binding to target elements and phosphorylation-dependent and -independent transcriptional activation. *J Bacteriol* 186: 2028–2037, 2004.
91. Geng H, Zhu Y, Mullen K, Zuber CS, and Nakano MM. Characterization of ResDE-dependent *fnr* transcription in *Bacillus subtilis*. *J Bacteriol* 189: 1745–1755, 2007.
92. Geng H, Zuber P, and Nakano MM. Regulation of respiratory genes by ResD-ResE signal transduction system in *Bacillus subtilis*. *Methods Enzymol* 422: 448–464, 2007.
93. Giardina G, Castiglione N, Caruso M, Cutruzzola F, and Rinaldo S. The *Pseudomonas aeruginosa* DNR transcription factor: light and shade of nitric oxide-sensing mechanisms. *Biochem Soc Trans* 39: 294–298, 2011.
94. Giardina G, Rinaldo S, Johnson KA, Di Matteo A, Brunori M, and Cutruzzola F. NO sensing in *Pseudomonas aeruginosa*: structure of the transcriptional regulator DNR. *J Mol Biol* 378: 1002–1015, 2008.
95. Gilberthorpe NJ and Poole RK. Nitric oxide homeostasis in *Salmonella typhimurium*: roles of respiratory nitrate reductase and flavohemoglobin. *J Biol Chem* 283: 11146–11154, 2008.
96. Gilles-Gonzalez MA and Gonzalez G. Heme-based sensors: defining characteristics, recent developments, and regulatory hypotheses. *J Inorg Biochem* 99: 1–22, 2005.
97. Goddard AD, Moir JW, Richardson DJ, and Ferguson SJ. Interdependence of two NarK domains in a fused nitrate/nitrite transporter. *Mol Microbiol* 70: 667–681, 2008.
98. Gonzalez PJ, Correia C, Moura I, Brondino CD, and Moura JJ. Bacterial nitrate reductases: Molecular and biological aspects of nitrate reduction. *J Inorg Biochem* 100: 1015–1023, 2006.
99. Govantes F, Orjalo AV, and Gunsalus RP. Interplay between three global regulatory proteins mediates oxygen regulation of the *Escherichia coli* cytochrome *d* oxidase (*cydAB*) operon. *Mol Microbiol* 38: 1061–1073, 2000.
100. Granados-Baeza MJ, Gomez-Hernandez N, Mora Y, Delgado MJ, Romero D, and Girard L. Novel reiterated *Fnr*-type proteins control the production of the symbiotic terminal oxidase *cbb₃* in *Rhizobium etli* CFN42. *Mol Plant-Microbe Interact* 20: 1241–1249, 2007.
101. Green J, Crack JC, Thomson AJ, and LeBrun NE. Bacterial sensors of oxygen. *Curr Opin Microbiol* 12: 145–151, 2009.
102. Green J and Paget MS. Bacterial redox sensors. *Nat Rev Microbiol* 2: 954–966, 2004.
103. Green J, Scott C, and Guest JR. Functional versatility in the CRP-FNR superfamily of transcription factors: FNR and FLP. *Adv Microb Physiol* 44: 1–34, 2001.
104. Gyan S, Shiohira Y, Sato I, Takeuchi M, and Sato T. Regulatory loop between redox sensing of the NADH/NAD⁺ ratio by Rex (YdiH) and oxidation of NADH by NADH dehydrogenase Ndh in *Bacillus subtilis*. *J Bacteriol* 188: 7062–7071, 2006.
105. Hanke A and Strous M. Climate, fertilization, and the nitrogen cycle *J Cosmol* 8: 1838–1845, 2010.
106. Happ HN, Braatsch S, Broschek V, Osterloh L, and Klug G. Light-dependent regulation of photosynthesis genes in *Rhodobacter sphaeroides* 2.4.1 is coordinately controlled by photosynthetic electron transport via the PrrBA two-component system and the photoreceptor AppA. *Mol Microbiol* 58: 903–914, 2005.
107. Hartsock A and Shapleigh JP. Identification, functional studies, and genomic comparisons of new members of the NnrR regulon in *Rhodobacter sphaeroides*. *J Bacteriol* 192: 903–911, 2010.
108. Hauser F, Pessi G, Friberg M, Weber C, Rusca N, Lindemann A, Fischer HM, and Henneke H. Dissection of the *Bradyrhizobium japonicum* NifA+s⁵⁴ regulon, and identification of a ferredoxin gene (*fdxN*) for symbiotic nitrogen fixation. *Mol Genet Genomics* 278: 255–271, 2007.
109. Hayatsu M, Tago K, and Saito M. Various players in the nitrogen cycle: diversity and functions of the microorganisms involved in nitrification and denitrification. *Soil Sci Plant Nutr* 54: 33–45, 2008.
110. Hendriks J, Oubrie A, Castresana J, Urbani A, Gemeinhardt S, and Saraste M. Nitric oxide reductases in bacteria. *Biochim Biophys Acta* 1459: 266–273, 2000.
111. Heurlier K, Thomson MJ, Aziz N, and Moir JW. The nitric oxide (NO)-sensing repressor NsrR of *Neisseria meningitidis* has a compact regulon of genes involved in NO synthesis and detoxification. *J Bacteriol* 190: 2488–2495, 2008.
112. Hino T, Matsumoto Y, Nagano S, Sugimoto H, Fukumori Y, Murata T, Iwata S, and Shiro Y. Structural basis of biological N₂O generation by bacterial nitric oxide reductase. *Science* 330: 1666–1670, 2010.
113. Isabella VM, Lapek JD Jr., Kennedy EM, and Clark VL. Functional analysis of NsrR, a nitric oxide-sensing Rrf2 repressor in *Neisseria gonorrhoeae*. *Mol Microbiol* 71: 227–239, 2009.
114. Jepson BJ, Marietou A, Mohan S, Cole JA, Butler CS, and Richardson DJ. Evolution of the soluble nitrate reductase: defining the monomeric periplasmic nitrate reductase subgroup. *Biochem Soc Trans* 34: 122–126, 2006.
115. Jepson BJ, Mohan S, Clarke TA, Gates AJ, Cole JA, Butler CS, Butt JN, Hemmings AM, and Richardson DJ. Spectropotentiometric and structural analysis of the periplasmic nitrate reductase from *Escherichia coli*. *J Biol Chem* 282: 6425–6437, 2007.
116. Jones KM, Kobayashi H, Davies BW, Taga ME, and Walker GC. How rhizobial symbionts invade plants: the *Sinorhizobium-Medicago* model. *Nat Rev Microbiol* 5: 619–633, 2007.
117. Jones MR, Richardson DJ, McEwan AG, Jackson JB, and Ferguson SJ. *In vivo* redox poisoning of the cyclic electron transport system of *Rhodobacter capsulatus* and the effects of the auxiliary oxidants, nitrate, nitrous oxide and trimethylamine N-oxide, as revealed by multiple short flash excitation *Biochem Biophys Acta* 1017: 209–216, 1990.
118. Jormakka M, Richardson D, Byrne B, and Iwata S. Architecture of NarGH reveals a structural classification of Mo-bisMGD enzymes. *Structure* 12: 95–104, 2004.
119. Justino MC, Vicente JB, Teixeira M, and Saraiva LM. New genes implicated in the protection of anaerobically grown *Escherichia coli* against nitric oxide. *J Biol Chem* 280: 2636–2643, 2005.
120. Kaplan S, Eraso J, and Roh JH. Interacting regulatory networks in the facultative photosynthetic bacterium, *Rhodobacter sphaeroides* 2.4.1. *Biochem Soc Trans* 33: 51–55, 2005.
121. Kawakami T, Kuroki M, Ishii M, Igarashi Y, and Arai H. Differential expression of multiple terminal oxidases for aerobic respiration in *Pseudomonas aeruginosa*. *Environ Microbiol* 12: 1399–1412, 2010.
122. Kim YJ, Ko IJ, Lee JM, Kang HY, Kim YM, Kaplan S, and Oh JI. Dominant role of the *cbb₃* oxidase in regulation of photosynthesis gene expression through the PrrBA system in *Rhodobacter sphaeroides* 2.4.1. *J Bacteriol* 189: 5617–5625, 2007.
123. Kommineni S, Yukl E, Hayashi T, Delepine J, Geng H, Moenne-Loccoz P, and Nakano MM. Nitric oxide-sensitive and -insensitive interaction of *Bacillus subtilis* NsrR with a

- ResDE-controlled promoter. *Mol Microbiol* 78: 1280–1293, 2010.
124. Körner H, Sofia HJ, and Zumft WG. Phylogeny of the bacterial superfamily of Crp-Fnr transcription regulators: exploiting the metabolic spectrum by controlling alternative gene programs. *FEMS Microbiol Rev* 27: 559–592, 2003.
 125. Kraft B, Strous M, and Tegetmeyer HE. Microbial nitrate respiration—Genes, enzymes and environmental distribution. *J Biotechnol* 155: 104–117, 2011.
 126. Kulajta C, Thumfart JO, Haid S, Daldal F, and Koch HG. Multi-step assembly pathway of the *cbb₃*-type cytochrome *c* oxidase complex. *J Mol Biol* 355: 989–1004, 2006.
 127. Kullik I, Fritsche S, Knobel H, Sanjuan J, Hennecke H, and Fischer HM. *Bradyrhizobium japonicum* has two differentially regulated, functional homologs of the *s⁵⁴* gene (*rpoN*). *J Bacteriol* 173: 1125–1138, 1991.
 128. Laguri C, Phillips-Jones MK, and Williamson MP. Solution structure and DNA binding of the effector domain from the global regulator PrrA (RegA) from *Rhodobacter sphaeroides*: insights into DNA binding specificity. *Nucleic Acids Res* 31: 6778–6787, 2003.
 129. Laguri C, Stenzel RA, Donohue TJ, Phillips-Jones MK, and Williamson MP. Activation of the global gene regulator PrrA (RegA) from *Rhodobacter sphaeroides*. *Biochemistry* 45: 7872–7881, 2006.
 130. Laratta WP, Choi PS, Tosques IE, and Shapleigh JP. Involvement of the PrrB/PrrA two-component system in nitrite respiration in *Rhodobacter sphaeroides* 2.4.3: evidence for transcriptional regulation. *J Bacteriol* 184: 3521–3529, 2002.
 131. Laratta WP and Shapleigh JP. Site-directed mutagenesis of NnrR: a transcriptional regulator of nitrite and nitric oxide reductase in *Rhodobacter sphaeroides*. *FEMS Microbiol Lett* 229: 173–178, 2003.
 132. Lazazzera BA, Beinert H, Khoroshilova N, Kennedy MC, and Kiley PJ. DNA binding and dimerization of the Fe-S-containing FNR protein from *Escherichia coli* are regulated by oxygen. *J Biol Chem* 271: 2762–2768, 1996.
 133. Lindemann A, Koch M, Pessi G, Muller AJ, Balsiger S, Hennecke H, and Fischer HM. Host-specific symbiotic requirement of BdeAB, a RegR-controlled RND-type efflux system in *Bradyrhizobium japonicum*. *FEMS Microbiol Lett* 312: 184–191, 2010.
 134. Lindemann A, Moser A, Pessi G, Hauser F, Friberg M, Hennecke H, and Fischer HM. New target genes controlled by the *Bradyrhizobium japonicum* two-component regulatory system RegSR. *J Bacteriol* 189: 8928–8943, 2007.
 135. Liu HP, Takio S, Satoh T, and Yamamoto I. Involvement in denitrification of the *napKEFDABC* genes encoding the periplasmic nitrate reductase system in the denitrifying phototrophic bacterium *Rhodobacter sphaeroides* f. sp. *denitrificans*. *Biosci Biotechnol Biochem* 63: 530–536, 1999.
 136. Liu X and De Wulf P. Probing the ArcA-P modulon of *Escherichia coli* by whole genome transcriptional analysis and sequence recognition profiling. *J Biol Chem* 279: 12588–12597, 2004.
 137. Malpica R, Franco B, Rodriguez C, Kwon O, and Georgellis D. Identification of a quinone-sensitive redox switch in the ArcB sensor kinase. *Proc Natl Acad Sci U S A* 101: 13318–13323, 2004.
 138. Malpica R, Sandoval GR, Rodriguez C, Franco B, and Georgellis D. Signaling by the arc two-component system provides a link between the redox state of the quinone pool and gene expression. *Antioxid Redox Signal* 8: 781–795, 2006.
 139. Mao L, Mackenzie C, Roh JH, Eraso JM, Kaplan S, and Resat H. Combining microarray and genomic data to predict DNA binding motifs. *Microbiology* 151: 3197–3213, 2005.
 140. Marietou A, Richardson D, Cole J, and Mohan S. Nitrate reduction by *Desulfovibrio desulfuricans*: a periplasmic nitrate reductase system that lacks NapB, but includes a unique tetraheme *c*-type cytochrome, NapM. *FEMS Microbiol Lett* 248: 217–225, 2005.
 141. Martinez-Espinosa RM, Cole JA, Richardson DJ, and Watmough NJ. Enzymology and ecology of the nitrogen cycle. *Biochem Soc Trans* 39: 175–178, 2011.
 142. Martínez-Espinosa RM, Dridge EJ, Bonete MJ, Butt JN, Butler CS, Sargent F, and Richardson DJ. Look on the positive side! The orientation, identification and bioenergetics of ‘Archaeal’ membrane-bound nitrate reductases. *FEMS Microbiol Lett* 276: 129–139, 2007.
 143. Matsuda Y, Inamori K, Osaki T, Eguchi A, Watanabe A, Kawabata S, Iba K, and Arata H. Nitric oxide-reductase homologue that contains a copper atom and has cytochrome *c*-oxidase activity from an aerobic phototrophic bacterium *Roseobacter denitrificans*. *J Biochem* 131: 791–800, 2002.
 144. Mesa S, Bedmar EJ, Chanfon A, Hennecke H, and Fischer HM. *Bradyrhizobium japonicum* NnrR, a denitrification regulator, expands the FixLJ-FixK₂ regulatory cascade. *J Bacteriol* 185: 3978–3982, 2003.
 145. Mesa S, Hauser F, Friberg M, Malaguti E, Fischer HM, and Hennecke H. Comprehensive assessment of the regulons controlled by the FixLJ-FixK₂-FixK₁ cascade in *Bradyrhizobium japonicum*. *J Bacteriol* 190: 6568–6579, 2008.
 146. Mesa S, Hennecke H, and Fischer HM. A multitude of CRP/FNR-like transcription proteins in *Bradyrhizobium japonicum*. *Biochem Soc Trans* 34: 156–159, 2006.
 147. Mesa S, Reutimann L, Fischer HM, and Hennecke H. Posttranslational control of transcription factor FixK₂, a key regulator for the *Bradyrhizobium japonicum*-soybean symbiosis. *Proc Natl Acad Sci U S A* 106: 21860–21865, 2009.
 148. Mesa S, Ucurum Z, Hennecke H, and Fischer HM. Transcription activation *in vitro* by the *Bradyrhizobium japonicum* regulatory protein FixK₂. *J Bacteriol* 187: 3329–3338, 2005.
 149. Mills PC, Rowley G, Spiro S, Hinton JC, and Richardson DJ. A combination of cytochrome *c* nitrite reductase (NrfA) and flavorubredoxin (NorV) protects *Salmonella enterica* serovar Typhimurium against killing by NO in anoxic environments. *Microbiology* 154: 1218–1228, 2008.
 150. Minchin FR, James EK, and Becana M. Oxygen diffusion, production of reactive oxygen and nitrogen species, and antioxidants in legume nodules. In: *Nitrogen-Fixing Leguminous Symbioses*, edited by Dilworth MJ, James EK, Sprent JI, and Newton WE. New York: Springer Science, 2008, pp. 321–362.
 151. Moore CM, Nakano MM, Wang T, Ye RW, and Helmann JD. Response of *Bacillus subtilis* to nitric oxide and the nitrosating agent sodium nitroprusside. *J Bacteriol* 186: 4655–4664, 2004.
 152. Morales G, Ugidos A, and Rojo F. Inactivation of the *Pseudomonas putida* cytochrome *o* ubiquinol oxidase leads to a significant change in the transcriptome and to increased expression of the CIO and *cbb₃*-1 terminal oxidases. *Environ Microbiol* 8: 1764–1774, 2006.
 153. Moreno-Vivián C, Cabello P, Martinez-Luque M, Blasco R, and Castillo F. Prokaryotic nitrate reduction: molecular properties and functional distinction among bacterial nitrate reductases. *J Bacteriol* 181: 6573–6584, 1999.

154. Moreno-Vivián C and Ferguson SJ. Definition and distinction between assimilatory, dissimilatory and respiratory pathways. *Mol Microbiol* 29: 664–666, 1998.
155. Nakano MM. Induction of ResDE-dependent gene expression in *Bacillus subtilis* in response to nitric oxide and nitrosative stress. *J Bacteriol* 184: 1783–1787, 2002.
156. Nakano MM, Geng H, Nakano S, and Kobayashi K. The nitric oxide-responsive regulator NsrR controls ResDE-dependent gene expression. *J Bacteriol* 188: 5878–5887, 2006.
157. Nakano MM, Hoffmann T, Zhu Y, and Jahn D. Nitrogen and oxygen regulation of *Bacillus subtilis* *nasDEF* encoding NADH-dependent nitrite reductase by TnrA and ResDE. *J Bacteriol* 180: 5344–5350, 1998.
158. Nakano MM and Zuber P. Anaerobic growth of a “strict aerobe” (*Bacillus subtilis*). *Annu Rev Microbiol* 52: 165–190, 1998.
159. Nakano MM, Zuber P, Glaser P, Danchin A, and Hulett FM. Two-component regulatory proteins ResD-ResE are required for transcriptional activation of *fnr* upon oxygen limitation in *Bacillus subtilis*. *J Bacteriol* 178: 3796–3802, 1996.
160. Nellen-Anthamatten D, Rossi P, Preisig O, Kullik I, Babst M, Fischer HM, and Hennecke H. *Bradyrhizobium japonicum* FixK₂, a crucial distributor in the FixLJ-dependent regulatory cascade for control of genes inducible by low oxygen levels. *J Bacteriol* 180: 5251–5255, 1998.
161. Nelson KE, Weinel C, Paulsen IT, Dodson RJ, Hilbert H, Martins dos Santos VA, Fouts DE, Gill SR, Pop M, Holmes M, Brinkac L, Beanan M, DeBoy RT, Daugherty S, Kolonay J, Madupu R, Nelson W, White O, Peterson J, Khouri H, Hance I, Chris Lee P, Holtzapple E, Scanlan D, Tran K, Moazzes A, Utterback T, Rizzo M, Lee K, Kosack D, Moestl D, Wedler H, Lauber J, Stjepandic D, Hoheisel J, Straetz M, Heim S, Kiewitz C, Eisen JA, Timmis KN, Dusterhoft A, Tummler B, and Fraser CM. Complete genome sequence and comparative analysis of the metabolically versatile *Pseudomonas putida* KT2440. *Environ Microbiol* 4: 799–808, 2002.
162. Nienaber A, Huber A, Gottfert M, Hennecke H, and Fischer HM. Three new NifA-regulated genes in the *Bradyrhizobium japonicum* symbiotic gene region discovered by competitive DNA-RNA hybridization. *J Bacteriol* 182: 1472–1480, 2000.
163. Nikel PI, Zhu J, San KY, Mendez BS, and Bennett GN. Metabolic flux analysis of *Escherichia coli* *creB* and *arcA* mutants reveals shared control of carbon catabolism under microaerobic growth conditions. *J Bacteriol* 191: 5538–5548, 2009.
164. Nilavongse A, Brondijk TH, Overton TW, Richardson DJ, Leach ER, and Cole JA. The NapF protein of the *Escherichia coli* periplasmic nitrate reductase system: demonstration of a cytoplasmic location and interaction with the catalytic subunit, NapA. *Microbiology* 152: 3227–3237, 2006.
165. Oh JI and Kaplan S. The *cbb₃* terminal oxidase of *Rhodobacter sphaeroides* 2.4.1: structural and functional implications for the regulation of spectral complex formation. *Biochemistry* 38: 2688–2696, 1999.
166. Oh JI and Kaplan S. Redox signaling: globalization of gene expression. *EMBO J* 19: 4237–4247, 2000.
167. Oh JI, Ko IJ, and Kaplan S. The default state of the membrane-localized histidine kinase PrrB of *Rhodobacter sphaeroides* 2.4.1 is in the kinase-positive mode. *J Bacteriol* 183: 6807–6814, 2001.
168. Oh JI, Ko IJ, and Kaplan S. Reconstitution of the *Rhodobacter sphaeroides* *cbb₃*-PrrBA signal transduction pathway *in vitro*. *Biochemistry* 43: 7915–7923, 2004.
169. Oldroyd GE and Downie JA. Coordinating nodule morphogenesis with rhizobial infection in legumes. *Annu Rev Plant Biol* 59: 519–546, 2008.
170. Olmo-Mira MF, Gavira M, Richardson DJ, Castillo F, Moreno-Vivian C, and Roldan MD. NapF is a cytoplasmic iron-sulfur protein required for Fe-S cluster assembly in the periplasmic nitrate reductase. *J Biol Chem* 279: 49727–49735, 2004.
171. Overton TW, Whitehead R, Li Y, Snyder LA, Saunders NJ, Smith H, and Cole JA. Coordinated regulation of the *Neisseria gonorrhoeae*-truncated denitrification pathway by the nitric oxide-sensitive repressor, NsrR, and nitrite-insensitive NarQ-NarP. *J Biol Chem* 281: 33115–33126, 2006.
172. Pagels M, Fuchs S, Pane-Farre J, Kohler C, Menschner L, Hecker M, McNamarra PJ, Bauer MC, von Wachenfeldt C, Liebeke M, Lalk M, Sander G, von Eiff C, Proctor RA, and Engelmann S. Redox sensing by a Rex-family repressor is involved in the regulation of anaerobic gene expression in *Staphylococcus aureus*. *Mol Microbiol* 76: 1142–1161, 2010.
173. Pawlik G, Kulajta C, Sachelaru I, Schroder S, Waidner B, Hellwig P, Daldal F, and Koch HG. The putative assembly factor CcoH is stably associated with the *cbb₃*-type cytochrome oxidase. *J Bacteriol* 192: 6378–6389, 2010.
174. Pearson IV, Page MD, van Spanning RJ, and Ferguson SJ. A mutant of *Paracoccus denitrificans* with disrupted genes coding for cytochrome *c₅₅₀* and pseudoazurin establishes these two proteins as the *in vivo* electron donors to cytochrome *cd₁* nitrite reductase. *J. Bacteriol* 185: 6308–6315, 2003.
175. Peña-Sandoval GR and Georgellis D. The ArcB sensor kinase of *Escherichia coli* autophosphorylates by an intramolecular reaction. *J Bacteriol* 192: 1735–1739, 2010.
176. Pereira MM, Santana M, and Teixeira M. A novel scenario for the evolution of haem-copper oxygen reductases. *Biochim Biophys Acta* 1505: 185–208, 2001.
177. Pereira MM, Sousa FL, Verissimo AF, and Teixeira M. Looking for the minimum common denominator in haem-copper oxygen reductases: towards a unified catalytic mechanism. *Biochim Biophys Acta* 1777: 929–934, 2008.
178. Peters A, Kulajta C, Pawlik G, Daldal F, and Koch HG. Stability of the *cbb₃*-type cytochrome oxidase requires specific CcoQ-CcoP interactions. *J Bacteriol* 190: 5576–5586, 2008.
179. Pitcher RS and Watmough NJ. The bacterial cytochrome *cbb₃* oxidases. *Biochim Biophys Acta* 1655: 388–399, 2004.
180. Pohlmann A, Cramm R, Schmelz K, and Friedrich B. A novel NO-responding regulator controls the reduction of nitric oxide in *Ralstonia eutropha*. *Mol Microbiol* 38: 626–638, 2000.
181. Pomowski A, Zumft WG, Kroneck PM, and Einsle O. N₂O binding at a [4Cu:2S] copper-sulphur cluster in nitrous oxide reductase. *Nature* 477: 234–237, 2011.
182. Poole RK. Nitric oxide and nitrosative stress tolerance in bacteria. *Biochem Soc Trans* 33: 176–180, 2005.
183. Poole RK and Cook GM. Redundancy of aerobic respiratory chains in bacteria? Routes, reasons and regulation. *Adv Microb Physiol* 43: 165–224, 2000.
184. Potter L, Angove H, Richardson D, and Cole J. Nitrate reduction in the periplasm of gram-negative bacteria. *Adv Microb Physiol* 45: 51–112, 2001.
185. Preisig O, Zufferey R, Thöny-Meyer L, Appleby CA, and Hennecke H. A high-affinity *cbb₃*-type cytochrome oxidase terminates the symbiosis-specific respiratory chain of *Bradyrhizobium japonicum*. *J Bacteriol* 178: 1532–1538, 1996.

186. Price CE and Driessen AJ. Biogenesis of membrane bound respiratory complexes in *Escherichia coli*. *Biochim Biophys Acta* 1803: 748–766, 2010.
187. Pullan ST, Gidley MD, Jones RA, Barrett J, Stevanin TM, Read RC, Green J, and Poole RK. Nitric oxide in chemostat-cultured *Escherichia coli* is sensed by Fnr and other global regulators: unaltered methionine biosynthesis indicates lack of S nitrosation. *J Bacteriol* 189: 1845–1855, 2007.
188. Puri-Taneja A, Schau M, Chen Y, and Hulett FM. Regulators of the *Bacillus subtilis* *cydABCD* operon: identification of a negative regulator, CcpA, and a positive regulator, ResD. *J Bacteriol* 189: 3348–3358, 2007.
189. Ranson-Olson B, Jones DF, Donohue TJ, and Zeilstra-Ryalls JH. *In vitro* and *in vivo* analysis of the role of PrrA in *Rhodobacter sphaeroides* 2.4.1 *hemA* gene expression. *J Bacteriol* 188: 3208–3218, 2006.
190. Ranson-Olson B and Zeilstra-Ryalls JH. Regulation of the *Rhodobacter sphaeroides* 2.4.1 *hemA* gene by PrrA and FnrL. *J Bacteriol* 190: 6769–6778, 2008.
191. Rauhamäki V, Bloch DA, Verkhovsky MI, and Wikstrom M. Active site of cytochrome *cbb₃*. *J Biol Chem* 284: 11301–11308, 2009.
192. Ravishankara AR, Daniel JS, and Portmann RW. Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st century. *Science* 326: 123–125, 2009.
193. Reents H, Munch R, Dammeyer T, Jahn D, and Hartig E. The Fnr regulon of *Bacillus subtilis*. *J Bacteriol* 188: 1103–1112, 2006.
194. Rey FE and Harwood CS. FixK, a global regulator of microaerobic growth, controls photosynthesis in *Rhodospseudomonas palustris*. *Mol Microbiol* 75: 1007–1020, 2010.
195. Richardson D, Felgate H, Watmough N, Thomson A, and Baggs E. Mitigating release of the potent greenhouse gas N₂O from the nitrogen cycle - could enzymic regulation hold the key? *Trends Biotechnol* 27: 388–397, 2009.
196. Richardson D and Ferguson SJ. The influence of carbon-substrate on the activity of the periplasmic nitrate reductase in aerobically grown *Thiosphaera pantotropha*. *Arch Microbiol* 157: 535–537, 1992.
197. Richardson DJ. Bacterial respiration: a flexible process for a changing environment. *Microbiology* 146: 551–571, 2000.
198. Richardson DJ. Redox complexes of the nitrogen cycle. In: *Nitrogen Cycling in Bacteria*, edited by Moir JWB. Norfolk, UK: Caister Academic Press, 2011, pp. 23–39.
199. Richardson DJ, Berks BC, Russell DA, Spiro S, and Taylor CJ. Functional, biochemical and genetic diversity of prokaryotic nitrate reductases. *Cell Mol Life Sci* 58: 165–178, 2001.
200. Richardson DJ, King GF, Kelly DJ, MacEwan AG, Ferguson SJ, and Jackson JB. The role of auxiliary oxidants in maintaining redox balance during phototrophic growth of *Rhodobacter capsulatus* on propionate or butyrate. *Arch Microbiol* 150: 131–137, 1988.
201. Richardson DJ, van Spanning RJ, and Ferguson SJ. The prokaryotic nitrate reductases. In: *Biology of the Nitrogen Cycle*, edited by Bothe H, Ferguson SJ, and Newton WE. The Netherlands: Elsevier, 2007, pp. 21–35.
202. Rinaldo S, Arcovito A, Giardina G, Castiglione N, Brunori M, and Cutruzzola F. New insights into the activity of *Pseudomonas aeruginosa* *cd₁* nitrite reductase. *Biochem Soc Trans* 36: 1155–1159, 2008.
203. Rinaldo S and Cutruzzola F. Nitrite reductases in denitrification. In: *Biology of the Nitrogen Cycle*, edited by Bothe H, Ferguson SJ, and Newton WE. The Netherlands: Elsevier, 2007, pp. 37–56.
204. Robles EF, Sanchez C, Bonnard N, Delgado MJ, and Bedmar EJ. The *Bradyrhizobium japonicum* *napEDABC* genes are controlled by the FixLJ-FixK₂-NnrR regulatory cascade. *Biochem Soc Trans* 34: 108–110, 2006.
205. Rodgers KR and Lukat-Rodgers GS. Insights into heme-based O₂ sensing from structure-function relationships in the FixL proteins. *J Inorg Biochem* 99: 963–977, 2005.
206. Rodionov DA, Dubchak IL, Arkin AP, Alm EJ, and Gelfand MS. Dissimilatory metabolism of nitrogen oxides in bacteria: comparative reconstruction of transcriptional networks. *PLoS Comput Biol* 1: e55, 2005.
207. Roldán MD, Sears HJ, Cheesman MR, Ferguson SJ, Thomson AJ, Berks BC, and Richardson DJ. Spectroscopic characterization of a novel multiheme *c*-type cytochrome widely implicated in bacterial electron transport. *J Biol Chem* 273: 28785–28790, 1998.
208. Rolfe MD, Ter Beek A, Graham AI, Trotter EW, Asif HM, Sumo S, Sanguinetti G, Teixeira de Mattos J, Poole RK, and Green J. Transcript profiling and inference of *Escherichia coli* K-12 ArcA activity across the range of physiologically relevant oxygen concentrations. *J Biol Chem* 286: 10147–10154, 2011.
209. Salmon K, Hung SP, Mekjian K, Baldi P, Hatfield GW, and Gunsalus RP. Global gene expression profiling in *Escherichia coli* K12. The effects of oxygen availability and FNR. *J Biol Chem* 278: 29837–29855, 2003.
210. Salmon KA, Hung SP, Steffen NR, Krupp R, Baldi P, Hatfield GW, and Gunsalus RP. Global gene expression profiling in *Escherichia coli* K12: effects of oxygen availability and ArcA. *J Biol Chem* 280: 15084–15096, 2005.
211. Satterlee JD, Suquet C, Bidwai AK, Erman JE, Schwall L, and Jimenez R. Mass instability in isolated recombinant FixL heme domains of *Bradyrhizobium japonicum*. *Biochemistry* 47: 1540–1553, 2008.
212. Schau M, Chen Y, and Hulett FM. *Bacillus subtilis* YdiH is a direct negative regulator of the *cydABCD* operon. *J Bacteriol* 186: 4585–4595, 2004.
213. Schreiber K, Krieger R, Benkert B, Eschbach M, Arai H, Schobert M, and Jahn D. The anaerobic regulatory network required for *Pseudomonas aeruginosa* nitrate respiration. *J Bacteriol* 189: 4310–4314, 2007.
214. Sciotti MA, Chanfon A, Hennecke H, and Fischer HM. Disparate oxygen responsiveness of two regulatory cascades that control expression of symbiotic Genes in *Bradyrhizobium japonicum*. *J Bacteriol* 185: 5639–5642, 2003.
215. Sears HJ, Sawers G, Berks BC, Ferguson SJ, and Richardson DJ. Control of periplasmic nitrate reductase gene expression *napEDABC* from *Paracoccus pantotrophus* in response to oxygen and carbon substrates. *Microbiology* 146 (Pt 11): 2977–2985, 2000.
216. Sears HJ, Spiro S, and Richardson DJ. Effect of carbon substrate and aeration on nitrate reduction and expression of the periplasmic and membrane-bound nitrate reductases in carbon-limited continuous cultures of *Paracoccus denitrificans* Pd1222. *Microbiology* 143: 3767–3774, 1997.
217. Shalel-Levanon S, San KY, and Bennett GN. Effect of oxygen, and ArcA and FNR regulators on the expression of genes related to the electron transfer chain and the TCA cycle in *Escherichia coli*. *Metab Eng* 7: 364–374, 2005.
218. Shapleigh JP. Oxygen control of nitrogen oxide respiration, focusing on α -proteobacteria. *Biochem Soc Trans* 39: 179–183, 2011.

219. Sharma V, Wikstrom M, and Kaila VR. Redox-coupled proton transfer in the active site of cytochrome *cbb₃*. *Biochim Biophys Acta* 1797: 1512–1520, 2010.
220. Sickmier EA, Brekasis D, Paranaawithana S, Bonanno JB, Paget MS, Burley SK, and Kielkopf CL. X-ray structure of a Rex-family repressor/NADH complex insights into the mechanism of redox sensing. *Structure* 13: 43–54, 2005.
221. Simon J, van Spanning RJ, and Richardson DJ. The organisation of proton motive and non-proton motive redox loops in prokaryotic respiratory systems. *Biochim Biophys Acta* 1777: 1480–1490, 2008.
222. Smith MS. Nitrous oxide production by *Escherichia coli* is correlated with nitrate reductase activity. *Appl Environ Microbiol* 45: 1545–1547, 1983.
223. Sousa EH, Tuckerman JR, Gonzalez G, and Gilles-Gonzalez MA. A memory of oxygen binding explains the dose response of the heme-based sensor FixL. *Biochemistry* 46: 6249–6257, 2007.
224. Spector MP, Garcia del Portillo F, Bearson SM, Mahmud A, Magut M, Finlay BB, Dougan G, Foster JW, and Pallen MJ. The *rpoS*-dependent starvation-stress response locus *stiA* encodes a nitrate reductase (*narZYWV*) required for carbon-starvation-inducible thermotolerance and acid tolerance in *Salmonella typhimurium*. *Microbiology* 145: 3035–3045, 1999.
225. Spiro S. Nitric oxide-sensing mechanisms in *Escherichia coli*. *Biochem Soc Trans* 34: 200–202, 2006.
226. Spiro S. Regulators of bacterial responses to nitric oxide. *FEMS Microbiol Rev* 31: 193–211, 2007.
227. Stewart V. Biochemical Society Special Lecture. Nitrate- and nitrite-responsive sensors NarX and NarQ of proteobacteria. *Biochem Soc Trans* 31: 1–10, 2003.
228. Stewart V and Bledsoe PJ. Fnr-, NarP- and NarL-dependent regulation of transcription initiation from the *Haemophilus influenzae* Rd *napF* (periplasmic nitrate reductase) promoter in *Escherichia coli* K-12. *J Bacteriol* 187: 6928–6935, 2005.
229. Stock AM, Robinson VL, and Goudreau PN. Two-component signal transduction. *Annu Rev Biochem* 69: 183–215, 2000.
230. Suharti, Heering HA, and de Vries S. NO reductase from *Bacillus azotoformans* is a bifunctional enzyme accepting electrons from menaquinol and a specific endogenous membrane-bound cytochrome *c₅₅₁*. *Biochemistry* 43: 13487–13495, 2004.
231. Swem DL and Bauer CE. Coordination of ubiquinol oxidase and cytochrome *cbb₃* oxidase expression by multiple regulators in *Rhodobacter capsulatus*. *J Bacteriol* 184: 2815–2820, 2002.
232. Swem DL, Swem LR, Setterdahl A, and Bauer CE. Involvement of SenC in assembly of cytochrome *c* oxidase in *Rhodobacter capsulatus*. *J Bacteriol* 187: 8081–8087, 2005.
233. Swem LR, Elsen S, Bird TH, Swem DL, Koch HG, Myllykallio H, Daldal F, and Bauer CE. The RegB/RegA two-component regulatory system controls synthesis of photosynthesis and respiratory electron transfer components in *Rhodobacter capsulatus*. *J Mol Biol* 309: 121–138, 2001.
234. Swem LR, Gong X, Yu CA, and Bauer CE. Identification of a ubiquinone-binding site that affects autophosphorylation of the sensor kinase RegB. *J Biol Chem* 281: 6768–6775, 2006.
235. Swem LR, Kraft BJ, Swem DL, Setterdahl AT, Masuda S, Knaff DB, Zaleski JM, and Bauer CE. Signal transduction by the global regulator RegB is mediated by a redox-active cysteine. *EMBO J* 22: 4699–4708, 2003.
236. Takaya N. Dissimilatory nitrate reduction metabolisms and their control in fungi. *J Biosci Bioeng* 94: 506–510, 2002.
237. Thorndycroft FH, Butland G, Richardson DJ, and Watmough NJ. A new assay for nitric oxide reductase reveals two conserved glutamate residues form the entrance to a proton-conducting channel in the bacterial enzyme. *Biochem J* 401: 111–119, 2007.
238. Tolla DA and Savageau MA. Phenotypic repertoire of the FNR regulatory network in *Escherichia coli*. *Mol Microbiol* 79: 149–165, 2011.
239. Torres MJ, Bueno E, Mesa S, Bedmar EJ, and Delgado MJ. Emerging complexity in the denitrification regulatory network of *Bradyrhizobium japonicum*. *Biochem Soc Trans* 39: 284–288, 2011.
240. Tosques IE, Shi J, and Shapleigh JP. Cloning and characterization of *nmrR*, whose product is required for the expression of proteins involved in nitric oxide metabolism in *Rhodobacter sphaeroides* 2.4.3. *J Bacteriol* 178: 4958–4964, 1996.
241. Tseng CP, Albrecht J, and Gunsalus RP. Effect of microaerophilic cell growth conditions on expression of the aerobic (*cyoABCDE* and *cydAB*) and anaerobic (*narGHJI*, *frdABCD*, and *dmsABC*) respiratory pathway genes in *Escherichia coli*. *J Bacteriol* 178: 1094–1098, 1996.
242. Tucker NP, D'Autreaux B, Yousafzai FK, Fairhurst SA, Spiro S, and Dixon R. Analysis of the nitric oxide-sensing non-heme iron center in the NorR regulatory protein. *J Biol Chem* 283: 908–918, 2008.
243. Tucker NP, Le Brun NE, Dixon R, and Hutchings MI. There's NO stopping NsrR, a global regulator of the bacterial NO stress response. *Trends Microbiol* 18: 149–156, 2010.
244. Ugidos A, Morales G, Rial E, Williams HD, and Rojo F. The coordinate regulation of multiple terminal oxidases by the *Pseudomonas putida* ANR global regulator. *Environ Microbiol* 10: 1690–1702, 2008.
245. Van Alst NE, Sherrill LA, Iglewski BH, and Haidaris CG. Compensatory periplasmic nitrate reductase activity supports anaerobic growth of *Pseudomonas aeruginosa* PAO1 in the absence of membrane nitrate reductase. *Can J Microbiol* 55: 1133–1144, 2009.
246. van Spanning RJ. Structure, function, regulation and evolution of the nitrite and nitrous oxide reductases: denitrification enzymes with a b-propeller fold. In: *Nitrogen Cycling in Bacteria*, edited by Moir JWB. Norfolk, UK: Caister Academic Press, 2011, pp. 135–161.
247. van Spanning RJ, Delgado MJ, and Richardson DJ. The nitrogen cycle: denitrification and its relationship to N₂ fixation. In: *Fixation in Agriculture, Forestry, Ecology and the Environment*, edited by Werner D and Newton WE. The Netherlands: Elsevier, 2005, pp. 277–342.
248. van Spanning RJ, Richardson DJ, and Ferguson SJ. Introduction to the biochemistry and molecular biology of denitrification. In: *Biology of the Nitrogen Cycle*, edited by Bothe H, Ferguson SJ, and Newton WE. The Netherlands: Elsevier, 2007, pp. 83–93.
249. Veldman R, Reijnders WN, and van Spanning RJ. Specificity of FNR-type regulators in *Paracoccus denitrificans*. *Biochem Soc Trans* 34: 94–96, 2006.
250. von Wachenfeldt C and Hederstedt L. Respiratory cytochromes, other heme-proteins and heme biosynthesis. In: *Bacillus subtilis and Its Close Relatives: from Genes to Cells*, edited by Hoch JA, Losick R, and Sonenshein AL. Washington DC: American Society for Microbiology Publications, 2001, pp. 163–179.

251. Wang E, Ikonen TP, Knaapila M, Svergun D, Logan DT, and von Wachenfeldt C. Small-angle X-ray scattering study of a Rex family repressor: conformational response to NADH and NAD⁺ binding in solution. *J Mol Biol* 408: 670–683, 2011.
252. Watmough NJ, Field SJ, Hughes RJ, and Richardson DJ. The bacterial respiratory nitric oxide reductase. *Biochem Soc Trans* 37: 392–399, 2009.
253. Willett J, Smart JL, and Bauer CE. RegA control of bacteriochlorophyll and carotenoid synthesis in *Rhodobacter capsulatus*. *J Bacteriol* 189: 7765–7773, 2007.
254. Williams HD, Zlosnik JE, and Ryall B. Oxygen, cyanide and energy generation in the cystic fibrosis pathogen *Pseudomonas aeruginosa*. *Adv Microb Physiol* 52: 1–71, 2007.
255. Winstedt L, Yoshida K, Fujita Y, and von Wachenfeldt C. Cytochrome *bd* biosynthesis in *Bacillus subtilis*: characterization of the *cydABCD* operon. *J Bacteriol* 180: 6571–6580, 1998.
256. Wood NJ, Alizadeh T, Bennett S, Pearce J, Ferguson SJ, Richardson DJ, and Moir JW. Maximal expression of membrane-bound nitrate reductase in *Paracoccus* is induced by nitrate via a third FNR-like regulator named NarR. *J Bacteriol* 183: 3606–3613, 2001.
257. Wood NJ, Alizadeh T, Richardson DJ, Ferguson SJ, and Moir JW. Two domains of a dual-function NarK protein are required for nitrate uptake, the first step of denitrification in *Paracoccus pantotrophus*. *Mol Microbiol* 44: 157–170, 2002.
258. Wu J and Bauer CE. RegB/RegA, a global redox-responding two-component system. *Adv Exp Med Biol* 631: 131–148, 2008.
259. Wu J and Bauer CE. RegB kinase activity is controlled in part by monitoring the ratio of oxidized to reduced ubiquinones in the ubiquinone pool. *MBio* 1, pii: e00272–10, 2010.
260. Wunsch P and Zumft WG. Functional domains of NosR, a novel transmembrane iron-sulfur flavoprotein necessary for nitrous oxide respiration. *J Bacteriol* 187: 1992–2001, 2005.
261. Yan A and Kiley PJ. Techniques to isolate O₂-sensitive proteins: [4Fe-4S]-FNR as an example. *Methods Enzymol* 463: 787–805, 2009.
262. Yoon SS, Karabulut AC, Lipscomb JD, Hennigan RF, Lyman SV, Groce SL, Herr AB, Howell ML, Kiley PJ, Schurr MJ, Gaston B, Choi KH, Schweizer HP, and Hassett DJ. Two-pronged survival strategy for the major cystic fibrosis pathogen, *Pseudomonas aeruginosa*, lacking the capacity to degrade nitric oxide during anaerobic respiration. *EMBO J* 26: 3662–3672, 2007.
263. Yukl ET, Elbaz MA, Nakano MM, and Moenne-Loccoz P. Transcription factor NsrR from *Bacillus subtilis* senses nitric oxide with a 4Fe-4S cluster. *Biochemistry* 47: 13084–13092, 2008.
264. Zhang X and Hulet FM. ResD signal transduction regulator of aerobic respiration in *Bacillus subtilis*: *ctaA* promoter regulation. *Mol Microbiol* 37: 1208–1219, 2000.
265. Zufferey R, Preisig O, Hennecke H, and Thony-Meyer L. Assembly and function of the cytochrome *cbb₃* oxidase subunits in *Bradyrhizobium japonicum*. *J Biol Chem* 271: 9114–9119, 1996.
266. Zumft WG. Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 61: 533–616, 1997.
267. Zumft WG. Nitric oxide reductases of prokaryotes with emphasis on the respiratory, heme-copper oxidase type. *J Inorg Biochem* 99: 194–215, 2005.
268. Zumft WG and Kroneck PM. Respiratory transformation of nitrous oxide (N₂O) to dinitrogen by Bacteria and Archaea. *Adv Microb Physiol* 52: 107–227, 2007.

Address correspondence to:

Dr. María J. Delgado
Estación Experimental del Zaidín
CSIC
C/Profesor Albareda, 1
18008 Granada
Spain

E-mail: mdelgado@eez.csic.es

Date of first submission to ARS Central, May 09, 2011; date of final revised submission, November 16, 2011; date of acceptance, November 18, 2011.

Abbreviations Used

Arc = aerobic respiration control
*cd₁*Nir = *cd₁* type nitrite reductase
 CIO = cyanide insensitive oxidase
 Crp = cAMP receptor protein
 CuNir = Cu-type nitrite reductase
 Cys = cysteine
 Cyt = cytochrome
 DMSO = dimethyl sulfoxide
 ETC = electron transport chain
 Fnr = fumarate and nitrate reductase regulatory protein
 HCO = heme-copper oxidase
 His = histidine
 H-T-H = helix-turn-helix motif
 ITC = isothermal titration calorimetry
 MK = menaquinone
 MKH₂ = menahydroquinone
 NDH = NADH dehydrogenase
 Nap = periplasmic nitrate reductase
 Nar = membrane-bound nitrate reductase
 Nir = nitrite reductase
 NnrR = nitrite and nitric oxide reductase regulator
 Nor = nitric oxide reductase
 Nos = nitrous oxide reductase
 NrfA = cytochrome *c* nitrite reductase
 PMF = proton motive force
 RNAP = RNA polymerase
 SDH = succinate dehydrogenase
 TM = transmembrane
 UQ = ubiquinone
 UQH₂ = ubihydroquinone