CORRESPONDENCE

Clinical Practice Guideline: Uncomplicated Urinary Tract Infections

by Prof. Dr. med. Florian M.E. Wagenlehner, Prof. Dr. med. Udo Hoyme, Dr. med. Martin Kaase, Prof. Dr. med. Reinhard Fünfstück, Prof. Dr. med. Dr. h.c. Kurt G. Naber, Dr. med. Guido Schmiemann in volume 24/2011

Doubts and Questions Remain

It is vitally necessary, as well as commendable, to observe the development of resistance patterns in uncomplicated urinary tract infections and to develop recommendations and treatment strategies on that basis. The publication of the S3 guideline, however, still leaves doubts and questions. A key message of this guideline is the classification of three antibiotic substances as first-line treatments for uncomplicated cystitis: fosfomycin trometamol, nitrofurantoin, and pivmecillinam.

Fosfomycin trometamol is currently available commercially only in the form of one preparation. This preparation is licensed only for women in a restricted age range. If this preparation were to be prescribed to a 67-year-old woman, for example, this would constitute off-label use. This would mean that the treating physician is at risk of penalties from the health insurers. The guidelines do not include a warning in this respect.

The two other preparations do not offer an alternative.

How can a guideline for first-line treatment recommend a substance whose use should be restricted, according to product information? Nitrofurantoin must be given only if more effective and lower-risk antibiotics cannot be used. In my experience, resistance in longer-term therapy or after multiple administrations is clearly above 20%. In my opinion, this rules out nitrofurantoin too.

Pivmecillinam is the third recommended option, but this preparation is not available in Germany.

Some of the guideline's recommendations are thus not practical. The lack of guideline implementation in routine clinical practice has been the subject of much criticism. Perhaps better implementation would be possible if, when developing guidelines, attention is given to whether these are applicable for basic care such as is provided by general practices.

DOI: 10.3238/arztebl.2012.0079a

REFERENCES

1. Wagenlehner FME, Hoyme U, Kaase M, et al.: Clinical practice guideline: uncomplicated urinary tract infections. Dtsch Arztebl Int 2011; 108(24): 415–23.

Dr. med. Joachim Liebendörfer
Ostelsheim
Liebendoerfer.Ostelsheim@t-online.de

Conflict of interest statement

Dr Liebendörfer holds shares in Sanofi-Aventis and GSK.

The Reality of Healthcare in Germany Was Partially Ignored

I welcome guidelines based on high-level evidence. However, this article seems to me to be problematic in parts. The German recommendations ultimately reflect part of the current guideline from the European Association of Urology (EAU), “Urological Infections” (www.uroweb.org). The German recommendations for empirical treatment of uncomplicated cystitis replicate this guideline without modifications, which results in substantial compatibility problems within Germany's healthcare services.

In addition to fosfomycin trometamol, the substances recommended as first-line treatments are nitrofurantoin and pivmecillinam. In Germany nitrofurantoin is indicated only in cases in which other antibiotics or chemotherapeutic drugs that are more effective or with a lower risk-profile, cannot be used (Rote Liste [Germany's national formulary] 2011). The risk of severe pulmonary complications is probably rather low—in the Anglo-American healthcare systems, the risk seems to be estimated less seriously than in Germany. Still, it is amazing that the substance ranges among the “top 3” recommendations.

The recommendation for pivmecillinam is entirely unreasonable as this substance is not available in Germany.

There was obviously disagreement regarding the recommendation of cotrimoxazole/trimethoprim. This recommendation, in a setting where local resistance patterns are known, is also part of the EAU guideline, but it seems that in Germany, general practitioners had to ensure its inclusion into the guideline. Was a consensus not possible? The substances are mostly well tolerated and not expensive, and if the clinical symptoms do not improve after initial treatment, patients could still be switched to a different substance. Some 75% of cases of cystitis due to *E coli* would be “caught” by using cotrimoxazole/trimethoprim, and it is likely that a proportion of the cystitis cases due to *E coli* strains with resistance to cotrimoxazole would be self limiting, with a spontaneous cure rate of 30–50%. In spite of these problems I wish to thank the authors for setting out their guideline. They have prompted me to think/read up on a subject in more detail, which will certainly influence my routine prescribing behavior.

DOI: 10.3238/arztebl.2012.0079b

REFERENCES

1. Wagenlehner FME, Hoyme U, Kaase M, et al.: Clinical practice guideline: uncomplicated urinary tract infections. Dtsch Arztebl Int 2011; 108(24): 415–23.

Dr. med. Jörg Wefer, FEBU
Urologische Gemeinschaftspraxis, Oldenburg
joerg.wefer@googlemail.com

Conflict of interest statement

The author declares that no conflict of interest exists.

Authors' Objectivity Is in Doubt

The guideline on uncomplicated urinary tract infections recommends fosfomycin as first-line treatment—an antibiotic that first entered the market in the 1980s as a back-up antibiotic for multiresistant Staphylococci, and which up to the present time can be used for just this purpose in MRSA patients. In *E coli* infections, it has resistance rates of below 10%—in other words, an important, still highly effective antibiotic.

What will happen to this substance if it is now used in cases of uncomplicated cystitis—now, that a preparation for oral administration has been available for several years? Presumably, resistance patterns will take a similar course as they did for quinolones, which in the 1980s were the only oral antibiotic used in *Pseudomonas* infections; were used widely, especially for urology patients; and nowadays have a resistance rate of 25–30%.

Is there a compelling reason to use fosfomycin widely in primary care this early on, opening the doors to the development of resistance?

The physicians who are most often confronted with uncomplicated cystitis (general practitioners, pediatricians, gynecologists) have expressed a rather disapproving attitude via their specialist societies (see also FRAUENARZT 2/11).

In spite of this the guideline has been compiled. Three of the authors reported having received funding from Pierre-Fabre, the manufacturers of Monuril® (fosfomycin trometamol). Professor Wagenlehner reported ties to 19 pharmaceutical manufacturers, including Pierre-Fabre.

In a scenario of such close ties to the drug industry I ask myself whether the recommendations for the use of antibiotics were actually derived from objective and independent assessments, such as physicians and patients expect from a guideline.

DOI: 10.3238/arztebl.2012.0080a

REFERENCES

1. Wagenlehner FME, Hoyme U, Kaase M, et al.: Clinical practice guideline: uncomplicated urinary tract infections. Dtsch Arztebl Int 2011; 108(24): 415–23.

Dr. med. Swale Swalve-Bordeaux
Eckernförde
swalve-bordeaux@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Prevalence Rises in Postmenopausal Women

The S3 guidelines aim to promote the rational use of antimicrobial drugs. The authors did not mention the rational use of local vaginal administration of estriol. This mildly effective metabolite of estradiol has a shorter receptor binding time by a factor of 7 and thus bears neither cancer risks nor thrombosis risks.

Urinary tract infections (UTIs) are common after the menopause, and one in four women older than 65

is affected. A notable rise in the prevalence of dysuria, imperative urinary urge, and pollakisuria can biologically be explained with increasing urogenital atrophy and even colpitis senilis as a result of long-term estrogen deficiency. Vaginal pathogens can easily reach the bladder via the urinary meatus. Insufficient urethral closure due to estrogen deficiency is a further contributing factor. The effect is compounded by moderate fluid intake in older women (especially those with urinary incontinence) and thus insufficient flushing by means of bladder voiding.

Local vaginal estriol treatment prompts proliferation of the urethral and vesical mucosa (as can be detected on microscopic slides of vaginal epithelia showing cellular structures such as before the menopause, after only a few days. Six out of 10 women are found to have a reappearance of lactobacilli in their vaginal secretions, such as during their fertile years (when given placebo, this rate fell to 0 out of 10) (10). The production of lactic acid in an acidic vaginal milieu protects against pathogens.

Estriol as a metabolite of estradiol leads to better vascularization of the urogenital region, but climacteric problems are not eliminated. Estriol with its proliferation promoting effect has preventive benefits in recurrent UTIs after antibiotic treatment.

In a randomized double blind study (2), susceptibility to infection was reduced by means of local estriol to 0.5 episodes per year, compared with 5.9 in women of the same age. This benefit for estriol was confirmed in a placebo controlled study (3).

One pessary of estriol costs 70 cents (initially it should be administered twice weekly, later once weekly). Risks and side effects have not been seen in 40 years.

This is fully consistent with the objective of the UTI guidelines: fewer exposures to antibiotic agents.

DOI: 10.3238/arztebl.2012.0080b

REFERENCES

1. Yoshimura T, et al.: Short term oral estriol treatment restores normal premenopausal vaginal flora to elderly women. Maturitas 2001; 39: 253–7.
2. Raz R, et al.: A controlled trial of intravaginal estriol in postmenopausal women with recurrent urinary tract infections. N Engl J Med 1993; 329: 753–6.
3. Dessole S, et al.: Efficacy of low-dose intravaginal estriol on urogenital aging in postmenopausal women. Menopause 2004; 11: 49–56.
4. Wagenlehner FME, Hoyme U, Kaase M, et al.: Clinical practice guideline: uncomplicated urinary tract infections. Dtsch Arztebl Int 2011; 108(24): 415–23.

Prof. Dr. med J. Matthias Wenderlein
Universität Ulm
wenderlein@gmx.de

Conflict of interest statement

Professor Wenderlein has received honoraria for preparing scientific continuing medical educational events from Kade-Berlin

Is This how Times Are Changing?

What does uncomplicated really mean, and why does such an infection require treatment with different antibiotics? In my surprise at reading the clinical guideline I reached for my bookshelf for *Antibiotika Therapie* by C Simon und W Stille (7th edition), which dates back to 1989, in order to establish details about the older urinary tract antibiotics that seem to be recommended for first-line therapy these days. The book says that mecillinam (as ester pivmecillinam, p 68) is a highly effective broad spectrum penicillin, and that fosfomycin was contraindicated during pregnancy (p 204) and is associated with the development of secondary resistance in vitro and in vivo. As far as the chemotherapeutic substance nitrofurantoin is concerned, I wish to quote verbatim from the book (p 256) in German, paraphrasing in English: "Indications: when weighing up the advantages and disadvantages of nitrofurantoin it is vital to substantially reduce the indications of nitrofurantoin and recommend this preparation only as a back-up chemotherapeutic substance for particular forms of urinary tract infections ... incorrect indications: Nitrofurantoin should not be used any longer to treat uncomplicated urinary tract infections in women...". The contraindications include pregnancy, owing to the drug's mutagenicity. For further explanations see the 11th edition (2). Is this how times are changing? Furthermore, I wish to express unease at almost all authors' multifarious ties to the pharmaceutical industry.

DOI: 10.3238/arztebl.2012.0081a

REFERENCES

1. Wagenlehner FME, Hoyne U, Kaase M, et al.: Clinical practice guideline: uncomplicated urinary tract infections. Dtsch Arztebl Int 2011; 108(24): 415–23.
2. Stille W, Brodt HR, Groll AH, Just-Nübling G: Antibiotika-Therapie: Klinik und Praxis der antiinfektiösen Behandlung. Stuttgart, Germany: Schattauer-Verlag 2005; S 234–7.

Dr. med. Ulrike Grundmann
 Fachärztin für Innere Medizin
 Naturheilverfahren
 Marktplatz 32–34
 56727 Mayen

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

The clinical S3 guideline for uncomplicated urinary tract infections (UTIs) deals with the therapy of uncomplicated UTIs, not their prophylaxis. Professor Wenderlein's comment that in postmenopausal patients, local vaginal prophylactic administration of estradiol can prevent recurring UTIs is correct; however, it does not relate to acute treatment but to prophylaxis.

Wefer points out the S3 guideline's similarity to the current guideline of the European Association of Urology. Since all quality guidelines can be developed

according to the same rules of evidence based medicine, similar recommendations in international publications (1) are a quality indicator rather than a problem. The important pillars that this S3 guideline is based on (development of resistance, collateral damage, study data, side effects, dosages) are near-identical in the comparable, international, high-quality guidelines.

No consensus was reached with regard to assessing trimethoprim. This is primarily because of the different assessments of existing resistance limits. In Germany, the resistance rate of *E coli* to cotrimoxazole/trimethoprim is 30% (2).

The S3 guideline applies to all German speaking countries, which is why we included pivmecillinam into the therapy, which is available in Austria, for example, and constitutes a suitable substance for treating uncomplicated cystitis, for the reasons set out in the guideline.

With regard to Liebendörfer's comment that nitrofurantoin is not therapeutic alternative, we might say: the resistance rate for nitrofurantoin has not risen in recent years and has been reported as below 10% in all studies. Liebendörfer probably means that in breakthrough infections during long-term therapy with nitrofurantoin, resistance rates above 20% have been observed. This is possible, and we would not argue with this. For this reason, in every patient requiring antibiotic therapy, previous antibiotic therapy needs to be considered; this is also the case for previous therapy with cotrimoxazole and fluoroquinolones. The possible side effects of nitrofurantoin and the restricted indication in the product information are openly dealt with in the S3 guideline. However, cotrimoxazole can occasionally have severe adverse effects (Hopf et al., 1981), as can trimethoprim, albeit more rarely (Bijl et al., 1998).

Grundmann compares the current guideline with a 1989 book on antibiotic therapy by Simon and Stille. However, much has changed in the intervening 22 years. In 1989, resistance to fluoroquinolones did not exist in uncomplicated UTIs and was thus not perceived as a problem. This has changed fundamentally in the meantime. For this reason, fluoroquinolones are not recommended as first-line therapy in acute cystitis but only in acute uncomplicated pyelonephritis. Fosfomycin trometamol, by contrast, is recommended as first-line treatment for uncomplicated UTIs because of its currently favorable resistance profile, its low potential for inducing collateral damage, and the high quality evidence from studies (meta-analysis). Swalve-Bordeaux points out the possible rise in resistance if fosfomycin trometamol is given more often. But the development of resistance is an unavoidable problem associated with antibiotic treatment, which we can try to reduce as much as possible while remaining aware that it cannot be prevented altogether. For this reason, every prescription for an antibiotic substance needs to be analyzed critically with regard to indication and sequelae, as we have to make the most of the substances that are currently at our disposal.

In response to our much mentioned ties to pharmaceutical companies: these include primarily activities such as studies and lectures, without preference for any particular company. This is reflected in the fact that almost all manufacturers that deal with antibiotic substances were listed, which ultimately shows the lack of bias among the scientists involved in the guideline. All vital recommendations for antibiotic treatment are carried by all participants in the guideline team, independently of any existing potential competing interests.

DOI: 10.3238/arztebl.2012.0081b

REFERENCES

1. Gupta K, Hooton TM, Naber KG, et al.: International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: A 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *CID* 2011; 52: e103-e120.
2. Wagenlehner FME, Wagenlehner C, Savov O, Gualco L, Schito G, Naber KG: Klinik und Epidemiologie der unkomplizierten Zystitis bei Frauen. *Deutsche Ergebnisse der ARESC-Studie. Urologe* 2010; 49: 253–261.
3. Hopf U, Bernstein LS: Nebenwirkungen von Co-trimoxazol. In: Naber KG, Adam D (eds.): *Chemotherapeutika im Vergleich. Trimethoprim/Tetroxoprim-Sulfonamid-Kombinationen*. Gustav Fischer, Stuttgart, New York 1981: 177–81.
4. Bijl AM, Van der Klauw MM, Van Vliet AC, Stricker BH: Anaphylactic reactions associated with trimethoprim. *Clin Exp Allergy* 1998; 28: 510–12.
5. Wagenlehner FME, Hoyme U, Kaase M, et al.: Clinical practice guideline: uncomplicated urinary tract infections. *Dtsch Arztebl Int* 2011; 108(24): 415–23.

Prof. Dr. med. Florian M.E. Wagenlehner

Klinik und Poliklinik für Urologie, Kinderurologie und Andrologie Universitätsklinikum Gießen und Marburg GmbH, Standort Gießen
Wagenlehner@aol.com

Conflict of interest statement

Professor Wagenlehner has received travel and hotel expenses and lecture honoraria from the following companies: Astellas, Bayer-Vital, Calixa, Cerexa, Cernelle, Cubist, GSK, MerLion, OM-Pharma, Janssen-Cilag, Johnson and Johnson, Lilly, Pharmacia, Pierre-Fabre, Rosen-Pharma, Sanofi-Aventis, Strathmann, Zambon, and Serag Wiessner.