Edaravone reduces mitochondrial damage due to reperfusion injury following leg ischemia in rats

Mitsuhiro Yamamura MD FICA, Yuji Miyamoto MD, Masataka Mitsuno MD, Hiroe Tanaka MD, Masaaki Ryomoto MD, Shinya Fukui MD, Yoshiteru Yoshioka MD

M Yamamura, Y Miyamoto, M Mitsuno, et al. Edaravone reduces mitochondrial damage due to reperfusion injury following leg ischemia in rats. *Int J Angiol* 2010;19(4):e129-e131.

PURPOSE: Free radicals have been implicated in reperfusion injury. It was shown that the free radical scavenger edaravone might suppress reperfusion injury in rat extremities. The present study aimed to elucidate how edaravone suppresses reperfusion injury, focusing on its effect on the mitochondrial structure and glycogen storage in the lower extremity muscles.

METHODS: Sixteen male Lewis rats (mean $\pm$ SD) weight 497 ± 32 g) were divided into two groups and injected with either 3.0 mg/kg of edaravone (edaravone group, $n=8$ rats) or the same dose of saline (control group, $n=8$ rats) into the peritoneal cavity. The rat reperfusion injury models were created by clamping the bilateral common femoral arteries for 5 h, then declamping. The muscles were harvested more than 5 h after the start of reperfusion. The mitochondrial damage,

defined as mitochondrial swelling, was examined using a transmission electron microscope at $\times 30,000$ original magnification ($n=3$ for each rat). Glycogen storage, defined as a positive periodic acid-Schiff stain area, was examined using computerized densitometry ($n=5$ sections for each rat).

RESULTS: The mitochondria in the control group demonstrated marked swelling (mean mitochondrial size = $0.169 \pm 0.059 \mu\text{m}^2$). However, the mitochondria in the edaravone group had significantly less swelling (mean mitochondrial size = $0.102 \pm 0.036 \mu\text{m}^2$; $P < 0.01$). The mean percentage of positive periodic acid-Schiff stain was also significantly higher in the edaravone group than in the control group ($51.7 \pm 6.8\%$ versus $7.3 \pm 2.1\%$; $P < 0.01$).

CONCLUSION: The results suggested that edaravone reduces mitochondrial damage due to reperfusion injury, resulting in a high level of glycogen storage.

Key Words: *Edaravone; Free radical; Reperfusion injury*

Free radicals cause lower extremity reperfusion injury and myoneuropathic metabolic syndrome (MNMS) (1). Edaravone (Radicut, Mitsubishi Tanabe Pharma Co, Japan; 3-methyl-1-phenyl-2-pyrazoline-5-one, $\text{C}_{10}\text{H}_{10}\text{N}_2\text{O}$, molecular weight = 174.2) is the first free radical scavenger that was introduced for the treatment of acute cerebral infarction (2). We have shown that edaravone might suppress reperfusion injury in rat models (3,4). However, the mechanisms demonstrating how edaravone suppresses reperfusion injury remain unknown. Therefore, in the present study, we investigated the protective effects of edaravone against reperfusion injury, focusing on mitochondrial structure and glycogen storage in the lower extremity muscles.

METHODS

Rat reperfusion injury model

Sixteen male Lewis rats (mean $\pm$ SD) weight 497 ± 32 g) were divided into two groups. Before the operation, they were injected with either 3.0 mg/kg of edaravone (edaravone group, $n=8$ rats) or the same dose of saline (control group, $n=8$ rats) into the peritoneal cavity. General anesthesia was induced by intraperitoneal administration of 30 mg/kg of sodium pentobarbital (Nembutal, Abbott Laboratories, USA). The rat reperfusion injury model creation was performed under a microscope (OME-J&N J73507R, Olympus, Japan) as previously described (3,4). First, both deep femoral arteries and epigastric arteries were ligated to completely remove the collateral blood supply. The bilateral common femoral arteries were then clamped for 5 h and declamped.

All animal procedures complied with the criteria of the Hyogo College of Medicine Animal Research Committee, Hyogo, Japan (No B-09-010).

Transmission electron microscopic study

After 5 h, the lower extremity muscles were immediately harvested and cut transversely with a diamond knife to isolate muscle fibres ($n=3$ for each rat). The mitochondrial structure was then evaluated using a transmission electron microscope (TEM; JEM-1220, Nippon Denshi Co, Japan) at $\times 30,000$ original magnification.

The mitochondrial damage was defined as mitochondrial swelling (measured as mitochondrial size in μm^2), and was determined using computerized densitometry (Image J, version 1.37, National Institutes of Health, USA; <http://rsb.info.nih.gov>). Three fresh lower extremity muscles were also harvested before reperfusion injury and used as the TEM reference group (normal mitochondria). Among the control group and edaravone group, 15 mitochondria in each rat were measured and compared.

Periodic acid-Schiff stain study

Five sections in the control group and edaravone group were stained using the periodic acid-Schiff (PAS) method to examine glycogen storage as the result of mitochondrial activity. The positive PAS stain area was also measured three times on different fields using computerized densitometry (Image J). On Image J densitometry, the presence of between 125 and 250 over-dots was defined and calculated as the positive area.

Department of Cardiovascular Surgery, Hyogo College of Medicine, Nishinomiya-city, Hyogo, Japan

Correspondence and reprints: Dr Mitsuhiro Yamamura, Department of Cardiovascular Surgery, Hyogo College of Medicine, 1-1 Mukogawa-cho, Nishinomiya-city, Hyogo 663-8501, Japan. Telephone 81-798-45-6851, fax 81-798-45-6853, e-mail yamachan@hcm-mail.hyo-med.ac.jp

Figure 1) Mitochondria in transmission electron microscopic sections for the reference group (no reperfusion injury), control group and edaravone group (original magnification $\times 30,000$). Each arrow indicates a mitochondrion

Figure 2) The comparison of mitochondrial swelling according to mitochondrial size (control group versus edaravone group)

Glycogen storage was expressed as the percentage positive PAS stain area relative to the total area (positive PAS stain area/total area $\times 100\%$).

Statistical analysis

All results were expressed as mean $\pm$ SD. The data were statistically analyzed using unpaired *t* tests (StatView J-4.5 for Macintosh, Abacus Concept, USA). $P < 0.05$ was considered to indicate statistical significance.

RESULTS

There were no operative deaths in either group.

TEM study

Representative TEM sections after reperfusion injury are shown in Figure 1. The mitochondria in the control group after reperfusion injury demonstrated marked swelling. Additionally, the mitochondria in the control group had lost many cristae. However, there was much less mitochondrial swelling in the edaravone group. Furthermore, the mitochondria in the edaravone group maintained normal cristae.

A statistical comparison of mitochondrial damage is shown in Figure 2. The mean mitochondrial size before the operation

Figure 3) Periodic acid-Schiff stained sections (control group and edaravone group, original magnification $\times 200$). The purple staining indicates positive periodic acid-Schiff stain areas

Figure 4) Comparison of glycogen storage (control group versus edaravone group). (+) Positive; PAS Periodic acid-Schiff stain

was believed to be $0.067 \pm 0.016 \mu\text{m}^2$ (reference group). The mean mitochondrial swelling in the edaravone group was significantly less than that in the control group ($0.102 \pm 0.036 \mu\text{m}^2$ versus $0.169 \pm 0.059 \mu\text{m}^2$; $P < 0.01$). Edaravone reduced mitochondrial swelling.

PAS stain study

Representative muscle sections stained with PAS are shown in Figure 3. The purple staining indicated a positive PAS stain area. In the edaravone group, the muscles were stained in a 'marble stone' pattern and the staining was more intense than in the control group.

A statistical analysis of the positive PAS stain areas is shown in Figure 4. The glycogen storage in the edaravone group was significantly higher than that in the control group ($51.7 \pm 6.8\%$ versus $7.3 \pm 2.1\%$; $P < 0.01$). Edaravone also reduced glycogen loss from reperfusion injury.

DISCUSSION

In 1960, Haimovici (5) reported two postoperative cases of MNMS. Now, it is well known that free radicals cause lower extremity reperfusion injury and MNMS (1). In experimental studies (6,7), there have been some reports of successful suppression of reperfusion injury by free radical scavengers, but none of the studies have translated into an effective clinical

application. Therefore, we began a study to investigate how to suppress reperfusion injury (3). In rat models, we have shown that edaravone suppressed reperfusion injury (4). Recently, we have shown that edaravone might protect against tissue damage in the lungs and kidneys induced by MNMS (8). In vitro, edaravone was reported to reduce mitochondrial damage in homogenate made from ischemic incubated rat brain (9). Although some free radical scavengers were effective in suppressing reperfusion injury in several reports (6,7), there are few studies that demonstrate how free radical scavengers affect the mitochondria.

Using TEM at $\times 30,000$ original magnification, we can focus on the mitochondria. At first, we investigated whether reperfusion injury causes marked mitochondrial swelling in lower extremity muscles. We then found that edaravone reduced this mitochondrial swelling. In addition, we examined glycogen storage as a result of mitochondrial activity after reperfusion injury. Using PAS staining, we also found the muscles in the edaravone group maintained significantly higher levels of glycogen storage. Therefore, we confirmed that edaravone can suppress reperfusion injury in rat models by preventing mitochondrial damage.

In the present study, we preoperatively injected 3.0 mg/kg of edaravone into the peritoneal cavity of rats. We used peritoneal injection instead of intravenous injection to avoid any surgical stress to the vascular endothelial cells at the injection site. Edaravone has been commercially available for 10 years and the optimal intravenous dose for an adult is 30 mg twice daily. The dose from our injection was much higher than the

current clinical dose. Edaravone can be administered during clamping or after declamping. Regarding the timing of administration, edaravone may be administered after declamping in clinical settings. However, we do not know whether edaravone is still effective after declamping. Further studies are needed to confirm the optimal dose and timing of edaravone administration in a clinical setting.

CONCLUSION

Our results suggest that the free radical scavenger edaravone can suppress reperfusion injury after leg ischemia and reduce mitochondrial damage due to reperfusion injury, resulting in a high level of glycogen storage. Further studies are required to determine the precise effect before clinical use.

ACKNOWLEDGEMENTS: This study was supported by a Grant-in-Aid for Researchers, Hyogo College of Medicine, 2010. The authors thank Mitsubishi Tanabe Pharma Co for kindly donating edaravone. They also thank Ryoji Hariyama (Surgical Pathology, Hyogo College of Medicine), Ritsuko Fujimoto, Hirotsugu Kubo (Research Facility for Common Use, Hyogo College of Medicine) and Atsuhiko Kihara (The Institute of Experimental Sciences, Hyogo College of Medicine) for their technical support.

DISCLOSURE: This study was presented at the 2010 Vascular Annual Meeting in Boston, USA (June 2010), and at the 52nd Annual World Congress of the International College of Angiology in Kentucky, USA (October 2010).

REFERENCES

- Haimovici H. Metabolic complications of acute arterial occlusions and skeletal muscle ischemia: Myoneuropathic-metabolic syndrome. In: Haimovici H, ed. *Haimovici's Vascular Surgery*, 4th edn. Blackwell Science Inc, 1996:509-30.
 - Watanabe T, Morita I, Nishi H, Murota S. Preventive effect of MCI-186 on 15-HPETE induced vascular endothelial cell injury in vitro. *Prostaglandins Leukotrienes Essent Fatty Acids* 1988;33:81-7.
 - Yamamura M, Miyamoto Y, Mitsuno M, et al. Suppression of rat lower extremity postoperative reperfusion injury with edaravone. *Int J Angiol* 2006;15:34-6.
 - Yamamura M, Miyamoto Y, Mitsuno M, et al. Edaravone suppresses postoperative reperfusion injury in the rat lower extremity: An immunohistological study. *Int J Angiol* 2007;16:17-9.
 - Haimovici H. Arterial embolism with acute massive ischemic myopathy and myoglobinuria: Evaluation of a hitherto unreported syndrome with report of two cases. *Surgery* 1960;47:739-47.
 - Beyersdorf F, Matheis G, Kruger S, et al. Avoiding reperfusion injury after limb revascularization: Experimental observations and recommendations for clinical application. *J Vasc Surg* 1989;9:757-66.
 - Watarai T. The study of oxygen radical scavenger in MNMS with ascorbic acid as oxygen radical scavenger (in Japanese with English abstract). *Jpn J Cardio Vasc Surg* 1990;20:5-10.
 - Yamamura M, Miyamoto Y, Mitsuno M, et al. Edaravone protects against tissue damage in the lung and kidney induced by myoneuropathic metabolic syndrome. *Int J Angiol* 2010;19:38-40.
 - Watanabe T, Satoshi Y, Egawa M, Nishi H. Protective effects of MCI-186 on cerebral ischemia: Possible involvement of free radical scavenging and antioxidant action. *J Pharmacol Exp Ther* 1994;268:1597-604.
-