Dinucleotide repeat polymorphism at the D14S45 locus

J.A.Luty and M.Litt*

Oregon Health Sciences University, Portland, OR 97201, USA

Source and Description: Phage XL30 was from a flow sorted X-chromosome specific library LAOXNL01. DNA sequences flanking a (GT)₂₂ repeat within Sau3A subclone XL30E of this phage (EMBL accession no. X58705) were used to design PCR primers.

PCR Primers:

80718: TCCGTGAGTGATATCTTCTC

80719: CAGCATTACACAGGTACCAA

Polymorphism: Nine allelic fragments were resolved on DNA sequencing gels. Lengths of allelic fragments (nt) were: A1 = 95, A2 = 93, A3 = 91, A4 = 89, A5 = 87, A6 = 85, A7 = 83, A8 = 81, A9 = 79. Alleles found in four CEPH mothers were: 202-A5, A6; 4502-A4, A6; 10402-A6, A8; 1329402-A6, A9.

Frequencies: Allele frequencies in 48 unrelated Caucasians were:

A1 = .021 A2 = .010

A3 = .083 A4 = .062

A5 = .165 A6 = .412

A7 = .093 A8 = .124

A9 = .031

PIC = 0.74; heterozygosity in 57 unrelated Caucasians was 79%.

Chromosomal Localization and Mendelian Inheritance: Linkage analysis in 16 CEPH families against locus D14S1 (probe pAW101, localized to 14q32.32-q32.33 (1)) gave a maximum LOD score of 9.4 at $\theta = 0.15$. Mendelian inheritance was observed in all cases.

PCR Conditions: PCR reactions are carried out in a total volume of 25 μ l containing: 25 ng genomic DNA, 5 pmole of each primer, 1.5 mM MgCl₂, 200 μ M dNTPs, 50 mM Tris-Cl, 20 mM (NH₄)₂SO₄, pH 9, 0.25 mM spermidine and 0.6 units of Replisase™ (NEN/Dupont). Amplification was performed in 96 well plates in a Biosycler™ oven (Bios Corp., New Haven, CT) for 32 cycles. We used a 4 temperature cycle with denaturation at 93°C for 60 seconds, pre-annealing at 53°C for 10 sec, annealing at 48°C for 60 seconds and extension at 72°C for 15 seconds. A 3 minute ramp was programmed for the transition from 53°C to 48°C; all other ramps were programmed to be as brief as possible. PCR products were resolved on DNA sequencing gels, capillary-blotted onto Hybond N+™ nylon membranes (Amersham) and revealed by probing with 5' [³²P] labeled (CA)₁₀ oligomer.

Acknowledgement: This work was supported by NIH Grant HG00022 and by a grant from the Retinitis Pigmentosa Foundation Fighting Blindness.

Reference: 1) Cox, D.W. and Donlon, T.A. (1989) *Cytogenet. Cell Genet.* **51**, 280–298.

Dinucleotide repeat polymorphism at the D11S534 locus

X.Y.Hauge, G.A.Evans¹ and M.Litt*

Oregon Health Sciences University, Portland, OR 97201

and ¹Salk Institute, La Jolla, CA 92138, USA

Source and Description of Clone: Cosmid 30,1 was from a human chromosome 11q-specific library (1). p30,1–1, a Sau 3A subclone of this cosmid in the vector pTZ18u, was partially sequenced (EMBL accession no. X59147) and the sequences flanking a (AC)₂₀ repeat were used to design PCR primers.

PCR Primers:

(# 91249) –5'-ATATGGAACTCTCCGTACT-3'

(# 92005) –5'-GCAACCATGGAGAGTCTGGA-3'

Polymorphism: Allelic fragments were detected on 6% denaturing polyacrylamide sequencing gels. Lengths (nt) were: A1 = 244, A2 = 242, A3 = 240, A4 = 238, A5 = 236, A6 = 234, A7 = 232, A8 = 230, A9 = 228. The parents of CEPH family 1332 had the following genotypes: 133201-A5, A6; 133202-A3, A8.

Frequencies: From 58 unrelated European Caucasians.

A1 = .008 A2 = .05

A3 = .15 A4 = .34

A5 = .10 A6 = .28

A7 = .03 A8 = .03

A9 = .012

The PIC calculated from these frequencies is 0.74.

Chromosomal Localization: Linkage analysis in 9 CEPH families with the MspI RFLP identified by the probe pHBI59 (D11S146) gave a maximum LOD score of 11.88 at $\theta = 0$. The locus D11S146 has been mapped to chromosome 11q13 (2).

Mendelian Inheritance: Mendelian inheritance was observed in 9 informative CEPH families with a total of 70 children.

PCR Conditions: PCR was performed in a total volume of 12.5 μ l containing 25 ng genomic DNA, 2.5 pmole of each primer, 1.5 mM MgCl₂, 200 μ M dNTPs, 50 mM KCl, 5 mM Tris Cl, pH 8.3, 0.3 units of Taq polymerase (Perkin- Elmer/Cetus) and 0.01% gelatin. Primer # 91249 was 5' end labeled with ³²P. Amplification was for 30 cycles with denaturation at 94°C for 1 min, annealing at 55°C for 1 min and extension at 74°C for 1 min. Amplified products were resolved on DNA sequencing gels and detected by autoradiography.

Acknowledgement: This work was supported by NIH Grant HG00022 and by a grant from the Retinitis Pigmentosa Foundation Fighting Blindness.

References: 1) Lichter, P., Tang, C.C., Call, K., Hermanson, G., Evans, G.A., Housman, D. and Ward, D.C. (1990) *Science* **24**, 64–69. 2) Julier, C., Nakamura, Y., Lathrop, M., O'Connell, P., Leppert, M., Litt, M., Mohandas, T., Lalouel, J. and White, R. (1990) *Genomics* **7**, 335–345.

* To whom correspondence should be addressed

* To whom correspondence should be addressed