Published in final edited form as:

Tetrahedron Lett. 2011 October 5; 52(40): 5170–5172. doi:10.1016/j.tetlet.2011.07.125.

Fluoro-substituted ketones from nitriles using acidic and basic reaction conditions

Erum Raja and Douglas A. Klumpp*

Department of Chemistry and Biochemistry, Northern Illinois University, DeKalb, IL 60115

Abstract

Fluoro-substituted aliphatic nitriles are shown to undergo the Houben-Hoesch reactions with arenes in $\text{CF}_3\text{SO}_3\text{H}$ to give fluoro-substituted ketones in good yields. The fluorine substituents appear to enhance the reactivities of the nitriles (and the nitrilium ion intermediates) compared to similar aliphatic nitriles. Fluoro-substituted ketones are also shown to be accessible through the reactions of organometallic reagents and fluoro-substituted nitriles.

Keywords

Houben-Hoesch reaction; superacid; nitrilium ion; organofluorine

1. Introduction

The Houben-Hoesch reaction is an acid-catalyzed reaction of nitriles with aromatic compounds leading to aryl ketones.¹ It has previously been shown that nitriles having electron withdrawing groups can form fairly reactive electrophiles in the Houben-Hoesch reaction. This has been the basis for synthetic methodologies leading to halogen-substituted ketones,² aminoketones,³ *N*-heterocyclic ketones,³ and other products.⁴ Likewise, Shudo and Ohwada have described a superacid-promoted Houben-Hoesch reaction in which dicationic species are formed and these superelectrophilic conversions are shown give good yields aryl ketones.⁵

Although chlorinated alkylnitriles have been shown to provide good yields of the corresponding chlorinated aryl ketones,⁶ similar conversions with fluorinated alkylnitriles have not been extensively studied, the only example being the reaction of activated arenes with trifluoroacetonitrile (catalyzed by ZnCl_2/HCl).⁷ However, α -fluorinated ketones are useful synthetic intermediates⁸ and a direct route to these compounds would have significant value. Fluorine-substituted ketones are often prepared via electrophilic fluorinations and by other routes.^{9,10} In the following Letter, we describe a direct route to fluorinated ketones using the Houben-Hoesch reaction and fluorinated nitriles. The superacid-promoted chemistry is also shown to compliment synthetic reactions of organometallic reagents and fluorinated nitriles.

2. Results and Discussion

A variety of fluoro-substituted aliphatic nitriles are commercially available and convenient synthetic methods can provide ready access to these substrates.¹¹ For example, fluoroacetonitrile and difluoroacetonitrile were reacted with *p*-chloroanisole in the presence of CF₃SO₃H and the respective ketone (**1** and **2**) were formed in good yields (Figure 1). A series of arenes were reacted with mono-, di-, and trifluoroacetonitrile (Table 1). With fluoroacetonitrile and benzene, α -fluorinated acetophenone (**3**) is prepared in 89% isolated yield. This result indicates that the electrophilic intermediate is fairly reactive, because a similar reaction with acetonitrile produces only a moderate amount of acetophenone. Electrophilic activation by fluorine is well known, as carbonyl groups and carboxonium ions show greater electrophilic reactivities when substituted by fluorine atom(s).¹² Reactions with aryl ethers, a dialkyl-substituted benzene, and a phenol also provide the products (**4–6**) from the Houben-Hoesch reaction. Unexpectedly, the yields for these conversions are lower than that with benzene, despite the use of activated arenes (i.e., stronger nucleophiles). This may be the result of partial protonation of the activated arenes, as phenols and aryl ethers are known to be protonated in superacid.¹³

Similar reactions utilizing difluoroacetonitrile also provide the Houben-Hoesch products (**7–10**) in good yields. In general, the product yields were higher than those obtained with fluoroacetonitrile. This is likely due to enhanced electrophilic activation from the second fluorine substituent. We also examined the reactions of trifluoroacetonitrile. At room temperature, this substance is a gas so it was condensed at -78°C and dissolved in CH₂Cl₂. The resulting solution was reacted with arenes (dissolved in CH₂Cl₂) and CF₃SO₃H at 0°C and allowed to warm to room temperature. The conversions using trifluoroacetonitrile were generally good, producing the trifluoromethyl ketones in high yields. Although 1,4-dimethoxybenzene does provide the expected ketone product (observed in solution by GCMS), isolation of the product leads to massive decomposition or polymerization. With the good conversion to α,α,α -trifluoroacetophenone (**11**), it is again clear that trifluoroacetonitrile forms a highly electrophilic intermediate with CF₃SO₃H. Interestingly, acetonitriles are known for their tendencies to form triazines in strong acid, but no triazines are formed with the fluoro-substituted acetonitriles.

Besides fluoro-substituted acetonitriles, several other types of nitriles gave ketone products in fair to good yields. The fluoro-substituted benzyl cyanide leads to ketone **16** (eq 1). Although benzylic fluoro-substituents are known to form carbocations in superacid,^{3a} there is no evidence for such chemistry in this system. The dinitrile (**17**) provides the 1,6-diketone from reaction with benzene in superacid (eq 2). Houben-Hoesch

(1)

(2)

reactions of aliphatic dinitriles (without fluoro-substituents) have been studied previously and reaction at both nitrile groups has been somewhat rare.¹⁴ Likewise, a perfluorinated ketone (**19**) is formed from the perfluorooctanenitrile (eq 3). This conversion is considered representative for these Houben-Hoesch reactions. Since superacid is necessary for these reactions with arenes, it suggests either partial or complete protonation of the nitrile group (eq 4). There is evidence for diprotonated, superelectrophilic nitriles being involved in some Houben-Hoesch reactions,⁵ however such intermediates are unlikely in the present case. It is expected that the fluoro-substituents reduce that basicity of the nitrile, so diprotonation would be extremely difficult.

(3)

(4)

Although these Houben-Hoesch reactions have not been optimized, good yields have been obtained of the fluorine-substituted ketones. In a typical procedure, the nitrile (1 mmol) is dissolved in 1 mL of the arene nucleophile to which is added 0.5 mL of triflic acid (5.6 mmol). For solid arenes, CHCl_3 may be used as a co-solvent. The solution is stirred at 50°C overnight and it is then poured over ice. The products are extracted into chloroform and isolated using standard methods. Other acids systems (H_2SO_4 , $\text{CF}_3\text{CO}_2\text{H}$, and sulfated zirconia) were also examined in the reaction of fluoroacetonitrile with benzene, but only superacidic $\text{CF}_3\text{SO}_3\text{H}$ provided a useful yield of the α -fluoroacetophenone. In Whalley's report of the Houben-Hoesch reaction with trifluoroacetonitrile, highly activated arenes (resorcinol and alkylated derivatives) gave the corresponding ketones from reactions with ZnCl_2 and HCl .⁷ This suggests that weaker acids systems can effectively promote the Houben-Hoesch reaction with fluoro-substituted nitriles, but it is only effective with activated aromatic compounds. Also noteworthy, several previous studies have utilized $\text{CF}_3\text{SO}_3\text{H}$ as an acid catalyst to give Houben-Hoesch products from activated arenes and non-fluorinated aliphatic nitriles.¹⁵

The superacid-promoted Houben-Hoesch reactions provide a complimentary set of products to those from organometallic reactions. For example, phenyllithium and 3-fluorophenylmagnesium bromide provide the α -fluoroacetophenones (**3** and **20**). Product **20** is notable because it possesses *meta*-oriented

(6)

substituents, a pattern not normally accessible by electrophilic aromatic substitution. We have also found aliphatic Grignard reagents provide the fluoro-substituted ketone (**21** and **22**) in reasonable yields (eqs 7–8). Organometallic reactions with nitriles are well-known as a route to ketones of varied structures.¹⁶ Previous studies have shown that aliphatic nitriles often require forcing conditions or hydrocarbon solvents to favor organometallic addition.^{16a} However, the fluoro-substituted nitriles provide the expected ketone products without the need of

(7)

(8)

unusual solvent combinations or other reagents.

Conclusion

Fluoro-substituted aliphatic nitriles have been shown to react with arenes in superacid-promoted Houben-Hoesch reactions. The reaction provides a convenient route to mono-, di-, and trifluoromethyl ketones. Two important factors appear to be involved: the superacidic $\text{CF}_3\text{SO}_3\text{H}$ more readily forms the required nitrilium ions and the fluoro-substituents enhance the electrophilic reactivities of the nitrilium ions. The fluoro-substituted aliphatic nitriles also react efficiently with organometallic reagents to afford the fluoro-substituted ketones.¹⁷

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the support of the National Science Foundation (CHE-0749907) and the NIH-National Institute of General Medical Sciences (GM085736-01A1).

References and notes

1. a Hoesch K. Ber. Dtsch. Chem. Ges. 1915; 48:1122. b Houben J. Ber. Dtsch. Chem. Ges. 1926; 59:2878. c Smith, MB.; March, J. March's Advanced Organic Chemistry. 6th Ed.. Wiley; NY: 2007. p. 732-733. d Mullins, RJ.; O'Reilly, MC. Name Reactions for Carbocyclic Ring Formations. Li, JJ.,

- editor. Wiley; NY: 2010. p. 675-687.e Ruske, W. Friedel-Crafts and Related Reactions. Olah, GA., editor. Vol. Volume III. Interscience; New York: 1964. p. 383-497.
2. a Booth BL, Noori GFM. J. Chem. Soc. Perkin Trans I. 1980;2894.b Sanchez-Viesca F, Gomez MR, Berros M. Org. Prep. Proc. Int. 2004; 36:135.
3. a Raja EK, Klumpp DA. Tetrahedron. 2011; 67:4494. [PubMed: 21666850] b Earley JV, Gilman NW. Synth. Commun. 1985; 15:1271.c Chang MN, Biftu T, Bouton DA, Finke PE, Hammond ML, Pessolano AA, Zambias RA, Bailey P, Goldenberg M, Rackham A. Eur. J. Med. Chem., Chim. Ther. 1986; 21:363.
4. a Neill AB, Amstutz ED. J. Am. Chem. Soc. 1951; 73:3687.B Kobayashi Y, Nakatani T, Tanaka R, Okada M, Torii E, Harayama T, Kimachi T. Tetrahedron. 2011; 67:3457.
5. Sato Y, Yato M, Ohwada T, Saito S, Shudo K. J. Am. Chem. Soc. 1995; 117:3037.
6. a Toyoda T, Sasakura K, Sugasawa T. J. Org. Chem. 1981; 46:189.b Sugasawa T, Adachi M, Sasakura K, Kitagawa A. J. Org. Chem. 1979; 44:578.c Pei T, Chen C.-y. Dormer PG, Davies IW. Angew. Chem. Int. Ed. 2008; 47:4231.
7. Walley WB. J. Chem. Soc. 1951:665.
8. a Kwiatkowski P, Beeson TD, Conrad JC, MacMillan DWC. J. Am. Chem. Soc. 2011; 133:1738.b Gao Y, Twamley B, Shreeve JM. Org. Biomol. Chem. 2009; 7:1716. [PubMed: 19343262] c Belanger, E.; Cantin, K.; Messe, O.; Tremblay, M.; Paquin, J-F. 2007. p. 1034
9. a Singh RP, Shreeve JM. Synthesis. 2002:2561.b Singh RP, Shreeve JM. Accts. Chem. Res. 2004; 37:31.c Singh RP, Meshri DT, Shreeve JM. Adv. Org. Synth. 2006; 2:291.d Peng W, Shreeve JM. J. Org. Chem. 2005; 70:5760-5763. [PubMed: 15989369]
10. a Kirk KL. Org. Process Res. Dev. 2008; 12:305-321.b Moore PR, Swallow S, Gouverneur V. Chem. Soc. Rev. 2008; 37:320-330. [PubMed: 18197348] c Hagmann WK. J. Med. Chem. 2008; 51:4359-4369. [PubMed: 18570365]
11. a Grunewald GL, Seim MR, Lu J, Makboul M, Criscione KR. J. Med. Chem. 2006; 49:2929.b Fukuhara T, Hara S. Synlett. 2009:198.c Lui, N.; Heinrich, J-D.; Wollner, T.; Pazenok, S. U.S. Patent no. 20100312002 p. 13113CAN 154Gitel PO, Spiridonova TG, Yakubovich AY. Zhur. Obsh. Khimii. 1966; 36:871.
12. a Kray WD, Rosser RW. J. Org. Chem. 1977; 42:1186.b Ohwada T, Yamagata N, Shudo K. J. Am. Chem. Soc. 1991; 113:1364.c Rusanov AL, Chebotarev VP, Lovkov SS. Russ. Chem. Rev. 2008; 77(6):547.d Diaz AM, Zolotukhin M, Fromine S, Salcedo S, Manero O, Cedillo G, Velasco VM, Guzman MT, Fritsch D, Khalizov AF. Macromol. Rapid Commun. 2007; 28:183.e Zolotukhin M, Fromine S, Salcedo S, Khalilov L. Chem. Commun. 2004:1030.
13. Olah, GA.; Prakash, GKS.; Molnar, A.; Sommer, J. Superacid Chemistry. 2nd Ed.. Wiley & Sons; New York: 2009.
14. p. 411-414.
15. a Yato M, Ohwada T, Shudo K. J. Am. Chem. Soc. 1991; 113:691.b Amer MI, Booth BL, Noori GFM, Proenca MFJRP. J. Chem. Soc., Perkin Trans. I. 1983:1075.
16. a Canonne P, Foscolos GB, Lemay G. Tetrahedron Lett. 1980; 21:155.b Weiberth FJ, Hall SS. J. Org. Chem. 1987; 52:3901. and references cited there within.
17. Some fluorinated compounds are known to be highly toxic, so work with these compounds should only be done by properly trained personnel.

Figure 1.
Superacid-promoted reactions of fluoro-substituted acetonitriles.

Table 1

Products and yields from the reactions of fluoro-substituted acetonitriles.

Products from CH_2FCN , $\text{CF}_3\text{SO}_3\text{H}$, arene:

3 (89%)	4 (66%)

5 (64%)	6 (89%)
Products from CHF_2CN , $\text{CF}_3\text{SO}_3\text{H}$, arene:

7 (91%)	8 (85%)

9 (80%)	10 (90%)
Products from CF_3CN , $\text{CF}_3\text{SO}_3\text{H}$, arene:

11 (88%)	12 (0%)

13 (91%)	15 (98%)