Published in final edited form as:

Mov Disord. 2012 March ; 27(3): 463–464. doi:10.1002/mds.24050.

The Normal Parkin Sequence

Yong Ren¹, Xiaojun Liu¹, Suzanne Lesage², Miao Cai³, Jiali Pu³, Baorong Zhang³, Alexis Brice², and Jian Feng^{1,*}

¹Department of Physiology and Biophysics, State University of New York at Buffalo, 124 Sherman Hall, Buffalo, NY 14214, USA

²Université Pierre et Marie Curie-Paris6, Centre de Recherche de l'Institut du Cerveau et de la Moelle épinière, UMR-S975, Paris, France; Inserm, U975, Paris, France; Cnrs, UMR 7225, Paris, France; AP-HP, Hôpital de la Salpêtrière, Department of Genetics and Cytogenetics, F-75013, Paris, France

³Department of Neurology, Second Affiliated Hospital, School of Medicine, Zhejiang University, 88 Jiefang Road, Hangzhou, Zhejiang 310009, China

Keywords

Parkinson's disease; parkin; sequence; mitochondria

The recruitment of parkin to mitochondria in response to mitochondrial membrane depolarization¹ is an important aspect of parkin biology that has attracted extensive research². We found that parkin with the sequence (AB009973.1) reported in the original discovery of the gene³ was not recruited to mitochondria following CCCP treatment, while parkin with the current Genbank Reference Sequence (NM_004562.2) was robustly recruited to mitochondria under the same condition. The only difference between the two sequences is nucleotide 768, which is C in AB009973.1 (encoding for proline at amino acid position 223) and T in NM_004562.2 (for serine at 223). As shown in Fig. 1, HeLa cells were transfected with FLAG-tagged parkin (with either P or S at amino acid 223). Cells were treated without or with CCCP (10 μ M) for 3 hr and immunostained for FLAG and Tom20. CCCP induced a strong mitochondrial recruitment of FLAG-parkin (223S), but no significant recruitment of FLAG-parkin (223P). This was confirmed using GFP-parkin with either P or S at position 223 (Fig. 1C). Similar results were also obtained in COS7 or HEK293 cells using FLAG-, Myc-, HA- or GFP-tagged human parkin with S or P at amino acid 223 (data not shown).

To test which version of parkin sequence is correct, we analyzed genomic DNA of 2102 individuals from Europe, China and the United States. None of them had C at this position; all were T. Briefly, genomic DNA from 231 normal subjects and 294 Parkinson's disease patients in China, as well as genomic DNA from 19 normal subjects and 18 Parkinson's disease patients in the United States were amplified by PCR using two primers in the introns flanking exon 6 of parkin (CTTGTCCTAAAGAGATTGTTACTGTGG and GGCTCGTGTGGCAGAACAAATATTGGG). The PCR product (318bp) should contain a single BsrI site (CCAGT) if AB009973.1 is correct. None of these samples could be cut by

*Correspondence to: Jian Feng, Department of Physiology and Biophysics, State University of New York at Buffalo, 124 Sherman Hall, Buffalo, NY 14214, USA. Phone: 1-716-829-2345, Fax: 1-716-829-2699, jianfeng@buffalo.edu.

Author Roles: Y.R., X.L., S.L., M.C. and J.P. performed the experiments, which were designed by J.F. The manuscript was written by J.F., A.B., and B.Z.

Relevant conflicts of interest/financial disclosures: Nothing to report.

BsrI, while a positive control PCR product containing a BsrI site was cut at the same condition. Using a BLAST search of DNA sequences from 1540 individuals (1290 Parkinson's disease patients and 250 healthy controls), mostly of French origin (~70%), we found that all of them had T at position 768; no one had C. Thus, the amino acid sequence reported in AB009973.1 (Pro223) is wrong; it should be Serine instead, as reported in NM_004562.2. This important error needs to be corrected, as it may affect other aspects of parkin biology, in addition to mitochondrial recruitment.

Acknowledgments

The work was supported by NIH grant NS061856 and NSFC grant 30928006 (to J.F.), and International Cooperation Project (2011C14026) of Zhejiang Province Department of Science and Technology (to B.Z).

References

1. Narendra D, Tanaka A, Suen DF, et al. Parkin is recruited selectively to impaired mitochondria and promotes their autophagy. *J Cell Biol.* 2008; 183:795–803. [PubMed: 19029340]
2. Youle RJ, Narendra DP. Mechanisms of mitophagy. *Nat Rev Mol Cell Biol.* 2011; 12:9–14. [PubMed: 21179058]
3. Kitada T, Asakawa S, Hattori N, et al. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature.* 1998; 392:605–608. [PubMed: 9560156]

Fig. 1. Parkin sequence error has important functional consequence

A–B, HeLa cells transfected with FLAG-parkin (223P) (**A**) or FLAG-parkin (223S) (**B**) were treated with CCCP (10 μ M for 3 hr) and co-stained for FLAG and the mitochondria marker Tom20. **C**, Percentage of cells with mitochondrial recruitment of transfected GFP-parkin or FLAG-parkin with 223P or 223S. *, $p < 0.001$, vs. the preceding bar, $n = 8–10$ views from 4 independent experiments, with each view having 70–80 cells.