Two RFLPs in the human protein 4.1 gene (EL1)

T.K.Tang¹, K.-b.Tam¹ and S.Chien^{1, 2}

¹Institute of Biomedical Sciences, Academia Sinica, Taipei 11529, Taiwan, ROC and ²AMES Department, University of California at San Diego, La Jolla, CA 92093, USA

Source/Description: pTM-1 (1) and pLym (2) are two cDNA clones containing a 1 kb and 2.7 kb EcoRI inserts of the protein 4.1 gene which were isolated from a human T cell leukemia cDNA library.

Polymorphism: Two allele co-dominant polymorphisms identified with HindIII (A1 = 4.5 kb, A2 = 4.0 kb) by the pTM-1 clone and with BglII (B1 = 22 kb, B2 = 17 kb) by the pLym clone.

Frequency: Studied in 16 (A allele) and 23 (B allele) unrelated Chinese.

A1: 0.44

A2: 0.56

B1: 0.24

B2: 0.76

Not Polymorphic For: PvuII, EcoRI, BamHI

Chromosome Localization: Assigned to chromosome 1p33→1p34.2 by non-radioactive *in situ* hybridization (3).

Mendelian Inheritance: Co-dominant segregation shown in three families, 20 individuals (A allele) and one family, 9 individuals (B allele).

Probe Availability: Available from Dr T.K.Tang.

Other Comments: The co-dominant segregation of HindIII polymorphism was also observed in a large white family (4).

Acknowledgements: We thank C.-J. Chang Tang and C.-M. Wang for technical assistance. This work was supported by NSC grant 80-0203-B001-03 and Academia Sinica, ROC.

References: 1) Tang *et al.* (1988) *Proc. Natl. Acad. Sci. USA* **85**, 3713-3717. 2) Tang *et al.* (1990) *J. Cell Biol.* **110**, 617-624. 3) Tang *et al.* (1991) *Cytogenet. and Cell Genet.* **57**, 119. 4) Costa *et al.* (1990) *N. Engl. J. Med.* **323**, 1046-1050.

AhaII polymorphism in human X-linked proteolipid protein gene (PLP)

J.A.Trofatter, V.M.Pratt, S.R.Dlouhy* and M.E.Hodes
Indiana University School of Medicine, Indianapolis, IN, USA

Source/Description: The polymerase chain reaction (PCR) and *AhaII* digestion were used to detect a common polymorphism within the open reading frame (ORF) of the human proteolipid protein (PLP) gene. This is a T→C transition in exon 4 that results in the formation of an *AhaII* restriction site. It is a single-base silent substitution (GAT→GAC) in aspartate codon 202.

PCR Primers: 5'-TCTCGAATTCCTCATGTCAATCATTTT-3' and 5'-TCTCGAATTCGCACCCGTACCCTAACTC-3'.

Polymorphism: Two hundred ng of amplified exon 4 DNA from each individual was digested with *AhaII* and size-fractionated on an 8% nondenaturing polyacrylamide gel. Amplified exon 4 was 227 bp long Allele B1 (Fig. 1, lane 4). Presence of the *AhaII* site resulted in production of fragments that are 183 and 44 bp in length Allele B2 (the 183-bp fragment is shown in Fig. 1).

Frequency: Studied in 58 unrelated individuals.

B1 (*AhaII* site absent) = 0.74

B2 (*AhaII* site present) = 0.26

Observed heterozygosity = 33%

Mendelian Inheritance: X-linked recessive inheritance demonstrated in three Caucasian families.

Other Comments: PCR mixtures (100 μ l) contained approximately 500 ng of genomic DNA, 600 ng of each primer, 200 μ M each dNTP, 10 mM Tris (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 0.01% gelatin and 2.5 units *Taq* polymerase (Cetus). Cycling conditions were: cycle set 1 94°C 1 min, 46°C 1 sec, 68°C 1.25 min for 4 cycles; cycle set 2 94°C 1 min, 58°C 5 sec, 72°C 1.25 min, for 31 cycles.

PLP is tightly linked to Pelizaeus-Merzbacher disease (PMD) at Xq21.33-q22 (1). A linked *MspI* RFLP outside *PLP* has been described (2). The mutation presented in the current study is the first silent mutation in the *PLP* ORF.

References: 1) Trofatter, J.A. *et al.* (1989) *Proc. Natl. Acad. Sci. USA* **86**, 9427-9430. 2) Wu, J.-S. *et al.* (1987) *Nucl. Acids Res.* **15**, 1882.

Figure 1. Lane 1, ϕ X174 marker; lane 2, hemizygous male with *AhaII* site; lane 3, heterozygous female; lane 4, hemizygous male lacking *AhaII*.

* To whom correspondence should be addressed