Trainees must seek simulated practice to maximise their surgical training. Existing tendon repair models have significant limitations. The dental roll lacks the translucency of the drinking straw. Silicon rods and pig trotters are excellent models; however, both are inconvenient, expensive and offer no feedback on handling technique. Furthermore, these models are easier to repair than human tendon through their size and handling characteristics. On drinking straws, tendon repair was picked up well by novices but impeccable stitch-handling is required for a 'perfect' repair. It is offered as an adjunct to existing models, offering specific technical benefits and the convenience to be used at any time, at home or at work.

A new surgical approach to remove subscapular exostoses in children

Christopher Brown, Rachel Buckingham
Nuffield Orthopaedic Centre, Oxford, UK

CORRESPONDENCE TO

Christopher Brown, Nuffield Orthopaedic Centre, Windmill Road, Oxford OX3 7LD, UK
E: cnbrown@doctors.org.uk

Figure 1 Longitudinal incision splitting the scapular apophysis.

Figure 2 Cobb elevator developing the subperiosteal plane of the inner table.

Figure 3 The base of the exostosis has been divided.

Figure 4 Excised exostosis with cartilaginous cap intact.

BACKGROUND

Exostoses (osteochondromas) are benign, cartilage-capped bone tumours. Subscapular exostoses can cause scapulothoracic crepitus, winging of the scapula and bursa formation.¹ They may be solitary or occur in the autosomal dominant condition hereditary multiple exostosis (HME).² Excision has traditionally been performed through a parascapular approach, dividing muscle fibres to gain extraperiosteal access to the inner table of the scapula. We propose that splitting the scapular apophysis and sub-periosteal dissection causes less soft tissue damage and is a simpler approach to remove these lesions in children.

TECHNIQUE

A skin incision is made over the medial border of the scapula. The subcutaneous tissues are sharply dissected to reach the scapular apophysis which is then split longitudinally with a scalpel in a similar way to splitting the iliac apophysis in the anterior approach to the hip. A Cobb elevator is used to raise the sub-periosteal plane of the inner table of the scapula. This dissection is continued to the base of the exostosis which is exposed using bone levers and cut with an osteotome. The lesion is then detached from the soft tissues and removed in its entirety (Figs 1–4). The apophysis is repaired with 1-vicryl sutures.

DISCUSSION

Splitting of the iliac apophysis is commonly performed to gain access to the paediatric hip. This produces minimal muscle damage and ease

of repair. We have found similar advantages to splitting the scapular apophysis in order to remove a subscapular exostosis. To our knowledge, this approach has not previously been described.

REFERENCES

1. Mohsen MS, Moosa NK, Kumar P. Osteochondroma of the scapula associated with winging and large bursa formation. *Med Princ Pract (Switzerland)* 2006; **15**: 387–90.
2. Solomon L. Hereditary multiple exostosis. *J Bone Joint Surg Br* 1963; **45**: 292–4.

Trapeziectomy made simpler (and quicker)

Stuart Goudie, Alberto Gregori, Michael Dowd

Department of Orthopaedics, Hairmyres Hospital, East Kilbride, UK

CORRESPONDENCE TO

Stuart Goudie, E: stuartgoudie@nhs.net

Figure 1 Drilling across the bulk of the trapezium.

Figure 2 Tap is inserted and used for leverage and retraction.

Figure 3 Complete trapezium is delivered attached to the tap.

BACKGROUND

Trapeziectomy for thumb carpal-metacarpal joint arthritis yields good results in terms of improvement of pain and physical function.¹ Traditional piecemeal trapeziectomy can be time-consuming. We describe a quick and simple way to remove the trapezium in one piece.

TECHNIQUE

A standard approach to the trapezium is made with protection of the radial nerve, the trapezium being exposed with a longitudinal incision in the carpometacarpal ligament.²

A 2.5mm drill is used to drill across the bulk of the trapezium (Fig. 1). A cortical 3.5mm tap is then inserted and used for leverage and retraction (Fig. 2) while soft tissue is dissected from around the bone using a scalpel, allowing delivery of the complete trapezium attached to the tap (Fig. 3).

DISCUSSION

We have found this method for excision of the trapezium to be slick and effective and to have good results. We have seen a time saving of around 15 minutes per procedure. We cannot claim credit for this technique, it being shown to us by one of the authors who learnt it in the US.

REFERENCES

1. Wajon A, Carr E, Edmunds I, Ada L. Surgery for thumb (trapeziometacarpal joint) osteoarthritis. *Cochrane Database Syst Rev* 2009; **4**: CD004631.
2. Wheelless CR. *Wheelless' Textbook of Orthopaedics*. Brooklandville, Maryland, US: Data Trace Internet Publishing; 2009. http://www.wheellessonline.com/ortho/cmc_arthritis_resection_arthroplasty (cited February 2011).