continuous suture up the wound to the proximal end. Once the proximal end is reached no knot is tied. Instead, the suture is simply turned back on itself and the continuous suture is continued back down the wound until the suture has passed two-thirds of the way down to approximately the level of the patella. At this point two bites are taken in the soft tissue to one side of the wound at 180 degrees to one another in order to lock the suture. The suture is then simply cut flush with the tissue. The superficial wound is closed in a conventional manner with a 2-0 braided polyglycolide suture (Vicryl) to approximate the wound edges and, finally, with clips to the skin.³

DISCUSSION

This provides a solid and reliable approximation of the deep tissues without any knot. This has been used by the authors for two years and has had led to excellent results with no cases of wound dehiscence.

REFERENCES

1. van Rijssel EJ, Brand R, Admiraal C *et al*. Tissue reaction and surgical knots: the effect of suture size, knot configuration, and knot volume. *Obstet Gynecol* 1989; **74**: 64–68.
2. Mahabir RC, Christensen B, Blair GK, Fitzpatrick DG. Avoiding stitch abscesses in subcuticular skin closure: the L-stitch. *Can J Surg* 2003; **46**: 223–224.
3. Khan RJ, Fick D, Yao F *et al*. A comparison of three methods of wound closure following arthroplasty: a prospective, randomised, controlled trial. *J Bone Joint Surg Br* 2006; **88**: 238–242.

Control of postoperative bleeding following a procedure for prolapse and haemorrhoids: a novel technique

A Bagul, S Jones, C Usmani

Department of Colorectal Surgery, Dorset County Hospital, Dorchester, UK

CORRESPONDENCE TO

A Bagul, E: anil.bagul@nhs.net

BACKGROUND

The majority of bleeding can be controlled by direct compression and using a tamponade effect. In an emergency situation, a common effective method is application of direct pressure by hand. This approach is not suitable for post-haemorrhoidectomy bleeding. Published data highlight a variable rate of postoperative bleeding, from 6% to 25%.^{1,2} We describe a novel technique for providing localised pressure, utilising materials that are freely available on most surgical wards or in the emergency department.

TECHNIQUE

A well lubricated large bore, 3-way catheter is introduced carefully above the staple line. The catheter balloon is inflated with 30ml of sterile water and pulled backwards against the staple line; the balloon acts as a tamponade. Two 100ml metronidazole infusion bottles are secured by suture to the first catheter port. The second port is connected to a catheter bag and the third port spigotted. The two attached metronidazole bottles are left hanging over the bed side, maintaining constant uniform traction. The advantage of the catheter channel is to drain any emerging bleeding

Figure 1 The catheter channel in situ

above the staple line and this is left in situ for 24 hours.

DISCUSSION

We have not come across this technique for control of postoperative procedure for prolapse and haemorrhoids (PPH) bleeding being described in the literature. It is effective, quick and practical. We have used this method in six patients who developed post-PPH bleeding. The haemorrhage was controlled in five patients; one patient needed to go back to theatre. In our experience it is an effective technique in controlling post-PPH haemorrhage.

REFERENCES

1. Finco C, Sarzo G, Savastano S *et al*. Stapled haemorrhoidopexy in fourth degree haemorrhoidal prolapse: is it worthwhile? *Colorectal Dis* 2006; **8**: 130–134.
2. Slawik S, Kenefick N, Greenslade GL, Dixon AR. A prospective evaluation of stapled haemorrhoidopexy/rectal mucosectomy in the management of 3rd and 4th degree haemorrhoids. *Colorectal Dis* 2007; **9**: 352–356.

TECHNICAL TIPS

New technique of optimal size skin incision for laparoscopic port insertion

Bynvant Sandhu, Ravi Marudanayagam, Mike Hallissey

Upper Gastrointestinal Surgical Unit, Queen Elizabeth Hospital, Birmingham, UK

CORRESPONDENCE TO

Ravi Marudanayagam, Upper Gastrointestinal Surgical Unit, Queen Elizabeth Hospital, University Hospital of Birmingham NHS Foundation Trust, Edgbaston, Birmingham B15 2TH, UK
E: ravim1973@yahoo.co.in

The optimal-sized skin incision for laparoscopic port insertion facilitates easy manoeuvrability of the port while reducing potential air leaks and results in smaller scars.

After infiltration of local anaesthetic, the port, without the trocar, is pushed against the anterior abdominal wall at the site of proposed port