Published in final edited form as:

J Med Chem. 2012 January 26; 55(2): 717–724. doi:10.1021/jm201157c.

Chemistry and Behavioral Studies Identify Chiral Cyclopropanes as Selective $\alpha 4\beta 2$ -Nicotinic Acetylcholine Receptor Partial Agonists Exhibiting an Antidepressant Profile

Hankun Zhang[†], Werner Tückmantel[§], J. Brek Eaton[‡], Po-wai Yuen[§], Li-Fang Yu[†], Krishna Mohan Bajjuri[†], Allison Fedolak[§], Daguang Wang[§], Afshin Ghavami[§], Barbara Caldarone[§], Neil E. Paterson[§], David A. Lowe[§], Daniela Brunner[§], Ronald J. Lukas[‡], and Alan P. Kozikowski^{†,*}

[†]Drug Discovery Program, Department of Medicinal Chemistry and Pharmacognosy, University of Illinois at Chicago, 833 South Wood Street, Chicago, Illinois 60612

[‡]Division of Neurobiology, Barrow Neurological Institute, 350 West Thomas Road, Phoenix, Arizona 85013

[§]PsychoGenics, Inc., 765 Old Saw Mill River Road, Tarrytown, New York 10591

Abstract

Despite their discovery in the early 20th century and intensive study over the last twenty years, nicotinic acetylcholine receptors (nAChRs) are still far from being well understood. Only a few chemical entities targeting nAChRs are currently undergoing clinical trials, and even fewer have reached the marketplace. In our efforts to discover novel and truly selective nAChR ligands, we designed and synthesized a series of chiral cyclopropane-containing $\alpha 4\beta 2$ -specific ligands that display low nanomolar binding affinities and excellent subtype selectivity, while acting as partial agonists at $\alpha 4\beta 2$ -nAChRs. Their favorable antidepressant-like properties were demonstrated in the classical mouse forced swim test. Preliminary ADMET studies and broad screening towards other common neurotransmitter receptors were also carried out to further evaluate their safety profile and eliminate their potential off-target activity. These highly potent cyclopropane ligands possess superior subtype selectivity compared to other $\alpha 4\beta 2$ -nAChR agonists reported to date, including the marketed drug varenicline, and therefore may fully satisfy the crucial prerequisite for avoiding adverse side effects. These novel chemical entities could potentially be advanced to the clinic as new drug candidates for treating depression.

Introduction

Over the past two decades, nicotinic acetylcholine receptors (nAChRs) have been investigated with the goal of developing drugs that can potentially treat a variety of nervous system disorders such as Alzheimer's disease, Parkinson's disease, schizophrenia, pathological pain, nicotine addiction, and depression.^{1–4} In vertebrates, nAChRs are pentameric ligand-gated ion channel proteins that are composed of seventeen known homologous subunits ($\alpha 1$ – $\alpha 10$, $\beta 1$ – $\beta 4$, γ , δ , and ϵ) that are expressed widely throughout the central and peripheral nervous systems (CNS and PNS) and neuromuscular junctions. They broadly participate in physiological and pathophysiological processes by modulating the

*To whom correspondence should be addressed. Phone: +1-312-996-7577. Fax: +1-312-996-7107. kozikowa@uic.edu.

SUPPORTING INFORMATION. Experimental details for synthesis of all compounds shown, and procedures for *in vitro* functional studies, and detailed broad screening data. This material is available free of charge via the internet at <http://pubs.acs.org>.

synaptic release of neurotransmitters such as dopamine (DA), serotonin (5-HT), glutamate (Glu), acetylcholine (ACh), and γ -aminobutyric acid (GABA) that are all involved in the aforementioned diseases.

There are twelve nAChR subunits expressed in the nervous system ($\alpha 2$ – $\alpha 10$ and $\beta 2$ – $\beta 4$), and different combinations of subunits allow the assembly of many functional pentamers, though the actual number of functional pentamers expressed is far less than the theoretical number of possible combinations. The predominant form of nAChRs in the CNS are heteromeric $\alpha 4\beta 2^*$ -nAChR complexes characterized by high-affinity ACh binding and slow desensitization (the asterisk denotes the possible integration of other subunits into the pentamer). Homomeric $\alpha 7$ -nAChRs, which are typified by low ACh affinity and fast activation, are the other major component in the brain. Ganglionic $\alpha 3\beta 4^*$ nAChRs play a dominant role in the sensory and autonomic ganglia as well as in subpopulations of neurons in the brain, and are frequently associated with adverse side effects such as emesis and nausea.^{5, 6} Less abundant in the brain overall, but nevertheless concentrated in dopaminergic, pleasure-reward centers putatively involved in mood and drug dependence, are $\alpha 6^*$ -nAChRs.

It is now well established that the $\alpha 4\beta 2^*$ -nAChRs have an essential role in mediating nicotine's rewarding properties,^{7, 8} and it is hypothesized that they are also responsible for the antidepressant effects of nicotinic agents. This notion is supported by the findings that knock-out mice lacking the nAChR $\beta 2$ subunit do not show any behavioral antidepressant response to mecamylamine or amitriptyline, and that nAChR $\alpha 4$ subunit knock-in mice exhibit increased anxiety.^{9–11} Furthermore, social defeat, a behavioral model of depression in rodents, produced a robust increase in the expression of the nAChR $\beta 2$ subunits in the brain.¹² In addition, nicotinic ligands targeting $\alpha 4\beta 2^*$ -nAChRs may likewise be used to treat neuropathic pain or attention deficit hyperactivity disorder (ADHD).⁴ Since there is a great deal of conservation between the primary structures of the nAChR subtypes, the design of ligands selective for $\alpha 4\beta 2$ -nAChRs over $\alpha 3\beta 4^*$ -nAChRs provides a challenge, but not one that is insurmountable.

Among the natural nicotinic ligands and a number of synthetic small molecules that have been pharmacologically tested as agents targeting brain $\alpha 4\beta 2$ -nAChRs, only a small fraction have been advanced to preclinical studies, and even fewer have made it to clinical trials. Abbott Laboratories developed ABT-089 (**1**), an $\alpha 4\beta 2$ -nAChR partial agonist that recently underwent clinical trials for the treatment of pediatric ADHD. Whereas this drug was found to be safe and well tolerated, it showed no significant difference from placebo in terms of efficacy.¹³ Another compound, TC-5214 (**2**) from Targacept, the *S*-enantiomer of mecamylamine, is now in Phase III development as an adjunctive therapy for major depressive disorder (MDD).¹⁴ Pfizer's varenicline (**3**), an $\alpha 4\beta 2$ -nAChR partial agonist and a full agonist at $\alpha 7$ - and $\alpha 3\beta 4^*$ -nAChR, is at present the most successful synthetic small molecule on the market for smoking cessation pharmacotherapy.¹⁵ The emergence of compound **3** lends support to the use of $\alpha 4\beta 2$ -nAChR partial agonists as clinical drugs to treat nervous system diseases. However, peripheral and central side effects of compound **3**, such as nausea, gastrointestinal symptoms, changes in mood, and, perhaps, suicidal ideation are most likely due to its insufficient subtype selectivity,⁴ indicating that the nicotinic arena is still rife with both opportunities and challenges.

Rational Design and Synthesis of Chiral Cyclopropane nAChR Ligands

There is still a need for antidepressants that exhibit fewer side effects, act pharmacologically in new ways, and that have a faster onset of action compared to currently available therapeutics. In pursuit of this goal, our group has identified sazetidine-A (**4**) as a highly

potent $\alpha 4\beta 2$ -nAChR partial agonist with excellent selectivity over $\alpha 3\beta 4^*$ -nAChRs.^{16, 17} Compound **4** has been shown to possess extremely promising antidepressant and anxiolytic effects in rodent studies, including nicotine-like effects in drug discrimination studies.^{18–20} In addition, analgesic effects of compound **4**, without any neurological side effects, have been reported using the rat formalin model.²¹ However, the potential metabolic liability of the acetylenic bond in compound **4**, which may be oxidized to generate a labile, highly reactive oxirene, thereby possibly giving rise to toxicity, discouraged further advancement of this compound down the drug discovery pipeline.²² Novel ligands were, therefore, designed to avoid the acetylene function while maintaining the important pharmacophoric elements of compound **4**. For various reasons, we considered replacement of the acetylene by a small and rigid cyclopropane ring. Cyclopropanes widely occur in both natural products and synthetic, biologically active compounds.^{23, 24} A cyclopropane ring in place of the acetylene group would not only function as a spacer but also might be directly involved in the ligand-receptor binding interaction. The rigid structure of the cyclopropane would endow the ligand with a unique restricted conformation in which the functional groups display a particular arrangement and might more effectively interact with the amino acid residues of the binding site of the target receptor. Moreover, the chiral cyclopropane could also be used to modify the areas of space accessible to the side-chain hydroxyl, with the goal to optimize its hydrogen bonding interactions.²⁵ In our previous studies on analogs of compound **4**, we found that a side chain length of between 4 and 6 carbon atoms was optimal for biological activity.²⁶ As our first goal, we chose to synthesize a cyclopropane ligand bearing a four-carbon side chain counting along the shortest path from the pyridine ring to the hydroxyl group.

The syntheses of the chiral cyclopropane ligands **12a**, **13a–17a**, and **19a** are described in Scheme 1. 3,5-Dibromopyridine (**5**) underwent Br displacement with benzyl alcohol, followed by a Heck reaction with *n*-butyl acrylate using a recently described, phosphine-free protocol²⁷ to afford the α,β -unsaturated ester **6**. Conversion of the ester group to the Weinreb amide²⁸ using a standard procedure and subsequent Corey-Chaykovsky cyclopropanation gave the racemic mixture of cyclopropanes **7**, which were then reduced to the corresponding alcohols in two steps, followed by chiral resolution on a ChiralPak® AD column to give alcohols **8a** and **8b** in gram quantities with essentially 100% ee values.^{29, 30} The absolute configuration of the alcohol **8a** was determined by the X-ray crystallography of its derivative **9a**, which was obtained by subsequent oxidation and coupling with a chiral Evans oxazolidinone.

The optically pure alcohol **8a** was subjected to standard Swern oxidation, Wittig reaction, and hydroboration to obtain the chain-extended terminal alcohol **10a**. Successive acylation of the alcohol, removal of the benzyl group, and Mitsunobu reaction to install the azetidine moiety furnished the intermediate **11a** after removal of the isobutyrate group. The intermediate **11a** was then converted to a carbamate using various amines or phenyl isocyanate. Removal of the Boc group from **11a** or the carbamate intermediates gave the desired products **12a** and **13a–17a**. The methyl ether analog **19a** was prepared by a similar procedure in which the methoxy group was introduced as a substituent in the Wittig reagent, and unsaturation removed by catalytic hydrogenation. Compounds **12b** and **19b**, diastereoisomers of **12a** and **19a**, were synthesized from **8b** by the same sequence of steps.

***In Vitro* Radioligand Binding and Functional Studies**

The K_i values of all the synthesized cyclopropane compounds were evaluated by [³H]epibatidine binding competition assays at seven heterologously expressed rat nAChR subtypes. As shown in Table 1, compound **12a** with a (*1S,2R*)-configured cyclopropane ring exhibited sub-nanomolar binding affinity for both the $\alpha 4\beta 2$ - (K_i = 0.1 nM) and $\alpha 4\beta 2^*$ - (K_i =

0.5 nM) nAChRs, thus having K_i values similar to those found for compound **4**. Compound **12a** was approximately 7-fold more potent than its diastereoisomer **12b** at both the $\alpha 4\beta 2$ - and $\alpha 4\beta 2^*$ -nAChRs (**12b**: $K_i = 0.7$ and 3.7 nM, respectively). Similarly, the (*1R,2S*)-configured ether analog **19b** was less active than **19a** at both the $\alpha 4\beta 2$ - ($K_i = 0.6$ vs. 0.1 nM) and $\alpha 4\beta 2^*$ -nAChR subtype ($K_i = 6.2$ vs. 0.3 nM). All four of these compounds demonstrated good selectivity for nAChRs containing $\beta 2$ subunits ($\alpha 2\beta 2$ -, $\alpha 3\beta 2$ -, $\alpha 4\beta 2$ -, and $\alpha 4\beta 2^*$ -nAChRs), which are associated with the regulation of dopamine release in the nucleus accumbens,³¹ over nAChRs containing $\beta 4$ subunits ($\alpha 3\beta 4$ -, $\alpha 2\beta 4$ -, and $\alpha 4\beta 4$ -nAChRs). Of considerable importance in the identification of therapeutically useful nicotinic ligands is the selectivity away from the $\alpha 3\beta 4$ -nAChR subtype. The $\alpha 3\beta 4$ -/ $\alpha 4\beta 2$ -nAChR K_i ratios of the two cyclopropane analogs featuring the (*1S,2R*)-configuration, **12a** and **19a**, were 65,200 and 100,000, respectively, which are much higher than that of nicotine ($\alpha 3\beta 4$ -/ $\alpha 4\beta 2$ -nAChR = 53) or even compound **4** ($\alpha 3\beta 4$ -/ $\alpha 4\beta 2$ -nAChR = 24,000). Collectively, these outcomes clearly suggest that the (*1S,2R*)-configuration of the cyclopropane ring improves subtype selectivity by conferring the proper orientation to the side chain, thereby improving upon the compounds' affinity for the $\alpha 4\beta 2$ -nAChR.

Next, we derivatized the hydroxyl group with a variety of carbamate groups, a common functional group in medicinal chemistry that has been successfully employed in the design of other selective $\alpha 4\beta 2$ -nAChR ligands, in our efforts to further optimize the side chain.³² The carbamate analogs **13a–17a**, which contain the preferred (*1S,2R*)-configured cyclopropane ring, also exhibited sub-nanomolar to low nanomolar binding affinities at both the $\alpha 4\beta 2$ - and $\alpha 4\beta 2^*$ -nAChRs (Table 1). Their binding preference for the $\beta 2$ -over the $\beta 4$ -containing nAChRs was consistent with that of the corresponding alcohols and ethers. For both the $\alpha 3\beta 2$ - and $\alpha 4\beta 2^*$ -nAChRs, binding affinities gradually decreased as the size of substituents at the carbamate nitrogen increased.

Additionally, radioligand competition binding assays revealed the cyclopropane ligands tested (**12a–17a**, **19a**, **12b**, and **19b**) to have very low affinity for $\alpha 7$ -nAChRs. $10\text{ }\mu\text{M}$ concentrations of test ligand inhibited binding of 10 nM ^3H -epibatidine by a maximum of 30% (**16a**) with other compounds showing less inhibition or no inhibition of radioligand binding at all (data not shown).

For functional studies, all compounds were tested in $^{86}\text{Rb}^+$ ion flux assays using SH-EP1-h $\alpha 4\beta 2$ cells, which heterologously and stably express human $\alpha 4\beta 2$ -nAChRs assembled from individual subunits.^{33, 34} SH-SY5Y and TE671/RD cells were used to assess activities of tested compounds at human $\alpha 3\beta 4^*$ - or $\alpha 1\beta 1\gamma\delta$ -nAChRs, respectively.^{35, 36} All of the cyclopropane ligands had agonist activity at $\alpha 4\beta 2$ -nAChRs with EC_{50} values $< 50\text{ nM}$ (Table 2). Consistent with the radioligand binding studies, compounds **12a** and **19a** bearing the (*1S,2R*)-configured cyclopropane ring exhibited about 3-fold higher potencies than their diastereoisomers **12b** and **19b** (EC_{50} : 10.2 vs 34.6 nM ; 17.5 vs 43.1 nM). The functional agonism in the carbamate series tended to decline stepwise with the increasing size of the terminal substituents of the side chain. All compounds were found to functionally inactivate the response of the $\alpha 4\beta 2$ -nAChRs to a full agonist at IC_{50} values similar to the agonism EC_{50} values (Table 2). All tested ligands had neither agonist nor antagonist activity at ganglionic $\alpha 3\beta 4^*$ - or muscle-type $\alpha 1\beta 1\gamma\delta$ -nAChRs even at the highest concentration ($10\text{ }\mu\text{M}$) tested.

In the functional agonism studies, the efficacies of the tested compounds were determined in a mixed population of high sensitivity (HS) and low sensitivity (LS) $\alpha 4\beta 2$ -nAChRs. The efficacy values at the HS $\alpha 4\beta 2$ -nAChRs were extrapolated using compound **4** defined as a full agonist at the HS $\alpha 4\beta 2$ -nAChR with 100% efficacy (see Supporting Information for

more details).²⁰ All of the tested ligands were found to be partial agonists at HS $\alpha 4\beta 2$ -nAChRs with efficacy values ranging from 60 to 92%.

In Vivo Behavioral Pharmacology

In order to assess the antidepressant effects of selected compounds *in vivo*, we used the mouse forced swim test,³⁷ an assay in which mice are placed into a beaker of water and the time the mouse spends passively floating in the water (immobility) is recorded. Most traditional anti-depressants decrease the amount of time the mouse spends immobile. Mice were administered the most potent compounds **12a**, **13a**, and **19a**, or the selective serotonin reuptake inhibitor, sertraline, as a positive control (20 mg/kg) (Figure 2).

All of the three tested compounds exhibited antidepressant-like effects at the minimal dose of 10 mg/kg (compound **13a**) or 3 mg/kg (compounds **12a** and **19a**). Receptor occupancy (RO) studies were also performed to quantify the relationship between drug concentration at the receptor and the observed antidepressant effects.³⁸ When tested at a dose of 10 mg/kg, both the compounds **12a** and **19a** showed very high levels of *ex vivo* receptor occupancy (85 to 95%) at the $\beta 2^*$ receptors, whereas the carbamate analog **13a** showed only approximately 65% occupancy (Figure 3). These RO findings are consistent with their observed antidepressant potencies, with compounds **12a** and **19a** being more potent and compound **13a** being less potent (Figure 2).

Broad Screening and Preliminary ADMET Studies

Apart from assessment of ligand interactions with nAChRs, a broad-ranging screen was carried out for compounds **12a**, **13a**, and **19a** to determine their off-target binding at fifty-three other neurotransmitter receptors and transporters that are widely distributed throughout the CNS. The PDSP broad screening studies indicated that none of the three tested compounds showed significant interactions with other neurotransmitter receptors and transporters (see Supporting Information for more details).

Compounds **12a**, **13a**, and **19a** were further tested in preliminary ADMET assays.³⁹ When incubated with human or mouse liver microsomes, at least 80% of compound **12a**, 98% of compound **13a**, or 84% of compound **19a** remained unchanged after 1 h incubation at 1 μ M. In the presence of compounds **12a**, **13a**, or **19a** at concentrations up to 10 μ M, none of the CYP isoforms tested (CYP1A, CYP2C9, CYP2C19, CYP2D6, and CYP3A) showed more than 25% inhibition, indicating minimal adverse drug-drug interactions, with the exception of compound **13a** and **19a**, which displayed about 80% and 70% inhibition of the CYP1A2 isoform, respectively. The plasma protein binding of compound **13a** was investigated using human and mouse plasma (CD-1) at 10 μ M. A mean percent binding of 8.4 and 23.8, respectively, was observed. Lastly, automated patch-clamp electrophysiology was employed to measure the inhibitory interactions between the test compounds (**12a**, **13a**, and **19a**) and hERG K⁺ channels using CHO cell lines in three test concentrations (0.1, 1, and 10 μ M). At the highest concentration, compounds **12a**, **13a**, and **19a** exhibited 19.1%, 16.6%, and 13.7% inhibition of tail current, respectively, indicating minimal potential for hERG-related cardiovascular toxicity.

In order to further explore the metabolic stability of these cyclopropane ligands, compound **19a** was selected for full mouse *in vivo* pharmacokinetic (PK) studies. The plasma and brain concentrations of compound **19a** in male CD-1 mice after a single oral gavage administration at a dose of 5 mg/kg were measured. Compound **19a** possessed a reasonable half-life in brain ($t_{1/2}$ = 150 min) as well as in plasma ($t_{1/2}$ = 144 min). The concentration of compound **19a** reached a value of 133 ng/mL ($C_{\max}$ -brain) in 30 min ($T_{\max}$ -brain) in brain, and of 359 ng/mL ($C_{\max}$ -plasma) in 10 min ($T_{\max}$ -plasma) in plasma. The brain to plasma ratio

of compound **19a** was found to be 0.37 ($C_{\text{max-brain}}/C_{\text{max-plasma}}$) indicating acceptable CNS penetration. In contrast, brain exposure levels of compound **4** were relatively low (~ 3 ng/g at 1 mg/kg or ~ 10 ng/g at 3 mg/kg) when measured 15 min after administration and were at or below detection level at later time points.³⁸

Conclusion

In summary, a series of chiral cyclopropane analogs of the lead structure, compound **4**, were identified as highly potent, $\alpha 4\beta 2$ -selective nAChR partial agonists. In order to avoid possible issues relating to the metabolic instability of the acetylene bond, a rigid cyclopropane ring was introduced in its place. The cyclopropane ring is also virtuous because of its ability to direct the orientation of the side chain in a manner that improves subtype selectivity for $\alpha 4\beta 2$ -nAChRs. The best compounds, **12a**, **13a**, and **19a**, exhibited sub-nanomolar to low-nanomolar binding affinity for both $\alpha 4\beta 2$ - and $\alpha 4\beta 2^*$ -nAChRs with negligible interaction with $\alpha 3\beta 4$ -nAChRs. In functional studies, these ligands acted as highly potent, partial agonists at HS $\alpha 4\beta 2$ -nAChRs and were totally inactive at both ganglionic $\alpha 3\beta 4^*$ - or muscle-type $\alpha 1\beta 1\gamma\delta$ -nAChRs. Compounds **12a**, **13a**, and **19a** were found to display antidepressant-like properties in the mouse forced swim test, associated with high levels of $\beta 2^*$ receptor occupancy. Furthermore, our findings that these three compounds lack any significant off-target activities and show favorable ADMET profiles commend these chiral cyclopropane ligands as potential drug candidates for the treatment of depression.

Experimental Section

General

All chemicals were purchased from Sigma-Aldrich or Chem-Impex, and solvents were used as obtained from Fisher Scientific or Sigma-Aldrich without further purification. Anhydrous THF and CH_2Cl_2 were obtained by distillation over sodium wire or CaH_2 , respectively. All non-aqueous reactions were run under an argon atmosphere with exclusion of moisture from reagents, and all reaction vessels were oven-dried. The progress of reactions was monitored by TLC on SiO_2 . Spots were visualized by their quenching of the fluorescence of an indicator admixed to the SiO_2 layer, or by dipping into I_2/SiO_2 mixture. Products were purified by column chromatography on 230–400 mesh SiO_2 . Proton and carbon NMR spectra were recorded at spectrometer frequencies of 400 MHz and 100 MHz, respectively. NMR chemical shifts were reported in δ (ppm) using the δ 7.26 signal of CHCl_3 (^1H NMR), the δ 4.80 signal of HDO (^1H NMR), and the δ 77.23 signal of CDCl_3 (^{13}C NMR) as internal standards. ^{13}C NMR spectra in D_2O were not adjusted. Optical rotation was detected on an Autopol IV automatic polarimeter. Mass spectra were measured in the ESI mode at an ionization potential of 70 eV with an LC-MS MSD (Hewlett Packard). The final compounds were purified by preparative HPLC, which was carried out on an ACE 5 AQ column (150×20 mm), with detection at 254 and 280 nm on a Shimadzu SPD-10A VP detector; flow rate = 17.0 mL/min; gradient of 0 to 50% methanol in water (both containing 0.05 vol% of CF_3COOH) in 30 min. Purities of final compounds ($> 98\%$) were established by both elemental analysis and by analytical HPLC, which was carried out on an Agilent 1100 HPLC system with a Synergi $4 \mu\text{m}$ Hydro-RP 80A column, with detection at 254 or 280 nm on a variable wavelength detector G1314A; flow rate = 1.4 mL/min; gradient of 0 to 100% methanol in water (both containing 0.05 vol% of CF_3COOH) in 18 min. See Supporting Information for detailed experimental procedures and NMR spectral data (^1H and ^{13}C) of all intermediates.

3-[(2(S)-Azetidinyl)methoxy]-5-[(1S,2R)-2-(2-hydroxyethyl)cyclopropyl]pyridine Trifluoroacetate (12a)

^1H NMR (D_2O): δ 8.33 (s, 1H), 8.23 (s, 1H), 7.85 (s, 1H), 4.98 (m, 1H), 4.53 (d, J = 4.0 Hz, 2H), 4.17-4.07 (m, 2H), 3.71 (t, J = 6.4 Hz, 2H), 2.70 (q, J = 8.4 Hz, 2H), 1.98 (m, 1H), 1.68 (q, J = 6.8 Hz, 2H), 1.33 (m, 1H), 1.20-1.14 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 155.8, 146.3, 132.0, 128.0, 125.3, 115.9 (TFA), 67.1, 60.8, 58.2, 43.3, 35.1, 21.9, 19.8, 19.4, 16.3; $[\alpha]_{\text{D}}^{20}$ = +36.5 (c 0.40, MeOH); Anal. Calcd for $\text{C}_{14}\text{H}_{20}\text{N}_2\text{O}_2 \cdot 2.15\text{CF}_3\text{COOH} \cdot 0.5\text{H}_2\text{O}$: C, 43.74; H, 4.64; F, 24.39; N, 5.57. Found: C, 43.55; H, 4.42; F, 24.38; N, 5.52.

3-[(2(S)-Azetidinyl)methoxy]-5-[(1R,2S)-2-(2-hydroxyethyl)cyclopropyl]pyridine Trifluoroacetate (12b)

^1H NMR (400 MHz, D_2O): δ 8.32 (s, 1H), 8.22 (s, 1H), 7.84 (s, 1H), 4.97 (m, 1H), 4.52 (d, J = 4.0 Hz, 2H), 4.16-4.06 (m, 2H), 3.70 (t, J = 6.4 Hz, 2H), 2.69 (q, J = 8.4 Hz, 2H), 1.97 (m, 1H), 1.67 (q, J = 6.8 Hz, 2H), 1.32 (m, 1H), 1.17-1.12 (m, 2H); ^{13}C NMR (100 MHz, D_2O): δ 162.3 (TFA), 155.8, 146.3, 132.0, 127.9, 125.3, 115.9 (TFA), 67.1, 60.8, 58.3, 43.3, 35.1, 21.9, 19.8, 19.4, 16.3; $[\alpha]_{\text{D}}^{20}$ = -49.4 (c 0.17, MeOH); Anal. Calcd for $\text{C}_{14}\text{H}_{20}\text{N}_2\text{O}_2 \cdot 2\text{CF}_3\text{COOH} \cdot 0.15\text{H}_2\text{O}$: C, 45.13; H, 4.69; F, 23.79; N, 5.85. Found: C, 45.10; H, 4.67; F, 23.90; N, 5.84.

3-[(2(S)-Azetidinyl)methoxy]-5-[(1S,2R)-2-[2-(N-methylcarbamoyloxy)ethyl]cyclopropyl]pyridine Trifluoroacetate (13a)

^1H NMR (D_2O): δ 8.35 (s, 1H), 8.23 (s, 1H), 7.84 (s, 1H), 4.98 (m, 1H), 4.52 (d, J = 3.6 Hz, 2H), 4.27-4.05 (m, 4H), 2.70 (q, J = 8.4 Hz, 2H), 2.62 (s, 3H), 1.99 (m, 1H), 1.88 (m, 1H), 1.65 (m, 1H), 1.34 (m, 1H), 1.16 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 159.2, 156.2, 146.5, 132.3, 128.2, 125.7, 115.9 (TFA), 67.5, 64.8, 58.6, 43.6, 32.6, 26.5, 22.3, 20.2, 19.8, 16.4; $[\alpha]_{\text{D}}^{20}$ = +43.3 (c 0.18, MeOH); Anal. Calcd for $\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}_3 \cdot 1.95\text{CF}_3\text{COOH} \cdot 1.8\text{H}_2\text{O}$: C, 42.67; H, 5.14; F, 19.84; N, 7.50. Found: C, 42.77; H, 4.99; F, 19.86; N, 7.35.

3-[(2(S)-Azetidinyl)methoxy]-5-[(1S,2R)-2-[2-(N,N-dimethylcarbamoyloxy)ethyl]cyclopropyl]-pyridine Trifluoroacetate (14a)

^1H NMR (D_2O): δ 8.31 (s, 1H), 8.19 (s, 1H), 7.80 (s, 1H), 4.94 (m, 1H), 4.48 (d, J = 4.0 Hz, 2H), 4.20-4.03 (m, 4H), 2.79 (s, 6H), 2.66 (q, J = 8.4 Hz, 2H), 1.95 (m, 1H), 1.83 (m, 1H), 1.65 (m, 1H), 1.33 (m, 1H), 1.11 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 157.8, 155.9, 146.2, 131.9, 127.9, 125.3, 115.9 (TFA), 67.1, 65.0, 58.2, 43.2, 35.1, 32.2, 21.9, 19.8, 19.4, 16.3; $[\alpha]_{\text{D}}^{20}$ = +31.3 (c 0.07, MeOH); Anal. Calcd for $\text{C}_{17}\text{H}_{25}\text{N}_3\text{O}_3 \cdot 2.15\text{CF}_3\text{COOH} \cdot 1.55\text{H}_2\text{O}$: C, 43.18; H, 5.15; F, 20.68; N, 7.09. Found: C, 43.35; H, 5.12; F, 20.51; N, 6.89.

3-[(2(S)-Azetidinyl)methoxy]-5-[(1S,2R)-2-[2-(N-cyclopropylcarbamoyloxy)ethyl]cyclopropyl]-pyridine Trifluoroacetate (15a)

^1H NMR (D_2O): δ 8.34 (s, 1H), 8.22 (s, 1H), 7.84 (s, 1H), 4.98 (m, 1H), 4.52 (d, J = 4.0 Hz, 2H), 4.19-4.05 (m, 4H), 2.69 (q, J = 8.4 Hz, 2H), 2.37 (m, 1H), 1.97 (m, 1H), 1.85 (m, 1H), 1.65 (m, 1H), 1.31 (m, 1H), 1.14 (m, 2H), 0.63 (m, 2H), 0.39 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 159.3, 155.9, 146.2, 131.9, 128.0, 125.4, 115.9 (TFA), 67.2, 64.4, 58.2, 43.3, 32.1, 21.9, 21.8, 19.8, 19.4, 16.1, 5.3, 5.2; $[\alpha]_{\text{D}}^{20}$ = +40.6 (c 0.68, MeOH); Anal. Calcd for $\text{C}_{18}\text{H}_{25}\text{N}_3\text{O}_3 \cdot 1.95\text{CF}_3\text{COOH} \cdot 0.7\text{H}_2\text{O}$: C, 46.44; H, 5.05; F, 19.62; N, 7.42. Found: C, 46.19; H, 4.74; F, 19.42; N, 7.16.

3-[(2(S)-Azetidiny)methoxy]-5-[(1S,2R)-2-[2-(1-pyrrolidinylcarbonyloxy)ethyl]cyclopropyl]-pyridine Trifluoroacetate (16a)

^1H NMR (D_2O): δ 8.32 (s, 1H), 8.20 (s, 1H), 7.81 (s, 1H), 4.95 (m, 1H), 4.49 (d, J = 4.0 Hz, 2H), 4.20-4.04 (m, 4H), 3.22 (m, 4H), 2.67 (q, J = 8.8 Hz, 2H), 1.94 (m, 1H), 1.87-1.76 (m, 5H), 1.66 (m, 1H), 1.33 (m, 1H), 1.11 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 156.1, 155.9, 146.2, 131.9, 128.0, 125.3, 115.9 (TFA), 67.2, 64.7, 58.2, 45.5, 43.2, 32.2, 24.3, 21.9, 19.8, 19.4, 16.3; $[\alpha]_{\text{D}}^{20}$ = +39.7 (c 0.66, MeOH); Anal. Calcd for $\text{C}_{19}\text{H}_{27}\text{N}_3\text{O}_3 \cdot 2.25\text{CF}_3\text{COOH} \cdot 0.45\text{H}_2\text{O}$: C, 46.26; H, 4.98; F, 21.02; N, 6.89. Found: C, 46.11; H, 4.81; F, 20.95; N, 6.91.

3-[(2(S)-Azetidiny)methoxy]-5-[(1S,2R)-2-[2-(N-phenylcarbamoyloxy)ethyl]cyclopropyl]pyridine Trifluoroacetate (17a)

^1H NMR (D_2O): δ 8.02 (s, 1H), 7.96 (s, 1H), 7.58 (s, 1H), 7.20 (t, J = 8.0 Hz, 2H), 7.07 (d, J = 7.6 Hz, 2H), 7.01 (t, J = 7.2 Hz, 1H), 4.41 (m, 1H), 4.21 (m, 1H), 4.10 (m, 3H), 3.97 (m, 1H), 2.59 (m, 2H), 2.02 (m, 1H), 1.83 (m, 1H), 1.45 (m, 1H), 1.28 (m, 1H), 1.04 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 155.5, 154.9, 145.9, 137.2, 131.4, 128.6, 127.4, 125.0, 123.1, 118.1, 115.9 (TFA), 66.8, 64.1, 58.0, 43.2, 32.4, 21.7, 19.9, 19.8, 16.3; $[\alpha]_{\text{D}}^{20}$ = +36.4 (c 0.91, MeOH); Anal. Calcd for $\text{C}_{21}\text{H}_{25}\text{N}_3\text{O}_3 \cdot 2.15\text{CF}_3\text{COOH} \cdot 1.0\text{H}_2\text{O}$: C, 48.19; H, 4.66; F, 19.43; N, 6.66. Found: C, 48.16; H, 4.51; F, 19.48; N, 6.51.

3-[(2(S)-Azetidiny)methoxy]-5-[(1S,2R)-2-(2-methoxyethyl)cyclopropyl]pyridine Trifluoroacetate (19a)

^1H NMR (D_2O): δ 8.35 (s, 1H), 8.24 (s, 1H), 7.86 (s, 1H), 5.00 (m, 1H), 4.54 (d, J = 4.0 Hz, 2H), 4.18-4.09 (m, 2H), 3.61 (t, J = 6.4 Hz, 2H), 3.36 (s, 3H), 2.71 (q, J = 8.4 Hz, 2H), 2.00 (m, 1H), 1.74 (m, 2H), 1.34 (m, 1H), 1.21-1.15 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 155.9, 146.2, 132.0, 128.0, 125.4, 115.9 (TFA), 71.5, 67.1, 58.2, 57.3, 43.3, 32.2, 22.0, 19.8, 19.4, 16.2; $[\alpha]_{\text{D}}^{20}$ = +40.0 (c 0.20, MeOH); Anal. Calcd for $\text{C}_{15}\text{H}_{22}\text{N}_2\text{O}_2 \cdot 2.05\text{CF}_3\text{COOH} \cdot 0.5\text{H}_2\text{O}$: C, 45.42; H, 5.00; F, 23.13; N, 5.55. Found: C, 45.43; H, 4.90; F, 23.05; N, 5.57.

3-[(2(S)-Azetidiny)methoxy]-5-[(1R,2S)-2-(2-methoxyethyl)cyclopropyl]pyridine Trifluoroacetate (19b)

^1H NMR (D_2O): δ 8.31 (s, 1H), 8.20 (s, 1H), 7.82 (s, 1H), 4.96 (m, 1H), 4.50 (d, J = 4.0 Hz, 2H), 4.14-4.05 (m, 2H), 3.55 (t, J = 6.4 Hz, 2H), 3.31 (s, 3H), 2.67 (q, J = 8.4 Hz, 2H), 1.96 (m, 1H), 1.69 (m, 2H), 1.28 (m, 1H), 1.16-1.10 (m, 2H); ^{13}C NMR (D_2O): δ 162.3 (TFA), 155.8, 146.2, 131.9, 127.9, 125.3, 115.9 (TFA), 71.4, 67.1, 58.3, 57.3, 43.3, 32.2, 21.9, 19.8, 19.4, 16.2; $[\alpha]_{\text{D}}^{20}$ = -42.9 (c 0.84, MeOH); Anal. Calcd for $\text{C}_{15}\text{H}_{22}\text{N}_2\text{O}_2 \cdot 2.05\text{CF}_3\text{COOH} \cdot 0.6\text{H}_2\text{O}$: C, 45.26; H, 5.02; F, 23.05; N, 5.53. Found: C, 45.22; H, 4.88; F, 22.91; N, 5.47.

In Vitro Studies

$[\text{}^3\text{H}]$ Epibatidine competition studies: For experimental details, please refer to the PDSP web site <http://pdsp.med.unc.edu/> for all nAChR subtypes except $\alpha 7$. For assay at $\alpha 7$, membrane preparations from SH-EP1 cells heterologously expressing human $\alpha 7$ nAChRs were used to test 10 μM concentrations of test ligands in competition with 10 nM $[\text{}^3\text{H}]$ epibatidine with an experimental protocol similar to that utilized by the PDSP.^{33, 34}

Cell lines and culture

Cell lines naturally or heterologously expressing specific, functional, human nAChR subtypes were used. The human clonal cell line TE671/RD naturally expresses human muscle-type $\alpha 1^*$ -nAChRs, containing $\alpha 1$, $\beta 1$, γ , and δ subunits, with function detectable

using $^{86}\text{Rb}^+$ efflux assays.³⁶ The human neuroblastoma cell line SH-SY5Y naturally expresses autonomic $\alpha 3\beta 4^*$ -nAChRs, containing $\alpha 3$, $\beta 4$, probably $\alpha 5$, and sometimes $\beta 2$ subunits, and also displays function detectable using $^{86}\text{Rb}^+$ efflux assays.³³ SH-SY5Y cells also express homopentameric $\alpha 7$ -nAChRs; however, their function is not detected in the $^{86}\text{Rb}^+$ efflux assay under the conditions used. SH-EP1 human epithelial cells stably transfected with cDNAs separately encoding human $\alpha 4$ or $\beta 2$ subunits (SH-EP1-h $\alpha 4\beta 2$ cells) have been established and characterized with both ion flux and radioligand binding assays.³⁴ These cells thus express a mixture of so-called “high sensitivity” (HS) $\alpha 4\beta 2$ -nAChRs, having the presumed subunit ratios of 2:3 $\alpha 4$: $\beta 2$ and comparatively high sensitivity to nicotinic agonists, and “low sensitivity” (LS) $\alpha 4\beta 2$ -nAChRs presumably having a 3:2 ratio of $\alpha 4$: $\beta 2$ subunits at which nicotinic agonists have lower observed potency.¹⁷

TE671/RD, SH-SY5Y, and transfected SH-EP1 cell lines were maintained as low passage number (1–26 from our frozen stocks) cultures to ensure stable expression of native or heterologously expressed nAChRs as previously described. Cells were passaged once a week by splitting just-confluent cultures 1/300 (TE671/RD), 1/10 (SH-SY5Y) or 1/40 (transfected SH-EP1) in serum-supplemented medium to maintain log-phase growth.

General Procedures for Behavioral Studies

Animals

BALB/cJ male mice (8–10 weeks old at testing) were obtained from Jackson Laboratory (Bar Harbor, ME, USA). Mice were housed four to a cage in a colony room maintained at 22 ± 2 °C on a 12 h light–dark cycle. All animal experiments were conducted in accordance with the NIH Guide for the Care and Use of Laboratory Animals and the PsychoGenics Animal Care and Use Committee.

Drugs

Compounds **12a**, **13a**, and **19a** were synthesized according to procedures described in the text, and sertraline was purchased from Toronto Research Chemicals (Ontario, Canada). All compounds were dissolved in injectable water and administered by oral gavage (PO) in a volume of 10 mL/kg.

Mouse forced swim test

Procedures were based on those previously described. Mice were individually placed into clear glass cylinders (15 cm tall $\times$ 10 cm diameter, 1 L beakers) containing 23 ± 1 °C water 12 cm deep (approximately 800 mL). Mice were administered vehicle, the SSRI sertraline (10 or 20 mg/kg; IP or PO) as a positive control, or compounds **12a** (PO), **13a** (PO), and **19a** (PO). Thirty minutes after compound administration, mice were placed in the water, and the time the animal spent immobile was recorded over a 6 min trial. Immobility was defined as the postural position of floating in the water.

Statistical Analysis

Data were analyzed with Analysis of Variance (ANOVA) with Treatment Group (Vehicle, Sertraline, compounds **12a**, **13a**, and **19a**) as the between group variable and total time immobile in sec (over the 6 min trial) as the dependent variable. Significant main effects were followed up with the post hoc Fisher’s PLSD test.

$\beta 2^*$ -nAChR *ex vivo* receptor occupancy

Compounds **12a**, **13a**, and **19a** (10 mg/kg) or water were administered via intraperitoneal injection 30 min before brain collection (the same time point as in forced swim testing) for

analysis of $\beta 2$ -nAChR occupancy in the thalamus (for compound **12a** and **19a**, n=6; for compound **13a**, n=4) as described before.³⁸

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Abbreviations

nAChR(s)	nicotinic acetylcholine receptor(s)
CNS	central nervous system
PNS	peripheral nervous system
DA	dopamine
5-HT	serotonin
Glu	glutamate
ACh	acetylcholine
GABA	γ -aminobutyric acid
ADHD	attention deficit hyperactivity disorder
MDD	major depressive disorder
ee	enantiomeric excess
HS	high-sensitivity
LS	low-sensitivity
ADMET	absorption, distribution, metabolism, excretion and toxicity
hERG	human ether-a-go-go-related gene
PK	pharmacokinetic
TFA	trifluoroacetic acid

Acknowledgments

This research was supported by award No. U19MH085193 from the National Institute of Mental Health. The Phoenix research component was also supported in part by the Barrow Neurological Foundation, and work by this team was conducted in part in the Charlotte and Harold Simensky Neurochemistry of Alzheimer's Disease Laboratory. We thank the PDSP program for assistance through providing the binding affinity data.

REFERENCES

1. Bunnelle WH, Dart MJ, Schrimpf MR. Design of ligands for the nicotinic acetylcholine receptors: the quest for selectivity. *Curr. Top. Med. Chem.* 2004; 4:299–334. [PubMed: 14754449]
2. Jensen AA, Frolund B, Liljefors T, Krogsgaard-Larsen P. Neuronal nicotinic acetylcholine receptors: structural revelations, target identifications, and therapeutic inspirations. *J. Med. Chem.* 2005; 48:4705–4745. [PubMed: 16033252]
3. Romanelli MN, Gratteri P, Guandalini L, Martini E, Bonaccini C, Gualtieri F. Central nicotinic receptors: structure, function, ligands, and therapeutic potential. *Chem Med Chem.* 2007; 2:746–767. [PubMed: 17295372]
4. Taly A, Corringer PJ, Guedin D, Lestage P, Changeux JP. Nicotinic receptors: allosteric transitions and therapeutic targets in the nervous system. *Nat. Rev. Drug Discov.* 2009; 8:733–750. [PubMed: 19721446]

5. Gotti C, Clementi F, Fornari A, Gaimarri A, Guiducci S, Manfredi I, Moretti M, Pedrazzi P, Pucci L, Zoli M. Structural and functional diversity of native brain neuronal nicotinic receptors. *Biochem. Pharmacol.* 2009; 78:703–711. [PubMed: 19481063]
6. de Filippi G, Mogg AJ, Phillips KG, Zwart R, Sher E, Chen Y. The subtype-selective nicotinic acetylcholine receptor positive allosteric potentiator 2087101 differentially facilitates neurotransmission in the brain. *Eur. J. Pharmacol.* 2010; 643:218–224. [PubMed: 20624387]
7. Picciotto MR, Zoli M, Rimondini R, Lena C, Marubio LM, Pich EM, Fuxe K, Changeux JP. Acetylcholine receptors containing the $\beta 2$ subunit are involved in the reinforcing properties of nicotine. *Nature.* 1998; 391:173–177. [PubMed: 9428762]
8. Tapper AR, McKinney SL, Nashmi R, Schwarz J, Deshpande P, Labarca C, Whiteaker P, Marks MJ, Collins AC, Lester HA. Nicotine activation of $\alpha 4^*$ receptors: sufficient for reward, tolerance, and sensitization. *Science.* 2004; 306:1029–1032. [PubMed: 15528443]
9. Rabenstein RL, Caldarone BJ, Picciotto MR. The nicotinic antagonist mecamylamine has antidepressant-like effects in wild-type but not $\beta 2$ - or $\alpha 7$ -nicotinic acetylcholine receptor subunit knockout mice. *Psychopharmacology (Berl).* 2006; 189:395–401. [PubMed: 17016705]
10. Caldarone BJ, Harist A, Cleary MA, Beech RD, King SL, Picciotto MR. High-affinity nicotinic acetylcholine receptors are required for antidepressant effects of amitriptyline on behavior and hippocampal cell proliferation. *Biol. Psychiatry.* 2004; 56:657–664. [PubMed: 15522249]
11. Labarca C, Schwarz J, Deshpande P, Schwarz S, Nowak MW, Fonck C, Nashmi R, Kofuji P, Dang H, Shi W, Fidan M, Khakh BS, Chen Z, Bowers BJ, Boulter J, Wehner JM, Lester HA. Point mutant mice with hypersensitive $\alpha 4$ nicotinic receptors show dopaminergic deficits and increased anxiety. *Proc. Natl. Acad. Sci. U. S. A.* 2001; 98:2786–2791. [PubMed: 11226318]
12. Kroes RA, Burgdorf J, Otto NJ, Panksepp J, Moskal JR. Social defeat, a paradigm of depression in rats that elicits 22-kHz vocalizations, preferentially activates the cholinergic signaling pathway in the periaqueductal gray. *Behav. Brain Res.* 2007; 182:290–300. [PubMed: 17452055]
13. Wilens TE, Gault LM, Childress A, Kratochvil CJ, Bensman L, Hall CM, Olson E, Robieson WZ, Garimella TS, Abi-Saab WM, Apostol G, Saltarelli MD. Safety and efficacy of ABT-089 in pediatric attention-deficit/hyperactivity disorder: results from two randomized placebo-controlled clinical trials. *J. Am. Acad. Child Adolesc. Psychiatry.* 2011; 50:73–84. e1. [PubMed: 21156272]
14. Lippiello PM, Beaver JS, Gatto GJ, James JW, Jordan KG, Traina VM, Xie J, Bencherif M. TC-5214 (S-(+)-mecamylamine): a neuronal nicotinic receptor modulator with antidepressant activity. *CNS Neurosci. Ther.* 2008; 14:266–277. [PubMed: 19040552]
15. Rollema H, Coe JW, Chambers LK, Hurst RS, Stahl SM, Williams KE. Rationale, pharmacology and clinical efficacy of partial agonists of $\alpha 4\beta 2$ nACh receptors for smoking cessation. *Trends Pharmacol. Sci.* 2007; 28:316–325. [PubMed: 17573127]
16. Xiao Y, Fan H, Musachio JL, Wei ZL, Chellappan SK, Kozikowski AP, Kellar KJ. Sazetidine-A, a novel ligand that desensitizes $\alpha 4\beta 2$ nicotinic acetylcholine receptors without activating them. *Mol. Pharmacol.* 2006; 70:1454–1460. [PubMed: 16857741]
17. Kozikowski AP, Eaton JB, Bajjuri KM, Chellappan SK, Chen Y, Karadi S, He R, Caldarone B, Manzano M, Yuen PW, Lukas RJ. Chemistry and pharmacology of nicotinic ligands based on 6-[5-(azetidin-2-ylmethoxy)pyridin-3-yl]hex-5-yn-1-ol (AMOP-H-OH) for possible use in depression. *Chem Med Chem.* 2009; 4:1279–1291. [PubMed: 19569163]
18. Turner JR, Castellano LM, Blendy JA. Nicotinic partial agonists varenicline and sazetidine-A have differential effects on affective behavior. *J. Pharmacol. Exp. Ther.* 2010; 334:665–672. [PubMed: 20435920]
19. Paterson NE, Min W, Hackett A, Lowe D, Hanania T, Caldarone B, Ghavami A. The high-affinity nAChR partial agonists varenicline and sazetidine-A exhibit reinforcing properties in rats. *Prog. Neuropsychopharmacol. Biol. Psychiatry.* 2010; 34:1455–1464. [PubMed: 20708056]
20. Zwart R, Carbone AL, Moroni M, Bermudez I, Mogg AJ, Folly EA, Broad LM, Williams AC, Zhang D, Ding C, Heinz BA, Sher E. Sazetidine-A is a potent and selective agonist at native and recombinant $\alpha 4\beta 2$ nicotinic acetylcholine receptors. *Mol. Pharmacol.* 2008; 73:1838–1843. [PubMed: 18367540]
21. Cucchiari G, Xiao Y, Gonzalez-Sulser A, Kellar KJ. Analgesic effects of Sazetidine-A, a new nicotinic cholinergic drug. *Anesthesiology.* 2008; 109:512–519. [PubMed: 18719450]

22. Chen H, Chen W, Gan LS, Mutlib AE. Metabolism of (S)-5,6-difluoro-4-cyclopropylethynyl-4-trifluoromethyl-3, 4-dihydro-2(1H)-quinazolinone, a non-nucleoside reverse transcriptase inhibitor, in human liver microsomes. Metabolic activation and enzyme kinetics. *Drug Metab. Dispos.* 2003; 31:122–132. [PubMed: 12485961]
23. Salaun J, Baird MS. Biologically active cyclopropanes and cyclopropenes. *Curr. Med. Chem.* 1995; 2:511–542.
24. Kozhushkov SI, Schill H. Three-Membered-Ring-Based Molecular Architectures. *Chem. Rev.* 2006; 106:4926–4996. [PubMed: 17165680]
25. Watanabe M, Hirokawa T, Kobayashi T, Yoshida A, Ito Y, Yamada S, Orimoto N, Yamasaki Y, Arisawa M, Shuto S. Investigation of the bioactive conformation of histamine H₃ receptor antagonists by the cyclopropylic strain-based conformational restriction strategy. *J. Med. Chem.* 2010; 53:3585–3593. [PubMed: 20397706]
26. Liu J, Yu LF, Eaton JB, Caldarone B, Cavino K, Ruiz C, Terry M, Fedolak A, Wang D, Ghavami A, Lowe DA, Brunner D, Lukas RJ, Kozikowski AP. Discovery of Isoxazole Analogues of Sazetidine-A as Selective $\alpha 4\beta 2$ -Nicotinic Acetylcholine Receptor Partial Agonists for the Treatment of Depression. *J. Med. Chem.* 2011; 54:7280–7288. [PubMed: 21905669]
27. Cui X, Zhou Y, Wang N, Liu L, Guo Q-X. *N*-Phenylurea as an inexpensive and efficient ligand for Pd-catalyzed Heck and room-temperature Suzuki reactions. *Tetrahedron Lett.* 2007; 48:163–167.
28. Toy PH, Dhanabalasingam B, Newcomb M, Hanna IH, Hollenberg PF. A Substituted Hypersensitive Radical Probe for Enzyme-Catalyzed Hydroxylations: Synthesis of Racemic and Enantiomerically Enriched Forms and Application in a Cytochrome P450-Catalyzed Oxidation. *J. Org. Chem.* 1997; 62:9114–9122.
29. The Charette protocol was also applied to the allylic alcohol corresponding to ester **6** to obtain alcohol **8a** with a modest ee of 77%. The major product was found to be identical with the (*S,S*)-isomer **8a**, an outcome which is consistent with the prediction using Charette's model.
30. Charette AB, Juteau H, Lebel H, Molinaro C. Enantioselective Cyclopropanation of Allylic Alcohols with Dioxaborolane Ligands: Scope and Synthetic Applications. *J. Am. Chem. Soc.* 1998; 120:11943–11952.
31. Exley R, Clements MA, Hartung H, McIntosh JM, Cragg SJ. $\alpha 6$ -containing nicotinic acetylcholine receptors dominate the nicotine control of dopamine neurotransmission in nucleus accumbens. *Neuropsychopharmacology.* 2008; 33:2158–2166. [PubMed: 18033235]
32. Hansen CP, Jensen AA, Christensen JK, Balle T, Liljefors T, Frolund B. Novel Acetylcholine and Carbamoylcholine Analogues: Development of a Functionally Selective $\alpha 4\beta 2$ Nicotinic Acetylcholine Receptor Agonist. *J. Med. Chem.* 2008; 51:7380–7395. [PubMed: 18989912]
33. Lukas, RJ.; Fryer, JD.; Eaton, JB.; Gentry, CL. Some methods for studies of nicotinic acetylcholine receptor pharmacology. In: Levin, ED., editor. *Nicotinic Receptors and the Nervous System*. Boca Raton: CRC Press; 2002. p. 3-27.
34. Eaton JB, Peng JH, Schroeder KM, George AA, Fryer JD, Krishnan C, Buhlman L, Kuo YP, Steinlein O, Lukas RJ. Characterization of human $\alpha 4\beta 2$ -nicotinic acetylcholine receptors stably and heterologously expressed in native nicotinic receptor-null SH-EP1 human epithelial cells. *Mol. Pharmacol.* 2003; 64:1283–1294. [PubMed: 14645658]
35. Lukas RJ, Norman SA, Lucero L. Characterization of Nicotinic Acetylcholine Receptors Expressed by Cells of the SH-SY5Y Human Neuroblastoma Clonal Line. *Mol. Cell. Neurosci.* 1993; 4:1–12. [PubMed: 19912902]
36. Lukas, RJ. Cell lines as models for studies of nicotinic acetylcholine receptors. In: Arneric, SP.; Brioni, JD., editors. *Neuronal Nicotinic Receptors: Pharmacology and Therapeutic Opportunities*. New York: Wiley-Liss, Inc; 1999. p. 81-97.
37. Porsolt RD, Bertin A, Jalfre M. Behavioral despair in mice: a primary screening test for antidepressants. *Arch. Int. Pharmacodyn. Ther.* 1977; 229:327–336. [PubMed: 596982]
38. Caldarone BJ, Wang D, Paterson NE, Manzano M, Fedolak A, Cavino K, Kwan M, Hanania T, Chellappan SK, Kozikowski AP, Olivier B, Picciotto MR, Ghavami A. Dissociation between duration of action in the forced swim test in mice and nicotinic acetylcholine receptor occupancy with sazetidine, varenicline, and 5-I-A85380. *Psychopharmacology (Berl.)* 2011; 217:199–210. [PubMed: 21487659]

39. The preliminary ADMET studies were carried out by Cerep, Inc.

ABT-089 (1)

Cognitive dysfunction, Phase II

TC-5214 (2)

Depression, Phase III

Varenicline (3)

Smoking cessation, On the market

Sazetidine-A (4)

Figure 1.
Selected examples of synthetic nAChR ligands

Figure 2.

Mouse forced swim data for compounds **12a** (A), **13a** (B), and **19a** (C). The selective serotonin reuptake inhibitor, sertraline, produced the expected decrease in immobility. (ANOVAs: $F(3,35) = 13.43$, $p < 0.001$ (A); $F(3,36) = 11.46$, $p < 0.001$ (B); $F(4,44) = 9.29$, $p < 0.001$ (C). *Fisher's PLSD post-hoc test: $ps < 0.05$ vs vehicle). All drugs were administered orally; $n = 9$ – 10 /group.

Figure 3.

Receptor occupancy studies of compounds **12a**, **13a**, and **19a** in mice showed a significant occupancy level. (*Mann-Whitney U; $p < 0.05$). All drugs were injected intraperitoneally; $n = 4-6$ /group.

Scheme 1.

^aReagents and conditions: (a) i. BuOH , NaH , DMF , rt; ii. *n*-butyl acrylate, 1% $\text{Pd}(\text{OAc})_2$, 2% PhNHCONH_2 , K_2CO_3 , 130°C ; (b) i. 2N NaOH , MeOH/THF (1:1), rt; ii. $\text{MeNH}_2^+\text{OMeCl}^-$, EDCI , DMAP , CH_2Cl_2 , rt; iii. $\text{Me}_3\text{S}(\text{O})^+\text{I}^-$, NaH , DMSO , rt; (c) i. DIBAL-H , THF , -78°C to -20°C ; ii. NaBH_4 , MeOH , rt; iii. Chiralpak AD, EtOH ; (d) i. $(\text{COCl})_2$, DMSO , Et_3N , CH_2Cl_2 , -78°C ; ii. NaClO_2 , KH_2PO_4 , 2-methyl-2-butene, *tert*-butanol/ H_2O , rt; (e) i. trimethylacetyl chloride, Et_3N , THF , -78°C to 0°C ; ii. (*S*)-(+)-4-phenyl-2-oxazolidinone lithium salt, THF , -78°C to 0°C ; (f) $(\text{COCl})_2$, DMSO , Et_3N , CH_2Cl_2 , -78°C ; (g) i. $\text{Ph}_3\text{P}=\text{CH}_2$, THF , 0°C ; ii. dicyclohexylborane, THF , 0°C to rt; iii. 30% H_2O_2 , 3N NaOH , 55°C ; (h) isobutyric anhydride, cat. DMAP , Et_3N , CH_2Cl_2 , rt; (i) i.

10% Pd/C, H₂, EtOAc/MeOH, rt; ii. 1-(*tert*-butoxycarbonyl)-(2*S*)-azetidinylmethanol, azodicarbonyldipiperidide (ADDP), P(*n*-Bu)₃, PhMe, 0 °C to rt; (j) NaOMe, MeOH, 40 °C; (k) 1,1'-carbonyldiimidazole, PhMe, THF, amine, rt; or phenyl isocyanate, PhMe, reflux; (l) CF₃COOH, CH₂Cl₂, rt; (m) Ph₃P=CHOCH₃, THF, 0 °C; (n) PtO₂, H₂, CH₂Cl₂, rt.

Table 1
Binding Affinities of Cyclopropane Ligands, Nicotine, and Sazetidine-A at Seven nAChR Subtypes

Compd.	K_i (nM) ^a						Selectivity
	$\alpha 2\beta 2$	$\alpha 2\beta 4$	$\alpha 3\beta 2$	$\alpha 3\beta 4$	$\alpha 4\beta 2$	$\alpha 4\beta 2^* b$	$\alpha 3\beta 4/ \alpha 4\beta 2$
12a	0.1	249 ^c	3.0±0.4	6,520	0.1	0.5±0.1	82.6±9
12b	0.5±0.1	65.0±7	17.0±7	1,040	0.7±0.1	3.7±0.6	29.0±4.3
13a	0.3±0.1	1,890	9.8±2.9	>10,000	0.6±0.1	1.7±0.2	441
14a	0.3±0.1	261	15.0±2.1	>10,000	0.4	2.1±0.3	281
15a	0.9±0.1	720	16.0±3	>10,000	0.7±0.1	2.4±0.3	234
16a	1.5±0.1	2,330	21.5±5.2	>10,000	0.8±0.1	4.0±0.3	162
17a	1.2±0.1	241	24.4±5.7	8,850	0.6±0.1	5.9±1.1	551
19a	0.1	236	2.4±0.4	>10,000	0.1	0.3	50.2±11
19b	0.5±0.1	405	20±7.3	>10,000	0.6±0.1	6.2±1.6	96.5±21
nicotine ^d	5.5	70	29	260	4.9	9.8	23
4 ^e				10,000	0.4	0.9	24,000

^a See Experimental Section.

^b $\alpha 4\beta 2^*$, prepared from rat forebrain.

^c SEM values are not provided for K_i values > 100 nM.

^d The binding data for nicotine are from the PDSP Assay Protocol Book (<http://pdsp.med.unc.edu/>).

^e The binding data for compound **4** were obtained from Reference 16.

Table 2

Sensitivities and Efficacies of Ligand Agonism and Inactivation at $\alpha 4\beta 2$ nAChRs^a

Compd.	Agonism			Desensitization	
	EC ₅₀ (nM)	Efficacy (%) ^b	Efficacy (%) HS ^c	IC ₅₀ (nM)	Efficacy (%) ^b
12a	10.2	21	92	9.4	63
12b	34.6	10	65	50.9	69
13a	15.7	17	77	18.2	85
14a	18.2	15	71	19.1	85
15a	23.3	27	80	19.6	82
16a	44.4	23	69	48.5	84
17a	48.9	24	78	50.2	87
19a	17.5	6	60	5.6	71
19b	43.1	8	62	50.4	75
nicotine ^d	290	88	--	430	93
4 ^d	5.8	55	100	4.8	63

^aSee Experimental Section.

^bThe efficacies were measured in a mixture of HS and LS $\alpha 4\beta 2$ -nAChRs.

^cThe efficacy values were extrapolated using compound **4** defined as a full agonist at the HS $\alpha 4\beta 2$ -nAChR (see Supporting Information for details).

^dResults for nicotine and compound **4** were obtained from Reference 17.