Published in final edited form as:

Int J Hyperthermia. 2012 ; 28(1): 9–18. doi:10.3109/02656736.2011.616182.

Effector CD8⁺ T cell IFN- γ production and cytotoxicity are enhanced by mild hyperthermia

Thomas A. Mace, Lingwen Zhong, Kathleen M. Kokolus, and Elizabeth A. Repasky¹

Department of Immunology, Roswell Park Cancer Institute, Buffalo, NY 14263

Abstract

Purpose—Clinical trials combining hyperthermia with radiation and/or chemotherapy for cancer treatment have resulted in improved overall survival and control of local recurrences. The contribution of thermally enhanced anti-immune function in these effects is of considerable interest, but not understood; studies on the fundamental effects of elevated temperature on immune effector cells are needed. The goal of this study is to investigate the potential of mild hyperthermia to impact tumor antigen-specific (Ag) effector CD8⁺ T cell functions.

Method—Pmel-1 Ag-specific CD8⁺ T cells were exposed to mild hyperthermia and tested for changes in IFN- γ production and cytotoxicity. Additionally, overall plasma membrane organization and the phosphorylation of signaling proteins were also investigated following heat treatment.

Results—Exposing effector Pmel-1 specific CD8⁺ T cells to mild hyperthermia (39.5°C) resulted in significantly enhanced Ag-specific IFN- γ production and tumor target cell killing compared to that seen using lower temperatures (33 and 37°C). Further, inhibition of protein synthesis during hyperthermia did not reduce subsequent Ag-induced IFN- γ production by CD8⁺ T cells. Correlated with these effects, we observed a distinct clustering of GM1⁺ lipid microdomains at the plasma membrane and enhanced phosphorylation of LAT and PKC θ which may be related to an observed enhancement of Ag-specific effector CD8⁺ T cell IFN- γ gene transcription following mild hyperthermia. However, mitogen-mediated production of IFN- γ , which bypasses T cell receptor activation with antigen, was not enhanced.

Conclusions—Antigen-dependent effector T cell activity is enhanced following mild hyperthermia. These effects could potentially occur in patients being treated with thermal therapies. These data also provide support for the use of thermal therapy as an adjuvant for immunotherapies to improve CD8⁺ effector cell function.

Keywords

T lymphocytes; fever; hyperthermia; cytotoxicity; interferon- γ

Introduction

Hyperthermia, used to elevate the temperature of tumors to between 40 and 45°C, is proving to be an important adjuvant in cancer therapy when combined with chemo and/or radiation therapy. More specifically, hyperthermia has been shown to improve clinical responses to

¹Correspondance to: Elizabeth A. Repasky, Ph.D., Professor of Oncology, Roswell Park Cancer Institute, Buffalo NY 14263, Phone: 716-845-3133, elizabeth.repasky@roswellpark.org.

Declaration of interests

The authors report no declarations of interest. This work was supported by grants NIH R01 CA135368-01A1, R01 CA071599-11, 2 T32 CA085183 Immunology Department Training Grant, and the Roswell Park Cancer Institute Core grant CA16056.

radiation in several clinical trials in patients with breast (1, 2), head and neck (3, 4), melanoma (5), cervical (6, 7), and brain (8, 9) cancers and improves response to chemoradiation therapy in patients with soft tissue sarcoma (10). Despite the growing documentation of the positive impact of hyperthermia as an adjuvant in cancer treatment, its use is still quite rare in comparison to other therapies. In part, this is due to a relative lack of new clinical trials that could document its effectiveness in various applications. However, the increased use of hyperthermia in cancer treatment may also be limited by a still-incomplete understanding of the precise mechanisms or cellular events by which temperature manipulation affects the tumor microenvironment or the efficacy of radiation or chemotherapy. Without more documentation of the complete range of cellular and/or molecular effects of temperature elevation, it is difficult to design the most effective clinical protocols.

Recently, there has been a growing appreciation for that fact that, due to physiological heat dissipating processes, the hyperthermic temperatures achieved in large regions of the tumor and surrounding area may be as mild as 39–40°C. Because this temperature range is not considered cytotoxic (at least not in treatment periods used in the clinic) the possibility exists that thermally sensitive activation of various cellular processes could be contributing to the beneficial effects seen with the addition of hyperthermia. One particularly attractive hypothesis is that the function of cells important for the anti-tumor immune system may be sensitive to thermal signals, possibility because of a natural, long-conserved sensitivity to elevated temperatures during fever. Based on a growing literature, it is apparent that the effect of hyperthermia on anti-tumor immunity is quite complex impacting various arms of the innate and adaptive immune response (11, 12). Excitingly, very recent studies in a clinically relevant animal model have demonstrated that adding hyperthermia to chemotherapy results in a curative therapy that is immunologically mediated (13). It has been widely reported that tumor infiltrating lymphocytes are essential for the inhibition of tumor growth and improved prognosis in various cancer types (14–19). Moreover, the specific immune cells located within tumors have been shown to be a favorable predictor of survival among colorectal cancer patients, even more accurate than histology (20). In particular, many of these observations indicate that the infiltration of CD8⁺ T cells into the tumor microenvironment is an immune mechanism benefiting patient survival. The effect of hyperthermia on effector CD8⁺ T cells is unknown in terms of antigen specific and T cell-dependent events. However, our lab has shown that hyperthermia enhances the rate of antigen-dependent contact hypersensitivity through a T cell-mediated response (21). This increased rate of hypersensitivity leads to an increased infiltration of lymphocytes accompanied by swelling at the site of antigen contact and blood vessel dilation. Further, our recent studies have indicated that increasing temperature enhances the early activation and differentiation of naïve CD8⁺ T cells into effector cells (Mace et al 2011, In press). Moreover, work by others (22) has shown that the differentiation of CD4⁺ T cells into T_H1 cells is accompanied by increased interferon gamma (IFN-γ) production and subsequent promotion of CD8⁺ T cell-mediated adaptive immune response. Hyperthermia trials have shown an effect on serum cytokine levels (23), but whether thermal treatment can directly influence IFN-γ production is also not known. However, since effector, but not naïve CD8⁺ T cells are able to directly kill tumor cells, it would be very important to determine whether hyperthermia could enhance the functions of this differentiated subset.

Here, we address the question of whether mild heat treatment can enhance tumor antigen-specific, effector CD8⁺ T cell functions, and in particular, their cytotoxicity and IFN-γ production. We also examined the effects of mild heating on the signaling pathway known to be engaged when CD8⁺ T cells are activated. Finally, we show that temperature induced effects on CD8⁺ T cell function are acting through events and signals mediated by the TCR in the plasma membrane. Collectively our data suggest that fever-range temperature can

enhance important CD8⁺ T cell functions and thereby may help to promote tumor control and improve long-term survival in patients receiving hyperthermia as part of their therapy.

Methods

Animals and cell lines

C57BL/6 mice were obtained from NCI (Bethesda, MA, USA). B6.Cg-*Thy1^a/Cy Tg* (TcraTcrb)8Rest/J (Pmel-1) mice were obtained from Jackson Laboratories (Bar Harbor, Maine). Mice were maintained in specific pathogen-free facilities and were treated in accordance with the guidelines established by the Animal Care and Use Committee at Roswell Park Cancer Institute (Buffalo, NY, USA). B16.F10 melanoma cells and EL-4 lymphoma cells were obtained from American Type Culture Collection (ATCC). Cells were cultured in 10% FBS, 10 mM L-glutamine, and 100 µg/ml penicillin/streptomycin in RPMI 1640 (Gibco).

Antibodies, peptide, and reagents

PE-conjugated anti-CD8 mAb (53–6.7), FITC-conjugated Thy1.1 mAb (OX-7), FITC-conjugated CD8 mAb (Ly-2), and PE-Cy5-conjugated CD8 (53–6.7) were purchased from BD Pharmingen (San Diego, CA). p-LAT was purchased from Millipore and p-PKCθ, and β-actin antibodies were purchased from Cell Signaling Technology (Boston, MA) for western blot analysis. gp100_{22–33} peptide was synthesized and purchased from JPT peptide (Berlin, Germany). FITC-Cholera Toxin B (CTxB) was purchased from Sigma (St. Louis, MO). Recombinant human IL-2 was purchased from R&D Systems (Minneapolis, MN).

In vitro generation of Pmel-1 effector CD8⁺ T cells

Pmel-1 splenocytes were cultured with 0.1 µg/ml of gp100_{25–33} peptide and 30 IU of rhIL-2 (R&D Systems) in 10% FBS, 2 mM 10 mM L-glutamine, and 100 µg/ml penicillin/streptomycin in RPMI (Gibco) for 5 days at 37°C. Cells were harvested and run over a ficoll gradient. Effector CD8⁺ T cell phenotype was confirmed by FACS analysis.

⁵¹Cr release assay for Ag-specific target cell killing

Target cells were labeled with 100–150 µCi of Na⁵¹CrO₄ (⁵¹Cr) (Perkin Elmer) at 37°C for 1 hour in FBS. After washing targets 3 times with 5% FBS in RPMI 1640 they were plated at 10⁴ ⁵¹Cr-labeled cells/well and mixed with effector Pmel-1 CD8⁺ T cells (incubated at indicated temperatures) in round bottom microtiter plates at indicated E:T ratios in triplicate. After 4 hours incubation at 37°C, plates were centrifuged for 5 min at 500 × g, and 50 µl of supernatant was collected and counted in a MicroBeta TriLux liquid scintillation counter (PerkinElmer). Percent-specific lysis was determined as (experimental release – spontaneous release)/(maximal release – spontaneous release) × 100.

ELISA

Supernatants were collected 18 hours after stimulating CD8⁺ T cells with C57BL/6 Ag-pulsed splenocytes. IFN-γ levels were measured in a sandwich ELISA using anti-IFN-γ capture mAb and biotin conjugated anti-IFN-γ mAb (XMG1.2) purchased from BD Pharmingen.

Cell Staining

To examine GM1 clustering, cells were adhered onto alcian blue coated cover-slips, washed with PBS, and fixed with 4% paraformaldehyde. Cells were stained with FITC-CTxB, visualized, and quantified by fluorescent microscopy.

Western Blot Analysis

Cells were collected by centrifugation and lysed in lysis buffer (20 mM Tris (pH 7.5), 120 mM NaCl, 100 mM NaF, 0.5% Nonionic P40) containing protease inhibitors (10 mM NaPPi, 4 mM PMSF, 1 µg/ml Leupeptine, 2 mM Benzamidine, and 10 µg/ml Aprotinin) and phosphatase inhibitors (200 µM sodium vanadate and 50 mM b-Glycerophosphate) for 30 min on ice. Lysates were centrifuged at $12,000 \times g$ for 10 min at 4°C and the supernatants were collected. Protein concentration in the supernatants was determined using the Pierce BCA protein assay kit. Lysates were boiled for 5 min in the presence of 1 × SDS-PAGE loading buffer and electrophoresed on a 10% SDS-polyacrylamide gel for 1.5 h at 100 V. The proteins were transferred to nitrocellulose and probed with indicated antibodies. The membrane was washed three times for 5 min with 0.1% Tween 20 and 1 × TBS. Detection was performed with the appropriate secondary Abs conjugated with HRP followed by ECL detection reagent (Pierce).

Quantitative RT-PCR Analysis

Total RNA was prepared from CD8⁺ T cells stimulated with Ag-pulsed splenocytes using RNeasy kit (Qiagen). cDNA was synthesized using oligo-dT primers and Superscript polymerase (Invitrogen). Quantitative real-time PCR was performed in an Applied Biosystems 7900HT Fast Real-Time PCR System using SYBR Green PCR kit from Applied Biosystems and specific primers to amplify 200 bp fragments from different genes analyzed. A threshold was set in the linear part of the amplification curve and the number of cycles needed to reach it was calculated for each gene. Normalization was performed using primers to GAPDH.

Statistical analysis

Results are expressed as mean $\pm$ standard deviation. Student's two-tailed t test was used for comparing experimental groups with p value < 0.05 considered significant.

Results

Ag-specific effector CD8⁺ T cell IFN- γ production is temperature sensitive

Previous work has shown that non-antigen specific T cell function has been enhanced with hyperthermia (24, 25). Moreover, our earlier work has demonstrated that elevated temperatures can enhance the differentiation of naïve CD8⁺ T cells into effector cells (Mace et al 2011, revised version under review). However, whether increased temperature affects Ag-specific, effector CD8⁺ T cell function is still unclear. Thus, we tested whether different physiological temperatures could regulate Ag-specific effector CD8⁺ T cell function, using antigen to achieve activation, rather than non-specific mitogens or crosslinking antibodies against surface receptors. Effector CD8⁺ T cells were generated from Pmel-1 splenocytes by stimulating cells for 5 days with 0.1 µg/ml gp100₂₅₋₃₃ peptide and 30 IU rIL-2. Effector CD8⁺ T cells were then incubated at 33, 37, or 39.5°C for 6 hours and re-stimulated with gp100₂₃₋₃₃ peptide-pulsed C57BL/6 splenocytes for 18 hours. First we examined production of IFN- γ because of the importance of this molecule in upregulating MHC class I expression on tumor cells, allowing them to be recognized by the immune system. We found that Ag-specific production of IFN- γ by CD8⁺ T cells is temperature sensitive with the highest production of IFN- γ observed with incubation at 39.5°C (Fig. 1A). Cells pre-incubated at 37°C had higher IFN- γ production than those incubated at 33°C. Re-stimulation of the CD8⁺ T cells with cognate peptide is still required as temperature alone does not result in production of IFN- γ . The temperature sensitive effect on IFN- γ production by CD8⁺ T cells is enhanced with a greater duration of heating (Fig. 1B). Pre-incubation at 39.5°C for 6 hours resulted in the highest level of IFN- γ production by effector CD8⁺ T cells. Consistent

with previously published results (26, 27), maximum effect on other immune cell subsets occurs by heating at 39.5°C for 6 hours. We next determined the range of Ag concentrations at which thermal enhancement of IFN- γ production could be observed. We incubated effector CD8⁺ T cells at 33, 37, and 39.5°C for 6 hours and then stimulated them with splenocytes pulsed with varying concentrations of gp100₂₅₋₃₃ peptide. The effect of higher temperatures on enhancing IFN- γ production is maximal at a peptide concentration of 0.1 to 0.01 μ g/ml. This enhancement is lost at higher peptide concentrations where IFN- γ production by cells incubated at 33 or 37°C is comparable to production by cells incubated at 39.5°C (Fig. 1C). Thus, if Ag stimulation is optimal, exposure to elevated temperature does not further enhance IFN- γ . When CD8⁺ T cells are stimulated at sub-optimal Ag concentrations, higher temperatures can enhance CD8⁺ T cell IFN- γ production.

Ag-specific effector CD8⁺ T cell cytotoxicity is temperature sensitive

Another important anti-tumor effector function of CD8⁺ T cells is cytotoxicity. A previous study has shown that Fas ligand specific killing is enhanced with hyperthermia (28), however little is known about how Ag specific T cell cytotoxicity is impacted by hyperthermia. To test Ag-specific CD8⁺ T cell cytotoxicity, effector Pmel-1 CD8⁺ T cells were generated *in vitro* by activating with C57BL/6 splenocytes pulsed with peptide for 5 days, and then effector cells were incubated at 33, 37, and 39.5°C for 6 hours and then co-incubated with B16.F10 (gp100 expressing) or EL-4 (gp100 negative) ⁵¹Cr labeled tumor cells. Similar to the effect of temperature on IFN- γ production, effector Pmel-1 CD8⁺ T cells incubated at higher temperatures displayed an enhanced cytotoxicity against B16.F10 cells (Fig. 2). Cytotoxicity was antigen specific as no lysis was observed when effector CD8⁺ T cells were co-incubated with chromium labeled EL-4 cells (Fig. 2). These results demonstrate that antigen specific CD8⁺ T cell effector responses are highly dependent on temperature. These observations suggest an important role for mildly elevated temperature (i.e. during a fever or hyperthermia) in enhancing a cellular mediated T cell response (IFN- γ production and cytotoxicity) to antigen.

Temperature dependent IFN- γ production does not depend on synthesis of new protein but does induce GM-1 clustering

To investigate the mechanisms by which increased temperature could affect IFN- γ production in effector CD8⁺ T cells, we first asked whether additional protein synthesis occurs during heating which could be mediating the thermal regulation of IFN- γ production when the CD8⁺ T cells are activated with Ag-pulsed splenocytes. CD8⁺ T cells were incubated with 10 μ M cycloheximide (to block protein synthesis) during incubation at 33, 37, and 39.5°C and then washed three times with pre-warmed media and subsequently stimulated with Ag-pulsed splenocytes from C57BL/6 mice. After incubation for 18 hours at 37°C supernatants were collected and analyzed by ELISA for IFN- γ . Overall, there is a slight decrease in IFN- γ production compared to DMSO controls as some cycloheximide may have remained in culture after subsequent washes. However, cells incubated at 39.5°C, even when co-cultured with cycloheximide during the heating process, still resulted in more IFN- γ secretion compared to cells incubated at 33 and 37°C (Fig. 3A). Thus, additional protein synthesis is not required for the temperature dependent effect on Ag-specific IFN- γ production by CD8⁺ T cells.

Recent work has shown that incubating naïve CD8⁺ T cells at 39.5°C for 6 hours induces the clustering of GM1⁺ cholesterol dependent (CD)-microdomains in the plasma membrane (Mace et al 2011, *In Press Journal of Leukocyte Biology*). We asked whether effector CD8⁺ T cells were similarly impacted thus changing the signaling required for IFN- γ production. Effector CD8⁺ T cells were incubated at 33, 37, and 39.5°C for 6 hours and GM1 was stained using CTxB to look at possible alterations in GM1 organization in the plasma

membrane. GM1 proteins appeared either diffuse or clustered depending on thermal treatment (Fig. 3B). Staining patterns were quantified by fluorescent microscopy and we observed a higher percentage of effector CD8⁺ T cells with GM1 clustering at 39.5°C compared to cells incubated at 33 or 37°C (Fig. 3C). This data suggests that temperature induced changes at the plasma membrane may be associated with the enhanced Ag-specific CD8⁺ effector function.

Increased temperature does not affect non-antigen specific CD8⁺ signaling that is independent of the TCR

Since elevated temperatures impact CD8⁺ T cell functions through altering the distribution of CD-microdomains, we hypothesized that CD8⁺ T cell activation through a stimulation signal bypassing the cellular membrane would not be influenced by temperature changes. Treatment with mitogen and ionophore (PMA and Ionomycin) bypasses the need for TCR engagement on the membrane. We did not observe differences in IFN- γ production between any of the groups when the effector CD8⁺ T cells were incubated at 33, 37, 39.5°C for 6 hours and then stimulated with PMA and ionomycin (Fig. 4). This indicates that the temperature dependent effect on CD8⁺ T cell function is acting through events and signals involving the plasma membrane.

Elevated physiological temperatures enhance CD8⁺ T cell TCR signaling

We next investigated how temperature could affect early signaling events following ligation of TCR during an Ag-specific CD8⁺ T cell response. Phosphorylation of PKC θ and LAT plays an important role in mediating TCR signaling (29–31) and previous work from our lab has shown that following hyperthermia, PKC reorganization and activity is enhanced in T cells (32). Therefore we asked how the phosphorylation of PKC θ and LAT might differ in antigen-stimulated Pmel-1 CD8⁺ T cells following pre-treatment with different temperatures. The effector Pmel-1 CD8⁺ T cells were incubated at 33, 37, and 39.5°C for 6 hours and then activated for 0–30 minutes at 37°C (Fig. 5–D). Lysates were analyzed by Western blot for PKC θ and LAT and densitometry was used to quantify the relative phosphorylated protein levels compared to control β -actin expression. The phosphorylation of LAT and PKC θ were enhanced after 5 minutes of Ag-stimulation when pre-incubated at 39.5°C compared to a delayed phosphorylation when cells were pre-incubated at 33 or 37°C prior to Ag-stimulation (Fig. 5B and 5D). These results suggest that TCR signaling is enhanced in a temperature dependent fashion and that higher temperatures increase the rate of activation. At later time points, the phosphorylation of LAT and PKC in the 37 and 39.5°C groups begins to decrease but remains higher in the 33°C pre-incubated cells. This suggests that signaling may not be occurring efficiently at lower temperatures (33°C).

Ag-specific effector CD8⁺ T cell IFN- γ gene transcription is regulated by temperature

To test whether these signaling effects resulted in enhanced IFN- γ gene transcription, CD8⁺ T cells were similarly incubated at 33, 37, and 39.5°C for 6 hours and re-stimulated with Ag-pulsed C57BL/6 splenocytes for 0 to 2 hours after which RNA was isolated. Ag-specific IFN- γ mRNA expression is regulated by temperature with higher fold message levels in cells incubated at 39.5°C (Fig. 6). Cells pre-incubated at 33°C showed a slower rate of IFN- γ transcription than cells incubated at 37 and 39.5°C temperatures. This demonstrates that mild hyperthermia can result in changes genes downstream of TCR signaling, thus resulting in the enhanced IFN- γ production observed at higher temperatures (Fig. 1).

Discussion

New strategies to enhance the activity of CD8⁺ T cells are critical for achieving long term protection from cancer recurrence. T cell infiltration of tumors has been documented to be

very important in overall survival, but substantial evidence exists that a suppressive microenvironment within the tumor limits the activation and function of CD8⁺ T cells. Indeed, many different factors and cell types within the tumor exert suppressive forces against a CD8⁺ T cell anti-tumor response. For example, factors such as IL-6, VEGF, MMPs, GM-CSF, IL-10 and many other cytokines and chemokines can be found within the tumor microenvironment and are responsible for the suppression of an anti-tumor immune response (33–35).

These factors and others promote the accumulation of cells including myeloid-derived suppressor cells (MDSCs), which suppress CD8⁺ T cell activity by secreting ROS, depleting arginine, and decreasing TCR associated signaling proteins (36–38). MDSCs also secrete anti-inflammatory cytokines including IL-10 and TGF- β which promote differentiation of T regulatory (T_{reg}) cells that also inhibit CD8⁺ T cell anti-tumor activity. A recent study suggested that infiltrating DCs can suppress CD8⁺ T cell function by inhibiting T cell L-Arginine metabolism in a spontaneous mammary tumor model leading to decreased T cell expansion and IFN- γ production ultimately resulting in anergy (39).

Since multiple factors and cell types within the tumor microenvironment can suppress T cell activity new therapies or adjuvants that could help to increase the activation and function of these cells could overcome this suppression would be important in generating an improved anti-tumor response. In this report, we document the ability of mild temperature elevation to significantly enhance activities (IFN- γ production and cytotoxicity) that would be important in tumor control, and we have associated these effects with changes in the organization of the plasma membrane and downstream signaling pathways, alterations which may be involved in the underlying mechanisms by which elevated temperature affects cellular function. Earlier studies in our laboratory have reported that heating mice *in vivo* enhances PKC activity and results in the aggregation of PKC and the cytoskeletal proteins spectrin and vimentin (27, 40). We have shown in this study that elevated temperature results in enhanced downstream signaling, with greater levels of phosphorylated PKC θ and LAT. Subsequent enhancement in IFN- γ gene transcription was observed when effector CD8⁺ T cells were incubated at higher temperatures and that these changes are also associated temporally with a thermally-induced reorganization of membrane domains.

It will now be important to determine if these thermally enhanced CD8⁺ T cell functions could play a role in controlling tumor growth *in vivo*. Previous studies have demonstrated general enhancement of immune activity against tumors following mild hyperthermia, but these studies have not studied a tumor antigen-dependent response. For example, in mice, fever-range hyperthermia has been shown to enhance anti-tumor immunity though both the innate and adaptive immune systems (41). Hyperthermia has also been shown to significantly enhance efficiency of heat shock protein vaccines (42). When combined with intratumoral DC injection, hyperthermia induces DC migration to the tumor draining lymph nodes and enhances the priming of CTLs in animal melanoma models (43). Clinically, Guo et. al. has shown that patients with advanced melanoma treated three times a week with local hyperthermia followed by intratumoral injections of immature DCs experienced significantly longer time to tumor progression ($p < 0.05$) (44). Furthermore, this study demonstrated that DC vaccination in combination with hyperthermia resulted in an increased infiltration of activated CD8⁺ cells into the tumor site accompanied by decreased infiltration of immune suppressive T_{reg} cells, possibly creating an environment for improved tumor control (44). Recently long-term anti-tumor immunity was achieved in murine models; treatment with oxaliplatin chemotherapy followed twenty-four hours later by six hour whole body hyperthermia was able to cure all primary and metastatic tumors in 50% of MTLn3 tumor-bearing rats (13).

We have shown that mild hyperthermia increases the clustering of GM-1⁺ regions within the plasma membrane (Fig. 3). These GM-1 regions contain important signaling molecules, such as TCR, Lck, LAT and many other proteins important in recognizing antigen presented by an APC or target cell (45). Formation of the immunological synapse between an APC and target cells requires the aggregation and localization of these signaling enriched GM-1 membrane domains to occur at this synapse (46). The hyperthermia induced increase in GM-1 clustering could potentially pre-condition effector T cells and allow them to react faster to antigen presentation and initiate killing mechanisms than at lower temperatures.

In summary, this study presents novel information describing the role of hyperthermia in improving tumor antigen specific, effector CD8⁺ T cell function. Our studies were preformed in Pmel-1 transgenic mice which carry a rearranged TCR specific for gp100 (47). While most prior work has been on non-specific antigen dependent immune responses, the gp100 protein is expressed by most melanoma cells making this an important model for future *in vivo* work on the antigen-specific immune response. Furthermore, the fundamental observations made here may expedite the design and implementation of new clinical protocols utilizing hyperthermia as an adjuvant immunotherapy.

Acknowledgments

We thank members of the Department of Flow and Image Cytometry Core Facility as well as the Roswell Park Cancer Institute Animal Resource Facility. We also thank Jeanne Prendergast for her assistance in the lab and Dr. Bonnie Hylander for her help in reviewing this manuscript.

References

1. Vernon CC, Hand JW, Field SB, Machin D, Whaley JB, van der Zee J, van Putten WL, van Rhoon GC, van Dijk JD, Gonzalez Gonzalez D, et al. Radiotherapy with or without hyperthermia in the treatment of superficial localized breast cancer: results from five randomized controlled trials. International Collaborative Hyperthermia Group. *Int J Radiat Oncol Biol Phys.* 1996 Jul 1; 35(4): 731–44. [PubMed: 8690639]
2. Jones EL, Oleson JR, Prosnitz LR, Samulski TV, Vujaskovic Z, Yu D, Sanders LL, Dewhirst MW. Randomized trial of hyperthermia and radiation for superficial tumors. *J Clin Oncol.* 2005 May 1; 23(13):3079–85. [PubMed: 15860867]
3. Datta NR, Bose AK, Kapoor HK, Gupta S. Head and neck cancers: results of thermoradiotherapy versus radiotherapy. *Int J Hyperthermia.* 1990 May-Jun; 6(3):479–86. [PubMed: 2198311]
4. Valdagni R, Amichetti M. Report of long-term follow-up in a randomized trial comparing radiation therapy and radiation therapy plus hyperthermia to metastatic lymph nodes in stage IV head and neck patients. *Int J Radiat Oncol Biol Phys.* 1994 Jan 1; 28(1):163–9. [PubMed: 8270437]
5. Overgaard J, Gonzalez Gonzalez D, Hulshof MC, Arcangeli G, Dahl O, Mella O, Bentzen SM. Randomised trial of hyperthermia as adjuvant to radiotherapy for recurrent or metastatic malignant melanoma. European Society for Hyperthermic Oncology. *Lancet.* 1995 Mar 4; 345(8949):540–3. [PubMed: 7776772]
6. Van der Zee J, Van Rhoon GC, Broekmeyer-Reurink MP, Kuijs AE, Van Den Berg AP, Treurniet-Donker D, Van Putten WL, Reinhold HS. Clinical experience with local hyperthermia in Rotterdam. *Adv Exp Med Biol.* 1990; 267:383–6. [PubMed: 2088056]
7. Harima Y, Nagata K, Harima K, Ostapenko VV, Tanaka Y, Sawada S. A randomized clinical trial of radiation therapy versus thermoradiotherapy in stage IIIB cervical carcinoma. *Int J Hyperthermia.* 2001 Mar–Apr; 17(2):97–105. [PubMed: 11252361]
8. Sneed PK, Stauffer PR, McDermott MW, Diederich CJ, Lamborn KR, Prados MD, Chang S, Weaver KA, Spry L, Malec MK, et al. Survival benefit of hyperthermia in a prospective randomized trial of brachytherapy boost +/- hyperthermia for glioblastoma multiforme. *Int J Radiat Oncol Biol Phys.* 1998 Jan 15; 40(2):287–95. [PubMed: 9457811]
9. Falk MH, Issels RD. Hyperthermia in oncology. *Int J Hyperthermia.* 2001 Jan-Feb; 17(1):1–18. [PubMed: 11212876]

10. Issels RD, Lindner LH, Verweij J, Wust P, Reichardt P, Schem BC, Abdel-Rahman S, Daugaard S, Salat C, Wendtner CM, et al. Neo-adjuvant chemotherapy alone or with regional hyperthermia for localised high-risk soft-tissue sarcoma: a randomised phase 3 multicentre study. *Lancet Oncol.* 2010 Jun; 11(6):561–70. [PubMed: 20434400]
11. Baronzio G, Gramaglia A, Fiorentini G. Hyperthermia and immunity. A brief overview. *In Vivo.* 2006 Nov–Dec; 20(6A):689–95. [PubMed: 17203747]
12. Peer AJ, Grimm MJ, Zynda ER, Repasky EA. Diverse immune mechanisms may contribute to the survival benefit seen in cancer patients receiving hyperthermia. *Immunol Res.* 2010 Mar; 46(1–3): 137–54. [PubMed: 19756410]
13. Rowe RW, Strebel FR, Proett JM, Deng W, Chan D, He G, Siddik Z, Bull JM. Fever-range whole body thermotherapy combined with oxaliplatin: a curative regimen in a pre-clinical breast cancer model. *Int J Hyperthermia.* 2010; 26(6):565–76. [PubMed: 20707651]
14. Naito Y, Saito K, Shiiba K, Ohuchi A, Saigenji K, Nagura H, Ohtani H. CD8+ T cells infiltrated within cancer cell nests as a prognostic factor in human colorectal cancer. *Cancer Res.* 1998 Aug 15; 58(16):3491–4. [PubMed: 9721846]
15. Diederichsen AC, Hjelmborg JB, Christensen PB, Zeuthen J, Fenger C. Prognostic value of the CD4+/CD8+ ratio of tumour infiltrating lymphocytes in colorectal cancer and HLA-DR expression on tumour cells. *Cancer Immunol Immunother.* 2003 Jul; 52(7):423–8. [PubMed: 12695859]
16. Fukunaga A, Miyamoto M, Cho Y, Murakami S, Kwarada Y, Oshikiri T, Kato K, Kurokawa T, Suzuoki M, Nakakubo Y, et al. CD8+ tumor-infiltrating lymphocytes together with CD4+ tumor-infiltrating lymphocytes and dendritic cells improve the prognosis of patients with pancreatic adenocarcinoma. *Pancreas.* 2004 Jan; 28(1):e26–31. [PubMed: 14707745]
17. Sato E, Olson SH, Ahn J, Bundy B, Nishikawa H, Qian F, Jungbluth AA, Frosina D, Gnjjatic S, Ambrosone C, et al. Intraepithelial CD8+ tumor-infiltrating lymphocytes and a high CD8+/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci U S A.* 2005 Dec 20; 102(51):18538–43. [PubMed: 16344461]
18. Piersma SJ, Jordanova ES, van Poelgeest MI, Kwappenberg KM, van der Hulst JM, Drijfhout JW, Melief CJ, Kenter GG, Fleuren GJ, Offringa R, et al. High number of intraepithelial CD8+ tumor-infiltrating lymphocytes is associated with the absence of lymph node metastases in patients with large early-stage cervical cancer. *Cancer Res.* 2007 Jan 1; 67(1):354–61. [PubMed: 17210718]
19. Yamada N, Oizumi S, Kikuchi E, Shinagawa N, Konishi-Sakakibara J, Ishimine A, Aoe K, Gemba K, Kishimoto T, Torigoe T, et al. CD8+ tumor-infiltrating lymphocytes predict favorable prognosis in malignant pleural mesothelioma after resection. *Cancer Immunol Immunother.* 2010 Oct; 59(10):1543–9. [PubMed: 20567822]
20. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A, Mlecnik B, Lagorce-Pages C, Tosolini M, Camus M, Berger A, Wind P, et al. Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science.* 2006 Sep 29; 313(5795):1960–4. [PubMed: 17008531]
21. Ostberg JR, Gellin C, Patel R, Repasky EA. Regulatory potential of fever-range whole body hyperthermia on Langerhans cells and lymphocytes in an antigen-dependent cellular immune response. *J Immunol.* 2001 Sep 1; 167(5):2666–70. [PubMed: 11509609]
22. Szabo SJ, Sullivan BM, Peng SL, Glimcher LH. Molecular mechanisms regulating Th1 immune responses. *Annu Rev Immunol.* 2003; 21:713–58. [PubMed: 12500979]
23. Atanackovic D, Nierhaus A, Neumeier M, Hossfeld DK, Hegewisch-Becker S. 41.8 degrees C whole body hyperthermia as an adjunct to chemotherapy induces prolonged T cell activation in patients with various malignant diseases. *Cancer Immunol Immunother.* 2002 Dec; 51(11–12): 603–13. [PubMed: 12439605]
24. Hanson DF. Fever and the immune response. The effects of physiological temperatures on primary murine splenic T-cell responses in vitro. *J Immunol.* 1993 Jul 1; 151(1):436–48. [PubMed: 8326136]
25. Smith JB, Knowlton RP, Agarwal SS. Human lymphocyte responses are enhanced by culture at 40 degrees C. *J Immunol.* 1978 Aug; 121(2):691–4. [PubMed: 150449]

26. Ostberg JR, Dayanc BE, Yuan M, Oflazoglu E, Repasky EA. Enhancement of natural killer (NK) cell cytotoxicity by fever-range thermal stress is dependent on NKG2D function and is associated with plasma membrane NKG2D clustering and increased expression of MICA on target cells. *J Leukoc Biol.* 2007 Nov; 82(5):1322–31. [PubMed: 17711975]
27. Di YP, Repasky EA, Subjeck JR. Distribution of HSP70, protein kinase C, and spectrin is altered in lymphocytes during a fever-like hyperthermia exposure. *J Cell Physiol.* 1997 Jul; 172(1):44–54. [PubMed: 9207924]
28. Cippitelli M, Fionda C, Di Bona D, Piccoli M, Frati L, Santoni A. Hyperthermia enhances CD95-ligand gene expression in T lymphocytes. *J Immunol.* 2005 Jan 1; 174(1):223–32. [PubMed: 15611244]
29. Shim EK, Jung SH, Lee JR. Role of two adaptor molecules SLP-76 and LAT in the PI3K signaling pathway in activated T cells. *J Immunol.* 2011 Mar 1; 186(5):2926–35. [PubMed: 21282515]
30. Huang SC, Tsai HF, Tzeng HT, Liao HJ, Hsu PN. Lipid raft assembly and Lck recruitment in TRAIL costimulation mediates NF-kappaB activation and T cell proliferation. *J Immunol.* 2011 Jan 15; 186(2):931–9. [PubMed: 21160038]
31. Ma JS, Haydar TF, Radoja S. Protein kinase C delta localizes to secretory lysosomes in CD8+ CTL and directly mediates TCR signals leading to granule exocytosis-mediated cytotoxicity. *J Immunol.* 2008 Oct 1; 181(7):4716–22. [PubMed: 18802074]
32. Wang XY, Ostberg JR, Repasky EA. Effect of fever-like whole-body hyperthermia on lymphocyte spectrin distribution, protein kinase C activity, and uropod formation. *J Immunol.* 1999 Mar 15; 162(6):3378–87. [PubMed: 10092792]
33. Bonecchi R, Locati M, Mantovani A. Chemokines and cancer: a fatal attraction. *Cancer Cell.* 2011 Apr 12; 19(4):434–5. [PubMed: 21481784]
34. Keeley EC, Mehrad B, Strieter RM. CXC chemokines in cancer angiogenesis and metastases. *Adv Cancer Res.* 2010; 106:91–111. [PubMed: 20399957]
35. Grivennikov SI, Karin M. Inflammatory cytokines in cancer: tumour necrosis factor and interleukin 6 take the stage. *Ann Rheum Dis.* 2011 Mar; 70(Suppl 1):i104–8. [PubMed: 21339211]
36. Bronte V, Serafini P, Mazzoni A, Segal DM, Zanovello P. L-arginine metabolism in myeloid cells controls T-lymphocyte functions. *Trends Immunol.* 2003 Jun; 24(6):302–6. [PubMed: 12810105]
37. Kusmartsev S, Nefedova Y, Yoder D, Gabrilovich DI. Antigen-specific inhibition of CD8+ T cell response by immature myeloid cells in cancer is mediated by reactive oxygen species. *J Immunol.* 2004 Jan 15; 172(2):989–99. [PubMed: 14707072]
38. Ezernitchi AV, Vaknin I, Cohen-Daniel L, Levy O, Manaster E, Halabi A, Pikarsky E, Shapira L, Baniyash M. TCR zeta down-regulation under chronic inflammation is mediated by myeloid suppressor cells differentially distributed between various lymphatic organs. *J Immunol.* 2006 Oct 1; 177(7):4763–72. [PubMed: 16982917]
39. Norian LA, Rodriguez PC, O'Mara LA, Zabaleta J, Ochoa AC, Cella M, Allen PM. Tumor-infiltrating regulatory dendritic cells inhibit CD8+ T cell function via L-arginine metabolism. *Cancer Res.* 2009 Apr 1; 69(7):3086–94. [PubMed: 19293186]
40. Di YP, Repasky E, Laszlo A, Calderwood S, Subjeck J. Hsp70 translocates into a cytoplasmic aggregate during lymphocyte activation. *J Cell Physiol.* 1995 Nov; 165(2):228–38. [PubMed: 7593200]
41. Burd R, Dziedzic TS, Xu Y, Caligiuri MA, Subjeck JR, Repasky EA. Tumor cell apoptosis, lymphocyte recruitment and tumor vascular changes are induced by low temperature, long duration (fever-like) whole body hyperthermia. *J Cell Physiol.* 1998 Oct; 177(1):137–47. [PubMed: 9731754]
42. Wang XY, Kazim L, Repasky EA, Subjeck JR. Characterization of heat shock protein 110 and glucose-regulated protein 170 as cancer vaccines and the effect of fever-range hyperthermia on vaccine activity. *J Immunol.* 2001 Jan 1; 166(1):490–7. [PubMed: 11123328]
43. Tanaka K, Ito A, Kobayashi T, Kawamura T, Shimada S, Matsumoto K, Saida T, Honda H. Intratumoral injection of immature dendritic cells enhances antitumor effect of hyperthermia using magnetic nanoparticles. *Int J Cancer.* 2005 Sep 10; 116(4):624–33. [PubMed: 15825167]

44. Guo J, Zhu J, Sheng X, Wang X, Qu L, Han Y, Liu Y, Zhang H, Huo L, Zhang S, et al. Intratumoral injection of dendritic cells in combination with local hyperthermia induces systemic antitumor effect in patients with advanced melanoma. *Int J Cancer*. 2007 Jun 1; 120(11):2418–25. [PubMed: 17294445]
45. Patel VP, Moran M, Low TA, Miceli MC. A molecular framework for two-step T cell signaling: Lck Src homology 3 mutations discriminate distinctly regulated lipid raft reorganization events. *J Immunol*. 2001 Jan 15; 166(2):754–64. [PubMed: 11145647]
46. Huse M, Quann EJ, Davis MM. Shouts, whispers and the kiss of death: directional secretion in T cells. *Nat Immunol*. 2008 Oct; 9(10):1105–11. [PubMed: 18800163]
47. Overwijk WW, Theoret MR, Finkelstein SE, Surman DR, de Jong LA, Vyth-Dreese FA, DelleMijn TA, Antony PA, Spiess PJ, Palmer DC, et al. Tumor regression and autoimmunity after reversal of a functionally tolerant state of self-reactive CD8+ T cells. *J Exp Med*. 2003 Aug 18; 198(4):569–80. [PubMed: 12925674]

Figure 1.

Ag-specific effector CD8⁺ T cell IFN- γ production is temperature sensitive. Effector Pmel-1 CD8⁺ T cells were generated by pulsing Pmel-1 splenocytes with 0.1 μ g/ml gp100₂₅₋₃₃ peptide (at 37°C) and 30 IU of rIL-2 for 5 days. **A**, Effector Pmel-1 CD8⁺ T cells were incubated at 33, 37, and 39.5°C for 6 hours and re-stimulated with C57BL/6 splenocytes pulsed with 0.1 μ g/ml gp100₂₅₋₃₃ peptide. Supernatants were removed after 18 hours of co-incubation at 37°C and analyzed for IFN- γ by ELISA. **B**, Effector Pmel-1 CD8⁺ T cells were incubated at 33, 37, or 39.5°C for 2 to 6 hours and activated with pulsed C57BL/6 splenocytes. Supernatants were analyzed for IFN- γ by ELISA 18 hours after pulsing. **C**, Effector Pmel-1 CD8⁺ T cells were incubated at 33, 37, or 39.5°C for 6 hours and stimulated with varying concentrations of gp100₂₅₋₃₃ peptide pulsed C57BL/6 splenocytes. Results are reported as the mean $\pm$ SD. These results are representative of two independent experiments ($p < 0.05$, # = vs. 37 and * = vs. 33°C).

Figure 2.

Mild hyperthermia enhances Ag-specific CD8⁺ cytotoxicity. Effector Pmel-1 CD8⁺ T cells were incubated at 33, 37, or 39.5°C for 6 hours and then co-incubated with Cr⁵¹ labeled EL-4 (gp100 negative) or B16.F10 (gp100 positive) target cells for 4 hours at 37°C. Supernatant was collected and percent lysis was determined by chromium-release by lysed target cells. Results are reported as the mean $\pm$ SD. These results are representative of two independent experiments.

Figure 3.

Mild hyperthermia does not affect new protein synthesis but induces GM1 clustering in the plasma membrane. A, Effector Pmel-1 CD8⁺ T cells were incubated at 33°, 37°, or 39.5°C for 6 hours in the presence of 10 μ m cycloheximide then stimulated with gp100₂₅₋₃₃ peptide pulsed C57BL/6 splenocytes for 18 hours. Supernatants were analyzed for IFN- γ production by ELISA. B, Effector cells were incubated at 33, 37, 39.5°C for 6 hours and adhered onto alcian blue coated coverslips, fixed with paraformaldehyde. Cells were stained with FITC-CTxB, visualized for clustering by fluorescent microscopy, and C, quantified. Results are reported as the mean $\pm$ SD. These results are representative of two independent experiments ($p < 0.05$, # = vs. 37 and * = vs. 33°C).

Figure 4.

Mild hyperthermia does not affect effector CD8⁺ T cell activation when TCR signaling is bypassed. Effector Pmel-1 CD8⁺ T cells were incubated at 33, 37, 39.5°C for 6 hours and activated with 0.5μg/ml of ionomycin and varying concentrations of PMA for 18 hours at 37°C. Supernatants were collected and analyzed for IFN-γ by ELISA. Results are reported as the mean ± SD. These results are representative of two independent experiments.

Figure 5.

Effector CD8⁺ T cell Ag-specific signaling is enhanced by mild hyperthermia Cells were incubated at 33, 37, or 39.5°C for 6 hours and stimulated with 0.1 µg/ml gp100₂₅₋₃₃ peptide pulsed C57BL/6 splenocytes at 37°C. Stimulation was varied between 0–30 minutes. A–D, Expression of phosphorylated LAT and PKCθ were determined by Western blot. (A, C) Western blots. (B, D) Densitometry was performed using total β-actin levels as background controls. These results are representative of two independent experiments.

Figure 6.

Effector CD8⁺ T cell IFN- γ transcription is regulated by temperature. Effector CD8⁺ T cells were incubated at 33, 37, 39.5°C for 6 hours and stimulated with 0.1 μ g/ml gp100₂₅₋₃₃ peptide pulsed C57BL/6 splenocytes for 0–2 hours at 37°C. RNA was isolated and cDNA was synthesized. mRNA levels were assessed by Real-time PCR. All message levels are relative to GAPDH controls and experimental gene expression is relative to cells activated with null peptide. Results are reported as the mean $\pm$ SD. These results are representative of two independent experiments ($p < 0.05$, # = vs. 37 and * = vs. 33°C).