Published in final edited form as:

J Zool (1987). 2012 February 1; 286(2): 102–109. doi:10.1111/j.1469-7998.2011.00856.x.

Estimating carnivoran diets using a combination of carcass observations and scats from GPS clusters

C.J. Tambling*,

Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa

Centre for African Conservation Ecology, Department of Zoology, Nelson Mandela Metropolitan University, Port Elizabeth, South Africa

S.D. Laurence,

Enviro-Insight, Murrayfield, Pretoria, South Africa

S.E. Bellan,

Department of Environmental Science, Policy and Management, University of California, Berkeley, California, USA

E.Z. Cameron,

Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa

School of Zoology, University of Tasmania, Hobart, Tasmania, Australia

J.T. du Toit, and

Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa

Department of Wildland Resources, Utah State University, Logan, Utah, USA

W.M. Getz

Mammal Research Institute, Department of Zoology and Entomology, University of Pretoria, Pretoria, South Africa

Department of Environmental Science, Policy and Management, University of California, Berkeley, California, USA

Abstract

Scat analysis is one of the most frequently used methods to assess carnivoran diets and Global Positioning System (GPS) cluster methods are increasingly being used to locate feeding sites for large carnivorans. However, both methods have inherent biases that limit their use. GPS methods to locate kill sites are biased towards large carcasses, while scat analysis over-estimates the biomass consumed from smaller prey. We combined carcass observations and scats collected along known movement routes, assessed using GPS data from four African lion (*Panthera leo*) prides in the Kruger National Park, South Africa, to determine how a combination of these two datasets change diet estimates. As expected, using carcasses alone under-estimated the number of feeding events on small species, primarily impala (*Aepyceros melampus*) and warthog (*Phacochoerus africanus*), in our case by more than 50% and thus significantly under-estimated the biomass consumed per pride per day in comparison to when the diet was assessed using

*Correspondent: Craig J. Tambling, Centre for African Conservation Ecology, Department of Zoology, Nelson Mandela, Metropolitan University, Port Elizabeth, South Africa. cjtambling@gmail.com.

carcass observations alone. We show that an approach that supplements carcass observations with scats that enables the identification of potentially missed feeding events increases the estimates of food intake rates for large carnivorans, with possible ramifications for predator-prey interaction studies dealing with biomass intake rate.

Keywords

African savanna; Kruger National Park; lion; *Panthera leo*; predation; prey consumption

Quantifying carnivoran diets is an essential step in investigating carnivoran ecology (Mills, 1992), and provides the basis for understanding population-level impacts that carnivorans may have on prey populations (Owen-Smith & Mason, 2005; Owen-Smith, 2008). Numerous techniques are available for the assessment of carnivoran diets (Mills, 1992), including highly invasive stomach content analysis (Smuts, 1979), moderately invasive continuous direct observations (Mills & Shenk, 1992) and non-invasive faecal analysis (Andheria, Karanth & Kumar, 2007) and carcass observation (Lehmann *et al.*, 2008). Each of the approaches is limited either due to inherent biases, or the feasibility and practicality of the technique.

Recent advances in Global Positioning System (GPS) technology permit the collection of animal location data at a scale that is sufficiently fine to provide a good approximation to the continuous movement path of individuals (Getz & Saltz, 2008). Despite this, a trade-off between the frequency of points collected and the total length of the movement path still exists owing to the limited battery life of most collars (Sand *et al.*, 2005). Obtaining GPS fixes at an hourly interval for female lions in the Kruger National Park allowed the development of models that increased the probability of locating carcasses at GPS clusters in time (Tambling *et al.*, 2010). However, small prey items might be entirely consumed by the predator (see Power, 2002), or scavengers might subsequently eliminate signs of some feeding sites. Similarly, the GPS approach has proved useful in dietary studies of mountain lions *Puma concolor* (Anderson & Lindzey, 2003), wolves *Canis lupus* (Sand *et al.*, 2005; Franke *et al.*, 2006) and leopards *Panthera pardus* (Martins *et al.*, 2011), but the bias towards large prey has remained largely intractable (but see Martins *et al.*, 2011).

The analysis of large numbers of carnivoran scats offers a non-invasive estimate of carnivoran diets. Despite widespread use, this method over-estimates the biomass, and under-estimates the numbers, of small species consumed (Karanth & Sunquist, 1995; Marker *et al.*, 2003), although correction factors are available for some species to correct for this bias (Ackerman, Lindzey & Hemker, 1984). In addition, a problem of autocorrelation occurs (Marucco, Pletscher & Boitani, 2008) when clusters of scats are produced by social carnivorans consuming prey items that are large in relation to their body size (e.g. wild dogs *Lycaon pictus*, lions *Panthera leo*, wolves). Despite the problems associated with both GPS cluster techniques and scat analysis, a combination of the two approaches can reduce possible pseudo-replication of correlated scats and identify small kills that are missed from direct observation (Marucco *et al.*, 2008; Martins *et al.*, 2011).

Although the most accurate method of quantifying lion diet composition is continuous observation of study animals, thereby allowing all predation activity to be directly observed (Mills & Shenk, 1992; Funston *et al.*, 1998), this approach is not always logistically feasible (Hayward *et al.*, 2009). Alternative methods, including scat and GPS approaches, have not been comprehensively tested; hence the bias inherent in results obtained through these methods is uncertain. Owing to their social nature, lions generally kill and consume prey items the same size as or larger than themselves (Radloff & du Toit, 2004; Hayward &

Kerley, 2005), resulting in the possibility of correlated clusters of scats from a single large carcass. In this study we investigated how the incorporation of scats into a GPS cluster approach might alter lion diet estimates, where diet estimates refer to both the composition and the total biomass of food consumed by lions.

MATERIALS AND METHODS

We conducted the study between April 2005 and May 2007, in the Kruger National Park, in a 1000km² area around the Satara rest camp (Figure 1, 31.77° E, 24.39° S) in the central region of the park. Rainfall in the area is highly seasonal, with the majority falling between October and March, when there is also an increase in average temperature (Venter, Scholes & Eckhardt, 2003). As a consequence faecal collection becomes more difficult during the rainy season as dung beetle (Coleoptera; Scarabaeinae) activity and rain reduce the number of collectable samples. The study area is an open tree savanna with a moderate to sparse shrub layer and dense grass layer, with marula (*Sclerocarya birrea*) and knob-thorn (*Senegalia nigrescens*) the dominant tree species, and red grass (*Themeda triandra*) and stinking grass (*Bothriocloa radicans*) the dominant grass species (Venter *et al.*, 2003). Our study area includes the northern component of the wildebeest (*Connochaetes taurinus*) and zebra (*Equus quagga*) migrations, and high densities of these species occur in the wet months (Gertenbach, 1983). Resident buffalo (*Syncerus caffer*), kudu (*Tragelaphus strepsiceros*), giraffe (*Giraffa camelopardalis*) and waterbuck (*Kobus ellipsiprymnus*) occur in large numbers, all combining as a prey base for a high lion population density (Ferreira & Funston, 2010). Across the park, impala (*Aepyceros melampus*) are the most abundant prey species and provide the bulk of food for lions and the other large predators (Mills & Biggs, 1993; Radloff & du Toit, 2004).

Five female lions from four prides were collared with GPS/GSM units (i.e. GPS with mobile phone capabilities to access data; African Wildlife Tracking) between April 2005 and May 2007. Lions were captured and collared by South African National Parks (SANParks) veterinarians using standard SANParks protocols (Smuts, Whyte & Dearlove, 1977). The fix schedule of the collars (one fix per hour at night [18:00–06:00] when the lions are most active (Hayward & Slotow, 2009), and three fixes during the day [06:00–18:00] when lions tend to rest) provides a set of locations used to construct a movement path over time for each collared lion. Along this movement path we identified spatio-temporal clusters of GPS locations (hereafter referred to as clusters), where a cluster is defined as consecutive (i.e. 2 or more) GPS co-ordinates within 100m of the previous GPS co-ordinate (see Tambling *et al.*, 2010). Investigated clusters were searched for carcasses and scats in a 50 m diameter around the GPS point, identified as the start of the cluster (accounting for GPS error - Tambling, 2010). If further GPS points identified as part of the cluster did not fall within the initial 50 m search diameter, new 50 m search diameters were created around those GPS positions. The duration of the search time was dependent on the vegetation cover of the area. Carcasses found at clusters were classified to species, age and sex based on the presence of identifiable material such as horns, jaws, bones and hair. We defined the number of feeding events for each species for each pride based on the observed carcasses found at clusters as $n_{\text{car.sp}}$. For all species specific analysis we removed carcasses that could not be identified to species level ($n = 3$).

Clusters (with and without observed carcasses) were searched for the presence of lion scats. Following collection, scats were washed under running water through a metal sieve to isolate all undigested distinguishable prey items (predominantly hair remains, but on some occasions hoof and horn remains as well). Hair was examined macroscopically (length, colour and texture) and microscopically (cross-section characteristics and cuticle-scale patterns) to identify prey species. Cross sections of hairs were made using the method

outlined in Douglas (1989) and cuticle scale imprints are obtained by investigating inverse imprints under a light microscope (van Kesteren, 2006). Both cross-section and cuticle-scale imprints were compared against reference collections to identify the species consumed (Keogh, 1983; Buys & Keogh, 1984). If multiple scats were found at GPS clusters, and if the same prey species occurred in more than one scat, we used the collection of scats containing remains of that prey species as a single sample unit to avoid over-representation of prey items from multiple scats (Marucco *et al.*, 2008).

Comparing frequency distributions of carcasses of prey species across the four prides revealed that two of the six pair-wise comparisons were significantly different. Therefore, we conducted analyses at the pride level and averaged results (mean $\pm$ SE) across prides ($n = 4$) to estimate the diet at the population level. Carcasses and scats found at each cluster occurred at known dates and times along the movement path of each pride, and these data were used to create a history of feeding events related to either a) a carcass alone, b) a scat alone or c) a combination of a carcass and a scat from the same feeding event. Since no estimate for transit times through the guts of lions exists (Breuer, 2005), based on cheetah (*Acinonyx jubatus*) transit times (48 – 111 hours: Marker *et al.*, 2003) we assumed a minimum of two days and a maximum of five days. The minimum and maximum gut transit times correspond to a high and low level of bias in terms of missed feeding events. The low bias assumes a long (120 hr) gut transit time, with all scats produced within five days of an observed carcass being the result of that carcass. The high bias assumes a 48 hr gut transit time with all scats produced within two days of an observed carcass being the result of that observed carcass. Therefore, if species A was consumed on day one, and three days later species A was found in the scats for the same pride of lions, we assumed that the scat originated from the observed carcass of species A for the low bias estimate, but that we had missed a feeding event of species A for the high bias estimate. For each pride we defined the number of missed feeding events for each species based on the inclusion of scats as $n_{\text{scat.sp}}$, and this value is calculated for both the low and high biases for missing feeding events.

To investigate the temporal relationship between observed carcasses and scats found for the same lion group following a feeding event, we tracked the presence and identity of scats found at subsequent clusters for each lion group for ten days after the observed feeding event. We then plotted the daily proportion of scats, containing the species identified at the feeding site, against the proportion of scats containing a different species over the ten-day period. Scats at a carcass site are a function of previous carcasses consumed and/or the observed carcass. Scats found at clusters not associated with a carcass are a function of the most recently observed carcass (within the range of days delineated by gut transit times) and/or possible missed feeding events that occurred once the lions had moved away from the last observed carcass. However, during the observed movement path for each lion pride, there exists a possibility that the same prey species contributes two consecutive feeding events, with either the first or second carcass not detected during cluster investigations. When this happens the scats found at a cluster could belong to a more recent feeding event of the same species that was missed, and thus our estimates for $n_{\text{scat.sp}}$ are an under-estimate. To quantify the possible magnitude of this under-estimation, we calculated how often the same species was found consecutively at clusters in a two-day window (average kill rate of female lions in Kruger National Park = 1.8 days, Funston *et al.*, 1998) as a proportion of all feeding events within a two-day window. We multiplied the proportion of same-species consecutive feeding events by $n_{\text{scat.sp}}$ for each prey species for each pride to estimate the number of missed feeding events per prey species per pride resulting from undetected consecutive feeding events, defined as $n_{\text{con.sp}}$. The final corrected number of individuals of each prey species consumed by each pride is then calculated as $n_{\text{sp}} = n_{\text{car.sp}} + n_{\text{scat.sp}} + n_{\text{con.sp}}$.

For each pride, as well as for all averaged prides, we used *G*-tests (Zar, 1999) to test if adding missed feeding events through scat-supplementation (using both the high- and low-bias levels of missed feeding events) changed the diet composition of lions as opposed to when the diet is estimated from observed carcasses alone. Using the log unit weight (taking into account the range of age and sex categories in the prey population) of each prey species (Coe, Cumming & Phillipson, 1976), we investigated if missed feeding events (using both high- and low-bias estimates) were related to the size of the prey species. We calculated the total biomass of each species consumed by each pride as the product of the unit weight of the species (Coe *et al.*, 1976), the proportion of edible meat for that species (see Funston *et al.*, 1998 Appendix 1) and the number of individuals of that species consumed by each pride. We used *G*-tests to assess whether the percent diet composition based on the biomass of each prey species consumed changes after implementing the scat-supplemented approach. Finally, using these biomass estimates, we investigated the difference in consumed biomass per pride per day between the three approaches (carcass observations, high bias of missing feeding events and low bias of missing feeding events) using ANOVA with a Tukey *post hoc* test to identify significant differences between approaches.

RESULTS

Between April 2005 and May 2007 we investigated 59.5% (1447 out of 2433) of all clusters identified, finding 236 carcasses (103, 70, 41 and 22 carcasses and scats in brackets per pride respectively) and 208 scats (91, 48, 50 and 19 scats per pride respectively). Clusters were investigated a median of 24 (range 0–331) days after lions occupied the clusters. Scats were found at clusters that were investigated a median of 16 (range 0–195) days after lions occupied the clusters. Scats were found both at carcass sites ($n = 47$) and at resting sites not associated with a carcass ($n = 161$). In 50% ($\pm 9\%$) of the scats found at carcass sites, the detected species was the same as the consumed species at that cluster (Figure 2). For all consecutive feeding events within the two-day window ($n = 60$), the same species was found consecutively 16% ($\pm 7\%$) of the time. Therefore, two thirds ($50\% + 16\% = 66\%$) of all scats found at carcass sites probably originated from a carcass fed on prior to the feeding event, where the scat was collected. Once lions moved away from a known carcass, 52%–66% of scats found at subsequent clusters in the following two days contained remains of the same species as that carcass and this level declined to ~20% in days 4–10 (Figure 2).

Numerically, the dominant prey species identified from carcasses located at clusters were zebra ($24.6\% \pm 4.3\%$), wildebeest ($18.8\% \pm 5.6\%$) and impala ($15.4\% \pm 7.1\%$; Table 1). The most important prey species in terms of percent consumed biomass were giraffe ($30.6\% \pm 9.5\%$), zebra ($21.8\% \pm 2.7\%$) and buffalo ($17.9\% \pm 6.2\%$; Table 1). Assuming gut transit times of two and five days respectively, and consecutive kills were from the same prey species in 16% of kills made within two days of each other, we estimated that we missed 147 and 111 feeding events ($n_{\text{scat.sp}} + n_{\text{con.sp}}$) for the high- and low-bias levels respectively. Therefore, we detected 383 and 347 feeding events in total for the high- and low-bias levels respectively. Assuming a high bias of missing carcasses, detected feeding events increased by 69% ($\pm 13\%$) and assuming a low bias of missing carcasses, detected feeding events increased by 52% ($\pm 7\%$). Following scat-supplementation, impala ($19.6\% \pm 5.1\%$ and $17.3\% \pm 3.6\%$ for the high- and low-bias estimates respectively) became numerically the second most consumed prey species, but no change in the order of importance was evident in terms of the consumed biomass (Table 1). Following scat supplementation, we found no significant changes in the estimated diet composition calculated from prey numbers ($G < 13.9$, $p > 0.08$ for all prides) or biomass ($G < 9.3$, $p > 0.42$ for all prides).

Investigating carcasses alone and prior to scat-supplementation, we found 62% ($\pm 10\%$) and 57% ($\pm 12\%$) less impala feeding events in comparison to when carcasses and scats were

combined, depending on whether we implemented a short or long gut transit times (Figure 3a,b). Similarly, without scat-supplementation, we detected 68% less warthog (*Phacochoerus africanus*) feeding events for both the short and long gut transit times (Figure 3a,b). For medium-sized species (100 – 300kg [2–2.5 on the log scale]), the failure to detect carcasses declined to 25% – 45%, whereas for larger species (> 300kg [> 2.5 on the log scale]) we increased our detection of feeding events by 10% – 20% (Figure 3a,b), by incorporating scats and carcasses. As a result there is a significant negative relationship between the log unit weight of each prey species and the percentage of missed feeding events between the two approaches of estimating diet for both the high bias of missing carcasses ($F_{1,9} = 48.5$, $R^2 = 0.84$, $p < 0.005$, Figure 3a) and the low bias of missing carcasses ($F_{1,9} = 70.5$, $R^2 = 0.89$, $p < 0.005$, Figure 3b).

There was a significant difference in the estimated biomass of meat consumed by the prides per day when diets were assessed using the three different approaches ($F_{2,9} = 7.2$, $p < 0.05$, Figure 4). Post hoc tests revealed that this difference was driven by the detection of a significantly higher biomass of consumed meat by lion prides per day, when calculating diet estimates assuming a high bias of missing feeding events (46.0 ± 3.3 kg/day/pride) as compared to estimating consumed biomass from carcasses alone (31.2 ± 2.6 kg/day/pride, $p < 0.05$, Figure 4). The same trend was evident when we calculated consumption rates based on a low bias of missed feeding events (41.0 ± 3.3 kg/day/pride), although the difference to the biomass consumed from carcasses alone was no longer statistically significant ($p = 0.079$, Figure 4).

DISCUSSION

Adequate knowledge of the movement paths of carnivorans determined by GPS (our study) or spoor tracking (Melville, Bothma & Mills, 2004; Marucco *et al.*, 2008) enables data from scats and carcasses to be combined chronologically to identify feeding events. This can be achieved through two different approaches: 1) by collecting scats and then using carcass observations to help define independent feeding events (Marucco *et al.*, 2008); 2) by locating carcasses and then collecting scats to reduce the bias against smaller species in the final diet estimates (current study, Jedrzejewski *et al.*, 2000; Martins *et al.*, 2011).

GPS tracking of large carnivorans results in datasets of carcasses located at clusters and these datasets are increasingly being used to estimate prey selection and feeding rates in large carnivorans (Knopff *et al.*, 2010). However, it must be recognized that prey species are under-represented in such data sets (Franke *et al.*, 2006). Suggestions to counter this bias include reducing the time between GPS fixes (Sand *et al.*, 2005; Webb, Hebblewhite & Merrill, 2008), using alternate movement metrics for model development (Webb *et al.*, 2008), increasing the number of clusters investigated in the field (Sand *et al.*, 2005; Knopff *et al.*, 2009), or collecting scats independently of carcass observations (Martins *et al.*, 2011). However, until now, none has suggested using scats collected at GPS clusters to adjust the bias.

Our results confirm previous GPS-based studies in the northern hemisphere on mountain lions (Anderson & Lindzey, 2003; Knopff *et al.*, 2009) and wolves (Sand *et al.*, 2005; Franke *et al.*, 2006; Webb *et al.*, 2008) that the use of carcasses alone under-estimates the presence of smaller prey species in the diet of large carnivorans. In the Kruger National Park, impala (45–50kg) and warthog (45–100kg) were under-represented by as much as 50%. Similarly, in Hwange National Park (HNP) small species (common duiker *Sylvicapra grimmia* [15–20kg] and reedbuck *Redunca arundinum* [30–70kg]) were considerably under-represented when diets were based on carcass observations estimated from GPS cluster investigation (van Kesteren, 2006).

Despite the considerable under-estimation of these smaller prey species, their inclusion did not significantly alter the estimated composition of the lion diet. This is contrary to findings that wolf diet estimates changed significantly once scat-supplementation was applied in an additive approach (Marucco *et al.*, 2008), but the size of the sample of observed carcasses (51 for wolves vs. 236 for lions) could be a key factor. The effect of small sample size will be exacerbated, if some prey species inhabit inaccessible regions (Marucco *et al.*, 2008). In our open-savanna study area, our diet estimate based on carcasses alone is based on a large sample size, such that detecting an additional 30–37 impala carcasses only resulted in a compositional increase of 1.8–4.2% in the diet.

We did, however, find a significant effect of missed kills on the estimation of biomass consumed by the lion prides. The biomass of food contributed by each prey species is important for the estimation of kill rates, which in turn is important for the conservation and management of carnivorans (Hebblewhite *et al.*, 2003). Predator kill rates are calculated by taking into account all kills over an extended monitoring period (Merrill *et al.*, 2010), but the majority of studies on African carnivorans do not estimate kill rates, because finding the majority of kills over long periods requires continuous observation (Funston *et al.*, 1998), which is often logistically not possible (Martins *et al.*, 2011). Combining data from scats and carcasses over periods, during which researchers are frequently (but not necessarily continuously) monitoring the study animals in the field, may provide a better chance to estimate kill rates for African carnivorans and as such allow the development of functional response curves that will increase the understanding of the link between predators and prey (Merrill *et al.*, 2010). Additionally, accurate knowledge of the biomass consumption by large carnivorans is beneficial for the development of carrying-capacity models (Hayward, O'Brien & Kerley, 2007).

Although this approach deals with correlated clusters of scats associated with social carnivorans, other aspects of sociality are important to consider. Scats could be produced by other individuals not associated with feeding sites, reducing the confidence in combining scats and feeding events into single sample units. Nevertheless, female lions in southern Africa have a high group fidelity (females remain together 87–94% of the time - Funston *et al.*, 1998; Tambling & Belton, 2009; Valeix *et al.*, 2009) and pride females in the study site have limited interaction with each other (C.J. Tambling Unpublished Data), reducing the chance of the scat samples being contaminated by interlopers. Males, on the other hand, join females intermittently (detected at 23 [9.7%] of female kills) and could contaminate clusters to cause an over-estimation of missed feeding events, although the use of genetic testing could verify the origin of scats (Marucco *et al.*, 2008).

Where direct continuous observation of lions is not possible, the use of the scat-supplemented approach, through a combination of GPS point investigation and serial collection of scats, results in much improved estimates of lion dietary consumption. In the case of social carnivorans with fission-fusion societies, genetic identification of individuals producing each scat can improve efforts to obtain a fine-scale representation of diets. Our study demonstrates how, as ecological methods become more diverse, there are increasing opportunities to combine and analyse collected data collected, using multiple techniques to yield less biased and more accurate results to advance large carnivoran research.

Acknowledgments

We thank South African National Parks (SANParks) for permission to conduct this research in the Kruger National Park. The SANParks veterinary service is thanked for the immobilisation of the lions during the study. Augusta Mabunda, Kutani Bulunga and Robert Dugtig are thanked for helping with the GPS follow-up field data collection and James Roxburgh is thanked for assistance with the scat analysis. This work was supported by NIH Grant GM083863 (WMG) and a PhD NRF Grant to CJT.

LITERATURE CITED

- Ackerman BB, Lindzey FG, Hemker TP. Cougar food habits in Southern Utah. *J. Wildl. Manage.* 1984; 48:147.
- Anderson CR, Lindzey FG. Estimating cougar predation rates from GPS location clusters. *J. Wildl. Manage.* 2003; 67:307.
- Andheria AP, Karanth KU, Kumar NS. Diet and prey profiles of three sympatric large carnivores in Bandipur Tiger Reserve, India. *J. Zool.* 2007; 273:169.
- Breuer T. Diet choice of large carnivores in northern Cameroon. *Afr. J. Ecol.* 2005; 43:181.
- Buyts D, Keogh HJ. Notes on the microstructure of hair of the Orycteropodidae, Elephantidae, Equidae, Suidae and Giraffidae. *S. Afr. J. Wildl. Res.* 1984; 14:111.
- Coe MJ, Cumming DH, Phillipson J. Biomass and production of large African herbivores in relation to rainfall and primary production. *Oecologia.* 1976; 22:341.
- Douglas RM. A new method of cross-sectioning hair of larger mammals. *S. Afr. J. Wildl. Res.* 1989; 19:73.
- Ferreira SM, Funston PJ. Estimating lion population variables: prey and disease effects in Kruger National Park, South Africa. *Wildl. Res.* 2010; 37:194.
- Franke A, Caelli T, Kuzyk G, Hudson RJ. Prediction of wolf (*Canis lupus*) kill-sites using hidden Markov models. *Ecol. Model.* 2006; 197:237.
- Funston PJ, Mills MGL, Biggs HC, Richardson PRK. Hunting by male lions: ecological influences and socioecological implications. *Anim. Behav.* 1998; 56:1333. [PubMed: 9933529]
- Gertenbach WPD. Landscapes of the Kruger National Park. *Koedoe.* 1983; 26:9.
- Getz WM, Saltz D. A framework for generating and analyzing movement paths on ecological landscapes. *PNAS.* 2008; 105:19066. [PubMed: 19060192]
- Hayward MW, Hayward GJ, Druce DJ, Kerley GIH. Do fences constrain predator movements on an evolutionary scale? Home range, food intake and movement patterns of large predators reintroduced to Addo Elephant National Park, South Africa. *Biodivers. Conserv.* 2009; 18:887.
- Hayward MW, Kerley GIH. Prey preferences of the lion (*PANTHERA LEO*). *J. Zool.* 2005; 267:309.
- Hayward MW, O'Brien J, Kerley GIH. Carrying capacity of large African predators: Predictions and tests. *Biol. Conserv.* 2007; 139:219.
- Hayward MW, Slotow R. Temporal partitioning of activity in large African carnivores: tests of multiple hypotheses. *S. Afr. J. Wildl. Res.* 2009; 39:109.
- Hebblewhite M, Paquet PC, Pletscher DH, Lessard RB, Callaghan CJ. Development and application of a ratio estimator to estimate wolf kill rates and variance in a multiple-prey system. *Wildl. Soc. Bull.* 2003; 31:933.
- Jedrzejewski W, Jedrzejewska B, Okarma H, Schmidt K, Zub K, Musiani M. Prey selection and predation by wolves in Bialowieza Primeval Forest, Poland. *J. Mammal.* 2000; 81:197.
- Karanth KU, Sunquist ME. Prey selection by tiger, leopard and dhole in tropical forests. *J. Anim. Ecol.* 1995; 64:439.
- Keogh HJ. A photographic reference system of the microstructure of the hair of southern African bovids. *S. Afr. J. Wildl. Res.* 1983; 13:89.
- Knopff KH, Knopff AA, Kortello A, Boyce MS. Cougar Kill Rate and Prey Composition in a Multiprey System. *J. Wildl. Manage.* 2010; 74:1435.
- Knopff KH, Knopff AA, Warren MB, Boyce MS. Evaluating Global Positioning System Telemetry Techniques for Estimating Cougar Predation Parameters. *J. Wildl. Manage.* 2009; 73:586.
- Lehmann MB, Funston PJ, Owen CR, Slotow R. Feeding behaviour of lions (*Panthera leo*) on a small reserve. *S. Afr. J. Wildl. Res.* 2008; 38:66.
- Marker LL, Muntifering JR, Dickman AJ, Mills MGL, MacDonald DW. Quantifying prey preferences of free-ranging Namibian cheetahs. *S. Afr. J. Wildl. Res.* 2003; 33:43.
- Martins Q, Horsnell WGC, Titus W, Rautenbach T, Harris S. Diet determination of the Cape Mountain leopards using global positioning system location clusters and scat analysis. *J. Zool.* 2011; 283:81.
- Marucco F, Pletscher DH, Boitani L. Accuracy of scat samples for carnivore diet analysis: wolves in the Alps as a case study. *J. Mammal.* 2008; 89:665.

- Melville HIAS, Bothma JdP, Mills MGL. Prey selection by caracal in the Kgalagadi Transfrontier Park. *S. Afr. J. Wildl. Res.* 2004; 34:67.
- Merrill E, Sand Hk, Zimmermann B, McPhee H, Webb N, Hebblewhite M, Wabakken P, Frair JL. Building a mechanistic understanding of predation with GPS-based movement data. *Philosophical Transactions of the Royal Society B: Biological Sciences.* 2010; 365:2279.
- Mills, MGL. A comparison of methods used to study food habits of large African carnivores. In: McCullough, DR.; Barrett, RH., editors. *Wildlife 2001: Populations.* London, United Kingdom: Elsevier Science Publishers LTD; 1992.
- Mills MGL, Biggs HC. Prey apportionment and related ecological relationships between large carnivores in Kruger National Park. *Symposium of the Zoological Society of London.* 1993; 65:253.
- Mills MGL, Shenk TM. Predator-prey relationships: the impact of lion predation on wildebeest and zebra populations. *J. Anim. Ecol.* 1992; 61:693.
- Owen-Smith N. Changing vulnerability to predation related to season and sex in an African ungulate assemblage. *Oikos.* 2008; 117:602.
- Owen-Smith N, Mason DR. Comparative changes in adult vs. juvenile survival affecting population trends of African ungulates. *J. Anim. Ecol.* 2005; 74:762.
- Power RJ. Prey selection of lions *Panthera leo* in a small, enclosed reserve. *Koedoe.* 2002; 45:67.
- Radloff FGT, du Toit JT. Large predators and their prey in a southern African savanna: a predator's size determines its prey size range. *J. Anim. Ecol.* 2004; 73:410.
- Sand H, Zimmermann B, Wabakken P, Andren H, Pedersen HC. Using GPS technology and GIS cluster analyses to estimate kill rates in wolf-ungulate ecosystems. *Wildl. Soc. Bull.* 2005; 33:914.
- Smuts GL. Diet of lions and spotted hyaenas assessed from stomach contents. *S. Afr. J. Wildl. Res.* 1979; 9:19.
- Smuts GL, Whyte IJ, Dearlove TW. A mass capture technique for lions. *East African Wildlife Journal.* 1977; 15:81.
- Tambling, CJ. *Zoology and Entomology: 174.* Pretoria: University of Pretoria; 2010. *Movement Analysis for Monitoring Predation by a Large Carnivore: Lions in Kruger National Park.*
- Tambling CJ, Belton L. Feasibility of using proximity tags to locate female lion *Panthera leo* kills. *Wildl. Biol.* 2009; 15:435.
- Tambling CJ, Cameron EZ, du Toit JT, Getz WM. A Comparison of Methods for Locating African Lion Kills Using Global Positioning System Movement Data. *J. Wildl. Manage.* 2010; 74:549.
- Valeix M, Loveridge AJ, Chamaillé-Jammes S, Davidson Z, Murindagomo F, Fritz H, MacDonald DW. Behavioural adjustments of African herbivores to predation risk by lions: Spatiotemporal variations influence habitat use. *Ecology.* 2009; 90:23. [PubMed: 19294909]
- van Kesteren, F. *Wildlife Conservation Research Unit.* Oxford: University of Oxford; 2006. *Analysis of the diet of African Lions (Panthera leo) in Hwange National Park, Zimbabwe.*
- Venter, FJ.; Scholes, RJ.; Eckhardt, HC. The Abiotic Template and its Associated Vegetation Patterns. In: du Toit, JT.; Rogers, KH.; Biggs, HC., editors. *The Kruger Experience: Ecology and Management of Savanna Heterogeneity:* 83. Washington DC, USA: Island Press; 2003.
- Webb NF, Hebblewhite M, Merrill EH. Statistical methods for identifying wolf kill sites using global positioning system locations. *J. Wildl. Manage.* 2008; 72:798.
- Zar, JH. *Biostatistical Analysis.* 4th edn.. New Jersey: Prentice Hall; 1999.

Figure 1.
Study area in the central region of the Kruger National Park, South Africa, scats located at GPS clusters, and kills located at GPS clusters

Figure 2.

Proportion of scats ($\pm$ SE) containing remains of the same prey species (grey) or a different species (black) as the focal kill (at the kill site [KS] and each day thereafter), based on the investigation of GPS clusters in the central region of the Kruger National Park, South Africa, between April 2005 and May 2007

Figure 3.
Relationship between the log unit mass of prey species and the estimated percent of missed kills, using the scat-supplemented approach in the central region of the Kruger National Park, South Africa, between April 2005 and May 2007 (A. – high bias of missing feeding events, B. – low bias of missing feeding events)

Figure 4. Daily food consumption by female lion groups in the Kruger National Park, South Africa, calculated using carcasses against carcasses and scats for short and long gut transit times.

Mean percent representation ($\pm$ SE) of prey species in the diet (numerical and biomass consumed) of lions in the central region of Kruger National Park, South Africa, between April 2005 and May 2007, utilising three approaches to estimate the diet. The uncorrected estimate uses only carcasses found at GPS clusters, whereas the scat-supplemented approach combines scats and kills observed at GPS clusters along a known movement trajectory for each collared lion. The high bias assumes a gut transit time of two days and the low bias assumes a gut transit time of five days, which are used to identify feeding events that have been missed.

Table 1

	Numerical diet estimate				Biomass diet estimate			
	Carcasses		Scat-supplemented		Carcasses		Scat-supplemented	
	Av.	High bias	Low bias	Low bias	Av.	High bias	Low bias	Low bias
Buffalo	10.1 (3.9)	7.8 (3.3)	7.8 (3.3)	17.9 (6.2)	15.3 (6.2)	16.5 (5.9)		
Elephant	0.2 (0.2)	0.1 (0.1)	0.1 (0.1)	1.6 (1.6)	1.1 (1.1)	1.2 (1.2)		
Giraffe	10.6 (3.9)	8.9 (3.4)	7.1 (2.3)	30.6 (9.5)	28.4 (9.4)	26.4 (8)		
Impala	15.4 (7.1)	19.6 (5.1)	17.3 (3.6)	3.9 (1.7)	5.4 (1.3)	5.5 (1.3)		
Kudu	7.4 (2.5)	7.9 (2.1)	7.2 (1.5)	4.8 (1.7)	5.6 (1.5)	5.8 (1.4)		
Warthog	2.2 (1.4)	4.4 (0.8)	3.8 (0.5)	0.5 (0.3)	1.3 (0.1)	1.3 (0.1)		
Waterbuck	7 (1.4)	9.1 (3.4)	7.5 (2.3)	5.4 (1.5)	8.6 (4.3)	7.5 (3.1)		
Wildebeest	18.8 (5.6)	17.8 (5)	16.4 (4.4)	11.6 (4.6)	12.2 (4.4)	12.2 (4.2)		
Zebra	24.6 (4.3)	20.9 (3.7)	20 (3.5)	21.8 (2.7)	20.6 (2.6)	22 (2.3)		
Ostrich	0.6 (0.6)	0.6 (0.3)	0.6 (0.3)	0.3 (0.3)	0.3 (0.2)	0.4 (0.2)		
Porcupine	0.6 (0.6)	1.4 (0.5)	1.4 (0.5)	0 (0)	0.1 (0)	0.1 (0)		
Unknown	2.5 (2.2)	1.5 (1.3)	1.5 (1.3)	1.6 (1.4)	1.1 (1)	1.2 (1.1)		