A new PstI polymorphism between VK21 and St14 in Xq28 (DXS523)

S.Kenwick⁺, P.Bridge¹, D.Lillicrap¹, J.Bainton and J.Gitschier^{*}

Department of Medicine and Howard Hughes Medical Institute, University of California, San Francisco, CA 94143, USA and ¹Department of Pathology, Queen's University, Kingston, Ontario K7L 3N6, Canada

Source/Description: pDK9.5 is a 900 bp BglII/HindIII single copy genomic DNA insert in BamHI/HindIII-cut pBLUESCRIPT SK⁺ (Stratagene). The insert is derived from a 15 kb λ Dash clone from a hybrid cell line (X3000-11) containing human Xq24-qter (1) (library prepared by Kenwick, unpublished). This sequence has been assigned the provisional HGM Workshop number of DXS523.

Polymorphism: pDK9.5 identifies a 2 allele polymorphism following digestion of genomic DNA with PstI. The alleles are represented by fragments of 6.5 kb (A1) and 5 kb (A2).

Frequency: Allele frequencies are 28% for A1 and 72% for A2 (in 103 chromosomes tested) giving a predicted heterozygosity of 40%.

Not Polymorphic For: BclI, BglII, HaeIII, MspI, and TaqI.

Chromosomal Localization: Hybridization to DNA from a panel of somatic cell hybrids reveals that pDK9.5 is located proximal to St14 (DXS52) and distal to VK21 (DXS296) in Xq28. Linkage data further indicates that pDK9.5 is proximal to St14. No recombination was detected with DX13 (DXS15), with a LOD score of 7.2 at Q = 0.

Mendelian Inheritance: Demonstrated in 20 multigeneration families with hemophilia A, fragile X, or adrenoleukodystrophy.

Other Comments: pDK9.5 hybridizes to a NotI fragment of 350 kb in genomic DNA, distinct from fragments identified by St14, 1A1, DX13, F8, G6PD, and hs7. The polymorphism is in linkage equilibrium with the PstI polymorphisms identified by 767 (DXS115) and 1A1 (DXS374) (2).

Acknowledgements: JG is an assistant investigator with the Howard Hughes Medical Institute. We thank D.Callen, M.Hors-Cayla, S. and D.Ledbetter, R.Nussbaum, M.Rocchi, H.-H.Ropers, C.Schwartz, G.Sutherland and S.Warren for providing hybrid cell lines.

References: 1)Nussbaum,R.L., Airhart,S.D. and Ledbetter,D.H. (1986) *Am. J. Med. Genet.* **23**, 457-466. 2)Patterson,M.N., Bell,M.V., Bloomfield,J., Flint,T., Dorkins,H., Thibodeau,S.N., Schaid,D., Bren,G., Schwartz,C.E., Wieringa,B., Ropers,H.-H., Callen,D.F., Sutherland,G., Froster-Iskenius,U., Vissing,H. and Davies,K.E. (1989) *Genomics* **4**, 570-578.

A TaqI polymorphism adjacent to the factor VIII gene (F8C)

S.Kenwick⁺, P.Bridge¹, D.Lillicrap¹, A.-E.Lehesjoki², J.Bainton and J.Gitschier^{*}

Department of Medicine and Howard Hughes Medical Institute, University of California, San Francisco CA 94143, USA, ¹Department of Pathology, Queen's University, Kingston, Ontario K7L 3N6, Canada and ²Department of Medical Genetics, University of Helsinki, Helsinki, Finland

Source/Description: p701.1 is a 1.3 kb BamHI/HindIII insert of genomic DNA in pGem4 (Promega) isolated by walking from the 5' end of the factor VIII gene.

Polymorphism: p701.1 identifies a 2 allele TaqI polymorphism with fragment sizes of 9.5 kb (H1) or 4.0 kb (H2).

Frequency: Allele frequencies of 72% (H1) and 28% (H2) are observed.

Not Polymorphic For: BclI, BglII, BstXI, HaeIII, MspI, PstI, and RsaI.

Chromosomal Localization: p701.1 is located 17 kb 5' to the first exon of the factor VIII gene in Xq28.

Mendelian Inheritance: Demonstrated in ten 2- and 3-generation families.

Other Comments: The polymorphism is in strong linkage disequilibrium with the BclI polymorphism described for the factor VIII gene (1). The 9.5 kb TaqI fragment is associated with the 0.9 kb allele of the BclI system and the 4 kb with the 1.17 kb BclI band. For 38 independent chromosomes only one discordant haplotype was observed in which the 4 kb TaqI fragment was associated with the 0.9 kb BclI fragment. The correlation coefficient (ρ) is 0.93 with $\chi^2 = 33$. p701.1 is a convenient probe to use in conjunction with St14 (DXS52) (2) as both reveal TaqI polymorphisms.

Acknowledgement: JG is an assistant investigator with the Howard Hughes Medical Institute.

References: 1)Gitschier,J., Drayna,D., Tuddenham,E.G.D., White,R.L. and Lawn,R.M. (1985) *Nature* **314**, 738-740. 2)Oberle,I., Drayna,D., Camerino,G., White,R. and Mandel,J.L. (1985) *Proc. Natl. Acad. Sci. USA* **82**, 2824-2828.

* To whom correspondence should be addressed

⁺Present address: University of Cambridge Clinical School, Department of Medicine, Addenbrooke's Hospital, Cambridge CB2 2QQ, UK