Published in final edited form as:

Neurosci Lett. 2012 March 14; 512(1): 48–52. doi:10.1016/j.neulet.2012.01.062.

Mice lacking *Wnt2b* are viable and display a postnatal olfactory bulb phenotype

Tadasuke Tsukiyama^{1,2,3} and Terry P. Yamaguchi¹

¹Center for Cancer Research, Cancer and Developmental Biology Laboratory, Cell Signaling in Vertebrate Development Section, NCI-Frederick, NIH, Frederick, Maryland 21702, USA

²Department of Biochemistry, Hokkaido University Graduate School of Medicine, Sapporo, Hokkaido 060-8638, Japan

Abstract

Wnts are secreted glycoproteins that play important roles in embryonic development. *Wnt2b* is transiently expressed in the primitive streak (PS) during gastrulation and in several organs during organogenesis. To determine the biological function of *Wnt2b* during mouse development, we established a conditional null allele of *Wnt2b*. Mice lacking *Wnt2b* were viable, fertile, and displayed a normal life span, however, the olfactory bulb in adult *Wnt2b* mutant mice was significantly reduced in length. Our results suggest that *Wnt2b* primarily plays a supportive role in gastrulation and organogenesis, functioning redundantly with canonical Wnts, such as *Wnt2*, in numerous tissues.

Keywords

Wnt; embryogenesis; brain; redundancy; olfactory bulb

Wnt proteins encode a large family of cysteine-rich secreted glycoproteins that play established roles in embryonic development and tumorigenesis [16, 17, 25]. Nineteen Wnt genes have been identified in murine and human genomes. These Wnts were initially functionally classified into two groups based on their activity in in vitro transformation assays and 2°-body axis induction assays in *Xenopus* embryos. Generally speaking, Wnt1-class Wnts (canonical Wnts), which include Wnt1, 3a, 8a and 8b, transform cells and induce the formation of an additional body axis when overexpressed in frog embryos, while Wnt5a class Wnts, such as Wnt4, 5a and 11, do not [5].

Wnt2b was first identified by degenerate RT-PCR and designated *Wnt13* [28]. *Wnt2b* binds the CRD domain of Frizzled4 and 5 to stabilize cytosolic β catenin, suggesting that *Wnt2b* is a canonical Wnt [11]. *Wnt2b* is transiently expressed in the mesoderm of the PS at mid-gastrulation stages only (E7–7.75) [25, 28] and is later expressed in numerous embryonic tissues including the optic vesicle (OpV)/retina, otic vesicle (OtV), kidney, lung, and limb bud, as well as the adult ovary [3, 7, 11, 14, 20, 28]. *Wnt2b* is also expressed in the

© 2012 Elsevier Ireland Ltd. All rights reserved.

³To whom correspondence should be addressed (contact address). Department of Biochemistry, Hokkaido University Graduate School of Medicine; N15, W7, Kita-ku, Sapporo, Hokkaido 060-8638, Japan, Phone: +81-11-706-5047, Fax: +81-11-706-5169, tsukit@med.hokudai.ac.jp.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

developing mouse brain at several developmental stages. It's expressed in the dorsal midline of the telencephalon and the mesencephalon at E8.5–9.5, cortical hem at E11.5–17.5, hippocampal fimbria at E15.5–17.5 and pineal body. Based on these observations, *Wnt2b* has been proposed to play a role in the patterning of the cerebral cortex however a requisite role for *Wnt2b* in these processes has not been rigorously tested.

A targeting vector for the *Wnt2b* locus was constructed as follows. A 5.9 kb genomic *XbaI* fragment for long arm 1 (LA1) and a 2.1 kb genomic *XbaI/EcoRI* fragment for LA2, which contains exon 2 and 3 for deletion, were inserted into *Sall* and *SmaI* sites of pML8 vector, respectively. A 1.2 kb genomic *EcoRI/KpnI* fragment for short arm (SA) was inserted into the *SacII* site of pK-11 vector, which contains a loxP-FRT-PGKNeo-FRT cassette for positive selection. An *XhoI/XbaI* fragment of LA1-loxP-LA2 isolated from pML8-LA1-LA2 was subcloned into *XhoI/XbaI* sites of pPGKtk, which contains a PGKtk cassette for negative selection. A *KpnI/SacI* fragment of loxP-FRT-PGKNeo-FRT-SA isolated from pK-11-SA was inserted into the *XbaI* site of pPGKtk-LA1-loxP-LA2. The targeting vector was linearized with *NotI* for transfection into ES cells. *NotI*-linearized *Wnt2b* conditional targeting vector DNA (80 µg) was electroporated into 8×10^7 of W9.5 ES cells (x2 pulses of 320 V/125 µF), and the cells were cultured on a layer of mitotically inactivated feeder cells carrying genomic PGKNeo gene. ES cell colonies that were resistant to 300 µg/ml of G418 and 1 µM gancyclovir were independently isolated. These clones of ES cells were screened by Southern blot analysis of *KpnI*-digested genomic DNA (15 µg) using a 370 bp 5' probe or *XbaI*-digested genomic DNA using a 540 bp 3' probe (Fig. 1A and B). *Wnt2b*^{WT/flox} ES clones were injected into albino C57BL/6N blastocysts to generate chimeras. Chimeric male mice that displayed a high contribution of agouti coat color were bred with C57BL/6N female mice to obtain a germ line-transmitted *Wnt2b*^{flox} allele. *Wnt2b*^{WT/flox} mice were then bred with βactin-Cre transgenic mice to delete exon 2 and 3 to generate the *Wnt2b*^Δ allele. *Wnt2b*^{WT/flox} and *Wnt2b*^{WT/Δ} mice were intercrossed to generate *Wnt2b*^{flox/flox} and *Wnt2b*^{Δ/Δ} mice, respectively. Genotypes were confirmed by PCR with allele-specific primers (KO1: 5'-GCC TCT CAC ACC AGC GTG TAA GAG-3', KO4: 5'-GTA ATT GAG TGG TCT CCA CC-3', Neo3': 5'-ATCAGC AGCCTC TGT TCC ACA TAC-3', PJL: 5'-TGC TAA AGC GCA TGC TCC AGA CTG-3', 2b-1: 5'-AGG AGT CTG CTC CTG ATT ACA GCC-3') to detect WT (200 bp fragment), flox (260 bp) or Δ (320 bp) allele (Fig. 1A and C).

Intercrosses of *Wnt2b*^{WT/flox} or *Wnt2b*^{WT/Δ} mice produced *Wnt2b*^{flox/flox} or *Wnt2b*^{Δ/Δ} (*Wnt2b* KO) mice in the expected Mendelian ratios (Table 1). *Wnt2b*^{Δ/Δ} mice appeared normal (Fig. 2A) and healthy during their lifespan (not shown). These results indicate that inactivation of *Wnt2b* did not affect body plan formation, despite *Wnt2b* being strongly and specifically expressed in the PS during gastrulation [28]. Indeed, several canonical Wnts, including *Wnt3*, *3a*, and *8a*, are co-expressed with *Wnt2b* in the primitive streak (PS) [25] and therefore could functionally compensate for the absence of *Wnt2b*. Only *Wnt3* and *Wnt3a* are required for A–P axis formation and extension [10, 15, 22, 25, 26], suggesting that they are the compensatory genes during gastrulation in *Wnt2b*^{Δ/Δ} animals. Intercrosses of *Wnt2b*^{Δ/Δ} mice led to viable *Wnt2b*^{Δ/Δ} progeny (Table 1), indicating that *Wnt2b*^{Δ/Δ} males and females are fertile and that there is no requirement for *Wnt2b* in the ovary where it is also expressed [20]. In addition, histological analyses indicated that limb, liver, kidney, eye and inner ear morphology was normal (not shown), despite strong *Wnt2b* expression in these tissues [3, 4, 8, 11, 12, 14, 18, 28](Fig. 2B, C).

We next examined the brain since *Wnt2b* is expressed there during neural development. Brains isolated from *Wnt2b*^{+/-Δ} and *Wnt2b*^{Δ/Δ} mice at 4 months of age were fixed with 10% formalin, dehydrated by ethanol and xylene and then embedded in paraffin to make sections for histochemical staining. Five-µm-thick sections were stained by hematoxylin and eosin.

Histochemical analysis indicated that there were no obvious patterning defects in the cerebral cortex or hippocampus (Fig. 3B). As in many other tissues, *Wnt2b* is coexpressed in the brain with several other canonical Wnts, particularly in the cortical hem, hippocampus and pineal gland [1, 21]. Surprisingly, we found that both the length (major axis) and width (minor axis) of the olfactory bulb (OfB) was significantly reduced (length, $p < 0.01$; width, $p < 0.05$, $n = 3$ in both *Wnt2b*^{WT/Δ} and *Wnt2b*^{Δ/Δ}, Student's *t*-test) in the *Wnt2b*^{Δ/Δ} brain (Fig. 3A–C). Small OfB's were observed in dissected, whole, adult *Wnt2b*^{Δ/Δ} brains, as well as in sagittal sections. To clarify whether the small OfB phenotype in adult *Wnt2b*^{Δ/Δ} mice arises during embryonic or postnatal development, we examined the size of the OfB in earlier-staged perinatal (P0, postnatal day 0) embryos. Contrary to the results observed in adult mice, significant size differences in OfBs isolated from perinatal *Wnt2b*^{+/-Δ} and *Wnt2b*^{Δ/Δ} mice were not observed (*Wnt2b*^{WT/Δ}; $n=4$ and *Wnt2b*^{Δ/Δ}; $n=7$), (Fig. 3D, E). These results suggest that the smaller OfB's observed in adult *Wnt2b*^{Δ/Δ} mice arises postnatally. We can conceive of several possible explanations for the small OfB phenotype. A small OfB could arise indirectly, for example due to a craniofacial developmental defect that physically constricts OfB development. Indeed, defects in nasal bone development have been observed in *Axin2* deficient mice in which Wnt/βcatenin signaling is perturbed [27]. Alternatively, a smaller OfB could be caused by impaired neuroblast cell migration into the OfB, or by impaired neurogenesis in the subventricular zone (SVZ). It is well known that the adult OfB is maintained through the recruitment of neuronal progenitor cells originating in the SVZ. These neuronal progenitors travel from the SVZ to the granule cell layer and glomerular layer of OfB along the rostral migratory stream (RMS) and differentiate into interneurons including granule cells or periglomerular cells [2, 24]. Wnt ligands secreted from the fimbrial neuroepithelium into the cerebrospinal fluid are thought to activate canonical Wnt signaling in the SVZ [13, 19]. In addition, it is also reported that loss of Sp8, which is a transcription factor expressing in postnatal SVZ and RMS and is a target gene of Wnt/βcatenin signaling leads the developmental defects in the generation and diversity of OfB interneurons [9] [23]. Although highly speculative, it is possible that loss of *Wnt2b* expression in fimbrial neuroepithelium leads to reduced *Wnt2b* in CSF and, consequently, reduced numbers of SVZ progenitors that populate the OfB. Clearly, additional studies are required to fully understand the underlying cause of the small OfB observed in adult *Wnt2b*^{Δ/Δ} mice.

We have previously reported that the combined inactivation of *Wnt2b* and *Wnt2* results in the complete loss of lung progenitor cells, providing irrefutable evidence that the closely related *Wnt2* can compensate for the absence of *Wnt2b* [6]. The present results showing that *Wnt2b* is not required at most of the sites at which it is expressed suggests that many other canonical Wnts can stand-in for *Wnt2b*, and that *Wnt2b* primarily plays a secondary, supportive role during embryogenesis.

Highlights

Wnt2b KO mice were generated to investigate the biological role of Wnt2b.
 Wnt2b KO mice appeared to be normal, healthy and reproductive.
 Wnt2b KO mice did not exhibit abnormalities in most organs.
 Wnt2b KO mice exhibited a short olfactory bulb phenotype.

Abbreviations

A-P anterior-posterior

PS	primitive streak
OpV	optic vesicle
OtV	otic vesicle
OfB	olfactory bulb
Hp	hippocampus
SVZ	subventricular zone

Acknowledgments

We thank Kristin K. Biris for laboratory management, Cheri A. Rhoderick for administrative assistance and Dr. Sawamoto for comments on the manuscript. We also thank Serguei Kozlov and Jaime Greear for embryonic engineering and maintenance of mice. This work was supported by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research and by a JSPS Research Fellowship for Japanese Biomedical and Behavioral Researchers at NIH.

References

1. Abu-Khalil A, Fu L, Grove EA, Zecevic N, Geschwind DH. Wnt genes define distinct boundaries in the developing human brain: implications for human forebrain patterning. *J Comp Neurol*. 2004; 474:276–288. <http://onlinelibrary.wiley.com/doi/10.1002/cne.20112/full>. [PubMed: 15164427]
2. Cayre M, Canoll P, Goldman JE. Cell migration in the normal and pathological postnatal mammalian brain. *Prog Neurobiol*. 2009; 88:41–63. <http://www.sciencedirect.com/science/article/pii/S0301008209000239>. [PubMed: 19428961]
3. Cho SH, Cepko CL. Wnt2b/beta-catenin-mediated canonical Wnt signaling determines the peripheral fates of the chick eye. *Development*. 2006; 133:3167–3177. <http://dev.biologists.org/content/133/16/3167.long>. [PubMed: 16854977]
4. Choo D, Ward J, Reece A, Dou H, Lin Z, Greinwald J. Molecular mechanisms underlying inner ear patterning defects in kreisler mutants. *Dev Biol*. 2006; 289:308–317. <http://www.sciencedirect.com/science/article/pii/S001216060500713X>. [PubMed: 16325169]
5. Du SJ, Purcell SM, Christian JL, McGrew LL, Moon RT. Identification of distinct classes and functional domains of Wnts through expression of wild-type and chimeric proteins in *Xenopus* embryos. *Molecular & Cellular Biology*. 1995; 15:2625–2634. <http://mcb.asm.org/cgi/reprint/15/5/2625?view=long&pmid=7739543>. [PubMed: 7739543]
6. Goss AM, Tian Y, Tsukiyama T, Cohen ED, Zhou D, Lu MM, Yamaguchi TP, Morrissey EE. Wnt2/2b and beta-catenin signaling are necessary and sufficient to specify lung progenitors in the foregut. *Dev Cell*. 2009; 17:290–298. <http://www.sciencedirect.com/science/article/pii/S1534580709002226>. [PubMed: 19686689]
7. Grove EA, Tole S, Limon J, Yip L, Ragsdale CW. The hem of the embryonic cerebral cortex is defined by the expression of multiple Wnt genes and is compromised in *Gli3*-deficient mice. *Development*. 1998; 125:2315–2325. <http://dev.biologists.org/content/125/12/2315.long>. [PubMed: 9584130]
8. Kawakami Y, Capdevila J, Buscher D, Itoh T, Rodriguez Esteban C, Izpisua Belmonte JC. WNT signals control FGF-dependent limb initiation and AER induction in the chick embryo. *Cell*. 2001; 104:891–900. <http://www.sciencedirect.com/science/article/pii/S0092867401002859>. [PubMed: 11290326]
9. Kawakami Y, Esteban CR, Matsui T, Rodriguez-Leon J, Kato S, Izpisua Belmonte JC. Sp8 and Sp9, two closely related buttonhead-like transcription factors, regulate *Fgf8* expression and limb outgrowth in vertebrate embryos. *Development*. 2004; 131:4763–4774. <http://www.sciencedirect.com/science/article/pii/S089662730600047X>. [PubMed: 15358670]
10. Kemp C, Willems E, Abdo S, Lambiv L, Leyns L. Expression of all Wnt genes and their secreted antagonists during mouse blastocyst and postimplantation development. *Dev Dyn*. 2005; 233:1064–1075. <http://onlinelibrary.wiley.com/doi/10.1002/dvdy.20408/full>. [PubMed: 15880404]

11. Kubo F, Takeichi M, Nakagawa S. Wnt2b controls retinal cell differentiation at the ciliary marginal zone. *Development*. 2003; 130:587–598. <http://dev.biologists.org/content/130/3/587.long>. [PubMed: 12490564]
12. Kubo F, Takeichi M, Nakagawa S. Wnt2b inhibits differentiation of retinal progenitor cells in the absence of Notch activity by downregulating the expression of proneural genes. *Development*. 2005; 132:2759–2770. <http://dev.biologists.org/content/132/12/2759.long>. [PubMed: 15901663]
13. Lehtinen MK, Zappaterra MW, Chen X, Yang YJ, Hill AD, Lun M, Maynard T, Gonzalez D, Kim S, Ye P, D'Ercole AJ, Wong ET, LaMantia AS, Walsh CA. The cerebrospinal fluid provides a proliferative niche for neural progenitor cells. *Neuron*. 2011; 69:893–905. <http://www.sciencedirect.com/science/article/pii/S0896627311000717>. [PubMed: 21382550]
14. Lin Y, Liu A, Zhang S, Ruusunen T, Kreidberg JA, Peltoketo H, Drummond I, Vainio S. Induction of ureter branching as a response to Wnt-2b signaling during early kidney organogenesis. *Dev Dyn*. 2001; 222:26–39. <http://onlinelibrary.wiley.com/doi/10.1002/dvdy.1164/full>. [PubMed: 11507767]
15. Liu P, Wakamiya M, Shea MJ, Albrecht U, Behringer RR, Bradley A. Requirement for Wnt3 in vertebrate axis formation. *Nat Genet*. 1999; 22:361–365. http://www.nature.com/ng/journal/v22/n4/full/ng0899_361.html. [PubMed: 10431240]
16. Logan CY, Nusse R. The Wnt signaling pathway in development and disease. *Annu Rev Cell Dev Biol*. 2004; 20:781–810. <http://www.annualreviews.org/doi/full/10.1146/annurev.cellbio.20.010403.113126>. [PubMed: 15473860]
17. Moon RT, Kohn AD, De Ferrari GV, Kaykas A. WNT and beta-catenin signalling: diseases and therapies. *Nat Rev Genet*. 2004; 5:691–701. <http://www.nature.com/nrg/journal/v5/n9/full/nrg1427.html>. [PubMed: 15372092]
18. Ober EA, Verkade H, Field HA, Stainier DY. Mesodermal Wnt2b signalling positively regulates liver specification. *Nature*. 2006; 442:688–691. <http://www.nature.com/nature/journal/v442/n7103/full/nature04888.html>. [PubMed: 16799568]
19. Pino D, Choe Y, Pleasure SJ. Wnt5a controls neurite development in olfactory bulb interneurons. *ASN Neuro*. 2011; 3:e00059. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3155196/>. [PubMed: 21539518]
20. Ricken A, Lochhead P, Kontogiannina M, Farookhi R. Wnt signaling in the ovary: identification and compartmentalized expression of wnt-2, wnt-2b, and frizzled-4 mRNAs. *Endocrinology*. 2002; 143:2741–2749. <http://endo.endojournals.org/content/143/7/2741.long>. [PubMed: 12072409]
21. Shimogori T, Banuchi V, Ng HY, Strauss JB, Grove EA. Embryonic signaling centers expressing BMP, WNT and FGF proteins interact to pattern the cerebral cortex. *Development*. 2004; 131:5639–5647. <http://dev.biologists.org/content/131/22/5639.full>. [PubMed: 15509764]
22. Takada S, Stark KL, Shea MJ, Vassileva G, McMahon JA, McMahon AP. Wnt-3a regulates somite and tailbud formation in the mouse embryo. *Genes Dev*. 1994; 8:174–189. <http://genesdev.cshlp.org/content/8/2/174.long>. [PubMed: 8299937]
23. Wacław RR, Allen ZJ 2nd, Bell SM, Erdelyi F, Szabo G, Potter SS, Campbell K. The zinc finger transcription factor Sp8 regulates the generation and diversity of olfactory bulb interneurons. *Neuron*. 2006; 49:503–516. <http://dev.biologists.org/content/131/19/4763.long>. [PubMed: 16476661]
24. Whitman MC, Greer CA. Adult neurogenesis and the olfactory system. *Prog Neurobiol*. 2009; 89:162–175. <http://www.sciencedirect.com/science/article/pii/S0301008209000999>. [PubMed: 19615423]
25. Yamaguchi TP. Heads or tails: Wnts and anterior-posterior patterning. *Curr Biol*. 2001; 11:R713–R724. <http://www.sciencedirect.com/science/article/pii/S0960982201004171>. [PubMed: 11553348]
26. Yoshikawa Y, Fujimori T, McMahon AP, Takada S. Evidence that absence of Wnt-3a signaling promotes neuralization instead of paraxial mesoderm development in the mouse. *Dev Biol*. 1997; 183:234–242. <http://www.sciencedirect.com/science/article/pii/S001216069798502X>. [PubMed: 9126297]

27. Yu HM, Jerchow B, Sheu TJ, Liu B, Costantini F, Puzas JE, Birchmeier W, Hsu W. The role of Axin2 in calvarial morphogenesis and craniosynostosis. *Development*. 2005; 132:1995–2005. <http://dev.biologists.org/content/132/8/1995.long>. [PubMed: 15790973]
28. Zakin LD, Mazan S, Maury M, Martin N, Guenet JL, Brulet P. Structure and expression of Wnt13, a novel mouse Wnt2 related gene. *Mech Dev*. 1998; 73:107–116. <http://www.sciencedirect.com/science/article/pii/S0925477398000409>. [PubMed: 9545553]

Figure 1. Generation of *Wnt2b* conditional knockout mice. (A) Schematic of gene targeting for the *Wnt2b* locus. B, *Bam*HI; E1, *Eco*RI; K, *Kpn*I; S, *Spe*I; X, *Xba*I. (B) Southern hybridization for *Wnt2b*-targeted ES cells. Both probes for the 5' locus and 3' locus were used for screening. (C) Genotyping for WT, flox and Δ alleles of the *Wnt2b* locus by PCR. Positions of primers to distinguish WT, flox and Δ alleles of the *Wnt2b* locus are indicated in A. All WT, flox and Δ alleles were typed with single PCR reaction (3-primer mix PCR) using primers KO1, KO4 and Neo3'.

Figure 2.

Normal appearance of *Wnt2b* mutant mice. (A) *Wnt2b*^{WT/Δ} (agouti coat color) and *Wnt2b*^{Δ/Δ} (albino coat color) mice are shown. (B–C) Eye (B) and limbs (C) of *Wnt2b*^{WT/WT}, *Wnt2b*^{WT/Δ} and *Wnt2b*^{Δ/Δ} mice are indicated. FL, forelimb; HL, hindlimb.

Figure 3.

Postnatal development of the brain in *Wnt2b* mutant mice. (A) Adult brains isolated from 4 month old *Wnt2b*^{WT/Δ} and *Wnt2b*^{Δ/Δ} mice are shown. Scale bar represents 5 mm. (B) 4 month old *Wnt2b*^{flx/flx} and *Wnt2b*^{Δ/Δ} brains were sectioned sagittally (5 μm) and stained with hematoxylin and eosin. (C) Measurements of adult OfBs (length, distance between solid white lines; widths, measured between two dashed lines in Fig. 3A) in *Wnt2b*^{WT/Δ} control (n=3) and *Wnt2b*^{Δ/Δ} (n=3) mice are indicated. (D) Brains isolated from perinatal *Wnt2b*^{WT/Δ} and *Wnt2b*^{Δ/Δ} mice are shown. Scale bar indicates 5 mm. (E) Measurements of OfBs isolated from perinatal *Wnt2b*^{WT/Δ} (n=4) and *Wnt2b*^{Δ/Δ} (n=7) mice are indicated. OfB, olfactory bulb; Hp, hippocampus; SVZ, subventricular zone.

Table 1

Wnt2b mutant mice were intercrossed with each other, and then the genotypes of offsprings obtained from the intercrossing were determined by PCR (shown in Fig. 1A and C) at 3 weeks of age. The numbers and ratios of each genotype are indicated.

Intercross	No. of pups (%)			Total
<i>Wnt2b</i> ^{WT/flox}	WT/WT 13 (29.5)	WT/flox 19 (43.2)	flox/flox 12 (27.2)	44
<i>Wnt2b</i> ^{WT/Δ}	WT/WT 30 (31.9)	WT/Δ 40 (42.6)	Δ/Δ 24 (25.5)	94
<i>Wnt2b</i> ^{Δ/Δ}	WT/WT 0 (0)	WT/Δ 0 (0)	Δ/Δ 14 (100)	14