CORRESPONDENCE

Epidemic Keratoconjunctivitis: The Current Situation and Recommendations for Prevention and Treatment

by Dr. med. Birthe Meyer-Rüsenberg, Dr. med. Ulrike Loderstädt, Prof. Dr. med. Gisbert Richard, Prof. Dr. rer. nat. Paul-Michael Kaulfers, Dr. med. Caroline Gesser in volume 27/2011

Prof. Dr. med. Dieter Schmidt
University Medical Center Freiburg, Germany
Dieter.Schmidt@uniklinik-freiburg.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Avoid Further Transmission

As far as the diagnosis is concerned, Wollensak (1965) said that infants and toddler—in contrast to adolescents and adults—often show signs of general illness, such as a raised temperature and fatigue. Conjunctival pseudomembranes may also occur in such children (1).

Regional (mostly preauricular) lymph nodes are palpable in 80–90% of cases in the initial days of the conjunctivitis (1).

Early-stage keratoconjunctivitis in adults is characterized by crimson-red and edematous, velvety appearing swelling of the plica semilunaris and the caruncle (1).

Typically, epidemic keratoconjunctivitis (EKC) does not lead to recurrence. Conjunctival hemorrhages of different extents are not rare in adenovirus infection. They are, however, not pathognomonic, since the rarer pneumococcal conjunctivitis can also cause local hemorrhages. Important diagnostic tools that are recommended in the early stages include Giemsa staining of conjunctival scrapings and a finding of a lymphocytic inflammation (3).

Occasionally, the clinical symptoms of EKC may mask herpes conjunctivitis (3). Since no causal treatment is known so far, the symptoms should be alleviated by using artificial tears, eye drops with unspecific effects for humidification of the eye (3). W Manz (2) mentioned cold compresses to alleviate pain, among others. In order to prevent further transmission, the recommended hygiene measures, especially in household settings, include strict separation of washing utensils, such as soap or towels from those of other residents.

DOI: 10.3238/arztebl.2012.0148a

Poor-Quality Data

The authors report in their article that in 2010, the Robert Koch-Institute (RKI) registered an increase in notified adenovirus infections of 300% compared with previous years. They correctly point out that this is but the tip of an iceberg.

A survey conducted at regional continuing medical educational events found that none of the ophthalmologists present had ever notified an adenovirus infection to the health authorities or had tried to confirm the virus by using a test. The health insurers do not cover the costs of the available rapid test, and the test result is hardly of any therapeutic relevance (any therapeutic recommendations are off-label only). Notifications would therefore come mainly from eye hospitals and federal states where a suspected diagnosis is notifiable.

The increase derived from the RKI data (1) is even less credible when one looks further back into the past. For 2005, 525 cases of infection were notified; for 2006, 138 cases; for 2007, 375 cases; and for 2008, 168 cases. The increase thus obviously relates only to the 169 cases notified in 2009 compared with the 455 cases notified in 2010.

To demand in view of these rather poor data that, because of possible transmission by contaminated or insufficiently disinfected contact lenses, only proven virucidal disinfectants for fitted lenses should be permitted—which is impossible, because of a lack of suitable or approved solutions—seems rather daring (3). However, this is exactly what the responsible authority in Baden-Württemberg has imposed on ophthalmological practices, citing recommendations from the RKI (2). Whether this impractical obligation, and its monitoring, is also directed at ophthalmological opticians, who fit and distribute more than half of all contact lenses, is not known.

DOI: 10.3238/arztebl.2012.0148b

REFERENCES

- Schmidt D: Photophobie bei Augenkrankheiten des Kindes- und Jugendalters. Verlag Dr. Köster, Berlin 2008.
- Schmidt D: Wilhelm Johann Baptist Manz (1833–1911), erster Lehrstuhlinhaber der Univ.-Augenklinik Freiburg (1868). Mitteilungen der JULIUS-HIRSCHBERG-GESELLSCHAFT 2004; 6: 180–1.
- Sundmacher R: Viruskonjunktivitis. Fortschr Ophthalmol 1986; 83: 257–62.
- Meyer-Rüsenberg B, Loderstädt U, Richard G, Kaulfers PM, Gesser C: Epidemic Keratoconjunctivitis—the current situation and recommendations for prevention and treatment. Dtsch Arztebl Int 2011; 108(27): 475–80.

REFERENCES

- www.rki.de: Epidemiologisches Bulletin Nr. 50, Aktuelle Statistik medienpflichtiger Infektionskrankheiten, Stand 20.12.2010.
- www.rki.de: Ratgeber für Ärzte/Merkblätter – Keratoconjunctivitis epidemica und andere Konjunktividen durch Adenoviren, Stand 18.03.2010.
- www.iho.de: IHO – Viruzidieliste/manuelle Instrumentendesinfektion, Stand 21.11.2010.
- Meyer-Rüsenberg B, Loderstädt U, Richard G, Kaulfers PM, Gesser C: Epidemic Keratoconjunctivitis—the current situation and recommendations for prevention and treatment. Dtsch Arztebl Int 2011; 108(27): 475–80.

Dr. med. Wolfgang Werner
Stuttgart
Wwerner-Augenarzt@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Hand Hygiene—Other Remedies Are Also Effective

Under the subheading “Hygienic measures in clinical practice”, the authors rightly point out the crucial importance of proper hand hygiene. It is not correct, however, that exclusively “virucidal” disinfectants can be used. Rapid, 30-second effectiveness against adenoviruses is sufficient, as is possible by using one of several commercially available hand disinfectants (color free and scent free). Complete elimination of all viruses in keeping with the regulations of the Robert Koch-Institute is not required and does not make sense since the preparations used for that purpose are often far less well tolerated.

DOI: 10.3238/arztebl.2012.0149a

REFERENCES

1. Meyer-Rüsenberg B, Loderstädt U, Richard G, Kaulfers PM, Gesser C: Epidemic Keratoconjunctivitis—the current situation and recommendations for prevention and treatment. *Dtsch Arztebl Int* 2011; 108(27): 475–80.

Prof. Dr. med. Markus Dettenkofer

Department of Environmental Health Sciences, University Medical Center Freiburg, Germany, markus.dettenkofer@uniklinik-freiburg.de

Conflict of interest statement

Professor Dettenkofer has received funding for a research project he initiated from B Braun, Sempach, Ecolab, and Schulze and Mayr.

In Reply:

First of all, we wish to thank our correspondents for their letters and our many colleagues who have contacted us directly with their questions and suggestions. All this underlines the importance and topicality of the subject.

Professor Schmidt has highlighted several issues. The information about the slightly changed disease course in children, in whom general symptoms predominate, is important for general practitioners who are expected to make the diagnosis. We also thank him for summarizing the most important symptoms, such as swelling of the plica, caruncle, and preauricular lymph nodes. We mention these in the paragraph entitled “Clinical features.” It is correct that EKC typically does not lead to recurrences. In theory, however, renewed infection with another of the three serotypes mentioned in our article is entirely possible.

In view of today’s clinical or practice routines, we do not think that a Giemsa stained conjunctival scrape, such as probably would have been very useful in 1986, should be recommended, since polymerase chain reaction (PCR) provides a safe and reliable result within less than four hours (1).

We are also grateful for the summary of the most important hygienic measures and symptomatic therapeutic measures. These can be found in our article under the headings “Treatment” and “Hygienic measures in clinical practice.”

We share Dr Werner’s views regarding the estimated numbers of undetected cases of EKC. The Robert Koch-Institute itself reported 489 notified cases for 2010. We are not aware of any explicit recommendation from the RKI with respect to disinfecting contact lenses. The RKI’s advice for physicians says that the fact that pathogens of EKC can be transmitted by means of contaminated ophthalmological instruments underlines the importance of careful preparation or use of non-contact instruments (for example, tonometers), which should be the preferred option wherever possible. This is also true for tolerability of the materials in the relevant agents and procedures (2). We have not seen to the exact requirement from the Baden-Württemberg authorities and can therefore not comment.

With regard to Professor Dettenkofer’s correspondence, we wish to point out that selecting the appropriate hand disinfectant is worth discussing. The poorer skin tolerability of hand disinfectants classed as fully virucidal is important and should not be ignored. However, we agree with Dettenkofer that other hand disinfectants that have proven effectiveness against adenoviruses are also suitable.

DOI: 10.3238/arztebl.2012.0149b

REFERENCES

1. Doerr HW: Medizinische Virologie: Grundlagen, Diagnostik, Prävention und Therapie viraler Erkrankungen. 2nd edition. Stuttgart: Thieme 2010; 639ff
2. www.rki.de; Keratoconjunctivitis epidemica und andere Konjunktividen durch Adenoviren. RKI-Ratgeber für Ärzte, 3/20
3. Meyer-Rüsenberg B, Loderstädt U, Richard G, Kaulfers PM, Gesser C: Epidemic Keratoconjunctivitis—the current situation and recommendations for prevention and treatment. *Dtsch Arztebl Int* 2011; 108(27): 475–80.

Dr. med. Birthe Meyer-Rüsenberg

Ophthalmology Department, University Medical Center Hamburg-Eppendorf, Germany, b.meyer-ruesenberg@uke.de

Conflict of interest statement

The author declares that no conflict of interest exists.