Published in final edited form as:

Mol Microbiol. 2012 March ; 83(6): 1210–1228. doi:10.1111/j.1365-2958.2012.08003.x.

SlrA/SlrR/SinR inhibits motility gene expression upstream of a hypersensitive and hysteric switch at the level of σ^D in *Bacillus subtilis*

Loralyn M. Cozy¹, Andrew M. Phillips¹, Rebecca A. Calvo¹, Ashley R. Bate², Yi-Huang Hsueh¹, Richard Bonneau², Patrick Eichenberger², and Daniel B. Kearns^{1,*}

¹Indiana University, Department of Biology, Bloomington, IN 47408

²New York University, Center for Genomics and Systems Biology, Department of Biology, 12 Waverly Place, 8th floor, New York, NY 10003

Abstract

Exponentially growing *Bacillus subtilis* cultures are epigenetically differentiated into two subpopulations in which cells are either ON or OFF for σ^D -dependent gene expression: a pattern suggestive of bistability. The gene encoding σ^D , *sigD*, is part of the 31-gene *fla/che* operon where its location at the 3' end, 25 kb away from the strong *P_{fla/che}* promoter, determines its expression level relative to a threshold. Here we show that addition of a single extra copy of the *slrA* gene in the chromosome inhibited σ^D -dependent gene expression. SlrA together with SinR and SlrR reduced *sigD* transcript by potentiating a distance-dependent decrease in *fla/che* operon transcript abundance that was not mediated by changes in expression from the *P_{fla/che}* promoter. Consistent with acting upstream of σ^D , SlrA/SinR/SlrR could be bypassed by artificial ectopic expression of *sigD* that hysteretically maintained for 20 generations by engaging the *sigD* gene at the native locus. SlrA/SinR/SlrR was also bypassed by increasing *fla/che* transcription and resulted a hypersensitive output in flagellin expression. Thus, flagellin gene expression demonstrated hypersensitivity and hysteresis and we conclude that σ^D -dependent gene expression is bistable.

Keywords

SlrA; SigD; FlgM; bistability; hysteresis

INTRODUCTION

Development is the differentiation of clonal cells into one or more specialized cell types by epigenetic and self-sustaining patterns of gene expression. While obvious within the three dimensional tissues of multicellular eukaryotes, development in bacteria is difficult to detect as it commonly manifests as freely-mixing differentiated subpopulations. Molecular techniques with single cell resolution have revealed bacterial development in the form of heterogeneity at the level of gene expression in which cells exist in either ON or OFF states for a variety of phenotypes (Dubnau and Losick, 2006). The maintenance of two distinct epigenetic subpopulations has been attributed to a regulatory phenomenon known as bistability (Ferrell, 2002). Bistability arises from the stochastic variation of key genetic regulators relative to a critical threshold (Swain et al., 2002; Kaern et al., 2005). Hypersensitivity of the regulator creates a molecular hair-trigger such that the system sharply transitions to the ON state once a threshold is exceeded. Hysteresis reinforces state

*Corresponding author: dbkearns@indiana.edu, Phone: 1-812-856-2523, Fax: 1-812-855-6705.

acquisition and resists switching to the opposite state by engagement of positive or negative regulatory feedback loops (Smits et al., 2006). Whereas hysteresis makes bistable systems robust by maintaining state choice over many generations, hypersensitivity governs initial entrance of a cell to the ON or OFF state (Ferrell, 2002).

Growing populations of *Bacillus subtilis* bifurcate into ON and OFF cell types with respect to the expression of flagellar genes at the level of the alternative sigma factor, σ^D (Kearns and Losick, 2005). Cells that have high levels of σ^D are ON for the expression of genes encoding the flagellar filament protein (Hag) and cell-separating autolysins (LytC and LytF); consequently the cells are motile and separated from one another (Mirel and Chamberlin, 1989; Lazarevic et al., 1992; Margot et al., 1999; Kearns and Losick, 2005; Chen et al., 2009). Cells that have a low level of σ^D are OFF for σ^D -dependent genes and grow in long, non-motile chains (Helmann et al., 1988; Márquez et al., 1990; Cozy and Kearns, 2010). Differentiation of the two subpopulations depends on whether transcription of the gene encoding σ^D , *sigD*, exceeds a critical threshold that depends on the native location of *sigD* at the end of the 31-gene, 25 kb *fla/che* operon (Figure 1; Márquez-Magaña and Chamberlin, 1994; Cozy and Kearns, 2010; Lim et al., 2011). Transcription of the *fla/che* operon is primarily directed by a strong σ^A -dependent promoter ($P_{fla/che}$) with a weaker, auxiliary σ^D -dependent promoter ($P_{D.3}$) upstream (Estacio et al., 1998; West et al., 2000). Isolation of ON and OFF subpopulations of a SwrA mutant (see below) revealed that both cell types experienced a gradual decrease in *fla/che* operon transcript abundance from the front to the back of the operon, but that ON cells had a greater amount of *fla/che* transcript initially (Cozy and Kearns, 2010). Thus, whether *sigD* expression exceeded a threshold appeared to depend on the combination of transcription magnitude of the *fla/che* operon, the gradual decrease in transcript abundance along the operon, and the location of the *sigD* gene.

A variety of proteins control σ^D -dependent gene expression (Figure 1). The activity of σ^D is held in check by the anti-sigma factor FlgM, and FlgM antagonism is relieved upon assembly of flagellar hook/basal body, encoded by genes forward in the *fla/che* operon (Hughes et al., 1993; Kutsukake, 1994; Barillà et al., 1994; Caramori et al., 1996). Furthermore, some genes activated by σ^D , such as those encoding flagellin and cell separating autolysins, are repressed by a heterodimer of two DNA binding proteins called SlrR and SinR (Chai et al., 2010a; Chai et al., 2010b). Finally, two poorly understood proteins SwrA and SwrB increase the number of cells ON for σ^D -dependent gene expression. SwrA is thought to act upstream of σ^D by activating expression from the $P_{fla/che}$ promoter and SwrB is thought to act downstream of σ^D by enhancing σ^D activity via an unknown mechanism (Werhane et al., 2004; Kearns and Losick, 2005). When both SwrA and SwrB are mutated, nearly all cells in the population are turned OFF (Kearns and Losick, 2005). How all of the various regulators combine to produce a biased population remains unclear.

The simultaneous maintenance of ON cells and OFF cells in a population is indicative of, but does not definitively support, the mechanism of bistability. Here we find that providing a single extra copy of the gene encoding SlrA, a small protein antagonist of SinR, abolishes expression of the entire σ^D regulon. We further show that the extra copy effect of SlrA requires SinR and SlrR and acts upstream of the *sigD* gene to inhibit *fla/che* transcript abundance after transcription initiation and that SlrA/SinR/SlrR may be bypassed by artificial *sigD* expression. We then use strains expressing an extra copy of *slrA* as tools to demonstrate that σ^D -dependent gene expression is both hypersensitive and hysteretic. We conclude that the switch governing the motile/non-motile state choice is bistable under the control of the σ^D alternative sigma factor.

RESULTS

SlrA inhibits flagellin expression by inhibiting σ^D protein accumulation

The motility alternative sigma factor, σ^D , is required for the expression of a regulon of genes including genes encoding late flagellum biosynthesis proteins and peptidoglycan-remodeling, cell-separating enzymes called autolysins (Márquez et al., 1990). The activity of σ^D can be monitored with the σ^D -dependent reporter of flagellin (*hag*) expression, *P_{hag}-lacZ*. Whereas wild type colonies with *P_{hag}-lacZ* were blue when grown on media containing X-gal, colonies simultaneously mutated for *swrA* and *swrB*, encoding the activators of σ^D -dependent gene expression SwrA and SwrB respectively, appeared pale blue to white (Fig. 2A) (Kearns and Losick, 2005). To find other regulators of σ^D -dependent gene expression, the *swrA swrB* double mutant background was mutagenized with transposons and screened for insertions that resulted in enhanced blue colony color. Fifteen of seventeen transposons that enhanced blue colony color disrupted either the DegS/DegU two-component system which activates the *P_{flgM}* promoter and expresses the anti- σ^D anti-sigma factor FlgM, or separated the *P_{flgM}* promoter from the DegU-phosphate binding site (Figure 1; Hsueh et al., 2011). The final two transposon insertions that restored blue colony color were found in the open reading frame, *slrA*, encoding SlrA, a small protein antagonist of the DNA binding protein and master regulator of biofilm formation, SinR (Figure 2A, 2B; Kobayashi, 2008; Chai et al., 2009). Despite the fact that mutation of *slrA* restored blue colony color to the *swrA swrB* double mutant on solid media containing X-gal, no increase in log phase β -galactosidase activity was observed in liquid media in the same genetic background (Fig. 2A). Expression of σ^D -dependent genes is heterogeneous in log phase cultures of *B. subtilis* however, and heterogeneity can obscure subpopulation level effects on gene expression (Kearns and Losick, 2005). To assess the effect of an *slrA* mutation in the *swrA swrB* background at the individual cell level, a fluorescent reporter for σ^D -dependent gene expression, *P_{hag}-GFP*, was integrated at an ectopic location in the chromosome (*lacA::P_{hag}-GFP*). Wild type populations were heterogeneous for *P_{hag}-GFP* expression, whereas simultaneous mutation of *swrA* and *swrB* resulted in a population that failed to express σ^D -dependent genes and grew as chains (Figure 3A, 3B). Mutation of *slrA* in the *swrA swrB* double mutant background also failed to reduce cell chaining and did not increase *P_{hag}-GFP* expression (Fig. 3C). Thus, in liquid-based assays, neither β -galactosidase enzymatic measurements nor GFP fluorescence microscopy explained the enhanced blue colony color observed in the original screen. We infer that the effect of the *slrA* mutation in a *swrA swrB* mutant background was either subtle, or had maximal effect at timepoints in the *B. subtilis* growth curve beyond our limited measurements at mid-log phase.

To further explore the reason that mutation of *slrA* appeared to restore *P_{hag}-lacZ* expression to the *swrA swrB* double mutant, *slrA* was mutated in backgrounds separately mutated for either *swrA* or *swrB* alone. Mutation of *swrB* decreased the number of cells in the population that expressed *P_{hag}-GFP* compared to wild type, and simultaneous mutation of *swrB* and *slrA* resulted in a population that resembled *swrB* alone (Fig 3D, 3E). Mutation of *swrA* decreased the number of cells in the population that expressed *P_{hag}-GFP* as well, but simultaneous mutation of *swrA* and *slrA* resulted in a population that resembled the wild type (Fig 3F, 3G). Thus, mutation of *slrA* increased the number of cells in the population that express *P_{hag}-GFP* in the absence of SwrA but not SwrB. We infer that the original screen to bypass the *swrA swrB* double mutant was stringent but sufficiently sensitive to detect the effect of mutating *slrA* on bypassing the absence of SwrA alone. We conclude that SlrA is an inhibitor of σ^D -dependent gene expression. We infer that SlrA acts in the same pathway as SwrA, and like SwrA, SlrA may act upstream of *sigD* expression.

To confirm that mutation of *slrA* was necessary for the increased frequency of σ^D -dependent gene expression, we genetically complemented the *slrA* mutation. Expression of *slrA* from its native promoter at the ectopic *amyE* site (*slrA*⁺) abolished expression of *P_{hag}-GFP* in *swrA* mutant background (Fig 3H) and reduced expression of *P_{hag}-lacZ* reporter relative to the *swrA swrB slrA* triple mutant (Fig 2A). In each case however, introduction of the (*slrA*⁺) complementation construct appeared to have a dominant effect as it reduced expression of the *P_{hag}* reporters below that of the either the *swrA* single mutant or the *swrA swrB* double mutant parental strains. Furthermore, introduction of the (*slrA*⁺) complementation construct as an extra copy into an otherwise wild type background dramatically reduced *P_{hag}-lacZ* reporter activity to render colonies white on media containing X-gal and abolish β -galactosidase activity in liquid culture (Fig. 2A). Cytologically, an extra copy of *slrA* also abolished *P_{hag}-GFP* expression and made the population grow constitutively as long chains that often formed braids (Fig. 3I, Fig. 4). We conclude that inhibition by SlrA is potent because a single extra copy of *slrA* in the chromosome was sufficient to dramatically diminish σ^D -dependent gene expression in wild type cells.

One way in which the single extra copy of *slrA* could cause reduced σ^D -dependent gene expression is through reduced σ^D protein accumulation. To assess σ^D protein levels, protein from whole cell lysates of wild type, a strain mutant for *sigD*, and a strain containing an extra copy of *slrA* were separated by SDS-PAGE and probed using anti- σ^D and anti- σ^A antibodies. Whereas the vegetative sigma factor, σ^A , was constant in all samples, σ^D protein accumulation was dramatically reduced in both the *sigD* mutant and *slrA* extra copy backgrounds (Fig. 5A). We conclude that one way in which SlrA inhibits σ^D -dependent gene expression is by reducing σ^D protein levels.

SlrA inhibits σ^D protein accumulation by inhibiting *fla/che* operon transcript levels through SinR/SlrR

To test if SlrA inhibited σ^D protein accumulation at the level of *sigD* gene expression, we created a strain background carrying three fluorescent reporters at three levels of the flagellar regulatory hierarchy (Figure 1). To monitor the 5' end of the *fla/che* operon, a transcriptional fusion between *P_{D-3P_{fla/che}}* promoter and the gene encoding red fluorescent protein (mCherry) was integrated at the ectopic *thrC* locus (*thrC::P_{D-3P_{fla/che}}*-mCherry). To monitor the 3' end of the *fla/che* operon where the *sigD* gene is located, the gene encoding cyan fluorescent protein (CFP) was integrated as a transcriptional fusion at the native site after the last gene of the operon (*fla/che Ω CFP*). To monitor σ^D -dependent gene expression, a transcriptional fusion between *P_{hag}* and the gene encoding yellow fluorescent protein (YFP) was integrated at the ectopic *lacA* locus (*lacA::P_{hag}-YFP*). Wild type cells uniformly expressed the *P_{D-3P_{fla/che}}* reporter, but heterogeneously expressed reporters for the 3' end of the operon and the σ^D -dependent *P_{hag}* reporter (Fig. 5B, Cozy and Kearns, 2010). When a single extra copy of *slrA* was introduced into this background, *P_{D-3P_{fla/che}}* expression was not reduced, but expression of the 3' end of the operon as well as *P_{hag}* were reduced below the level of detection (Figure 5B, *slrA*⁺). We conclude that SlrA lowers σ^D protein levels by decreasing *sigD* gene expression.

An extra copy of *slrA* appeared to reduce *sigD* gene expression within the *fla/che* operon downstream of the *P_{fla/che}* promoter. To determine where within the *fla/che* operon inhibition by SlrA was occurring, total RNA was purified from wild type cells and from cells with an extra copy of *slrA*, and quantitative reverse transcriptase PCR (qRT-PCR) was performed using primers specific to fifteen different genes along the operon. Cells with an extra copy of *slrA* showed reduced transcript abundance throughout the operon relative to wild type that was dependent on the distance from the *P_{fla/che}* promoter and culminated in a 20-fold reduction at the *sigD* gene. (Figure 5C). Consistent with a decrease in *fla/che* operon transcript levels, cells containing an extra copy of *slrA* also showed reduced levels of two

proteins encoded within the operon, FliG and FliY (Fig. 5A). It is noteworthy that the effect of *slrA* extra copy manifested as far forward in the operon as the first gene *flgB* which experienced a 5-fold decrease in transcript abundance (Figure 5C). The inhibitory effect of SlrA was not observed at the level of initiation at the *P_{fla/che}* promoter as an extra copy of the *slrA* gene in the chromosome had no discernable effect on the expression of an ectopically integrated *P_{fla/che}-lacZ* reporter fusion (Fig 5D). We conclude that SlrA inhibits transcript abundance of the entire *fla/che* operon following transcription initiation and in a manner dependent on the distance from the *P_{fla/che}* promoter. We further conclude that the decrease in *fla/che* transcript abundance accounts for the low levels *sigD* transcript, the low levels of σ^D protein, and inhibition of σ^D -dependent gene expression.

To search for genes under control of SlrA that might account for the decrease in *fla/che* operon transcript abundance, transcriptional profiling was conducted. RNA was purified from growing cells of wild type and from growing cells of a strain containing an extra copy of *slrA*, differentially labeled, and compared on an oligo-based genome-wide DNA microarray. Consistent with previous reports, SlrA not only inhibited expression of certain σ^D -dependent genes but appeared to inhibit the entire σ^D regulon and, as suggested above, was a potent inhibitor of the *fla/che* operon (Table 1, Table S1) (Kobayashi, 2008). Also consistent with previous reports, SlrA activated genes required for biofilm formation (Table 2, Table S1) (Kobayashi, 2008; Chai et al., 2009). We hypothesized that SlrA might diminish *fla/che* transcript abundance downstream of promoter initiation by activation of RNA turnover enzymes, activation of an RNA polymerase termination factor, or inhibition of an RNA polymerase elongation factor (Roberts et al., 2008; Silva et al., 2011). No proteins overtly associated with RNA stability or transcript elongation appeared to be under the control of SlrA. Perhaps, SlrA regulates RNA elongation/turnover at the post-transcriptional level or the factor that reduces *fla/che* operon transcript abundance may be encoded among the various proteins and small RNAs of unknown function under SlrA control.

To find components required to inhibit *fla/che* transcript levels that were epistatic to SlrA, a strain containing an extra copy of *slrA* and a *P_{hag}-lacZ* reporter that normally grows as white colonies on media containing X-gal was mutagenized with transposons and screened for colonies with increased blue colony color. Insertion in either the native or ectopic copy of *slrA* restored reporter expression to wild type levels (Fig 2A). In addition, insertions in *remA*, *remB*, *sinR* and *slrR* genes all restored reporter expression to a level equal to or greater than wild type (Fig 2A, Table 3). Mutations that bypassed the extra copy of *slrA* seemed to have in common a relationship to the transcription factor SlrR: SlrR was directly mutated in the screen, RemA and RemB both activate expression of the SlrR protein, and SinR forms a co-repressing heterodimer with SlrR to inhibit expression of autolysin genes (Figure 1) (Winkelman et al., 2009; Chai et al., 2010a). We focused our work on the epistasis between SlrA and SlrR because a relationship between SlrA, SinR, and SlrR has been previously reported and we inferred that SlrR was the most downstream component of all the genes identified in the bypass screen (Kobayashi, 2008; Chai et al., 2009).

To explore the epistatic relationship between *slrA* and *slrR* with regards to *fla/che* operon expression, an *slrR* mutation was introduced into a background with an extra copy of *slrA* and *fla/che* operon expression was examined cytologically using a strain containing three fluorophores in the motility hierarchy. Mutation of *slrR* restored expression of the *CFP* reporter integrated at the 3' end of the *fla/che* operon as well as the σ^D -dependent *P_{hag}-YFP* reporter (Fig. 5B). Furthermore, Western blot analysis revealed that mutation of *slrR* in the *slrA* extra-copy background restored protein accumulation of FliG, FliY, σ^D and flagellin (Hag) to near wild type levels (Fig. 5A). Consistent with the cytological and protein accumulation observations, quantitative measurement of *fla/che* operon transcript abundance

showed that mutation of *slrR* restored the *fla/che* operon expression to levels comparable to wild type (Fig. 5C). Although mutation of *slrR* restored *fla/che* operon transcript levels in the presence of an extra copy of *slrA*, no change in expression was detected using promoter fusions of $P_{D-3P_{fla/che}}$ to either mCherry or *lacZ* (Fig. 5B, 5D). We conclude that SlrA/SinR/SlrR inhibits *fla/che* operon transcript levels after the initiation of transcription from the $P_{D-3P_{fla/che}}$ promoter. Furthermore, we infer that SlrA/SinR/SlrR inhibits σ^D -dependent gene expression at two levels, by inhibiting *sigD* transcription post-initiation and by inhibiting the assembly of flagellar hook/basal body complexes (encoded within the *fla/che* operon) that antagonize the σ^D -antagonist FlgM (Figure 1).

σ^D is hysteretic for the activation of P_{hag} expression

All our observations suggested that SlrA inhibited P_{hag} expression upstream of σ^D protein production by reducing *fla/che* transcript levels. We wondered whether artificial expression of *sigD* was sufficient to bypass the inhibitory effect of *slrA*. An artificial expression construct was generated that fused *sigD* expression to the IPTG-inducible $P_{hyspank}$ promoter and was inserted at an ectopic locus (*thrC::P_{hyspank}-sigD*) of a strain containing an extra copy of *slrA* (*amyE::P_{slrA}-slrA*). In the absence of *sigD* induction, P_{hag} -YFP expression remained OFF and cells grew in long chains (Fig 6A). Induction of the *sigD* artificial expression construct restored YFP expression to a subpopulation and cells grew in short chains or single cells (Fig. 6B). Expression was restricted to a subpopulation because of the anti-sigma factor FlgM that inhibits σ^D activity. Mutation of *flgM* restored P_{hag} expression to all cells in the population whether or not inducer was added (Fig. 6C, 6D). We conclude that artificial expression of *sigD* bypassed the inhibitory effect of SlrA on P_{hag} expression.

Hysteresis is the epigenetic maintenance of a cell's regulatory state over many generations in the absence of a stimulus. We wondered whether we could use the fact that artificial expression of *sigD* could restore P_{hag} expression in the presence of an extra copy of *slrA* to test for hysteresis. To build a strain suitable to detect hysteresis of the OFF to ON transition, we created a *sigD* artificial expression construct with enhanced dependence on IPTG induction by mutating the *sigD* ribosome binding site away from consensus (*thrC::P_{hyspank}-^{crRBS}sigD*). The RBS-crippled ^{crRBS}*sigD* expression construct was introduced to a strain containing an extra copy of *slrA* (to increase the number of cells in the P_{hag} OFF state), a mutation in *flgM* (to eliminate regulation at the level of σ^D activity), and a P_{hag} -YFP reporter construct (to detect σ^D -dependent gene expression). The resulting strain was sensitive to artificial *sigD* induction as the population was largely OFF for P_{hag} -YFP expression in the absence of IPTG and all ON for P_{hag} -YFP expression in the presence of 1 mM IPTG (Fig. 7A).

To assay for hysteresis, cells grown in the presence of IPTG were washed, serially diluted in the absence of IPTG, and allowed to grow such that the cells would return to our standard measurement condition of 0.5 OD₆₀₀ after a defined number of generations. Cells that had acquired the flagellin ON state remained ON for at least 20 generations after inducer had been removed (Fig. 7A, "WT"). Furthermore, the maintenance of the ON state was dependent on the *sigD* gene at the native location, as mutation of native *sigD* resulted in loss of P_{hag} -YFP fluorescence within four generations after inducer removal (Fig 7A, Δ *sigD*). σ^D -dependent hysteresis of P_{hag} -YFP was also found to occur in strains mutated for *swrA* (Figure S1A) and in a subpopulation of a strain wild type for *flgM* (Figure S1B). Direct induction of a fluorescent reporter alone from a $P_{hyspank}$ promoter ($P_{hyspank}$ -GFP) did not display hysteresis (Figure S2). Finally, consistent with SlrA/SlrR/SinR acting upstream of σ^D , hysteretic activation of σ^D did not change the levels of either the SlrR or SinR proteins (Figure 7B). We conclude that activation of σ^D was hysteretic and that hysteresis was maintained by expression of the *sigD* gene within the *fla/che* operon after artificial induction of the ectopic construct was removed.

The expression of *sigD* depends on the *fla/che* promoter region. We hypothesized that deletion of the *fla/che* promoter region ($P_{D-3}P_{fla/che}$) would mimic deletion of the *sigD* gene with respect to hysteresis. To test the promoter requirement, we deleted a region encompassing the P_{D-3} promoter, the $P_{fla/che}$ promoter, and the DNA between them in the “WT” genetic background from Figure 7A. Upon induction of *sigD*, the population grew exclusively in the ON state, however after removal of inducer, outgrowth in media lacking IPTG revealed that the σ^D state of the population quickly reverted to the uninduced (OFF) phenotype ($\Delta P_{D-3}P_{fla/che}$, Fig 7A). In contrast, when only the σ^D -dependent P_{D-3} promoter was deleted, the population exhibited hysteresis comparable to that of “WT” (ΔP_{D-3} , Fig. 7A). We conclude that the $P_{fla/che}$ but not the P_{D-3} promoter was required for ON-state maintenance and that $P_{fla/che}$ is likely needed to provide a high level of operon expression such that *sigD* transcript levels rest above a threshold.

Activation of P_{hag} expression is hypersensitive to σ^D

While hysteresis stabilizes cells in one regulatory state or another, hypersensitivity provides for state acquisition in bistable systems (Ferrell, 2002). Hypersensitivity invokes a non-linear or sigmoidal output in response to linear increase in stimulus. To test for hypersensitivity, a strain was generated in which all cells began in the OFF state for σ^D -dependent gene expression and *sigD* was artificially induced. First, the strain contained an extra copy of *slrA* integrated at the ectopic *amyE* locus (*amyE::P_{slrA}-slrA*). Second, the strain was mutated for *flgM* to remove activity level regulation of σ^D (*ΔflgM*). Third, the native $P_{fla/che}$ promoter was replaced by an IPTG-inducible $P_{hyspank}$ promoter such that expression of the *fla/che* operon and therefore *sigD* expression could be controlled by linear variation in IPTG concentration ($\Omega P_{hyspank-fla/che}$). Fourth, a P_{hag} -GFP reporter was introduced at the ectopic *lacA* locus so that σ^D -dependent gene expression could be measured cytologically (*lacA::P_{hag}-GFP*).

In the absence of IPTG, the engineered strain for the hypersensitivity experiment was all OFF for P_{hag} expression. When grown in the presence of low levels of IPTG, cells induced P_{hag} expression heterogeneously and population level expression became homogeneously ON at high levels of IPTG (Figure 8A). GFP intensity was measured by pixel counting of individual cells at each level of IPTG induction and the population wide behavior was represented by the median fluorescence intensity (Figure 8C). We found that as the IPTG concentration increased linearly, σ^D -dependent gene expression output displayed a sigmoidal response with a Hill constant of 3.6 as derived by fitting the data to a standard binding curve (van Holde et al., 1998). As a control, the median fluorescence intensity of cells expressing GFP directly from an IPTG-inducible $P_{hyspank}$ promoter (*amyE::P_{hyspank}-GFP*) was largely linear with a Hill Constant of 1.5 (Figure 8B, 8C). We conclude that σ^D -dependent gene expression exhibits a nonlinear response to linear *fla/che* operon induction and we infer that σ^D displays hypersensitivity.

DISCUSSION

Fluorescence microscopy and single cell analysis have revealed remarkable heterogeneity at the level of gene expression in bacterial populations in which particular regulons are either ON or OFF in individuals. Population heterogeneity has been found for a variety of conditionally-advantageous, time-sensitive phenotypes including competence for DNA uptake, sporulation, biofilm formation, persistence to antibiotic treatment, nutrient uptake, and virulence factor synthesis (Mamaar et al., 2007; Chai et al., 2008; Veening et al., 2008; Siegele and Hu, 1997; Tolker-Neilsen et al., 1998; Rietsch and Mekalanos, 2006; Nakata et al., 2009; Balaban et al., 2004). Simultaneous maintenance of specialized cell types primes a subpopulation to immediately exploit a selective advantage in a harsh and fluctuating environment (Kussell and Leibler, 2005; Veening et al., 2008). One explanation for ON/

OFF bifurcated subpopulations is the epigenetic phenomenon of bistability, which invokes a gene expression switch with hypersensitivity to govern state acquisition relative to a threshold, and hysteresis to govern state stability often by engaging regulatory feedback loops (Ferrell, 2002). Here we study the heterogeneous expression of motility genes in *Bacillus subtilis* and demonstrate hypersensitivity and hysteresis in the system.

Motility in *B. subtilis* requires over 30 proteins to assemble flagella. Many proteins for early flagellar assembly are encoded in the massive 25 kb, 31 gene *fla/che* operon. The second to last gene in the *fla/che* operon, *sigD*, encodes σ^D an alternative sigma factor that directs RNA polymerase to express a regulon of proteins including late flagellar assembly proteins and autolysin enzymes involved in daughter cell separation after cell division. Whereas a only minority of cells in a wild type population is OFF for σ^D -dependent gene expression, the number of OFF cells increases dramatically in cells mutated for *SwrA*, a protein that appears to activate the *P_{fla/che}* promoter of the *fla/che* operon (Kearns and Losick, 2005). In previous work, ON and OFF subpopulations were isolated from a *swrA* mutant and separately analyzed for *fla/che* operon expression (Cozy and Kearns, 2010). Both cell types experienced a decrease in *fla/che* operon expression that depended on the distance from the *P_{fla/che}* promoter, and variations in *fla/che* transcript levels caused the *sigD* gene to be expressed either above or below a threshold. Here we show that introducing an extra copy of the gene *slrA* into the chromosome caused a distance-dependent decrease in *fla/che* transcript levels reminiscent of the *swrA* mutant. Furthermore, a cell doubly mutated for *swrA* and *slrA* restored the σ^D ON cells to wild type frequency, and we infer that the distance-dependent decrease in *fla/che* transcript originally observed in a *swrA* mutant was likely due to the action of SlrA.

SlrA function has been shown to be mediated through the paralogous DNA binding proteins SinR/SlrR (Kobayashi, 2008; Chai et al., 2009; Chai et al., 2010a). Consistent with this genetic organization, mutation of *slrR* was epistatic to an extra copy of *slrA* in all experiments we performed. SlrA interacts with SinR to derepress SlrR expression; SlrR in turn forms a heterodimer with SinR that acts downstream of σ^D and directly represses two of the vegetative autolysins (LytC and LytF) and flagellin (Hag) (Figure 1; Chai et al., 2010a). Here we show that SlrA/SinR/SlrR also act upstream of σ^D by inhibiting expression of the entire σ^D regulon by diminishing *fla/che* operon transcript levels and may be bypassed by artificial *sigD* expression (Figure 1, Table 1). SlrA/SinR/SlrR reduce *fla/che* operon transcript levels as early as the first gene in the *fla/che* operon but the effect does not appear to be mediated at the level of transcript initiation at the *P_{fla/che}* promoter. We infer that the effect must occur post-initiation and may involve control of a transcription elongation factor, a transcriptional terminator, or RNA turnover. No genes overtly associated with RNA management, however, were found to be under *slrA* control by transcriptome analysis or by a forward genetic bypass screen. The mechanism of the SlrA mediated *fla/che* operon transcript decrease is unknown but may be mediated by SlrA-transcriptionally-regulated proteins of unknown function or by essential proteins that are regulated at the functional level.

SlrA not only diminishes σ^D levels but also indirectly inhibits σ^D activity. Early inhibition of the *fla/che* transcript by SlrA reduced the amount of basal body proteins synthesized by the cell. Fully assembled basal bodies antagonize the FlgM anti-sigma factor that binds to σ^D and inhibits σ^D activity (Barillà et al., 1994; Caramori et al., 1996; Bertero et al., 1999). Thus, SlrA restricts basal body assembly and releases FlgM from its antagonist. By introducing a *flgM* mutation to eliminate the contribution of activity-level regulation on σ^D , we were able to bypass SlrA by artificial expression of the *sigD* gene integrated at ectopic site in the chromosome and demonstrate hysteresis in the system. When artificial induction of *sigD* was removed, *P_{hag}* expression remained in the ON state for over 20 generations that

depended on the *sigD* gene at the native locus. Hysteresis required the $P_{fla/che}$ promoter presumably to set a high level of *fla/che* transcription and express the native *sigD* gene. We hypothesize that the ON state was maintained by positive feedback at one or more σ^D -dependent promoters internal to the *fla/che* operon (West et al., 2000; Cozy and Kearns, 2010). By replacing the $P_{fla/che}$ promoter with an artificial IPTG inducible promoter, we were able to vary the amount of inducer linearly and show sigmoidal output in P_{hag} expression indicative of hypersensitivity in the system. Hypersensitivity often invokes cooperative protein-protein interactions that are difficult to explain in the context of a sigma factor. σ^D is unusual among the sigma factors, however, in that it binds to DNA in the absence of core RNA polymerase, and generates supershifted complexes that may indicate more than one protein bound at the promoter (Chen and Helmann, 1995; Bertero et al., 1999; Sevim et al., 2011).

We propose a model in which ON/OFF motility gene expression is governed by the amount of *fla/che* operon expression which determines the probability that *sigD* is transcribed to set σ^D protein levels relative to a threshold. SlrA/SinR/SlrR imposes resistance on the system upstream of *sigD* by reducing *fla/che* transcript levels and thus the probability of transcript completion, which in turn favors the acquisition of an OFF state. SlrA/SinR/SlrR may reinforce the OFF state by reducing basal body antagonism of FlgM, and by directly repressing certain genes under σ^D control. For example, direct repression of LytC may be particularly important for cell chaining as it is expressed from both σ^A -dependent and σ^D -dependent promoters (Lazarevic et al., 1992). SwrA, on the other hand, activates expression from the $P_{fla/che}$ promoter to increase *fla/che* operon transcript and increase the probability that σ^D protein exceeds a threshold. Hypersensitivity in σ^D output, perhaps by nucleation of σ^D protein at promoters, overrides SlrA/SinR/SlrR activity and provides for a sharp transition to the ON state. Once ON, σ^D hysteretically maintains its own expression to stabilize cells as single motile individuals. While the mechanisms that contribute to the observed hypersensitivity and hysteresis remain unknown, both properties support a model in which motility gene expression is bistable under the control of σ^D .

EXPERIMENTAL PROCEDURES

Strains and growth conditions

B. subtilis PY79, 3610 and their derivatives were grown at 37°C in Luria-Bertani (LB) (10g of tryptone, 5 g of yeast extract, 5 g of NaCl per liter) broth or LB plates supplemented with 1.5% Bacto agar. When appropriate, antibiotics were included at the following concentrations: 10 µg/mL tetracycline, 100 µg/mL spectinomycin, 5 µg/mL chloramphenicol, 5 µg/mL kanamycin and 1 µg/mL erythromycin plus 25 µg/mL lincomycin (MLS). Isopropyl β-D-thiogalactopyranoside (IPTG, Sigma) was added to liquid medium at indicated concentration when appropriate.

Strain construction

All constructs were first introduced into the domesticated strain PY79 by natural competence and then transferred to the 3610 background using SPP1-mediated generalized phage transduction (Yasbin and Young, 1974). All strains used in this study are listed in Table 4. All plasmids used in this study are listed in Supplemental Table S2. All primers used in this study are listed in Supplemental Table S3.

In-frame deletions—To generate the $\Delta sigD$ in frame marker-less deletion construct, the region upstream of *sigD* was PCR amplified using the primer pair 2019/2020 and digested with *EcoRI* and *XhoI*, and the region downstream of *sigD* was PCR amplified using the primer pair 2021/2022 and digested with *XhoI* and *BamHI*. The two fragments were then

simultaneously ligated into the *EcoRI* and *BamHI* sites of pMiniMAD which carries a temperature sensitive origin of replication and an erythromycin resistance cassette to generate pDP326 (Patrick and Kearns, 2008). The plasmid pDP326 was introduced to PY79 by single cross-over integration by transformation at the restrictive temperature for plasmid replication (37°C) using *mls* resistance as a selection. The integrated plasmid was then transduced into 3610. To evict the plasmid, the strain was incubated in 3ml LB broth at a permissive temperature for plasmid replication (22°C) for 14 hours, diluted 30-fold in fresh LB broth, and incubated at 22°C for another 8 hours. Dilution and outgrowth was repeated 2 more times. Cells were then serially diluted and plated on LB agar at 37°C. Individual colonies were patched on LB plates and LB plates containing *mls* to identify *mls* sensitive colonies that had evicted the plasmid. Chromosomal DNA from colonies that had excised the plasmid was purified and screened by PCR using primers 2019/2022 to determine which isolate had retained the *ΔsigD* allele.

To generate the *ΔslrR* in frame marker-less deletion construct, the region upstream of *slrR* was PCR amplified using the primer pair 1965/1966 and digested with *EcoRI* and *XhoI*, and the region downstream of *slrR* was PCR amplified using the primer pair 1967/1968 and digested with *XhoI* and *BamHI*. The two fragments were then simultaneously ligated into the *EcoRI* and *BamHI* sites of pMiniMAD which carries a temperature sensitive origin of replication and an erythromycin resistance cassette to generate pDP321. pDP321 was integrated into *B. subtilis* genome and evicted as described previously. Colonies were screened by PCR using primers 1965/1968 to determine which isolate had retained the *ΔslrR* allele.

To generate the $\Delta P_{D-3P_{fla/che}}$ marker-less deletion construct, the region upstream of $\Delta P_{D-3P_{fla/che}}$ was PCR amplified using the primer pair 1648/1649 and digested with *EcoRI* and *XhoI*, and the region downstream of $\Delta P_{D-3P_{fla/che}}$ was PCR amplified using the primer pair 2315/2316 and digested with *XhoI* and *BamHI*. The two fragments were then simultaneously ligated into the *EcoRI* and *BamHI* sites of pMiniMAD which carries a temperature sensitive origin of replication and an erythromycin resistance cassette to generate pDP310. pDP310 was integrated into *B. subtilis* genome and evicted as described previously. Colonies were screened by PCR using primers 1648/2316 and digesting the PCR product with *XhoI* to determine which isolate had retained the $\Delta P_{D-3P_{fla/che}}$ allele.

SUMO-FliG fusion protein expression vector—To generate the translational fusion of FliG to the SUMO his tag, a fragment containing *fliG* was amplified using 3610 as a template and the primer pair 882/883 and was digested with *SapI* and *XhoI*. The fragment was ligated into the *SapI* and *XhoI* sites of pTB146 containing ampicillin resistance cassette to create pKB43.

SUMO-FliY fusion protein expression vector—To generate the translational fusion of FliY to the SUMO his tag, a fragment containing *fliY* was amplified using 3610 as a template and the primer pair 1295/1296 and was digested with *SapI* and *XhoI*. The fragment was ligated into the *SapI* and *XhoI* sites of pTB146 containing ampicillin resistance cassette to create pDP288.

6His-Hag fusion protein expression vector—To generate the translational fusion of Hag to the 6His tag, a fragment containing *hag* was amplified using 3610 as a template and the primer pair 663/664 and was digested with *NheI* and *XhoI*. The fragment was ligated into the *NheI* and *XhoI* sites of pET28A containing kanamycin resistance cassette to create pJP4.

SlrA Complementation—The *slrA* region was amplified using 3610 genomic DNA as template and the primer pair 540/541 and was digested with *EcoRI* and *HindIII*. The

fragment was ligated into the *EcoRI/HindIII* sites of pDG364 containing a chloramphenicol resistance cassette to create pKB8.

IPTG inducible *sigD* constructs—The *sigD* gene was amplified using 3610 genomic DNA as template and the primer pair 374/375 and was digested with *SphI* and *NheI*. The fragment was ligated into the *SphI/NheI* sites of pLC135 containing a spectinomycin resistance cassette to create pLC136.

To generate the inducible *sigD* construct with a crippled ribosome binding site, the *sigD* gene was amplified using 3610 genomic DNA as template and the primer pair 1628/1973 and was digested with *SphI* and *NheI*. The fragment was ligated into the *SphI/NheI* sites of pDR111 containing a spectinomycin resistance cassette to create pKB144. Primer 1628 replaced sequence immediately upstream of the *sigD* start codon with the “AAAGGAAAAAAGCG” altered ribosome binding site sequence from the mutant *epsE* allele *loc2* (Guttenplan, 2010).

IPTG inducible *GFP*—The *GFP* gene was amplified using pDP155 plasmid DNA as template and the primer pair 378/1528 and was digested with *NheI* and *SphI*. The fragment was ligated into the *NheI/SphI* sites of pDP150 containing a erythromycin resistance cassette to create pAP11.

YFP transcriptional reporters—The *P_{hag}-YFP* fragment from pDP193 was amplified using the primer pair 380/563 and digested with *EcoRI* and *BamHI*. The fragment was ligated into the *EcoRI/BamHI* sites of pDR183 containing a mls resistance cassette to create pLC19. Alternatively, the *P_{hag}-YFP* fragment from pDP193 was digested with *EcoRI* and *BamHI* and ligated into the *EcoRI/BamHI* sites of pNC018 containing a tetracycline resistance cassette to create pLC20.

SPP1 phage transduction

To 0.2 ml of dense culture grown in TY broth (LB broth supplemented after autoclaving with 10 mM MgSO₄ and 100 μM MnSO₄), serial dilutions of SPP1 phage stock were added and statically incubated for 15 minutes at 37°C. To each mixture, 3 ml molten TY supplemented with 0.5% agar was added, poured atop fresh TY plates, and incubated at 37°C overnight. Top agar from the plate containing near confluent plaques was harvested by scraping into a 50 ml conical tube, vortexed, and centrifuged at 5,000 × g for 10 minutes. The supernatant was treated with 25 μg/ml DNase final concentration before being passed through a 0.45 μm syringe filter and stored at 4°C.

Microscopy

For phase contrast images, cells were grown to mid-log-phase, concentrated by centrifugation and 3 μL of culture was spotted on a glass microscope slide and immobilized with a poly-L-lysine-treated glass coverslip. Samples were observed under phase contrast at 1000x magnification using a Nikon eclipse 80i microcope and recorded with a Nikon CoolSNAP HQ² camera and Metamorph image capture software (Universal Imaging).

For fluorescence microscopy, 1.0 mL of broth culture was harvested at mid-log phase and washed once in T-Base buffer (20 g (NH₄)₂ SO₄, 140 g K₂HPO₄, 60 g KH₂PO₄, 10 g Sodium Citrate per liter). The suspension was pelleted and resuspended in 40 μL of T-Base buffer containing 1 μg/mL FM4–64 (Molecular Probes) and incubated for 5 min at room temperature in the dark. Three microliters of suspension were placed on a microscope slide and immobilized with a poly-L-lysine-treated coverslip. Fluorescence microscopy was performed with an Nikon eclipse 80i microcope and X-Cite lamp, recorded with a Nikon

CoolSNAP HQ² camera and Metamorph image capture software (Universal Imaging), and adjusted with Adobe Photoshop software.

FliG protein purification

The FliG protein expression vector pKB43 was transformed into *Rosetta gami E. coli*, grown to ~0.5 OD₆₀₀ in 500 ml of Terrific Broth, induced with 1 mM IPTG and grown overnight at 16°C. Cells were pelleted and resuspended in Lysis Buffer (50 mM Na₂HPO₄, 300 mM NaCl, 10 mM Imidazole) and lysed by treatment with BugBuster for 1 hour at room temperature. Lysed cells were ultracentrifuged at 35,000 rpm. Cleared supernatant was combined with Ni-NTA resin (Novagen) and incubated for 1 hour at 4°C. The bead/lysate mixture was poured onto a 1 cm separation column (BioRad), the resin was allowed to pack and was washed with Wash Buffer (50 mM Na₂HPO₄, 300 mM NaCl, 30 mM Imidazole). SUMO-FliG bound to the resin was then eluted using a stepwise elution of Wash Buffer with 50 mM – 500 mM Imidazole. Elutions were separated by SDS-PAGE and Coomassie stained to verify purification of the SUMO-FliG fusion. Purified SUMO-FliG was combined with Ubiquitin Ligase (protease) and Cleavage Buffer and incubated overnight at 4°C to cleave the SUMO tag from the FliG protein (Butt et al., 2005). The cleavage reaction was combined with Ni-NTA beads, incubated for 2 hours at 4°C, and centrifuged to pellet the resin. Supernatant was removed and dialyzed in to 50 mM Tris (pH 8.0), 150 mM NaCl, 10% Glycerol, 1mM DTT and stored at –20°C. Removal of the SUMO tag was verified by SDS-PAGE and Coomassie staining. Pooled fractions were dialyzed in to 50 mM Tris (pH 8.0), 150 mM NaCl, 10% Glycerol, 1mM DTT and stored at –20°C.

FliY protein purification

The FliY protein expression vector pDP288 was transformed into *Rosetta gami E. coli*, grown to ~0.5 OD₆₀₀ in 500 ml of Terrific Broth, induced with 1 mM IPTG and grown overnight at 16°C. Cells were pelleted and resuspended in Lysis Buffer (50 mM Na₂HPO₄, 300 mM NaCl, 10 mM Imidazole) and lysed by French Press. Lysed cells were ultracentrifuged at 35,000 rpm. Cleared supernatant was combined with Ni-NTA resin (Novagen) and incubated for 1 hour at 4°C. The bead/lysate mixture was poured onto a 1 cm separation column (BioRad), the resin was allowed to pack and was washed with Wash Buffer (50 mM Na₂HPO₄, 300 mM NaCl, 30 mM Imidazole). SUMO-SlrA bound to the resin was then eluted using a stepwise elution of Wash Buffer with 50 mM – 500 mM Imidazole. Elutions were separated by SDS-PAGE and Coomassie stained to verify purification of the SUMO-SlrA fusion. Purified SUMO-FliY was combined with Ubiquitin Ligase (protease) and Cleavage Buffer and incubated overnight at 4°C to cleave the SUMO tag from the FliY protein (Butt et al., 2005). The cleavage reaction was combined with Ni-NTA beads, incubated for 2 hours at 4°C, and centrifuged to pellet the resin. Supernatant was removed and dialyzed in to 50 mM Tris (pH 8.0), 150 mM NaCl, 10% Glycerol, 1mM DTT and stored at –20°C. Removal of the SUMO tag was verified by SDS-PAGE and Coomassie staining.

Hag protein purification

The Hag protein expression vector pJP4 was transformed into *Rosetta gami E. coli*, grown to ~0.5 OD₆₀₀ in 500 ml of Terrific Broth, induced with 1 mM IPTG and grown overnight at 16°C. Cells were pelleted and resuspended in Lysis Buffer (50 mM Na₂HPO₄, 300 mM NaCl, 10 mM Imidazole) and lysed by treatment with BugBuster for 1 hour at room temperature. Lysed cells were ultracentrifuged at 35,000 rpm. Cleared supernatant was combined with Ni-NTA resin (Novagen) and incubated for 1 hour at 4°C. The bead/lysate mixture was poured onto a 1 cm separation column (BioRad), the resin was allowed to pack and was washed with Wash Buffer (50 mM Na₂HPO₄, 300 mM NaCl, 30 mM Imidazole). His₆-Hag bound to the resin was then eluted using a stepwise elution of Wash Buffer with

50 mM – 500 mM Imidazole. Elutions were separated by SDS-PAGE and Coomassie stained to verify purification of the His₆-Hag fusion. Purified His₆-Hag was combined with Thrombin (protease) and Cleavage Buffer and incubated overnight at 25°C to cleave the His₆ tag from the Hag protein. The cleavage reaction was combined with Ni-NTA beads, incubated for 2 hours at 4°C, and centrifuged to pellet the resin. Supernatant was removed and dialyzed into 50 mM Tris (pH 8.0), 150 mM NaCl, 10% Glycerol, 1mM DTT and stored at –20°C.

FliG antibody preparation

1.7 mg of purified FliG protein was cut from SDS-PAGE gel and sent to Cocalico Biologicals Inc. for serial injection into a rabbit host for antibody generation. Anti-FliG serum was mixed with FliG-conjugated Affigel-10 beads and incubated overnight at 4°C. Beads were packed onto a 1 cm column (BioRad) and then washed with 100 mM glycine (pH 2.5) to release the antibody and immediately neutralized with 2M unbuffered Tris. Purification of the antibody was verified by SDS-PAGE. Purified anti-FliG antibody was dialyzed into 1x PBS, 50% glycerol and stored at –80°C.

FliY antibody preparation

1 mg of purified FliY protein was sent to Cocalico Biologicals Inc. for serial injection into a rabbit host for antibody generation. Anti-FliY serum was mixed with FliY-conjugated Affigel-10 beads and incubated overnight at 4°C. Beads were packed onto a 1 cm column (BioRad) and then washed with 100 mM glycine (pH 2.5) to release the antibody and immediately neutralized with 2M unbuffered Tris. Purification of the antibody was verified by SDS-PAGE. Purified anti-FliY antibody was dialyzed into 1x PBS, 50% glycerol and stored at –80°C.

Hag antibody preparation

1 mg of purified Hag protein was sent to Cocalico Biologicals Inc. for serial injection into a rabbit host for antibody generation. Anti-Hag serum was mixed with Hag-conjugated Affigel-10 beads and incubated overnight at 4°C. Beads were packed onto a 1 cm column (BioRad) and then washed with 100 mM glycine (pH 2.5) to release the antibody and immediately neutralized with 2M unbuffered Tris. Purification of the antibody was verified by SDS-PAGE. Purified anti-Hag antibody was dialyzed into 1x PBS, 50% glycerol and stored at –80°C.

Western Blotting

B. subtilis strains were grown in LB to mid-log-phase, 1 or 10 mL were harvested by centrifugation, and resuspended to OD₆₀₀ 100 in Lysis buffer (20 mM Tris pH 7.0, 10 mM EDTA, 1 mg/ml lysozyme, 10 µg/ml DNase I, 100 µg/ml RNase I) and incubated 30 minutes at 37°C. 10 µl of lysate was mixed with 2 µl 6x SDS loading dye. Samples were separated by 15% Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). The proteins were electroblotted onto nitrocellulose and developed with a 1:5,000 dilution (Anti-SigD, Anti-FliY), 1:10,000 dilution (Anti-FliG, Anti-SinR), or 1:40,000 dilution (Anti-Hag, Anti-SigA) of primary antibody and a 1:10,000 dilution secondary antibody (horseradish peroxidase-conjugated goat anti-rabbit immunoglobulin G). Immunoblot was developed using the Immun-Star HRP developer kit (Bio-Rad).

RNA Purification

(Modified from www.microarrays.org, performed as in Cozy and Kearns, 2010) 25 ml of cells were grown to midlog phase in LB broth at 37°C and either separated by density gradient centrifugation (see above) or left as a mixed culture. Cells were then combined with

25 ml cold methanol, pelleted and stored at -80°C . Cell pellets were then combined with 75°C LETS buffer (0.1 M LiCl, 0.01 M EDTA, 0.01 M Tris, 0.2% SDS) and 75°C acid phenol (pH 4.3) and glass beads. The mixture was vortexed and centrifuged $3,200 \times g$ for 10 minutes at 4°C . The resulting aqueous phase was removed and vortexed with phenol-chloroform (5:1, 75°C) and then centrifuged $3,200 \times g$ for 10 minutes at 4°C . The resulting aqueous phase was removed and combined with equal volumes of isopropanol, mixed and microfuged 25 minutes at 4°C to precipitate the RNA. Supernatant was removed from the pellet and the pellet was washed in cold 75% DEPC H_2O -EtOH and microfuged 5 minutes at 4°C . Supernatant was removed from the pellet and the pellet was resuspended in 20 μl of DEPC H_2O . Trizol reagent was added to the resuspended RNA, vortexed. Chloroform was then added, shaken and microfuged 15 minutes at 4°C . The resulting aqueous phase was removed and combined with equal volumes of isopropanol, mixed and microfuged 25 minutes at 4°C to precipitate the RNA. Supernatant was removed from the pellet and the pellet was washed in cold 75% DEPC treated H_2O -EtOH and microfuged 5 minutes at 4°C . Supernatant was removed from the pellet and the pellet was air dried for 5 minutes before it was resuspended in 80 μl of DEPC treated H_2O at 55°C .

Quantitative RT-PCR

Total RNA was isolated from late exponential-phase cultures as described above. Isolated RNA was DNase digested using the TURBO DNA Free Kit (Ambion). cDNA was reverse transcribed from each DNase-treated RNA sample using random dT primers of qScript cDNA SuperMix (Quanta Biosciences). Quantitative PCR was performed with specific primer pairs (1 μM) and either diluted cDNA template or DNase digested RNA template using SYBR Green SuperMix Low ROX (Quanta Biosciences) on the Stratagene MX3500 Pro thermocycler. Data were analyzed using the MXPro Stratagene software package. The following primer pairs were used: 1560/1561 (*flgB*), 2560/2561 (*flgC*), 2562/2563 (*flgE*), 1596/1597 (*fliF*), 2564/2565 (*fliG*), 1598/1599 (*flgE*), 1654/1655 (*fliK*), 1656/1657 (*fliY*), 1600/1601 (*fliZ*), 1658/1659 (*fliR*), 1602/1603 (*flhA*), 1604/1605 (*cheB*), 1662/1663 (*cheW*), 1562/1563 (*cheD*), 1448/1449 (*sigD*), 1450/1451 (*sigA*).

To quantify the amount of mRNA corresponding to each gene, cross-threshold (C_T) values generated using cDNA as template were compared against C_T values generated using a known concentration of chromosomal DNA as template. All primer pairs used in quantitative PCR amplify a 100 bp target in the chromosome. Assuming one copy of each gene per chromosome, the ratio of the concentration of target DNA to total chromosomal DNA is approximately 100bp/4,400,000bp. Chromosomal DNA was isolated from a 3610-derived strain that lacks the endogenous plasmid (McLoon *et al.*, 2011) using a standard protocol and treated with Ribonuclease A (Sigma) for 1 hour at 37°C . RNase-treated chromosomal DNA was purified using phenol:chloroform (1:1) extraction and resuspended in sterile MilliQ water. Quantitative PCR was performed as described above with specific primer pairs (1 μM) and diluted chromosomal DNA of known concentrations spanning 5 orders of magnitude (from 1×10^{-5} ng/ μl to 1×10^{-10} ng/ μl) as template (Figure S3). Data were analyzed using the MXPro Stratagene software package. The C_T values obtained were plotted against the amount of template DNA (ng) used in the quantitative PCR reaction using KaleidaGraph software. From these data, the equation of a line of best fit was calculated and used as a standard curve. The amount of RNA calculated for each gene in the *fla/che* operon was normalized to *sigA* RNA amounts in each genetic background tested.

Transcriptional profiling

Custom Agilent DNA microarrays were designed using the Agilent eArray application. The arrays include 15,744 60-mer probes which consist of the annotated protein coding genes of the *B. subtilis* str. 168 genome, as well as the known non-coding RNAs entered into NCBI

and reported in Rasmussen *et al.*, 2009 and Irnov *et al.*, 2010. Each protein coding gene includes 3 probes and the non-coding RNAs have either 2 or 3 probes.

B. subtilis strains 3610 and DS1578 were grown in LB at 37°C to an optical density of 1.0–1.1 at 600 nm and samples were immediately mixed with an equal volume of methanol (–20°C) and centrifuged to pellet cells. Cell pellets were stored at –80°C. RNA was isolated using a hot acid-phenol isolation procedure as described above. The sample is then treated with Qiagen's RNase-Free DNase in solution and the RNA cleanup protocol of Qiagen's RNeasy® kit was used. The quality of the RNA was checked by visualizing the integrity of the 23S and 16S rRNA bands on an agarose gel.

Labeled cDNA was generated from RNA using Agilent's Fairplay® III Microarray Labeling Kit. The Agilent Two-Color Microarray-Based Prokaryotic Analysis (FairPlay III Labeling) Protocol was used with the following changes: between 6–10 µg total RNA was used, the reverse transcription reaction was performed at 42°C for 2 hours, and the NHS-ester dye-coupling reaction to amino allyl dUTP (using GE Healthcare CyTM3 and CyTM5 monofunctional reactive dyes) was incubated for 90 minutes. Hybridizations were incubated at 65°C for 17 hours. Arrays were scanned by Agilent Technologies DNA Microarray Scanner with SureScan High-Resolution Technology.

Data was subjected to standard lowess normalization using Bioconductor and run in R was performed on the processed signal given by the Agilent software and the geometric mean of the probes was used to give the final value for each gene and nc-RNA.

Hysteresis

Cells were grown in LB supplemented with 0.1mM IPTG to midlog phase, pelleted and washed twice in 10 mL LB without IPTG. The washed pellet was resuspended in LB without IPTG to the starting optical density. Washed cells were then diluted 2^4 , 2^{10} or 2^{20} in 25 mL LB without IPTG and allowed to grow until reaching the starting optical density representing 4, 10 and 20 generations respectively.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank T. Bernhardt, S. Barry, K. Blair, A. Camp, S. Guttenplan, T. Kacmarczyk, D. Rudner, M. Winkler, J. Winkelman, R. Chen, S. Mukherjee, T. Redditt, and A. Zlotnick, for reagents and assistance. This work was supported by the NIH Training Grant (NIH T32 GM007757) to LC, the Women in Science Fellowship to LC, NIH grant GM092616 to PE, and NIH GM093030 to DBK.

References

- Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S. Bacterial persistence as a phenotypic switch. *Science*. 2004; 305:1622–1625. [PubMed: 15308767]
- Barillà D, Caramori T, Galizzi A. Coupling of flagellin gene transcription to flagellar assembly in *Bacillus subtilis*. *J Bacteriol*. 1994; 176:4558–4564. [PubMed: 8045886]
- Becskei A, Seraphin B, Serrano L. Positive feedback in eukaryotic gene networks: cell differentiation by graded to binary response conversion. *EMBO J*. 2001; 20:2528–2535. [PubMed: 11350942]
- Bendezu FO, Hale CA, Bernhardt TG, de Boer PA. RodZ (YfgA) is required for proper assembly of the MreB actin cytoskeleton and cell shape in *E. coli*. *EMBO J*. 2009; 28:193–204. [PubMed: 19078962]

- Bertero MG, Gonzales B, Tarricone C, Cecilian F, Galizzi A. Overproduction and characterization of the *Bacillus subtilis* anti-sigma factor FlgM. *J Biol Chem*. 1999; 274:12103–12107. [PubMed: 10207036]
- Caramori T, Barilla D, Nessi C, Sacchi L, Galizzi A. Role of FlgM in σ^D -dependent gene expression in *Bacillus subtilis*. *J Bacteriol*. 1996; 178:3113–3118. [PubMed: 8655488]
- Chai Y, Chu F, Kolter R, Losick R. Bistability and biofilm formation in *Bacillus subtilis*. *Mol Microbiol*. 2008; 67:254–263. [PubMed: 18047568]
- Chai Y, Kolter R, Losick R. Paralogous antirepressors acting on the master regulator of biofilm formation in *Bacillus subtilis*. *Mol Microbiol*. 2009; 74:876–887. [PubMed: 19788541]
- Chai Y, Norman T, Kolter R, Losick R. An epigenetic switch governing daughter cell separation in *Bacillus subtilis*. *Genes Dev*. 2010a; 24:754–765. [PubMed: 20351052]
- Chai Y, Kolter R, Losick R. Reversal of an epigenetic switch governing cell chaining in *Bacillus subtilis* by protein instability. *Mol Microbiol*. 2010b; 78:218–229. [PubMed: 20923420]
- Chatterjee A, Kaznessis YN, Hu W. Tweaking biological switches through a better understanding of bistability behavior. *Curr Opin Biotechnol*. 2008; 19:475–481. [PubMed: 18804166]
- Chen R, Guttenplan SB, Blair KM, Kearns DB. Role of the sigmaD-dependent autoysins in *Bacillus subtilis* population heterogeneity. *J Bacteriol*. 2009; 191:5775–5784. [PubMed: 19542270]
- Chen YF, Helmann JD. The *Bacillus subtilis* flagellar regulatory protein σ^D : overproduction, domain analysis, and DNA-binding properties. *J Mol Biol*. 1995; 249:743–753. [PubMed: 7602586]
- Chu F, Kearns DB, McLoon A, Chai Y, Kolter R, Losick R. A novel regulatory protein governing biofilm formation in *Bacillus subtilis*. *Mol Microbiol*. 2008; 68:1117–1127. [PubMed: 18430133]
- Cozy LM, Kearns DB. Gene position in a long operon governs motility development in *Bacillus subtilis*. *Mol Microbiol*. 2010; 76:273–285. [PubMed: 20233303]
- Doan T, Marquis KA, Rudner DZ. Subcellular localization of a sporulation membrane protein is achieved through a network of interactions along and across the septum. *Mol Microbiol*. 2005; 55:1767–1781. [PubMed: 15752199]
- Dubnau D, Losick R. Bistability in bacteria. *Mol Microbiol*. 2006; 61:564–572. [PubMed: 16879639]
- Estacio W, Anna-Arriola SS, Adedipe M, Márquez-Magaña LM. Dual promoters are responsible for transcription initiation of the *fla/che* operon in *Bacillus subtilis*. *J Bacteriol*. 1998; 180:3548–3555. [PubMed: 9657996]
- Ferrell JE. Self-perpetuating states in signal transduction: positive feedback, double negative feedback and bistability. *Curr Opin Chem Biol*. 2002; 6:140–148.
- Ferrell JE, Pomeroy JR, Kim SY, Trunnell NB, Xion W, Huang CF, Machleder EM. Simple, realistic models of complex biological processes: Positive feedback and bistability in a cell fate switch and a cell cycle oscillator. *FEBS Lett*. 2009; 583:3999–4005. [PubMed: 19878681]
- Guérout-Fleury AM, Frandsen N, Stragier P. Plasmids for ectopic integration in *Bacillus subtilis*. *Gene*. 1996; 180:57–61. [PubMed: 8973347]
- Guttenplan SB, Blair KM, Kearns DB. The EpsE flagellar clutch is bifunctional and synergizes with EPS biosynthesis to promote *Bacillus subtilis* biofilm formation. *PLOS Genet*. 2010; 6:e1001243. [PubMed: 21170308]
- Helmann JD, Márquez LM, Chamberlin MJ. Cloning, sequencing, and disruption of the *Bacillus subtilis* σ^{28} gene. *J Bacteriol*. 1988; 170:1568–1574. [PubMed: 2832368]
- Hughes KT, Gillen KL, Semon MJ, Karlinsey JE. Sensing structural intermediates in bacterial flagellar assembly by export of a negative regulator. *Science*. 1993; 262:1277–1280. [PubMed: 8235660]
- Imov I, Sharma C, Vogel J, Winkler W. Identification of regulatory RNAs in *Bacillus subtilis*. *Nuc Acid Res*. 2010; 38:6637–6651.
- Kaern M, Elston TC, Blake WJ, Collins JJ. Stochasticity in gene expression: From theories to phenotypes. *Nat Rev Genet*. 2005; 6:451–464. [PubMed: 15883588]
- Kearns DB, Losick R. Cell population heterogeneity during growth of *Bacillus subtilis*. *Genes Dev*. 2005; 19:3083–3094. [PubMed: 16357223]
- Kobayashi K. SlrR/SlrA controls the initiation of biofilm formation in *Bacillus subtilis*. *Mol Microbiol*. 2008; 69:1399–1410. [PubMed: 18647168]

- Kussell E, Leibler S. Phenotypic diversity, population growth, and information in fluctuating environments. *Science*. 2005; 309:2075–2078. [PubMed: 16123265]
- Kutsukake K. Excretion of the anti-sigma factor through a flagellar substructure couples flagellar gene expression with flagellar assembly in *Salmonella typhimurium*. *Mol Gen Genet*. 1994; 243:605–612. [PubMed: 8028576]
- Laurent M, Charvin G, Guespin-Michel J. Bistability and hysteresis in epigenetic regulation of the lactose operon. *Cell Mol Biol*. 2005; 51:583–594. [PubMed: 16359608]
- Lazarevic V, Margo P, Soldo B, Karamata D. Sequencing and analysis of the *Bacillus subtilis* *lytRABC* divergon: a regulatory unit encompassing the structural genes of the *N*-acetylmuramoyl-L-alanine amidase and its modifier. *J Gen Micro*. 1992; 138:1949–1961.
- Lim HN, Lee Y, Razika H. Fundamental relationship between operon organization and gene expression. *Proc Natl Acad Sci USA*. 2011; 108:10626–10631. [PubMed: 21670266]
- Maamar H, Dubnau D. Bistability in the *Bacillus subtilis* K-state (competence) system requires a positive feedback loop. *Mol Microbiol*. 2005; 56:615–624. [PubMed: 15819619]
- Margot P, Pagni M, Karamata D. *Bacillus subtilis* 168 gene *lytF* encodes a γ -D-glutamate-meso-diaminopimelate muropeptidase expressed by the alternative vegetative sigma factor, σ^D . *Microbiol*. 1999; 145:57–65.
- Márquez LM, Helmann JD, Ferrari E, Parker HM, Ordal GW, Chamberlin MJ. Studies of the σ^D -dependent functions in *Bacillus subtilis*. *J Bacteriol*. 1990; 172:3435–3443. [PubMed: 2111808]
- Marquez-Magana LM, Chamberlin MJ. Characterization of the *sigD* transcription unit of *Bacillus subtilis*. *J Bacteriol*. 1994; 176:2427–2434. [PubMed: 8157612]
- McLoon AL, Guttenplan SB, Kearns DB, Kolter R, Losick R. Tracing the domestication of a biofilm-forming bacterium. *J Bacteriol*. 2011; 193:2027–2034. [PubMed: 21278284]
- Mirel DB, Chamberlin MJ. The *Bacillus subtilis* flagellin gene (*hag*) is transcribed by the σ^{28} form of RNA polymerase. *J Bacteriol*. 1989; 171:3095–3101. [PubMed: 2498284]
- Nakata M, Köller T, Moritz K, Ribardo D, Jonas L, McIver KS, Sumitomo T, Terao Y, Kawabata S, Podbielski A, Kreikemeyer B. Mode of expression and functional characterization of FCT-3 pilus region encoded proteins in *Streptococcus pyogenes* serotype M49. *Infect Immun*. 2009; 77:32–44. [PubMed: 18852238]
- Patrick JE, Kearns DB. MinJ (YvjD) is a topological determinant of cell division in *Bacillus subtilis*. *Mol Microbiol*. 2008; 70:1166–1179. [PubMed: 18976281]
- Rasmussen S, Bjorn N, Jarner H. The transcriptionally active regions in the genome of *Bacillus subtilis*. *Mol Microbiol*. 2009; 76:1043–1057. [PubMed: 19682248]
- Rietsch A, Mekalanos JJ. Metabolic regulation of type III secretion gene expression in *Pseudomonas aeruginosa*. *Mol Microbiol*. 2006; 59:807–820. [PubMed: 16420353]
- Roberts JW, Shankar S, Filter JJ. RNA polymerase elongation factors. *Annu Rev Microbiol*. 2008; 62:211–233. [PubMed: 18729732]
- Serizawa M, Yamamoto H, Yamaguchi H, Fujita Y, Kobayashi K, Ogasawara N, Sekiguchi J. Systematic analysis of SigD-regulated genes in *Bacillus subtilis* by DNA microarray and Northern blotting analyses. *Gene*. 2004; 329:125–136. [PubMed: 15033535]
- Sevim E, Gaballa A, Beldüz AO, Helmann JD. DNA-binding properties of the *Bacillus subtilis* and *Aeribacillus pallidus* AC6 σ^D proteins. *J Bacteriol*. 2011; 193:575–579. [PubMed: 21097624]
- Siegele DA, Hu JC. Gene expression from plasmids containing the *araBAD* promoter at subsaturating inducer concentrations represents mixed populations. *Proc Natl Acad Sci USA*. 1997; 94:8168–8172. [PubMed: 9223333]
- Silva IJ, Saramengo M, Dressaire D, Domingues S, Viegas SC, Arraiano CM. Importance and key events of prokaryotic RNA decay: the ultimate fate of an RNA molecule. *WIREs RNA*. 2011; 2:818–836. [PubMed: 21976285]
- Smith TJ, Blackman SA, Foster SJ. Autolysins of *Bacillus subtilis*: multiple enzymes with multiple functions. *Microbiology*. 2000; 146:249–262. [PubMed: 10708363]
- Smits WK, Eschevins CC, Susanna KA, Bron S, Kuipers OP, Hamoan LW. Stripping *Bacillus*: ComK auto-stimulation is responsible for the bistable response in competence development. *Mol Microbiol*. 2005; 56:604–614. [PubMed: 15819618]

- Smits WK, Kuipers OP, Veening JW. Phenotypic variation in bacteria: the role of feedback regulation. *Nat Rev Microbiol.* 2006; 4:259–271. [PubMed: 16541134]
- Swain PS, Elowitz MB, Siggia ED. Intrinsic and extrinsic contributions to stochasticity in gene expression. *Proc Natl Acad Sci USA.* 2002; 99:12795–12800. [PubMed: 12237400]
- Tolker-Nielsen T, Holmstrøm K, Boe L, Molin S. Non-genetic population heterogeneity studied by *in situ* polymerase chain reaction. *Mol Microbiol.* 1998; 27:1099–1105. [PubMed: 9570396]
- van Holde, KE.; Johnson, WC.; Ho, PS. Principles of Physical Biochemistry. Prentice Hall; Upper Saddle River, NJ: 1998. Chemical equilibria involving macromolecules; p. 587-629.
- Veening JW, Smits WK, Kuipers OP. Bistability, epigenetics, and bet-hedging in Bacteria. *Annu Rev Microbiol.* 2008; 62:193–210. [PubMed: 18537474]
- Werhane H, Lopez P, Mendel M, Zimmer M, Ordal GW, Márquez-Magaña LM. The last gene of the *fla/che* operon in *Bacillus subtilis*, *ylxL*, is required for maximal σ^D function. *J Bacteriol.* 2004; 186:4025–4029. [PubMed: 15175317]
- West JT, Estacio W, Márquez-Magaña LM. Relative roles of the *fla/che* P_A , P_{D-3} , and P_{sigD} promoters in regulating motility and *sigD* expression in *Bacillus subtilis*. *J Bacteriol.* 2000; 182:4841–4848. [PubMed: 10940026]
- Winkelman JT, Blair KM, Kearns DB. RemA (YlzA) and RemB (YaaB) regulate extracellular matrix operon expression and biofilm formation in *Bacillus subtilis*. *J Bacteriol.* 2009; 191:3981–3991. [PubMed: 19363116]
- Yasbin RE, Young FE. Transduction in *Bacillus subtilis* by bacteriophage SPP1. *J Virol.* 1974; 14:1343–1348. [PubMed: 4214946]

Figure 1. The regulatory architecture governing σ^D -dependent gene expression

Figure 4: The regulatory architecture governing σ^{54} -dependent gene expression. Block arrows represent open reading frames. Bent arrows represent promoters. Dashed line indicates *fla/che* operon transcript. Straight arrows represent positive genetic interaction. T-bars represent negative genetic interaction. Gray lines indicate the region of the *fla/che* operon that encodes genes required for the assembly of the flagellar hook basal body. To support Fig 5B, three arrows representing genes encoding the fluorophores mCherry (*mcherry*), CFP (*cfp*), and YFP (*yfp*) are also indicated to show how and where each reporter fits into the regulatory hierarchy.

Figure 2. SlrA inhibits σ^D -dependent gene expression

A) The indicated genetic backgrounds were plated on media containing the chromogenic substrate X-gal, incubated overnight at 37°C and photographed. β -galactosidase assays of *P_{hag}-lacZ* transcriptional activity were conducted in the same genetic backgrounds as the colony pictures above each bar and expressed in Miller Units (Table S4). Error bars are the standard deviation of three replicates. The following strains were used to generate this panel: wild type (DS2746), *swrA swrB* (DS2748), *swrA swrB slrA* (DS9243), *swrA swrB slrA (slrA⁺)* (DS9244), *(slrA⁺)* (DS9242), *(slrA⁺) slrA* (DS7929), *(slrA⁺) remA* (DS7846), *(slrA⁺) remB* (DS7845), *(slrA⁺) sinR* (DS9331), *(slrA⁺) slrR* (DS7932). Note: an *epsH* mutation was introduced to the *(slrA⁺) sinR* strain in order to abolish excessive cell cohesion and permit β -galactosidase measurements. B) Map of the region surrounding *slrA*. Block arrows represent open reading frames. Bent arrows represent promoters. Black triangles represent the location of transposon insertion (DS3446 and DS3452). Double T-bar represents the region of DNA used for complementation of *slrA*, referred to as *(slrA⁺)*.

Figure 3. SlrA inhibits P_{hag} expression in a subpopulation of *swrA* cells

Exponentially growing populations containing a σ^D -dependent reporter *lacA::P_{hag}-GFP* in wild type (A, DS9294), *swrA swrB* (B, DS9344), *swrA swrB slrA* (C, DS9358), *swrB* (D, DS9343), *swrB slrA* (E, DS9357), *swrA* (F, DS9342), *swrA slrA* (G, DS9356), *swrA slrA* (*slrA*⁺) (H, DS9381), and (*slrA*⁺) (I, DS9334). Cells were observed using fluorescence microscopy. Membranes were false colored red and the *P_{hag}-GFP* reporter was false colored green. The percentage of ON cells from greater than 600 cells counted in the population is indicated in the upper right hand corner of the overlay panels. Scale bar is 2 μ m.

wild type

(slrA⁺)

Figure 4. An extra copy of *slrA* causes severe chaining

Phase microscopy of wild type (3610) and *(slrA⁺)* (DS1578) cells during exponential growth. Scale bar is 5 μ m.

Figure 5. SlrA inhibits *fla/che* operon transcript levels

A) Proteins from whole cell lysates of wild type (3610), *sigD* (DS6420), *(slrA⁺)* (DS1578) and *(slrA⁺) slrR* (DS3113) were separated by SDS-PAGE and probed by Western blot with anti-FliG, anti-FliY, anti- σ^D , anti-Hag or anti- σ^A primary antibody and indicated by a black triangle. Open triangle indicates a non-specific cross reacting band for anti- σ^D . B) Three transcriptional reporters *P_{D-3P_{fla/che}}*-mCherry (false color red), *fla/che::CFP* (false color blue), and *P_{hag}-YFP* (false color green), were introduced into wild type (DS3139), *(slrA⁺)* (DS6335) and *(slrA⁺) slrR* (DS7944) as indicated by the colored arrows in Figure 1. Cells were grown to exponential phase and observed using phase and fluorescence microscopy. Scale bar is 2 μ m. C) Total RNA was purified from wild type (3610, dashed line), *(slrA⁺)* (DS1578, open circles) and *(slrA⁺) slrR* (DS3113, closed circles) genetic backgrounds and used as a template for quantitative reverse transcriptase PCR. Dots on the graph represent transcript levels for (in operon order) *flgB*, *flgC*, *fliE*, *fliF*, *fliG*, *fliH*, *fliK*, *fliM*, *fliN*, *fliP*, *fliQ*, *fliR*, *fliS*, *fliT*, *fliU*, *fliV*, *fliW*, *fliX*, *fliY*, *fliZ*, *fliA*, *cheB*, *cheW*, *cheD*, and *sigD* at the indicated distances from the *P_{fla/che}* promoter. Transcript levels were quantified using chromosomal DNA to generate a standard curve and normalized to the expression of the *sigA* gene encoding the housekeeping sigma factor,

σ^A . D) $P_{D-3P_{fla/che}}$ expression is unchanged by an extra copy of *slrA* in the chromosome. The indicated genetic backgrounds were used for β -galactosidase assays of $P_{D-3P_{fla/che}}-lacZ$ transcriptional activity and expressed in Miller Units (Table S5). Error bars are the standard deviation of three replicates. The following strains were used to generate this panel: wild type (DS611), (*slrA*⁺) (DS3269), Δ *slrR* (DS8625), (*slrA*⁺) Δ *slrR* (DS8626).

Figure 6. Artificially expressed σ^D restores σ^D -dependent gene expression in the presence of an extra copy of *slrA*

Vertical panels A and B show fluorescence microscopy of DS7228 cells containing the σ^D -dependent reporter *P_{hag}-YFP*, the IPTG-inducible promoter fusion *P_{hyspank}-sigD* and the *slrA* complementation copy (*slrA*⁺). Vertical panels C and D show fluorescence microscopy of DS7546 cells with the same genotype as panels A and B but also includes a *flgM* mutation. Cells were grown to exponential phase in the absence (–) or presence (+) of 1 mM IPTG. Membrane stain was false colored red. *P_{hag}-YFP* was false colored green. Scale bar is 4 μ m.

Figure 7. σ^D expression is hysteretic for activation of P_{hag} expression

A) Fluorescence microscopy of cells containing the σ^D -dependent reporter P_{hag} -YFP (false colored green) and membrane stained (false colored red). Strains were grown to OD₆₀₀ 0.5 in LB with or without 1 mM IPTG. Cells grown with IPTG were then pelleted, washed in LB without IPTG, and separately serially diluted to make growth back to OD₆₀₀ 0.5 require 4, 10 or 20 generations. The following strains were used: DS7803, DS7804, DS8243, DS8129. Common genotypes are indicated along the top of each panel with differences indicated vertically. "WT" refers to the genetic background listed above each panel series. Percent population in the σ^D -ON state is written in the upper right hand corner of each panel as determined by counting over 600 cells. Scale bar is 4 μ m. B) Samples were taken from the hysteresis timecourse in Fig 9A of DS7803 and DS7804, whole cell lysates were

separated by SDS-PAGE and probed by Western blot with anti- σ^D , anti- σ^A , and anti-SinR primary antibodies. The “-” indicates samples grown in the absence of IPTG, “+” indicates cells grown in the presence of IPTG, and numbers indicate the number of generations grown after IPTG was removed. The position of the σ^D protein is indicated by a black triangle. Grey triangle indicates a non-specific cross reacting band for anti- σ^D . The anti-SinR antibody cross-reacts with SlrR (Chu et al., 2008); the position of SlrR is indicated by an open black triangle and the position of SinR is indicated by an open gray triangle. Each series represents a separate gel using the same samples.

Figure 8. Activation of P_{hag} expression is hypersensitive to induction of σ^D

A) Fluorescence microscopy of cells containing the σ^D -dependent reporter P_{hag} -GFP (false colored green) and membrane stained (false colored red). Strain DS9345 was grown to OD₆₀₀ 1.0 in LB in the presence of the indicated amount of IPTG. B) Fluorescence microscopy of cells containing an IPTG inducible $P_{hyspank}$ -GFP construct (false colored green) and membrane stained (false colored red). Strain DS5237 was grown to OD₆₀₀ 1.0 in LB in the presence of the indicated amount of IPTG. C) The fluorescence intensity of over 600 cells of strains DS9345 (closed circles) and DS5237 (open circles) was determined by pixel counting at each IPTG concentration and the median fluorescence intensity was plotted on the graph (Table S6). Hill constants were derived by fitting the data to a standard binding curve (van Holde et al., 1998).

Table 1

Genes inhibited by SlrA^a

Gene	Fold	Annotation
		(fla/che operon)
<i>flgB</i>	12.5	<i>fla/che</i> operon, flagellar rod
<i>flgC</i>	13.4	<i>fla/che</i> operon, flagellar rod
<i>fliE</i>	23.5	<i>fla/che</i> operon, flagellar basal body
<i>fliF</i>	15.9	<i>fla/che</i> operon, flagellar basal body
<i>fliG</i>	22.0	<i>fla/che</i> operon, flagellar rotor
<i>fliH</i>	27.5	<i>fla/che</i> operon, flagellar secretion
<i>fliI</i>	83.5	<i>fla/che</i> operon, flagellar secretion
<i>fliJ</i>	69.5	<i>fla/che</i> operon, flagellar secretion
<i>ylxF</i>	54.3	<i>fla/che</i> operon, unknown
<i>fliK</i>	65.6	<i>fla/che</i> operon, hook length regulator
<i>ylxG</i>	97.8	<i>fla/che</i> operon, putative hook cap
<i>flgE</i>	58.1	<i>fla/che</i> operon, hook protein
<i>ylzI</i>	79.6	<i>fla/che</i> operon, unknown
<i>fliL</i>	75.3	<i>fla/che</i> operon, unknown
<i>fliM</i>	71.8	<i>fla/che</i> operon, flagellar switch complex
<i>fliY</i>	55.9	<i>fla/che</i> operon, flagellar switch complex
<i>cheY</i>	11.1	<i>fla/che</i> operon, chemotaxis
<i>fliZ</i>	38.4	<i>fla/che</i> operon, unknown
<i>fliP</i>	32.4	<i>fla/che</i> operon, flagellar secretion
<i>fliQ</i>	59.6	<i>fla/che</i> operon, flagellar secretion
<i>fliR</i>	58.6	<i>fla/che</i> operon, flagellar secretion
<i>flhB</i>	37.1	<i>fla/che</i> operon, flagellar secretion
<i>flhA</i>	55.1	<i>fla/che</i> operon, flagellar secretion
<i>flhF</i>	31.3	<i>fla/che</i> operon, unknown
<i>ylxH</i>	40.2	<i>fla/che</i> operon, unknown
<i>cheB</i>	51.6	<i>fla/che</i> operon, chemotaxis
<i>cheA</i>	44.4	<i>fla/che</i> operon, chemotaxis
<i>cheW</i>	26.7	<i>fla/che</i> operon, chemotaxis
<i>cheC</i>	34.6	<i>fla/che</i> operon, chemotaxis
<i>cheD</i>	30.8	<i>fla/che</i> operon, chemotaxis
<i>sigD</i>	9.6	<i>fla/che</i> operon, σ^D sigma factor
<i>swrB</i>	7.0	<i>fla/che</i> operon, swarming motility
		(miscellaneous)
<i>cgeD</i>	5.3	glycosyltransferase, sporulation
<i>cgeE</i>	4.8	unknown, sporulation
<i>liaH</i>	5.2	unknown
<i>liaI</i>	6.0	putative permease
<i>maeN</i>	3.1	sodium/malate symporter

Gene	Fold	Annotation
<i>mtlD</i>	2.9	mannitol-1-phosphate 5-dehydrogenase
<i>mtlF</i>	4.0	mannitol PTS enzyme IIA
<i>pspA</i>	3.0	phage shock protein A
<i>rapB</i>	6.5	Rap phosphatase
<i>sdpA</i>	3.4	SDP bacteriocin secretion
<i>spo0M</i>	6.6	unknown
<i>ybfK</i>	3.4	carboxylesterase NP
<i>ydjG</i>	3.0	putative phage replication protein
<i>ydjH</i>	3.1	unknown
<i>ydjI</i>	3.3	putative flotillin
<i>yheJ</i>	4.0	unknown
<i>yjcM</i>	16.2	unknown, putative amidase
<i>yjfA</i>	2.5	unknown
<i>ykoW</i>	4.4	GGDEF/EAL containing protein
<i>yolA</i>	9.3	unknown
<i>yolB</i>	8.0	unknown
<i>ypfA</i>	3.4	unknown
<i>ypfB</i>	5.6	unknown
<i>yuaG</i>	4.4	putative flotillin
<i>yuaI</i>	4.3	putative acetyltransferase
<i>ythQ</i>	3.3	putative ABC transport permease
<i>ytxE</i>	3.6	flagellar-motor sodium channel
<i>yvbX</i>	5.6	putative peptidoglycan epimerase
<i>yvlA</i>	4.6	unknown
<i>yydF</i>	3.4	peptide controlling LiaRS sensor
		(σ^D-regulon)
<i>yvyF</i>	5.3	unknown (σ^D)
<i>flgM</i>	11.4	flagellar anti-sigma factor (σ^D)
<i>flgN</i>	10.7	flagellar hook chaperone (σ^D)
<i>flgL</i>	11.4	flagellar hook associated protein (σ^D)
<i>flgK</i>	13.3	flagellar hook associated protein (σ^D)
<i>yviE</i>	4.2	unknown (σ^D)
<i>hag</i>	157.4	flagellin (σ^D)
<i>yvyC</i>	35.3	unknown (σ^D)
<i>fliD</i>	20.2	flagellin cap (σ^D)
<i>fliS</i>	24.5	flagellin chaperone (σ^D)
<i>fliT</i>	21.2	flagellin cap chaperone (σ^D)
<i>smiA</i>	16.4	swarming motility inhibitor (σ^D)
<i>motA</i>	48.2	flagellar motor protein (σ^D)
<i>motB</i>	34.6	flagellar motor protein (σ^D)
<i>flhO</i>	5.8	flagellar distal rod protein (σ^D)

Gene	Fold	Annotation
<i>flhP</i>	7.4	flagellar distal rod protein (σ^D)
<i>yfmT</i>	82.3	putative dehydrogenase (σ^D)
<i>yfmS</i>	82.1	MCP (σ^D)
<i>mcpA</i>	24.5	MCP (σ^D)
<i>mcpB</i>	13.5	MCP (σ^D)
<i>mcpC</i>	39.4	MCP (σ^D)
<i>hemAT</i>	34.3	MCP (σ^D)
<i>tlpA</i>	6.9	MCP homolog (σ^D)
<i>tlpC</i>	9.9	MCP homolog (σ^D)
<i>yvaQ</i>	3.7	MCP homolog (σ^D)
<i>cheV</i>	6.0	chemotaxis (σ^D)
<i>epr</i>	6.8	extracellular protease (σ^D)
<i>pgdS</i>	15.2	poly-g-glutamate hydrolase (σ^D)
<i>ybdO</i>	4.8	unknown (σ^D)
<i>yjdB</i>	18.3	unknown (σ^D)
<i>ylqB</i>	15.4	unknown (σ^D)
<i>yscB</i>	8.6	unknown (σ^D)
<i>yxkC</i>	19.4	unknown (σ^D)
		(autolysin cluster)
<i>lytA</i>	2.7	unknown (σ^D)
<i>lytB</i>	14.5	autolysin facilitator protein (σ^D)
<i>lytC</i>	20.9	autolysin (σ^D)
<i>lytD</i>	6.9	autolysin (σ^D)
<i>lytF</i>	18.2	autolysin, cell separation (σ^D)
<i>iseA</i>	6.1	autolysin inhibitor protein
		(surfactin cluster)
<i>urfAA</i>	3.9	surfactin synthesis
<i>urfAB</i>	4.1	surfactin synthesis
<i>urfAC</i>	4.1	surfactin synthesis
<i>urfAD</i>	3.4	unknown
<i>comS</i>	3.9	MecA chaperone antagonist
<i>comK</i>	2.9	Transcription factor, competence
		(wapA cluster)
<i>wapA</i>	3.2	protease, wall-associated
<i>yxxG</i>	4.0	unknown
<i>yxiF</i>	3.8	unknown
<i>yxiG</i>	3.4	unknown
<i>yxiH</i>	3.1	unknown
<i>yxzG</i>	3.0	unknown
<i>yxiI</i>	2.6	unknown

Gene	Fold	Annotation
<i>yxiJ</i>	2.6	unknown
<i>yxiM</i>	2.5	unknown (sRNA)
<i>RNA61</i>	2.8	between <i>yxjI</i> and <i>yxjH</i>
<i>RNA62</i>	3.2	between <i>yxjH</i> and <i>yxjG</i>

^a Genes assigned to functional categories in bold. (σ^D) indicates genes previously identified as being expressed under the control of σ^D (Serizawa et al., 2004; Kearns and Losick, 2005)

Table 2Genes activated by SlrA^a

Gene	Fold	Annotation
(biofilm, polymer cluster)		
<i>epsA</i>	4.4	EPS synthesis, putative length regulator
<i>epsB</i>	4.2	EPS synthesis, putative tyrosine kinase
<i>epsC</i>	6.6	EPS synthesis, epimerase/dehydratase
<i>epsD</i>	8.1	EPS synthesis, glycosyltransferase
<i>epsE</i>	8.9	EPS synthesis and flagellar clutch
<i>epsF</i>	15.0	EPS synthesis, glycosyltransferase
<i>epsG</i>	14.3	EPS synthesis, unknown
<i>epsH</i>	12.7	EPS synthesis, glycosyltransferase
<i>epsI</i>	12.6	EPS synthesis, pyruvyltransferase
<i>epsJ</i>	16.9	EPS synthesis, glycosyltransferase
<i>epsK</i>	16.3	EPS synthesis, putative flippase
<i>epsL</i>	15.0	EPS synthesis, glycosyltransferase
<i>epsM</i>	15.0	EPS synthesis, acetyltransferase
<i>epsN</i>	12.4	EPS synthesis, aminotransferase
<i>epsO</i>	7.6	EPS synthesis, pyruvyltransferase
<i>slrR</i>	3.5	Regulator of EPS synthesis/motility
<i>tapA</i>	4.4	TasA anchoring protein
<i>sipW</i>	6.2	TapA/TasA signal peptidase
<i>tasA</i>	7.5	extracellular amyloid fiber
<i>pgsB</i>	15.1	poly- γ -glutamate synthase
<i>pgsC</i>	17.7	poly- γ -glutamate synthase
<i>pgsA</i>	14.9	poly- γ -glutamate synthase
<i>pgsE</i>	11.1	poly- γ -glutamate synthase
<i>bglC</i>	8.6	endo- β -1,4-glucanase
<i>pelB</i>	4.2	pectate lyase
<i>yoaJ</i>	7.5	putative extracellular endoglucanase
<i>slrA</i>	7.9*	Regulator of EPS synthesis/motility
(miscellaneous)		
<i>catD</i>	4.6	catechol-2,3-dioxygenase
<i>catE</i>	3.9	catechol 2,3-dioxygenase
<i>pksJ</i>	4.4	polyketide synthase
<i>ung</i>	2.8	uracil DNA-glycosylase
<i>yjiA</i>	5.1	unknown
<i>yraJ</i>	4.3	unknown
<i>yvkA</i>	3.0	putative multidrug efflux pump
<i>yvkB</i>	3.6	putative TetR-like transcription fractor
<i>yvkC</i>	5.5	putative pyruvate water dikinase
<i>ywmC</i>	10.0	unknown

Gene	Fold	Annotation
<i>ywmD</i>	5.9	unknown

^a*indicates that *slrA* activation is elevated in part due to the presence of the extra copy included in one strains for comparison.

Table 3

Transposon insertions that restored $P_{hag-lacZ}$ expression when an extra copy of *slrA* was present in the genome.

Gene	Function of gene product	Transposon insertion	Transposon insertion site
<i>slrA</i>	Inhibitor of motility gene expression	TnΩ7929	TAGTAACCT ¹
		TnΩ7930	TATGCCAAC
		TnΩ7931	TAAAAACTG
		TnΩ7848	TAGCTTCTT
<i>remA</i>	Activator of SlrR expression	TnΩ7846	TAGCCTAGT ²
<i>remB</i>	Activator of SlrR expression	TnΩ7845	TATTACTCT
		TnΩ7851	TACTCTCCC
<i>sinR</i>	Co-repressor with SlrR	TnΩ7933	TAGTTCTGA
<i>slrR</i>	Inhibitor of motility gene expression	TnΩ7932	TAATATGAA ¹

¹The transposon inserted between the start codon from the ribosome binding site.

²The transposon inserted upstream of the ribosome binding site and is likely polar on *remA* expression.

Table 4

Strains^a

Strain	Genotype
3610	Wild type
PY79	<i>swrA</i> ^{PY79} <i>sfp</i> ⁰ (laboratory strain)
DS611	<i>thrC</i> :: <i>P</i> _{fla/che} - <i>lacZ mls</i>
DS1578	<i>amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS2746	<i>lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet</i>
DS2748	<i>ΔswrA ΔswrB lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet</i>
DS3113	<i>amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat slrR</i> :: <i>tet</i>
DS3139	<i>thrC</i> :: <i>P</i> _{flache} - <i>mCherry spec lacA</i> :: <i>P</i> _{hag} - <i>YFP tet swrBΩCFP mls</i>
DS3269	<i>amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat thrC</i> :: <i>P</i> _{fla/che} - <i>lacZ mls</i>
DS3446	<i>swrA swrB lacA</i> :: <i>tet Phag-lacZ slrA</i> :: <i>TnYLB kan (TATGCCAAC)</i>
DS3452	<i>swrA swrB lacA</i> :: <i>tet Phag-lacZ slrA</i> :: <i>TnYLB kan (TATGAAAAC)</i>
DS5237	<i>thrC</i> :: <i>P</i> _{hyspank} - <i>GFP mls</i>
DS6335	<i>thrC</i> :: <i>P</i> _{flache} - <i>mCherry spec lacA</i> :: <i>P</i> _{hag} - <i>YFP tet swrBΩCFP mls amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS6356	<i>ΔflgM ΔswrA amyE</i> :: <i>P</i> _{hag} - <i>GFP cat thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls sigD</i> :: <i>tet</i>
DS6420	<i>ΔsigD</i>
DS6447	<i>ΔflgM ΔswrA amyE</i> :: <i>P</i> _{hag} - <i>GFP cat thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls</i>
DS7228	<i>amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP mls thrC</i> :: <i>P</i> _{hyspank} - <i>sigD spec</i>
DS7564	<i>ΔflgM amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} - <i>sigD mls</i>
DS7803	<i>ΔflgM amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls</i>
DS7804	<i>ΔflgM amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls ΔsigD</i>
DS7845	<i>remB</i> :: <i>TnYLB kan (TATTACTCT) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7846	< <i>remA</i> :: <i>TnYLB kan (TAGCCTAGT) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7848	<i>slrA</i> :: <i>TnYLB kan (TAGCTTCTT) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7851	<i>remB</i> :: <i>TnYLB kan (TACTCTCCC) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7929	<i>slrA</i> :: <i>TnYLB kan (TAGTAACCT) amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet</i>
DS7930	<i>slrA</i> :: <i>TnYLB kan (TATGCCAAC) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7931	<i>slrA</i> :: <i>TnYLB kan (TAAAACTG) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7932	<i>slrR</i> :: <i>TnYLB kan (TAATATGAA) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7933	<i>sinR</i> :: <i>TnYLB kan (TAGTTCTGA) lacA</i> :: <i>P</i> _{hag} - <i>lacZ tet amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7944	<i>ΔslrR thrC</i> :: <i>P</i> _{flache} - <i>mCherry spec lacA</i> :: <i>P</i> _{hag} - <i>YFP tet swrBΩCFP mls amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat</i>
DS7966	<i>ΔsigD amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls</i>
DS7967	<i>amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls</i>
DS8129	<i>ΔP_{D-3} amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls</i>
DS8243	<i>ΔP_{D-3} ΔP_{flache} amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat lacA</i> :: <i>P</i> _{hag} - <i>YFP tet thrC</i> :: <i>P</i> _{hyspank} ^(crRBS) - <i>sigD mls</i>
DS8625	<i>slrR</i> :: <i>tet thrC</i> :: <i>P</i> _{fla/che} - <i>lacZ mls</i>
DS8626	<i>amyE</i> :: <i>P</i> _{slrA} - <i>slrA cat slrR</i> :: <i>tet thrC</i> :: <i>P</i> _{fla/che} - <i>lacZ mls</i>

Strain	Genotype
DS8666	<i>AflgM amyE::P_{hyspank}-slrR cat thrC::P_{hyspank}-GFP mls</i>
DS8667	<i>AflgM ΔsigD amyE::P_{hyspank}-slrR cat thrC::P_{hyspank}-GFP mls</i>
DS9242	<i>amyE::P_{slrA}-slrA cat lacA::P_{hag}-lacZ tet</i>
DS9243	<i>ΔswrA ΔswrB slrA::TnYLB kan lacA::P_{hag}-lacZ tet</i>
DS9244	<i>ΔswrA ΔswrB slrA::TnYLB kan amyE::P_{slrA}-slrA cat lacA::P_{hag}-lacZ tet</i>
DS9294	<i>lacA::P_{hag}-GFP mls</i>
DS9331	<i>sinR::TnYLB kan epsH::Tn10 spec lacA::P_{hag}-lacZ tet amyE::P_{slrA}-slrA cat</i>
DS9334	<i>amyE::P_{slrA}-slrA cat lacA::P_{hag}-GFP mls</i>
DS9342	<i>ΔswrA lacA::P_{hag}-GFP mls</i>
DS9343	<i>ΔswrB lacA::P_{hag}-GFP mls</i>
DS9344	<i>ΔswrA ΔswrB lacA::P_{hag}-GFP mls</i>
DS9345	<i>AflgM amyE::P_{slrA}-slrA cat P_{fla/che}-ΩP_{hyspank}-fla/che spec lacA::P_{hag}-GFP mls</i>
DS9356	<i>ΔswrA slrA::TnYLB kan lacA::P_{hag}-GFP mls</i>
DS9357	<i>ΔswrB slrA::TnYLB kan lacA::P_{hag}-GFP mls</i>
DS9358	<i>ΔswrA ΔswrB slrA::TnYLB kan lacA::P_{hag}-GFP mls</i>
DS9381	<i>ΔswrA slrA::TnYLB kan amyE::P_{slrA}-slrA cat lacA::P_{hag}-GFP mls</i>

^a All strains are in the undomesticated 3610 genetic background unless otherwise indicated.