An MspI and PstI RFLP detected by probe pFMS22 – 1.4 [D20S22]

Frank-Michael Stolz and Ingo Hansmann
Institut für Humangenetik, Universität Göttingen,
Goßlerstr. 12d, 3400 Göttingen, FRG

Source/Description: The phage clone FMS22, containing a 0.8 and a 1.4 kb HindIII insert, was isolated from a human chromosome 20-specific gene library (Lawrence Livermore National Laboratory, California, LL20NS01, ATCC # 57712). The 1.4 kb HindIII fragment was subcloned into the HindIII site of pBR322, giving rise to pFMS22–1.4.

Polymorphisms: MspI identifies a three-allele polymorphism with bands at 4.5 (a), 2.6 (b) and 2.0 kb (c). Pst I detects two alleles with bands at 1.6 (y) and 1.1 kb (z) and an additional constant band at 1.2 kb. The DNA source is blood lymphocytes.

Frequencies:

MspI: a (4.5 kb) : 0.17	PstI : y (1.6 kb) : 0.18
b (2.6 kb) : 0.46	z (1.1 kb) : 0.82
c (2.0 kb) : 0.37	
Studied in 49 unrelated European Caucasians.	Studied in 60 unrelated European Caucasians.
PIC = 0.55	PIC = 0.25

Not Polymorphic For: HindIII (one band at 1.3 kb), RsaI (one band at 1.5 kb) and TaqI (one band at 5.9 kb) with panels of 5 or 6 unrelated individuals.

Chromosomal Localization: Localized to chromosome 20 using a panel of human/mouse and human/hamster somatic cell hybrids (kindly provided by Prof. K.-H.Grzeschik, Marburg, FRG).

Mendelian Inheritance: Co-dominant segregation was shown in a three-generation family.

Probe Availability: Available for collaboration.

Other Comments: For each polymorphism the distribution of homozygotes and heterozygotes was consistent with Hardy-Weinberg equilibria determined from the individual allele frequencies. However, the occurrence of an a-allele was found to be always linked to the occurrence of a y-allele. Accordingly, maximum likelihood estimation of the haplotype frequencies disclosed a significant linkage disequilibrium ($\chi^2 = 48.22$, 3df, $P < 0.001$) between the MspI and the PstI RFLP.

At low stringency further bands are visible from 1.2 to 11.0 kb with MspI and 1.7 to 11.6 kb with PstI, respectively.

Acknowledgements: We thank Dr Hans Neumeyer, Abt. Transfusionsmedizin, for providing us with blood samples, Pia Ballhausen for the preparation of DNA, Dr. Michael Krawczak for the maximum likelihood calculations and Christiane Kläehn for the photographic reproductions. This work was supported by the Deutsche Forschungsgemeinschaft, grant Ha 747/8 – 1 to I.H.

A rare BclI RFLP in the putative oncogene bcl-1 locus

P.Selvanayagam, L.C.Strong, G.F.Saunders
and B.Barlogie

The University of Texas M.D.Anderson Cancer Center, Departments of Hematology, Pediatrics, Biochemistry and Molecular Biology, Houston, TX 77030, USA

Source and Description of Clone: A genomic 0.6 kb SmaI–EcoRI fragment pRc8SmR, subcloned from the lambda clone Rc8, isolated from the tumor cell DNA of a CLL patient carrying a t(11;14) chromosomal translocation (Tsujimoto *et al.*, 1984).

Polymorphism: Restriction endonuclease BclI identified a 9 kb variant allele besides the normal 13 kb allele in normal cells of a patient with multiple myeloma.

Frequency: Only one out of 120 unrelated multiple myeloma patients and healthy individuals was heterozygous, exhibiting a frequency of 0.01 for the 9 kb allele.

Other Enzymes Tested and Negative: BamHI, EcoRI, HindIII, SstI, PvuII and BglII.

Chromosomal Localization: The probe pRc8SmR is derived from chromosome 11q13, as determined by karyotyping of the tumor cell line (Erickson *et al.*, 1984).

Mendelian Inheritance: Co-dominant segregation of the variant allele was observed in four informative matings covering three generations.

Probe Availability: Contact Y.Tsujimoto at the Wistar Institute, 36th and Spruce Street, Philadelphia, PA 19104, USA

Acknowledgement: This work was supported by CA 37161 from the National Cancer Institute, Bethesda, Maryland.

References: Tsujimoto, Y. *et al.* (1984) *Science* **224**, 1403–1406. Erickson, J. *et al.* (1984) *Proc. Natl. Acad. Sci. USA* **81**, 4144–4148.