Published in final edited form as:

Org Lett. 2012 March 16; 14(6): 1408–1411. doi:10.1021/ol3001365.

Enantioselective Total Synthesis of (*S*)-Bisoranjidiol, an Axially Chiral Bisanthraquinone

Erin E. Podlesny and Marisa C. Kozlowski*

Department of Chemistry, Roy and Diana Vagelos Laboratories, University of Pennsylvania, Philadelphia, Pennsylvania 19104, United States

Abstract

The first enantioselective total synthesis of the bisanthraquinone (*S*)-bisoranjidiol and an unnatural regioisomer has been accomplished. Key features of the synthesis include the asymmetric oxidative biaryl coupling of a hindered 8-substituted 2-naphthol, selective *para*-quinone formation, and regioselective tandem Diels-Alder/aromatization reactions.

(*S*)-Bisoranjidiol [(*S*)-**1**, Figure 1] is a photosensitizing bisanthraquinone capable of interacting with molecular oxygen to generate electronically excited singlet oxygen and the ground state radical anion upon irradiation with light.¹ These photosensitizing properties are responsible for the phototoxicity of the shrub *Heterophyllaea pustulata* (genera Rubiaceae), from which it was isolated in 2006.² Native to the Andean Mountains of Argentina and Bolivia, the plant poses a danger to grazing livestock. Ingestion of the plant and exposure to sunlight cause dermatitis, keratoconjunctivitis (which may lead to blindness), and behavioral changes, such as restlessness and photophobia.^{3,4} The photosensitizing properties of (*S*)-**1** make it an interesting natural product with potential applications in photodynamic therapy.⁵ Recently, it has been reported to be a good antibacterial agent⁶ and photodynamically active toward cancer cells.⁷ To date, no asymmetric syntheses of bisanthraquinones nor any synthesis of **1** have been achieved. In this letter, we report the first total synthesis of (*S*)-bisoranjidiol (**1**) via a route that would allow analog generation to optimize biological properties.

Bisoranjidiol (**1**) is a member of a larger class of axially chiral symmetrical bisanthraquinone natural products possessing a biaryl linkage at the 1,1'-positions (see Figure 1). Spectroscopic analyses and chemical methods established the structure and configuration of **1**.² Bisoranjidiol (**1**) is unique compared to the other bisanthraquinone natural products (**2**) as it lacks the 4,4'-hydroxyl groups and the *meta*-substitution pattern of the distal rings shared by the other congeners.

marisa@sas.upenn.edu.

Supporting Information Available: Experimental procedures, characterization data, NMR spectra, CD spectra, HPLC traces. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Prior syntheses of symmetrical bisanthraquinones possessing a 1,1'-biaryl linkage include biomimetic coupling of an anthraquinone or anthrone monomer,⁸ the Ullman reaction between brominated anthraquinones,⁹ and double addition of an isobenzofuran-1,3-dione to a biphenyl intermediate.¹⁰ However, the only bisanthraquinone natural products synthesized include skyrin (0.28 – 35% yield, R^{1,2} = H, R³ = Me; Figure 1) and hinakurin (≤ 58% yield, R¹ = H, R^{2,3} = Me; Figure 1), which were formed via biomimetic syntheses⁸ and an Ullman reaction, respectively.^{9a} The total syntheses of related bisanthraquinones, biphyscion, 2,2'-*epi*-cytoskyrin A, and rugulosin have also been reported, however, biphyscion contains a 2,2'- biaryl linkage¹¹ and the latter two possess a cage-like “skyrane” motif.¹²

Retrosynthetic analysis of (*S*)-**1** revealed a different, nonbiomimetic approach to this structural subtype that hinges on the much simpler prospect of forming a binaphthol via oxidative coupling rather than the more difficult oxidative coupling of an oxidation resistant anthraquinone. Specifically, a double Diels-Alder reaction between a binaphtho-*para*-quinone [(*S*)-**4a**] and vinyl ketene acetal **3**, followed by aromatization would readily reveal the core bisanthraquinone structure (Scheme 1). The Diels-Alder reaction of 1,4-naphthoquinones with vinylketene acetals, followed by aromatization is well known.¹³ Alkyl trimethylsilyl- or bis(trimethylsilyl)vinyl ketene acetals, also referred to as Brassard dienes,¹³ are typically employed in these types of reactions although less ornate dienes such as Danishefsky's diene¹⁴ have also been used.

The key binaphtho-*para*-quinone (*S*)-**4a** would, in turn, require a selective oxidation of (*S*)-**5**, a heretofore unexplored undertaking. Enantioselective oxidative binaphthol coupling of **7**, containing substitution at the 8-position, with our 1,5-diaza-*cis*-decalin copper catalyst,¹⁵ followed by ester and protecting group removal would provide key intermediate (*S*)-**6**. This asymmetric coupling reaction has been an integral part of the syntheses of several other axially or helically chiral naphthalene-based natural products from our group.^{16,17}

Coupling precursor **7** was effectively prepared from **8** via a Fischer esterification and selective benzylation of the less hindered phenol (Scheme 2). The enantioselective biaryl coupling of substrate **8** with (*R,R*)-**9** proceeded surprisingly smoothly even with the C8-substitution, which provides considerable steric hindrance to the coupling at C1. As a result, (*S*)-**6**¹⁸ was obtained in 62% yield and 87% ee (61% yield, 92% ee with (*S,S*)-**9**). After one trituration, material of >99% ee was readily obtained. To the best of our knowledge, the only reported enantioselective binaphthol coupling reaction with substitution in the hindered 8-position is the coupling of naphtho[2,1-*b*]furan-8-ol in 50% yield and 75% ee.¹⁹

After methylation of (*S*)-**6**, the esters were effectively removed in 63% yield via a three step sequence involving DIBALH reduction to the alcohol, IBX promoted oxidation to the aldehyde and decarbonylation with RhCl(PPh₃)₃ (Scheme 3). In this instance, this combination was found to be more efficient and reliable than decarboxylation. Hydrogenolysis of the benzyl groups followed by a selective salcomine catalyzed oxidation yielded the binaphtho-*para*-quinone (*S*)-**12** in 63% yield without loss of enantiopurity. Approximately 23% of an unsymmetrical binaphtho-*ortho,para*-quinone was also isolated. This side product could be avoided by using of (diacetoxyiodo)benzene, but yields were much lower.

With (*S*)-**12** in hand, the remaining key step was transformation of the binaphtho-*para*-quinone into a bisanthraquinone via regioselective tandem Diels-Alder/aromatization reactions. Initial investigations revealed that the use of Lewis acids, such as ZnCl₂, provided control over the regioselectivity of the cycloaddition, however, side reactions and decomposition of the quinone consistently resulted in low yields. Aside from Lewis acids,

halogens are often used as directing groups when working with vinyl ketene acetals and *para*-quinones.²⁰ Subsequent investigations on a monomeric system, indicated that halogen directing groups provided superior yields and complete regiocontrol in the cycloaddition reaction.

To install the desired bromine directing group, (*S*)-**12** was brominated with Br₂, followed by dehydrohalogenation in acetic acid (Scheme 4). This procedure provided a mixture of bromoquinones in 95% yield. The mixture was comprised of the 6,6', 6,7', and 7,7'-dibrominated regioisomers. Approximately equal ratios of the 6,6' and 6,7'-dibromoquinones [(*S*)-**4a** and (*S*)-**4b**; Scheme 4] were obtained, plus 10% of the 7,7'-dibromoquinone [(*S*)-**4c**]. The production of (*S*)-**4a** and (*S*)-**4b** was seen as an opportunity to produce both a significant amount of bisoranjidiol precursor [(*S*)-**13a**] and an unnatural bisanthraquinone regioisomer. Due to the directing ability of the halogen in the cycloaddition reaction, we expected the ratio of regioisomers **4a–c** to be reflected in the product bisanthraquinones. When diene **14**²¹ was used in a test reaction with a substrate analogous to **4a–c**,²² this result did occur, but interestingly, with diene **3**²³ and **4a–c**, the process was very selective for (*S*)-**13a**. It appeared that (*S*)-**4b** and (*S*)-**4c** were either not reacting as well as (*S*)-**4a** and/or were decomposing. To evaluate the reasons for the selectivity, the regioisomers (*S*)-**4a** and (*S*)-**4b** were separated and treated separately with diene **3**. The key intermediate to bisoranjidiol, (*S*)-**4a**, reacted completely with diene **3** in two days to afford the “out-out” bisanthraquinone (*S*)-**13a** in 80% yield, following aromatization over silica (Scheme 4). Dibromoquinone (*S*)-**4b**, on the otherhand, stalled at the monocycloadduct stage and required 5–6 days to reach completion, with noticeably more decomposition over the course of the reaction compared to (*S*)-**4a**. Aromatization over silica provided “out-in” (*S*)-**13b** in 46–71% yield. This was deprotected to give the regioisomer of bisoranjidiol (Scheme 4). The minor isomer (*S*)-**4c** mostly led to decomposition. Presumably, the formation of the “in” Diels-Alder adduct with the phenolic substituent disposed to the hindered bay region of the structure is highly disfavored on steric grounds.

Finally, deprotection of (*S*)-**13a** with excess BBr₃ provided (*S*)-bisoranjidiol, (*S*)-**1**, in 80% yield with no erosion of the enantiomeric excess (Scheme 4). Both the chemical shifts and splitting patterns from the ¹H and ¹³C NMR data of the synthetic material were in accord with the published spectroscopic data for the natural product.² Using an authentic sample of the natural product kindly provided by Dr. Cabrera,² we undertook chiral HPLC analysis to confirm the original stereochemical assignment as *S*.² Surprisingly, the authentic sample showed only 5% ee of the (*S*)-enantiomer. Since the ee is small, it is also possible that the natural sample is racemic. Concurrently, we discovered that the enantiopurity of bisoranjidiol eroded in MeOH (25 °C, t_{1/2} = 3.8 months; 80 °C, t_{1/2} = 1.8 h). As the isolation conditions involved refluxing solvent (benzene) and exposure to base and acid,² it is not unreasonable that significant erosion of the enantiopurity occurred during isolation and subsequent handling of the material.

In summary, the first total synthesis of (*S*)-bisoranjidiol has been completed in 12 steps and 4% overall yield from commercially available **8**. This synthesis also represents the most selective coupling of a 2-naphthol with substitution in the hindered 8-position, as well as the first enantioselective synthesis of a 1,1'-linked bisanthraquinone. This approach represents a more efficient entry into this natural product series and also permits flexibility in generation of further analogs for study.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support for this work was provided by the National Science Foundation (CHE-0911713). E.E.P. gratefully acknowledges the National Institutes of Health for both an NRSA (F31-GM093515) and Chemistry-Biology Interface (T32-GM071339) predoctoral fellowships. Partial instrumentation support was provided by the NIH for MS (1S10RR023444) and NMR (1S10RR022442). We also thank Dr. José Cabrera (Universidad Nacional de Córdoba, Argentina) for an authentic sample of bisoranjidiol. In addition, we acknowledge Dr. Barbara J. Morgan (University of Pennsylvania) for her contributions to the biaryl coupling reaction, as well as Dr. Ömer Koz (visiting researcher, Ege University, Turkey), and undergraduates Xaira Lopez Corcino (REU student), and Kenny Puk (University of Pennsylvania) for scale-up of racemic material.

References

- Comini LR, Núñez Montoya SC, Sarmiento M, Cabrera JL, Argüello GA. *J. Photochem. Photobiol., A*. 2007; 188:185–191.
- Núñez Montoya SC, Agnese AM, Cabrera JL. *J. Nat. Prod.* 2006; 69:801–803. [PubMed: 16724844]
- Núñez Montoya SC, Comini LR, Rumie Vittar B, Fernández IM, Rivarola VA, Cabrera JL. *Toxicon*. 2008; 51:1409–1415. [PubMed: 18513778]
- Aguirre DH, Neumann RA. *Med. Vet.* 2001; 18:487–490.
- (a) Dougherty TJ, Gomer CJ, Henderson BW, Jori G, Kessel D, Korbelik M, Moan J, Peng Q. *J. Natl. Cancer Inst.* 1998; 90:889–905. [PubMed: 9637138] (b) Dolmans DE, Fukumara D, Jain RK. *Nat. Rev. Cancer*. 2003; 3:380–387. [PubMed: 12724736] (c) Hamblin MR, Hasan T. *Photochem. Photobiol. Sci.* 2004; 3:436–450. [PubMed: 15122361] (d) Yano S, Hirohara S, Obata M, Hagiya Y, Ogura S, Ikeda A, Kataoka H, Tanaka M, Joh T. *J. Photochem. Photobiol., C*. 2011; 12:46–67.
- Comini LR, Núñez Montoya SC, Páez PL, Argüello GA, Albesa I, Cabrera JL. *J. Photochem. Photobiol., B*. 2011; 102:108–114. [PubMed: 20965744]
- Fernández I, Comini L, Nigra A, Núñez S, Rumie Vittar B, Cabrera J, Rivarola VA. *Biocell*. 2009; 33:A266.
- (a) Franck B, Chahin R, Eckert H-G, Langenberg R, Radtke V. *Angew. Chem. Int. Ed.* 1975; 14:819–820. (b) Banks HJ, Cameron DW, Raverty WD. *Aust. J. Chem.* 1976; 29:1509–1521. (c) Cameron DW, Edmonds JS, Raverty WD. *Aust. J. Chem.* 1976; 29:1535–1548.
- (a) Tanaka O, Kaneko C. *Pharm. Bull.* 1955; 3:284–286. [PubMed: 13289274] (b) Shibata S, Tanaka O, Kitagawa I. *Pharm. Bull.* 1955; 3:278–283. [PubMed: 13289273] (c) Tanaka O. *Chem. Pharm. Bull.* 1958; 2:203–208. [PubMed: 13537188] (d) Iio H, Zenfuku K, Tokoroyama T. *Tetrahedron Lett.* 1995; 36:5921–5924.
- Scholl R, Liese K, Michelson K, Grunewald E. *Ber.* 1910; 43:512–518.
- Hauser FM, Gauuan PJF. *Org. Lett.* 1999; 1:671–672. [PubMed: 10823198]
- (a) Nicolaou KC, Papageorgiou CD, Piper JL, Chadha RK. *Angew. Chem. Int. Ed.* 2005; 44:5846–5851. (b) Nicolaou KC, Lim YH, Piper JL, Papageorgiou CD. *J. Am. Chem. Soc.* 2007; 129:4001–4013. [PubMed: 17355133]
- Savard J, Brassard P. *Tetrahedron*. 1984; 40:3455–3464.
- Danishefsky S, Kitaharai T. *J. Am. Chem. Soc.* 1974; 96:7807–7808.
- (a) Li X, Yang J, Kozlowski MC. *Org. Lett.* 2001; 3:1137–1140. [PubMed: 11348178] (b) Kozlowski MC, Li X, Carroll PJ, Xu Z. *Organometallics*. 2002; 21:4513–4522. (c) Xie X, Phuan P-W, Kozlowski MC. *Angew. Chem. Int. Ed.* 2003; 42:2168–2170. (d) Mulrooney CA, Li X, DiVirgilio ES, Kozlowski MC. *J. Am. Chem. Soc.* 2003; 125:6856–6857. [PubMed: 12783524] (e) Li X, Hewgley JB, Mulrooney CA, Yang J, Kozlowski MC. *J. Org. Chem.* 2003; 68:5500–5511. [PubMed: 12839440] (f) Morgan BJ, Xie X, Phuan P-W, Kozlowski MC. *J. Org. Chem.* 2007; 72:6171–6182. [PubMed: 17629337] (g) Hewgley JB, Stahl SS, Kozlowski MC. *J. Am. Chem. Soc.* 2008; 130:12232–12233. [PubMed: 18710234]
- (a) Kozlowski MC, Morgan BJ, Linton EC. *Chem. Soc. Rev.* 2009; 38:3193–3207. [PubMed: 19847351] (b) DiVirgilio ES, Dugan EC, Mulroony CA, Kozlowski MC. *Org. Lett.* 2007; 9:385–388. [PubMed: 17249768] (c) O'Brien EM, Morgan BJ, Kozlowski MC. *Angew. Chem. Int. Ed.* 2008; 47:6877–6880. (d) Morgan BJ, Dey S, Johnson SW, Kozlowski MC. *J. Am. Chem. Soc.* 2009; 131:9413–9425. [PubMed: 19489582] (e) Morgan BJ, Mulrooney CA, O'Brien EM,

- Kozlowski MC. J. Org. Chem. 2010; 75:30–43. [PubMed: 19894745] (f) Morgan BJ, Mulrooney CA, Kozlowski MC. J. Org. Chem. 2010; 75:44–56. [PubMed: 19894744] (g) O'Brien EM, Morgan BJ, Mulrooney CA, Kozlowski MC. J. Org. Chem. 2010; 75:57–68. [PubMed: 19894741]
17. For a general review, see: Bringmann G, Gulder T, Gulder TAM, Breuning M. Chem. Rev. 2011; 111:563–639. [PubMed: 20939606]
18. The absolute configuration was assigned based upon analogy to other binaphthols formed by the diaza-*cis*-decalin copper catalyst. This coupling reaction has been demonstrated to be highly stereoregular. See references 15 and 16.
19. Upadhyay SP, Karnik AV. Tetrahedron Lett. 2007; 48:317–318.
20. For examples, see: (a) Cameron DW, Riches AG. Aust. J. Chem. 1999; 52:1165–1171. (b) Tietz LF, Gericke KM, Schuberth I. Eur. J. Org. Chem. 2007:4563–4577. (c) Danishefsky SJ, Uang BJ, Quallich G. J. Am. Chem. Soc. 1985; 107:1285–1293.
21. Tietz LF, Gericke KM, Singidi RR, Schuberth I. Org. Biomol. Chem. 2007; 5:1191–1200. [PubMed: 17406717]
22. The test reaction was carried out with a mixture of the “out”, “out-in”, and “in” bromoquinones, analogous to 4a–c, but containing 3,3'-methyl esters and 2,2'-hydroxyls
23. Kim GT, Wenz M, Park JI, Hasserodt J, Janda KD. Bioorg. Med. Chem. 2002; 10:1249–1262. [PubMed: 11886788]

(*S*)-Bisoranjidiol, (*S*)-1

"C4 Derivatives," 2

$R^1 = \text{H, Me, CO}_2\text{C}_5\text{H}_{11}$

$R^2 = \text{H, Me}$

$R^3 = \text{Me, OMe, CH}_2\text{OH}$

Figure 1.
Bisanthraquinone natural products.

Scheme 1.
Retrosynthetic Analysis

Scheme 2.
Synthesis of Biaryl

Abbreviations: IBX, *o*-iodoxybenzoic acid

Scheme 3.
Synthesis of Binaphtho-*para*-quinone

Scheme 4.
 Synthesis of (S) -Bisoranjidiol and an Unnatural Regioisomer